



Supplement of

Identification of typical ecohydrological behaviours using InSAR allows landscape-scale mapping of peatland condition

Andrew V. Bradley et al.

Correspondence to: Andrew V. Bradley (andrew.bradley1@nottingham.ac.uk)

The copyright of individual parts of the supplement might differ from the article licence.

5 Supplement

10

Supplement contents

1.1 Multichannel Singular Spectrum Analysis (MSSA)

15 1.2 Sub site and random polygon creation

1.3 Topographical Attributes

1.4 Hierarchical Cluster Analysis

Fig. S1. MSSA reconstructions showing the contribution and periodicity of the first 8 EOFs (1-8) to an example time series from a wet peatland location.

20 **Fig. S2.** Example of polygon definition using topographic context, clues to management and vegetation contrasts

Fig. S3. Hierarchical cluster trees and dominant group averages for the sub site polygons.

Table S1. Characteristics of the cluster groups for the sub-sites, showing the average value of the semi quantitative scale for each contributing plant functional type (PFT), shrub (S), sedge (Sg), *Sphagnum* (Sp), moss (M), grass (G), rush (R), forest (F).

25 **Table S2.** Characteristics of the cluster groups for the sub-sites, showing the average value of each contributing hydrological feature, pool (P), stream (S), drained (D).

Table S3. Characteristics of the cluster groups for the random sites, showing the average value of each contributing plant functional type (PFT), shrub (S), sedge (Sg), *Sphagnum* (Sp), moss (M), grass (G), rush (R), forest (F)

Table S4. Characteristics of the cluster groups for the random-sites, showing the average value of each contributing hydrological feature, pool (P), stream (S), drained (D).

30 **Table S5.** Cumulative summary of the percentage of pool locations containing a ‘soft’ peat classified pixel and the different proportions of hits within the 150m buffer zone when searching for pixels with a soft *Sphagnum* dominated class

1.1 Multichannel Singular Spectrum Analysis (MSSA).

Multichannel Singular Spectrum Analysis (MSSA) was used to extract the seasonal oscillations from the InSAR time series. Each timeseries consisted of 133 data points, each separated by a 12-day time step. MSSA was applied to the data using the SSA-MTM software package (SPECTRA, 2021; Ghil et al., 2002) and method (Vautard and Ghil, 1989). Using this method, the first 20 Empirical Orthogonal Functions (EOFs) of the timeseries were calculated with a window (L) of 31 time steps. The results show that the first 6 EOFs account for ~ 72% of variance within the time series. EOFs 1, 2, 3 and 4 are indicative of low frequency multi-annual climatic trends (with cycles of 48.8, 48.8, 26.0 and 26.0 months respectively) as opposed to annual bog breathing. EOFs 5 and 6, each with a period of 11.8 months, define the annual cycle that when reconstructed corresponds to the observed annual cycle of bog breathing (Fig. S1). If all 6 EOFs are used to reconstruct the time series, the low frequency trends can obscure the higher frequency annual cycles as shoulders or merged peaks, which cannot always be detected with the R maximum peak calculation method. Hence, we selected EOFs 5 and 6 to reconstruct the annual cycles for calculating key variables of peak timing and amplitude whilst removing long-term trends and higher frequency noise (Fig 1 main text).

1.2 Sub-site and random polygon creation.









An illustration of how the polygons were created, using the UK Ordnance Survey basemap, Google Earth imagery and annual peak timing (2016-17) is shown for sub-site 3 Munsary (Figure S3).

1.3 Topographic attributes.


The STRM DEM (Jarvis et al., 2008) was resampled to the resolution of the InSAR data using the aggregate tool in ESRI ArcGIS to give the altitude. Slope and aspect was then calculated on a pixel-by pixel basis using 'Surface' toolbox in ArcGIS. Data was then summarised for all the data points that fell within each polygon to give the topographic attributes.

1.4 Hierarchical cluster analysis (HCA)

Using the semi quantitative scores 0, 1, 2, 3, for the sub-sites and random sites the Plant Functional Type (PFT) and then the hydrology classes were grouped. The approach used hierarchical clustering (Hierarchical Cluster Analysis, HCA) to identify polygons with similar combinations in semi quantitative scores. An iterative k-means clustering test indicated that 8-10 classes was sufficient to explain the dissimilarity within the data and this guided the location of the threshold on the hierarchical cluster tree (Figure S4 (A- D)). Once the data had been split into clusters, for each cluster, the average score in the semi-quantitative scale of each PFT in the cluster was calculated. The average scores were ranked and the top three were used to define the name of the Plant Functional Groups. In the case of the hydrology, the top scoring group within each class determined the name (Tables S1 -S4).




65 **Fig. S1. MSSA reconstructions including the calculated periodicity (in months) of the first 8 EOFs (1-8) to an example time series from a wet peatland location. EOFs 1 to 4 are calculated to have periodicities greater than 12 months, 5 and 6 correspond to the annual cycle and 7 and 8 have a periodicity greater than 12 months. Axis x is date; axis-y is relative height in metres.**




70 **Fig. S2.** Example of polygon definition using topographic context, clues to management and vegetation contrasts from, (a) UK Ordnance Survey map (Digimap Licence) © Crown copyright 2017. Distributed under the Open Government Licence (OGL). (b) True colour image, © Google Earth, and (c) Peak timing for 2016-2017 aggregated to months, for sub-site 3 Munsary.


(a)



(b)



(c)



(d)




Fig. S3. Hierarchical cluster trees and dominant group averages for the sub site polygons. A) vegetation. B) hydrology, and random polygons. C) vegetation. D) hydrology. Horizontal line defines the threshold for the final clusters.

Table S1. Characteristics of the cluster groups for the sub-sites, showing the average value of the semi quantitative scale for each contributing Plant Functional Type (PFT), shrub (S), sedge (Sg), *Sphagnum* (Sp), moss (M), grass (G), rush (R), forest (F). Top three scores for each cluster are highlighted, leading to the plant functional group (PFG) name, including Little or Absent vegetation (LoA).

Cluster	n	S	Sg	Sp	M	G	R	F	PFG name	
1	34		2.0	2.1	2.9	1.1	0.0	0.0	0.0	Sp,S,Sg
2	27		2.4	2.1	1.3	1.0	1.0	1.6	0.0	S,Sg,R
3	10		1.9	1.5	0.9	1.1	2.0	1.7	0.0	G,S,R
4	1		1.0	2.0	1.0	0.0	1.0	3.0	0.0	R,S
5	4		1.0	1.0	0.0	1.0	2.5	2.3	0.0	G,R
6	2		0.0	0.0	0.0	0.0	0.0	0.0	0.0	LoA
7	3		0.0	0.0	0.0	1.0	0.0	0.0	3.0	F
8	12		2.4	2.3	2.0	1.2	1.4	0.9	0.0	S,Sg,Sp
9	26		2.5	2.4	1.0	1.8	0.1	0.2	0.0	S,Sg,M
10	12		2.8	2.1	0.9	2.0	1.0	1.0	0.0	S,Sg,M

90

*Rank 1, rank 2, rank3

95 **Table S2. Characteristics of the cluster groups for the sub-sites, showing the average value of each contributing hydrological feature, pool (P), stream (S), drained (D). Top three scores for each cluster are highlighted, with rank 1 forming the final class name.**

Cluster	n	P	S	D	HydroName
1	27	3.0	0.1	0.0	P
2	11	0.0	0.2	0.9	D
3	28	0.3	0.7	2.4	D
4	16	0.4	2.2	1.6	S
5	18	1.2	0.3	1.0	P
6	14	0.0	0.0	2.2	D
7	1	0.0	0.0	2.0	D
8	13	3.0	0.0	1.5	P
9	2	2.0	2.0	0.5	P
10	1.0	3.0	2.0	0.0	P

*Rank 1, rank 2, rank3

Table S3. Characteristics of the cluster groups for the random sites, showing the average value of the semi quantitative scale for each contributing plant functional type (PFT), shrub (S), sedge (Sg), *Sphagnum* (Sp), moss (M), grass (G), rush (R), forest (F). Top three scores for each cluster are highlighted, leading to the Plant Functional Group (PFG) name, including Little or Absent vegetation (LoA).

Cluster	n	S	Sg	Sp	M	G	R	F	PFG name
1	35	2.9	1.9	1.1	1.3	0.0	0.1	0.0	S,Sg,M
2	10	2.6	0.6	0.2	0.9	2.5	1.4	0.0	S,G,R
3	10	0.9	0.4	0.0	0.0	2.8	2.1	0.2	G,R,S
4	9	2.1	1.9	0.9	1.0	1.1	0.8	0.0	S,Sg,G
5	7	0.1	0.1	0.0	0.0	0.0	0.1	0.1	LoA
6	35	2.0	2.2	2.3	1.1	0.1	0.1	0.0	Sp,Sg,S
7	12	0.0	0.0	0.0	0.8	0.0	0.0	3.0	F
8	7	1.4	1.9	0.4	0.9	0.9	2.0	0.0	R,Sg,S
9	2.0	1.0	2.4	0.5	1.0	2.5	2.0	0.0	G,S,R

105

*Rank 1, rank 2, rank3

110 Table S4. Characteristics of the cluster groups for the random-sites, showing the average value of each contributing hydrological feature, pool (P), stream (S), drained (D). Top three scores for each cluster are highlighted, with rank 1 forming the final class name.

Cluster	n	P	S	D		Hydro name
1	48	0.0	0.0		1.4	D
2	9	2.7	0.0		1.0	P
3	23	3.0	0.0		0.0	P
4	12	0.0	1.3		1.0	S
5	11	0.5	0.0		0.0	P
6	3	1.0	2.0		1.3	S
7	4	1.1	0.0		1.0	P
8	2	2.0	0.0		0.0	P
9	1	3.0	2.0		1.0	P
10	6	0.2	2.0		0.0	S
11	4	0.0	2.0		2.8	D
12	1	0.0	3.0		1.0	S
13	1.0	0.0	3.1		3.0	S

*Rank 1, rank 2, rank 3

115 **Table S5. Cumulative summary of the percentage of pool locations containing a ‘soft’ peat classified pixel and the different proportions of hits within the 150m buffer zone when searching for pixels with a soft *Sphagnum* dominated class.**

Proportion of ‘soft’ peat class occurrences in each buffer zone	Number of buffer zones (n=328)	Proportion of buffer zones (%)	Cumulative proportion of buffer zones (%)
100.0%	40	12.2	12.2
80.1 – 99.9%	57	17.4	29.6
60.1 – 80.0%	64	19.5	49.1
40.1 – 60.0%	69	21.0	70.1
20.1 – 40.0%	59	18.0	88.1
0.1 – 20.0%	32	9.8	97.9
0%	7	2.1	100.0

References

- 120 SPECTRA: SSA-MTM Toolkit for Spectral Analysis, SSA-MTM Group, Department of Atmospheric Sciences, University of California, Los Angeles [code], <http://research.atmos.ucla.edu/tcd/ssa/guide/guide4.html> , last access: 2 May 2021, 2021
- Ghil, M., Allen, M. R., Dettinger, M. D., Ide, K., Kondrashov, D., Mann, M. E., Robertson, A. W., Saunders, A., Tian, Y., Varadi, F., and Yiou, P.: Advanced spectral methods for climatic time series, *Rev. Geophys.*, 40, 1–40, <https://doi.org/10.1029/2000RG000092>, 2002.
- Vautard, R., and Ghil, M.: Singular spectrum analysis in nonlinear dynamics, with applications to paleoclimatic time series. *Physica D: Nonlin. Phenom.*, 35, 395-424, [https://doi.org/10.1016/0167-2789\(89\)90077-8](https://doi.org/10.1016/0167-2789(89)90077-8), 1989.
- 125 Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled seamless SRTM data V4, CIAT – International Centre for Tropical Agriculture [data set], <http://srtm.csi.cgiar.org> (last access: 19 July 2019), 2008.