



Landslide-lake outburst floods accelerate downstream hillslope slippage

Wentao Yang¹, Jian Fang², and Jing Liu-Zeng³

¹Three-gorges Reservoir Area (Chongqing) Forest Ecosystem Research Station, School of Soil and Water Conservation, Beijing Forestry University, Beijing, 100083, China

²College of Urban and Environmental Sciences, Central China Normal University, Wuhan, 430079, China

³Institute of Surface–Earth System Science, Tianjin University, Tianjin, 300072, China

Correspondence: Wentao Yang (yang_wentao@bjfu.edu.cn)

Received: 22 February 2021 – Discussion started: 11 March 2021

Revised: 22 August 2021 – Accepted: 29 August 2021 – Published: 16 September 2021

Abstract. The Jinsha River, which has carved a 2–4 km deep gorge, is one of the largest SE Asian rivers. Two successive landslide-lake outburst floods (LLFs) occurred after the 2018 Baige landslides along the river. Using Sentinel-2 images, we examined the LLF impacts on downstream river channels and adjacent hillslopes over a 100 km distance. The floods increased the width of the active river channel by 54 %. Subsequently, major landslides persisted for 15 months in at least nine locations for displacements >2 m. Among them, three moving hillslopes ~80 km downstream from the Baige landslides slumped more than 10 m 1 year after the floods. Extensive undercuts by floods probably removed hillslope buttresses and triggered a deformation response, suggesting strong and dynamic channel–hillslope coupling. Our findings indicate that infrequent catastrophic outburst flooding plays an important role in landscape evolution. Persistent post-flood hillslope movement should be considered in disaster mitigation in high-relief mountainous regions.

1 Introduction

In eastern Tibet, the Jinsha, Mekong, and Salween rivers, three of the largest Asian rivers, flow parallel to each other and incise steep gorges, with a maximum of 6 km of ridge–valley relief, while draining off the tectonically active margin of the Tibetan Plateau (Brookfield, 1998; Hallet and Molnar, 2001; Larsen and Montgomery, 2012; Liu-Zeng et al., 2008). The topographically varied terrain carved by fluvial incision, in association with the monsoonal climate, fosters a biodiversity hotspot with one of the highest speciation rates in the world (Myers et al., 2000). Ongoing rapid incision in the upper reaches (e.g., Zhang et al., 2021) destabilizes hillslopes, leading to landslides, occasional river damming, and associated landslide-lake outburst floods (LLFs) (e.g., Fan et al., 2020; Korup and Tweed, 2007). The extreme discharge of LLFs often entrains large volumes of sediments and is a major driver of geomorphic change. However, observations are lacking of the catastrophic impacts of LLFs on hill-

slopes (Baker, 2001; Turzewski et al., 2019), hindering the understanding of the geomorphic processes associated with LLFs and the development of hazards and landscape evolution models.

The extremely large water flux of LLFs to the receiving rivers is the most obvious phenomenon and has significant social impacts. For example, in 2000 CE, a large rock avalanche dammed the Yigong River with an estimated maximum impoundment of >2 km³. The resulting flood led to a record river level rise at gauging stations ~500 km from the landslide and caused major damage to infrastructure (such as roads and bridges) and loss of life in India (Delaney and Evans, 2015). Recently, the Tangjiashan landslide-dammed lake, which resulted from the 2008 Wenchuan earthquake, posed serious threats to the city of Mianyang, which is inhabited by millions of people (Fan et al., 2012a). A more astonishing LLF in 1920 BCE reportedly flooded the downstream Yellow River and even changed ancient Chinese history (Wu et al., 2016).

In addition to causing socioeconomic havoc, LLFs have long been recognized as important geomorphic drivers of fluvial systems. Most work on the geomorphic impacts of outburst floods have focused on their efficiency in sediment transportation and channel erosion (Cook et al., 2018; Turzewski et al., 2019). Bank undercutting and parallel retreat are the most frequently reported consequences of LLF lateral erosion (Korup and Tweed, 2007). Landslides related to bank undercutting along flood routes are regarded as instantaneous impacts of outburst floods, which are often recognized by retrospective field reconnaissance or post-event image interpretations (Cook et al., 2018; Higaki and Sato, 2012). Intersecting with the LLF routes, these already occurring landslides are easy to recognize (Cook et al., 2018). In addition to these instantaneous landslides, LLFs may also disturb upper hillslopes adjacent to the river channel over longer periods, which are more difficult to recognize and often overlooked. Thus, there is a lack of observations of LLF impacts on these ongoing slope slippages.

Two landslides occurred on 10 October 2018 and 10 November 2018 near Baige along the Jinsha River. These successive landslides caused two mega-floods (the peak discharges of the floods were $\sim 10\,000$ and $\sim 31\,000\text{ m}^3\text{ s}^{-1}$, respectively) (Cai et al., 2020) on 12 October and 12 November 2018. In this study, the ongoing impacts of LLFs on hillslope instability are investigated through quantitative measurements of downstream slope movements related to two upstream LLFs along the upper Jinsha River.

2 Materials and methods

2.1 Two floods related to the Baige landslides in 2018

A slope near Baige village in the Tibet Autonomous Region, China, failed twice in 2018 (Ouyang et al., 2019). The first landslide occurred on 10 October 2018, fully blocking the Jinsha River and forming a dammed lake with a maximum storage of 0.29 km^3 of water (Fan et al., 2019). The lake started to drain naturally on 12 October 2018, leading to a first on-site flood wave with a peak discharge of $10\,000\text{ m}^3\text{ s}^{-1}$ (Zhong et al., 2020). The first flood discharge finished on 16 October 2018, and there were no reports of economic damage during the first flood (Fan et al., 2019).

On 3 November 2018, the same slope collapsed again, leading to a second landslide dam, which fully blocked the Jinsha River. After 8 November 2018, several excavators were deployed to construct spillways. The lake started to drain through the spillway on 12 November. The maximum water volume of the second lake was $>0.52\text{ km}^3$ (Fan et al., 2019). The second landslide lake caused a mega-flood with a peak on-site discharge of $31\,000\text{ m}^3\text{ s}^{-1}$ near Baige (Zhong et al., 2020). This flood caused significant damage to major roads, schools, buildings, and farms in the Diqing and Lijiang prefectures, Yunnan Province, $\sim 500\text{ km}$ downstream of the Baige landslides. Fortunately, no casualties were re-




Figure 1. Map of the Jinsha River study region. The black rectangles are moving slopes detected after the Baige floods. The red circles are new landslides after the floods. The yellow rectangles with black numbers 1 to 4 are map extents shown in Fig. 3a–d. The blue rectangles with the black number 5 show the map extent for Fig. 7. (a) Two Sentinel-2 images were taken before the Baige landslide in July 2018 (top) and after in November 2018 (bottom). (b) Longitudinal profile of the Jinsha River in this section from the ALOS World 3D-30 m DEM; the localities of new landslides and moving slopes are marked. The Jinsha River is digitized by the authors from Sentinel-2 images before the 2018 floods.

ported during these two LLFs due to effective and timely evacuations.

Hillslopes on both sides along the Jinsha River were investigated for the downstream 100 km long stretch from the Baige landslide (Fig. 1). The elevation of the region ranges from 2500 to 5400 m, with a mean value of 4100 m above sea level (a.s.l.). In the study reach, the Jinsha River cuts through the plateau and flows from north to south; the valley floor descends from 2900 to 2600 m a.s.l. (Fig. 1) over a distance of $\sim 80\text{ km}$.

2.2 Slope displacements derived from Sentinel-2 images

We used the coregistration of optically sensed images and correlation (COSI-Corr) method to derive subpixel horizontal slope deformation in Sentinel-2 image pairs (Leprince et al., 2007). The COSI-Corr method is a pixel-matching method. There are two correlator engines that perform the procedure: the frequency and statistical engines. The frequency correlator transforms the images into the Fourier domain and detects subpixel surface changes in the phase images, whereas the statistical correlator compares changes in the spatial domain (Leprince et al., 2007). The frequency correlator is more accurate in detecting surface changes than the latter correlator and is used in this work (Leprince et al., 2007). To detect surface deformation, the method uses two images at a time: an earlier reference image and a later target image, both of which are transformed to the frequency domain. Using the reference image, ground deformation can be derived in the target image.

To perform the frequency correlation via the COSI-Corr method, the initial and final window sizes are two major parameters to be defined. The final window is set no larger than the initial window size by default. For both windows, smaller sizes are sensitive to background noise, whereas larger window sizes often result in smooth results (Lacroix et al., 2018; Yang et al., 2020a). In this work, we used moderate window size combinations of 64 and 32 pixels for the initial and final window sizes.

To detect slope deformation before and after the Baige floods, we used an image pair on 13 November 2015 and 12 November 2018 and a second image pair on 12 November 2018 and 12 November 2019. Images in these two pairs cover a large area from the Baige landslide to a 100 km downstream region along both banks of the Jinsha River (Fig. 1). Both image pairs are from the same date in different years to minimize uncertainties by having similar solar zenith and azimuth angles (Yang, 2020; Yang et al., 2020a).

We further analyzed the Mindu section ~ 80 km from the Baige landslide, where more moving hillslopes were detected. In this section, we composed 16 image pairs to detect the time series of slope displacements after the Baige floods. The dates of the reference and target images in these 16 image pairs are shown in Table 1. In this work, four reference images from (3, 13, and 16) January and (12) February 2017 were used. The acquisition dates of these reference images are very close to ensure that no displacements occur among them because the slopes in this section are stable before October 2018 and have been moving since then (Fig. 3). We used the same stable zone near Mindu to correct the image shifts between the reference and the target images with Yang et al. (2020b). In addition, we used two Sentinel-2 images in the summer months of 15 July 2017 and 16 August 2019 to map the width of the active channel before and after the 2018 floods.

Table 1. Dates of images used in the Mindu section. Image pairs with stars are shown in Fig. 4. All image pairs are used in Fig. 5.

| Image pairs | Reference images | Target images |
|-------------|------------------|------------------|
| No. 1* | 3 January 2017 | 13 January 2018 |
| No. 2 | 12 February 2017 | 3 April 2018 |
| No. 3* | 12 February 2017 | 23 April 2018 |
| No. 4* | 3 January 2017 | 9 November 2018 |
| No. 5* | 3 January 2017 | 29 November 2018 |
| No. 6 | 3 January 2017 | 9 December 2018 |
| No. 7* | 16 January 2017 | 6 January 2019 |
| No. 8 | 12 February 2017 | 23 January 2019 |
| No. 9* | 12 February 2017 | 7 February 2019 |
| No. 10* | 12 February 2017 | 14 March 2019 |
| No. 11* | 12 February 2017 | 29 March 2019 |
| No. 12* | 12 February 2017 | 18 April 2019 |
| No. 13* | 3 January 2017 | 29 December 2019 |
| No. 14 | 13 January 2020 | 13 January 2020 |
| No. 15 | 12 February 2017 | 18 January 2020 |
| No. 16* | 12 February 2017 | 7 February 2020 |

Uncertainties in the derived surface deformation have several sources, such as digital elevation model (DEM) errors during orthorectification and solar angle differences between the reference and the target image, and they can be estimated as the mean and standard deviation of displacements in a manually selected stable area (Bontemps et al., 2018; Lacroix et al., 2018; Yang et al., 2020a). In this work, we used the stable area of Yang et al. (2020b) near the MD-2 landslide.

Although previous work has demonstrated the capability of using a similar method to detect surface deformation of a 1/20 maximum pixel size (Leprince et al., 2007; Stumpf et al., 2017), the smallest reliable displacement is 1/5 of the image pixel size (Yang, 2020). For Sentinel-2 images of 10 m spatial resolution, displacements that are larger than 2 m are reliable.

Similar to the method used by Cook et al. (2018), we measured active river channels by manually interpreting active river channels from false color composite Sentinel-2 images. Fresh bare land near riverbanks is a major feature for interpreting active river channels. Topographic information from Google Earth was also used as ancillary data during interpretation. The uncertainty associated with the manual measurements of active channel width is one pixel size of the optical imagery used (10 m in this work).

3 Results

3.1 Magnitudes of the 2018 Baige LLFs

The nearest downstream hydrological station is the Batang station, which is 190 km away from the Baige landslide. The multiyear average discharge measured at the station is $924 \text{ m}^3 \text{ s}^{-1}$ (Xiong et al., 2020). The recorded peak discharges at this station for the first and second Baige floods




Figure 2. Multiyear peak discharge and suspended sediment flux at the Batang station. For comparison, the discharges for the 2018 Baige floods were 7850 and 21 200 $\text{m}^3 \text{s}^{-1}$, respectively, and the suspended sediment fluxes were 21.6 and 42 kg m^{-3} , respectively. Note multiyear gaps in the available data, which are partly due to the remoteness of the station. Despite the gaps, any event with a discharge larger than those in 1954 would be recorded.




Figure 3. Moving slopes in four selected sections along the Jinsha River (a–d) before (November 2015–November 2018) and after (November 2018–November 2019) the Baige floods. The left panels (a1, b1, c1, d1) are pre-flood results, and the right panels (a2, b2, c2, d2) are post-flood results. MD is short for Mindu village. The background images are cloud-free grayscale Sentinel-2 images before and after the 2018 floods. The red circles indicate concurrent landslides, and the black squares are slope slippages.

were 7850 and 21200 $\text{m}^3 \text{s}^{-1}$, respectively. We collected 40-year records of annual peak discharges for the Batang station from 1953 to 2017 that represent background climatic discharges (Fig. 2). Peak discharges caused by the Baige floods were 1.3 and 3.6 times larger than the recorded maximum an-

nual peak discharge in 1954 (Duan et al., 2016). In addition, the measured suspended sediment discharges for the first and second Baige floods were 2.3 and 4.5 times larger than the maximum annual peak sediment discharge (9.33 kg m^{-3} in 1972), respectively.




Figure 4. Cumulative hillslope displacements near Mindu village at different times between early 2018 and early 2020 using the early 2017 image (either January or February) as the reference. Cumulative displacement (**a**, **b**) before the first Baige flood, (**c**) between the two floods, and (**d**–**j**) at different times between 29 November 2019 and 7 February 2020. Background images are respective target Sentinel-2 images of the end date for the calculation.




Figure 5. Cumulative displacements of three points (crosses in Fig. 4) on three slopes near Mindu village (two black vertical lines indicate the occurrence date of the Baige outburst floods in late October and November 2018). There were no displacement data for MD-2 during March 2019 and January 2020 due to low image quality.

Although the peak discharge of LLFs is expected to decrease as the flood propagates downstream (Cook et al., 2018; Schwanghart et al., 2016; Cenderelli et al., 2001), the discharges recorded 190 km downstream of the Baige landslide are still much higher than background climate values.

3.2 Post-flood slope movements along the Jinsha River

We identified at least nine post-flood moving slopes (Fig. 3) in the studied reach. Before the Baige floods in 2018 (November 2015–November 2018), very few slopes along the Jinsha River had displacements larger than 2 m, whereas widespread slope movements with displacements larger than 2 m near the riverbank occurred after the floods (November 2018–November 2019). All these newly emerged slope movements had larger deformations near the valley bottom and decreased progressively higher along the hillslopes.

Near Mindu village ~ 80 km from the Baige landslide, at least four hillslopes slipped, and their movements appear to be ongoing (d2 in Fig. 3). We showed displacements of these slopes in 10 target images from 13 January 2018 to 7 February 2020 (Fig. 4) relative to the base images in early 2017 (Table 1). No displacements (>2 m) were detected in the MD-1 (MD is short for Mindu village), MD-2, and MD-4 hillslopes before the floods until late October and November 2018 (Fig. 4a, b). The toes of these hillslopes began to deform after the floods (since 29 November 2018, Fig. 4c–g). The deformation started from riverbanks, became larger over time, and propagated upslope until the end of the study period (February 2020, Fig. 4g–j). In the MD-3 area, the hillslope deformation was less than 4 m from January 2017 to January 2018 (Fig. 4a) and less than 8 m by 9 November 2018 (Fig. 4c); however, an acceleration could be observed from November 2018 to January 2019 (Fig. 4d, e). There is some background noise on hillslopes and tributary catchments in Fig. 4, which is irrelevant to the 2018 floods.

We further selected three representative points, each on the MD-1, MD-2, and MD-3 landslides, and measured their cumulative movements and velocities on 16 different dates (Table 1) from January 2018 to February 2020 (Fig. 5). The points on the MD-1 and MD-2 landslides show similar temporal patterns. The movement at both points before November 2018 was minimal (<0.18 cm d $^{-1}$ from February 2017) compared to the 4.55 cm d $^{-1}$ movement from November 2018 to April 2019. Even though the 12 November 2018 image was acquired during the second flood on 12 November, there are no detected displacements from 2015 to 12 November 2018, suggesting that the first flood had little post-effect on slow deformations and cannot be detected in a short time. Cumulative displacements at the MD-1 and MD-2 points continued to increase from ~ 8 to ~ 12 m after the second flood. Significant acceleration of the slope movement was also found for the point on the MD-3 landslide. The MD-3 point moved 1.52 cm d $^{-1}$ (uncertainty of <0.26 cm d $^{-1}$) before 9 November 2018 and then increased




Figure 6. River widths before and after the Baige floods. The gray shading indicates an uncertainty of ± 1 pixel in the Sentinel-2 imagery.

to 4.84 cm d $^{-1}$ (uncertainty of <0.96 cm d $^{-1}$) from November 2018 to February 2019. No additional displacements were detected after February 2019, which is due to collapse of the slope at that time.

3.3 Concurrent landslides and channel expansion during the 2018 floods

Active channel width increased for almost all parts of the study area (Fig. 6). There was widespread lateral erosion of the Jinsha riverbank in the form of bank undercutting and parallel retreat. The successive LLFs increased the mean width of the active channel from 96.33 to 148.56 m. The river width expanded 52.2 m on average after the 2018 floods. Lateral erosion of the Jinsha riverbanks led to undercutting of slopes, which propagated up slopes and could have been the direct trigger for the expansion of landslides existing before the Baige LLFs. In addition, we observed a few new landslides (Fig. 7), which could have been concurrent with the 2018 LLFs. We further measured the area of these concurrent landslides and post-event moving slopes with deformation >2 m in January 2020. The areas of these concurrent landslides and moving slopes were 2.18×10^5 m 2 and 11.86×10^5 m 2 , respectively. The areal extent of the moving slopes (with deformation >2 m) was 5.4 times that of the concurrent landslides. Although most of these moving slopes and the concurrent landslides occurred in the reach that experienced a significant increase in river channel width, there seems to be no relation between the channel widening and the maximum measured displacements of the moving slopes (Fig. 8). This is because slope stability is determined by the integrity of bedrock in the first order (Gallen et al., 2015).

The influence of the 2018 floods on adjacent hillslopes could be larger than that reported here. In this work, the method we use can reliably track slope deformation >2 m (Stumpf et al., 2017; Yang, 2020), and we only monitored the downstream riverbanks for less than 2 years. It is possible that there could be more slow-moving slopes with cu-




Figure 7. Concurrent landslides related to the Baige floods. False composite Sentinel-2 images were acquired on 15 July 2017 (a) and 16 August 2019 (b). The spatial extent of this area is shown as rectangle 5 in Fig. 1.

mulative displacements < 2 m that were not detected by our method.

4 Discussion

4.1 Implications for mountain hazards

Outbursts of landslide lakes present severe flooding threats to downstream communities (Delaney and Evans, 2015; Ling and Evans, 2014; Dai et al., 2005; Fan et al., 2012a). Landslide dams can be formed at more diverse locations along the river (Liu et al., 2019) than other types of natural dams, such as glacier or moraine dams, which are usually located in highlands near river heads and are typically far from human communities (Cook et al., 2018). LLFs pose more serious threats to humans due to their closer proximity to densely populated regions.

The observations in this study show that the 2018 Baige landslides destabilized some major downstream hillslopes tens of kilometers away. If these hillslopes subsequently fail, they may cause further disruptions to the main channel, forming a domino effect of “landslide–LLF–landslide” hazard chains. For example, Yang et al. (2020b) found that the size of the moving MD-2 slope is larger than that of the Baige landslide, whereas the river directly below MD-2 is narrower than the river at Baige, indicating a high risk of blocking the channel if a valley-blocking landslide occurs. This possible process demonstrates the propagation of LLF hazards downstream, which has not been given enough attention in the past but should be considered in future disaster mitigation measures. The findings of this work also have important implications for the ongoing construction of the Sichuan–Tibet railway, which runs through the deepest gorges along the Jinsha, Mekong, and Salween rivers.

4.2 Possible mechanism for hillslope slippage

To form the landslide–LLF–landslide hazard chain, weak riverbank hillslopes may be an important prerequisite. In our study area, we found that slopes with tensile cracks deformed after the Baige floods (Fig. 9). These tensile cracks may be the joint result of interactions among active regional tectonics, weak rock types, large slope gradients, and intense precipitation. (1) The Jinshajiang fault zone is the main fault zone in this study area. GPS measurements show that the near–east–west shortening rate of the fault zone is 2–3 mm per year (Chen et al., 1998, 2000). However, no strong earthquakes have occurred in the past 100 years for this section of the fault zone. Despite this, faults with prolonged fault motion activity produce damage zones and crush rocks along the Jinsha suture, leading to high landslide susceptibility in this study area (Cao et al., 2016). (2) The strata in the study area mainly comprise Mesoproterozoic metamorphic rocks, namely gneiss, quartz schist, and metagranulite, which can be easily weathered to form layers with weak shear strength during rainfall. (3) The landscape of the study area is deeply incised with high slope gradients greater than the angle of repose for these rock types (20–30°, Larsen and Montgomery, 2012). (4) Greater than 60 % of the precipitation occurs from July to September. Monsoon-driven precipitation, characterized by intense storms with large-magnitude and short-duration rainfall, is another stimulus for landslides.

In addition to deformations on slopes with tensile cracks, we also found deformations on slopes that have no visible tensile cracks. However, both types of slopes had bank collapses adjoining the active river channel, although the former slope type resulted in a much larger area of deformation. The removal of buttressing by the 2018 LLF erosion may be attributed to hillslope slippage in downstream reaches, which is substantiated by our observation that detected deformations are larger near the riverbank and decrease in concentric ellipses upslope.

Figure 3 indicates the velocity of three slope deformations. All these slopes showed movement with increased velocity




Figure 8. Relation between measured maximum slope displacements, river width increase (a), and ratio of river width increase (to the pre-disturbed river width) (b).




Figure 9. Tensile cracks on major hillslopes before the 2018 Baige floods as shown by high-spatial-resolution images from © Google Earth.

immediately after the floods, indicating that the Baige floods probably accelerated slope slippage. For MD-1, the velocity decreased quickly after the floods, which may indicate the deterioration of the floods' effect over time. For MD-2, a similar deformation–time pattern before May 2019 is visible. The deformation continued at a lower velocity from May 2019 to January 2020. The deformation of slopes may be related to precipitation or river discharge. As we do not have daily or monthly hydrological data, we analyzed the GPM (Global Precipitation Measurement mission) monthly precipitation from November 2018 to February 2020 (Fig. 10). Although GPM is known to be imperfect in mountain regions, it is the best satellite-based rainfall estimation to date and is sufficient to reveal the precipitation pattern for this study (Fang et al., 2019). The 2019 rainy season lasted from March to October. The beginning of the rainy season in March may explain the acceleration of MD-1 and MD-2 in March. The deformation continued during the rainy season (March to October). However, the acceleration of MD-2 after 18 January 2020 does not agree with precipitation change and should be explained by other triggers.

4.3 Implications for geomorphic processes

Previous work on the influence of outburst floods on hillslope instability along major rivers is hypothetical and lacks direct observations (Cook et al., 2018; Higaki and Sato, 2012). Widespread active channel expansions and bank collapses after the Baige floods confirm that mega-floods are major drivers of landscape evolution (Baynes et al., 2015; Lamb et al., 2014). Our finding of Baige flood-accelerated slope slippage supports the claim that landsliding is a key process linking fluvial incision and tectonics in landscape evolution (Larsen and Montgomery, 2012). Mega-floods can cause both instant landslides in terms of bank collapses (Cook et al., 2018) and post-flood landslides by destabilizing hillslopes along fluvial channels that fail in later years. However, much focus has been placed on the immediate hazards resulting from LLFs, such as instantaneous downstream floods and inundations (Dai et al., 2005; Delaney et al., 2015; Fan et al., 2012a). The lack of observations of post-flood landslides




Figure 10. Monthly GPM precipitation near the MD-1, MD-2, and MD-3 slopes. The precipitation data were downloaded from the Google Earth Engine.

hinders our understanding of the long-term impacts of mega-floods on landslides, landscape evolution, and related risks. This work shows that in addition to immediate (or concurrent) landslides (Cook et al., 2018; Higaki and Sato, 2012), the impact of LLF lateral erosion could propagate uphill by disturbing adjacent hillslope stability for a prolonged period, illustrating the dynamic response of mountain hillslopes to channel incision.

Infrequent catastrophic floods could play an important role in landscape evolution (Cook et al., 2018). Our finding that the mean active channel width increased by 54.2 % after the 2018 LLFs indicates that rare catastrophic events could leave a disproportionate footprint on local landscapes. Compared to monsoonal discharges, LLFs can cause extensive bank erosion and efficiently expand active river channels by undercutting hillslope bases and causing bank retreat. This is consistent with previous findings that bedrock canyons on the surfaces of Mars and Earth are probably caused by infrequent catastrophic floods instead of uniform, steady erosion from background runoff (Baynes et al., 2015; Keisling et al., 2020; Lamb et al., 2014; Lamb and Fonstad, 2010; Larsen and Lamb, 2016; Malatesta et al., 2017).

However, quantitative assessment of the role of LLFs in long-term landscape evolution awaits future research. First, the relationship between the magnitude and frequency of LLFs should be constrained. The volume of impoundment water and downstream channel morphology, which is site-specific, determines the severity and magnitude of an LLF (Fan et al., 2012b). Landslide triggers, such as large-magnitude earthquakes and high-intensity storms, control the frequency of LLFs. Strong earthquakes can trigger numerous landslides within river systems and form many landslide-dammed lakes (Fan et al., 2012b), the outbursts of which could accelerate the flush of post-seismic landslide debris

(Croissant et al., 2017). In addition, the strength of hillslopes along the LLF pathway should also be constrained. Hillslopes composed of weaker bedrock or that have been repeatedly damaged by seismicity could have more slope erosion after the occurrence of LLFs. Despite these related but poorly understood questions, our observations indicate that LLFs could significantly improve the efficiency of regional erosion and can be an important internal variable that modulates regional erosion.

5 Conclusions

In October and November 2018, two large landslides occurred on the same slope near Baige village along the Jinsha River, leading to two successive LLFs downstream. In addition to three new landslides and widespread lateral erosion, we found that the LLFs destabilized at least nine hillslopes, which progressively deformed over the following year within 100 km downstream of the landslide. Bank undercutting and parallel retreat were prevalent after the Baige floods, which probably activated these slopes by removing their buttresses. Landslide hazards propagated to long-range downstream regions by releasing mega-floods to undercut hillslopes along river channels, leading to more destabilized hillslopes. Persistent monitoring of slope deformation along major rivers may be a viable way to detect possible hazards in remote mountain regions.

Compared to LLF-triggered concurrent landslides, subsequent slowly moving slopes are less easily recognized and thus often overlooked. Realizing that LLFs may lead to more hillslope slumping over a larger areal extent and for a prolonged period afterwards could help researchers obtain a holistic picture of LLF impacts and improve geomorphic models of landscape evolution. In the future, LLF contribu-

tions to hillslope erosion may be constrained by quantifying the magnitude–frequency relationship.

Data availability. The Sentinel-2 remote sensing data can be accessed from the Sentinel hub (<https://www.sentinel-hub.com/>; Sinergise Laboratory for geographical information systems, Ltd, 2021). The ALOS World 3D-30 m DEM is available on the JAXA website (<https://www.eorc.jaxa.jp/ALOS/en/aw3d30/index.htm>; Japan Aerospace Exploration Agency, 2021).

Author contributions. The paper was written by WY with major contributions by JF and JLZ. WY processed the data. WY, JF, and JLZ interpreted the results. All authors reviewed and approved the paper.

Competing interests. The authors declare that they have no conflict of interest.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. Sentinel-2 images are courtesy of the ESA. We acknowledge JAXA for the use of the ALOS World 3D-30 m DEM. We thank the CIT team for the use of the COSI-Corr.

Financial support. The Second Tibetan Plateau Scientific Expedition and Research Program (STEP, grant no. 2019QZKK0906) and the National Science Foundation of China (grant nos. 42030305, 41807500) jointly supported this work.

Review statement. This paper was edited by Jens Turowski and reviewed by two anonymous referees.

References

- Baker, V. R.: Water and the Martian landscape, *Nature*, 412, 228–236, <https://doi.org/10.1038/35084172>, 2001.
- Baynes, E. R. C., Attal, M., Niedermann, S., Kirstein, L. A., Dugmore, A. J., and Naylor, M.: Extreme flood erosion dominates canyon evolution, *P. Natl. Acad. Sci. USA*, 112, 2355–2360, <https://doi.org/10.1073/pnas.1415443112>, 2015.
- Brookfield, M. E.: The evolution of the great river systems of southern Asia during the Cenozoic India-Asia collision: rivers draining southwards, *Geomorphology*, 22, 285–312, [https://doi.org/10.1016/S0169-555X\(97\)00082-2](https://doi.org/10.1016/S0169-555X(97)00082-2), 1998.
- Bontemps, N., Lacroix, P., and Doin, M. P. Inversion of deformation fields time-series from optical images, and application to the long term kinematics of slow-moving landslides in Peru, *Remote Sens. Environ.*, 210, 144–158, <https://doi.org/10.1016/j.rse.2018.02.023>, 2018.
- Cai, Y., Cheng, H., Wu, S., Yang, Q., Wang, L., Luan Y., and Chen, Z.: Breaches of the Baige Barrier Lake: Emergency response and dam breach flood, *Sci. China Technol. Sci.*, 63, 1164–1176, <https://doi.org/10.1007/s11431-019-1475-y>, 2020.
- Cao, C., Wang, Q., Chen, J., Ruan, Y., Zheng, L., Song, S., and Niu, C.: Landslide Susceptibility Mapping in Vertical Distribution Law of Precipitation Area: Case of the Xulong Hydropower Station Reservoir, Southwestern China, *Water*, 8, 270, <https://doi.org/10.3390/w8070270>, 2016.
- Cenderelli, D. A. and Wohl, E. E.: Peak discharge estimates of glacial-lake outburst floods and “normal” climatic floods in the Mount Everest region, Nepal, *Geomorphology*, 40, 57–90, [https://doi.org/10.1016/S0169-555X\(01\)00037-X](https://doi.org/10.1016/S0169-555X(01)00037-X), 2001.
- Chen, Z., Zhang, X., Shen, F., Zhao, J., Liu, Y., Tang, W., Burchfiel, B. C., King, R. W., and Royden, L. H.: GPS Monitoring of Crustal Movement in Southwest China, *Chin. Sci. Bull.*, 44, 851–854, <https://doi.org/10.1007/BF02886164>, 1999.
- Chen, Z., Burchfiel, B. C., Liu, Y., King, R. W., Royden, L. H., Tang, W., Wang, E., Zhao, J., and Zhang, X.: Global Positioning System measurements from eastern Tibet and their implications for India/Eurasia intercontinental deformation, *J. Geophys. Res.-Earth*, 105, 16215–16227, <https://doi.org/10.1029/2000JB900092>, 2000.
- Cook, K. L., Andermann, C., Gimbert, F., Adhikari, B. R., and Hovius, N.: Glacial lake outburst floods as drivers of fluvial erosion in the Himalaya, *Science*, 362, 53–57, <https://doi.org/10.1126/science.aat4981>, 2018.
- Croissant, T., Lague, D., Steer, P., and Davy, P.: Rapid post-seismic landslide evacuation boosted by dynamic river width, *Nat. Geosci.*, 10, 680–684, <https://doi.org/10.1038/ngeo3005>, 2017.
- Dai, F. C., Lee, C. F., Deng, J. H., and Tham, L. G.: The 1786 earthquake-triggered landslide dam and subsequent dam-break flood on the Dadu River, southwestern China, *Geomorphology*, 65, 205–221, <https://doi.org/10.1016/j.geomorph.2004.08.011>, 2005.
- Delaney, K. B. and Evans, S. G.: The 2000 Yigong landslide (Tibetan Plateau), rockslide-dammed lake and outburst flood: Review, remote sensing analysis, and process modelling, *Geomorphology*, 246, 377–393, <https://doi.org/10.1016/j.geomorph.2015.06.020>, 2015.
- Duan, W., He, B., Nover, D., Fan, J., Yang, G., Chen, W., Meng, H., and Liu, C.: Floods and associated socioeconomic damages in China over the last century, *Nat. Hazards*, 82, 401–413, <https://doi.org/10.1007/s11069-016-2207-2>, 2016.
- Fan, X., Xu, Q., Alonso-Rodriguez, A., Subramanian, S. S., Li, W., Zheng, G., Dong, X., and Huang, R.: Successive landsliding and damming of the Jinsha River in eastern Tibet, China: prime investigation, early warning, and emergency response, *Landslides*, 16, 1003–1020, <https://doi.org/10.1007/s10346-019-01159-x>, 2019.
- Fan, X., Dufresne, A., Siva Subramanian, S., Strom, A., Hermanns, R., Tacconi Stefanelli, C., Hewitt, K., Yunus, A. P., Dunning, S., Capra, L., Geertsema, M., Miller, B., Casagli, N., Jansen, J. D., and Xu, Q.: The formation and impact of landslide dams – State of the art, *Earth-Sci. Rev.*, 203, 103116, <https://doi.org/10.1016/j.earscirev.2020.103116>, 2020.

- Fan, X., Tang, C. X., van Westen, C. J., and Alkema, D.: Simulating dam-breach flood scenarios of the Tangjiashan landslide dam induced by the Wenchuan Earthquake, *Nat. Hazards Earth Syst. Sci.*, 12, 3031–3044, <https://doi.org/10.5194/nhess-12-3031-2012>, 2012a.
- Fan, X., van Westen, C. J., Xu, Q., Gorum, T., and Dai, F.: Analysis of landslide dams induced by the 2008 Wenchuan earthquake, *J. Asian Earth Sci.*, 57, 25–37, <https://doi.org/10.1016/j.jseaes.2012.06.002>, 2012b.
- Fang, J., Yang, W., Luan, Y., Du, J., Lin, A., and Zhao, L.: Evaluation of the TRMM 3B42 and GPM IMERG products for extreme precipitation analysis over China, *Atmos. Res.*, 223, 24–38, <https://doi.org/10.1016/j.atmosres.2019.03.001>, 2019.
- Gallen, S. F., Clark, M. K., and Godt, J. W.: Coseismic landslides reveal near-surface rock strength in a high-relief, tectonically active setting, *Geology*, 43, 11–14, <https://doi.org/10.1130/G36080.1>, 2015.
- Hallet, B. and Molnar, P.: Distorted drainage basins as markers of crustal strain east of the Himalaya, *J. Geophys. Res.-Earth*, 106, 13697–13709, <https://doi.org/10.1029/2000JB900335>, 2001.
- Higaki, D. and Sato, G.: Erosion and sedimentation caused by glacial lake outburst floods in the Nepal and Bhutan Himalayas, *Global Environ. Res.*, 16, 71–76, 2012.
- Japan Aerospace Exploration Agency: ALOS Global Digital Surface Model “ALOS World 3D – 30 m” (AW3D30), available at: <https://www.eorc.jaxa.jp/ALOS/en/aw3d30/index.htm>, last access: 15 September 2021.
- Keisling, B. A., Nielsen, L. T., Hvidberg, C. S., Nuterman, R., and DeConto, R. M.: Pliocene–Pleistocene megafloods as a mechanism for Greenlandic megacanyon formation, *Geology*, 48, 737–741, <https://doi.org/10.1130/G47253.1>, 2020.
- Korup, O. and Tweed, F.: Ice, moraine, and landslide dams in mountainous terrain, *Quaternary Sci. Rev.*, 26, 3406–3422, <https://doi.org/10.1016/j.quascirev.2007.10.012>, 2007.
- Lacroix, P., Bièvre, G., Pathier, E., Kniess, U., and Jongmans, D.: Use of Sentinel-2 images for the detection of precursory motions before landslide failures, *Remote Sens. Environ.*, 215, 507–516, <https://doi.org/10.1016/j.rse.2018.03.042>, 2018.
- Larsen, I. J. and Lamb, M. P.: Progressive incision of the Channeled Scablands by outburst floods, *Nature*, 538, 229–232, <https://doi.org/10.1038/nature19817>, 2016.
- Larsen, I. J. and Montgomery, D. R.: Landslide erosion coupled to tectonics and river incision, *Nat. Geosci.*, 5, 468–473, <https://doi.org/10.1038/ngeo1479>, 2012.
- Lamb, M. and Fongstad, M.: Rapid formation of a modern bedrock canyon by a single flood event, *Nat. Geosci.*, 3, 477–481, <https://doi.org/10.1038/ngeo894>, 2010.
- Lamb, M. P., Mackey, B. H., and Farley, K. A.: Amphitheater-headed canyons formed by megaflooding, *P. Natl. Acad. Sci. USA*, 111, 57–62, <https://doi.org/10.1073/pnas.1312251111>, 2014.
- Leprince, S., Barbot, S., Ayoub, F., and Avouac, J.: Automatic and precise orthorectification, coregistration, and subpixel correlation of satellite images, application to ground deformation measurements, *IEEE T. Geosci. Remote*, 45, 1529–1558, <https://doi.org/10.1109/TGRS.2006.888937>, 2007.
- Ling, S. and Evans S. G.: GIS-based analysis of 1933 Diexi Landslides and dam breach on the Min River, Sichuan, China, *EGU General Assembly Conference Abstracts*, 27 April–2 May 2014, Vienna, Austria, Bibcode: 2014EGUGA..1612446L, 2014.
- Liu, W., Carling, P. A., Hu, K., Wang, H., Zhou, Z., Zhou, L., Liu, D., Lai, Z., and Zhang, X.: Outburst floods in China: A review, *Earth-Sci. Rev.*, 197, 102895, <https://doi.org/10.1016/j.earscirev.2019.102895>, 2019.
- Liu-Zeng, J., Taponnier, P., Gaudemer, Y., and Ding, L.: Quantifying landscape differences across the Tibetan plateau: Implications for topographic relief evolution, *J. Geophys. Res.-Earth*, 113, F04018, <https://doi.org/10.1029/2007JF000897>, 2008.
- Malatesta, L. C., Prancevic, J. P., and Avouac, J. P.: Autogenic entrenchment patterns and terraces due to coupling with lateral erosion in incising alluvial channels, *J. Geophys. Res.-Earth*, 122, 335–355, <https://doi.org/10.1002/2015JF003797>, 2017.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., and Kent, J.: Biodiversity hotspots for conservation priorities, *Nature*, 403, 853–858, <https://doi.org/10.1038/35002501>, 2000.
- Ouyang, C., An, H., Zhou, S., Wang, Z., Su, P., Wang, D., Chen, D., and She, J.: Insights from the failure and dynamic characteristics of two sequential landslides at Baige village along the Jinsha River, China, *Landslides*, 16, 1397–1414, <https://doi.org/10.1007/s10346-019-01177-9>, 2019.
- Schwanghart, W., Worni, R., Huggel, C., Stoffel, M., and Korup, O.: Uncertainty in the Himalayan energy–water nexus: estimating regional exposure to glacial lake outburst floods, *Environ. Res. Lett.*, 11, 074005, <https://doi.org/10.1088/1748-9326/11/7/074005>, 2016.
- Sinergise Laboratory for geographical information systems, Ltd.: Sentinel Hub, available at: <https://www.sentinel-hub.com/>, last access: 15 September 2021.
- Stumpf, A., Malet, J. P., and Delacourt, C.: Correlation of satellite image time-series for the detection and monitoring of slow-moving landslides, *Remote Sens. Environ.*, 189, 40–55, <https://doi.org/10.1016/j.rse.2016.11.007>, 2017.
- Turzewski, M. D., Huntington, K. W., and LeVeque, R. J.: The Geomorphic Impact of Outburst Floods: Integrating Observations and Numerical Simulations of the 2000 Yigong Flood, Eastern Himalaya, *J. Geophys. Res.-Earth*, 124, 1056–1079, <https://doi.org/10.1029/2018JF004778>, 2019.
- Wu, Q., Zhao, Z., Liu, L., Granger, D. E., Wang, H., Cohen, D. J., Wu, X., Ye, M., Bar-Yosef, O., Lu, B., Zhang, J., Zhang, P., Yuan, D., Qi, W., Cai, L., and Bai, S.: Outburst flood at 1920 BCE supports historicity of China’s Great Flood and the Xia dynasty, *Science*, 353, 579–582, <https://doi.org/10.1126/science.aaf0842>, 2016.
- Xiong, M., Li, J., and Chen, Y.: Runoff Trend and Natural Driving Force in the Upper Jinsha River, *Water Resour. Res.*, 9, 235–248, <https://doi.org/10.12677/JWRR.2020.93025>, 2020.
- Yang, W.: Selecting the best image pairs to measure slope deformation, *Sensors*, 20, 4721, <https://doi.org/10.3390/s20174721>, 2020.
- Yang, W., Wang, Y., Wang, Y., Ma, C., and Ma, Y.: Retrospective deformation of the Baige landslide using optical remote sensing images, *Landslides*, 17, 659–668, <https://doi.org/10.1007/s10346-019-01311-7>, 2020a.
- Yang, W., Liu, L., and Shi, P.: Detecting precursors of an imminent landslide along the Jinsha River, *Nat. Hazards Earth Syst. Sci.*, 20, 3215–3224, <https://doi.org/10.5194/nhess-20-3215-2020>, 2020b.

- Zhang, J., Yang, H., Liu-Zeng, J., Ge, Y., Wang, W., Yao, W., and Xu, S.: Reconstructing the incision of the Lancang River (Upper Mekong) in southeastern Tibet below its prominent knickzone using fluvial terraces and transient tributary profiles, *Geomorphology*, 376, 107551, <https://doi.org/10.1016/j.geomorph.2020.107551>, 2021.
- Zhong, Q., Chen, S., Wang, L., and Shan, Y.: Back analysis of breaching process of Baige landslide dam, *J. Hydrol.*, 17, 1681–1692, <https://doi.org/10.1007/s10346-020-01398-3>, 2020.