Answer to the reviews on "Tectonic and climatic controls on the Chuquibamba landslide (western Andes, southern Peru)" by A. Margirier et al.

A. Margirier et al.

Audrey.margirier@gmail.com

Comments given by the two referees are repeated in bold for clarification purposes. We join the corrected manuscript after our answer.

OVERALL COMMENTS:

Anonymous referee 1 (AR1):

1) "There are several additional statements the authors make that seem to go beyond what they can reasonably conclude from currently available data (although if more references would be available to back up those statements, then they may be fine). "

 \rightarrow We changed some of our statements in which we were over interpreting the available data. We reorganize the text by removing some sentences to focus our discussion on the studied Chuquibamba landslide and we add some references to reinforce them.

2) "The referencing and organization of the manuscript can be improved."

 \rightarrow We added the references cited by AR1.

 \rightarrow We changed the organization of the manuscript, and restructured the paragraphs according to the two anonymous referees point of view: we gathered the dispersed geomorphic description in the "geomorphological setting" part.

3) "The authors should clarify their statements or provide better support."

 \rightarrow Adding references, improving figures and re organizing the text makes it clearer.

Anonymous referee 2 (AR2):

1) "There is little clarity on the new data regarding their presentation and structure, and the scope thereof. I believe that a proper reorganization of information, with a good exploitation of the data, can highlight the understanding of landslides in this part of the Andes." \rightarrow We improved and reorganized the text according to the two anonymous referees comments: we corrected unclear sentences and we did gather the geomorphological descriptions in a single paragraph.

2) "The dispersion of geomorphological data in context, method and results, suggest that they were obtained in the research. If so, they should be showed as a result, placing in the geomorphological context only the framework that was already published and indicating the used geomorphologic methods. If you do not want to exploit this aspect, then place everything in the geomorphological setting."

 \rightarrow We gathered all our geomorphological data in the same part (geomorphological setting) and we made a better distinction between our data and those issued from the literature, citing sources.

3) "Whereas ages are the main results of the work, better treatment of the numbers is needed. The text ages and figures do not always agree among themselves and with the Table. It is important to correct this point to avoid confusion in future cites."

→ We checked in detail all our ages, either in the text, figures and the tables. In the text and the figures, we now only indicate the ages calculated with an erosion rate of 0.21 mm/ka. We took in account this point and corrected the ages from the figure 5. We do now highlight the ¹⁰Be ages by removing the geomorphological description from the results part.

4) The fact that large earthquakes do not trigger landslides is mentioned isolated in the summary, without linking it to the results. However, the results dedicated an entire paragraph indicating that landslides could be triggering by seismicity; without arguments supporting this possibility. I suggest improving this point in the results and summary.

→ We improved the discussion, rewriting some sentence to make it clearer, citing new references. But in the absence of historical seismic data for ~100 ka, it's not possible to discuss 150yrs seismic cycles. Finally, we cannot conclude on the seismic triggering of the last debris-flow during the Ouki event. The main outcome of this paper is that a wet event seems to be necessary to trigger a large landslide.

LINE-SPECIFIC ANSWERS:

AR2, p. 1130, l. 14: The point (i) in the abstract is very general. Be more specific.

 \rightarrow 0k, we took this remark in account and focused on this point.

AR1, p. 1130, l. 24: McPhillips et al. (2014, Nature Geoscience) seems to be an appropriate reference to add to this list (or alternatively to line 25), particularly as it relates to the possible seismic triggering of landslides.

 \rightarrow 0k, we had this reference line 25.

AR1 / AR2, p. 1132, l. 2-4: This unit is quite well dated by Thouret et al. (2007), so it wouldn't hurt to mention the specific age range that they report. / The Huaylillas Ignimbrite is not well defined; its probable age is not mentioned. Explain more.

 \rightarrow 0k, we add the age of the ignimbrite after Thouret et al. (2007).

AR1, p. 1132, l. 19: This sentence should be restructured, because there are many studies that have now documented climatic fluctuations on the Altiplano over those timescales; is there a reason you mention Placzek specifically? If so, please state why, otherwise provide at least a representative list of the paleoclimate studies that have been done close to the field area.

 \rightarrow Ok, we change the sentence and add some critical references. At first, we mentioned Placzek et al. (2006, 2013) specifically because this study was the first to document climate fluctuations for such a long period in our study area: they documented climate fluctuations during the last 130 ka.

AR1, p. 1132, l. 20-22: To be fair, the study by McPhillips et al. (2014, Nature Geoscience) should also be mentioned, because they reported a very different interpretation in one of the areas where Steffen was working.

 \rightarrow 0k, we add McPhillips et al. (2014).

AR1, p. 1132, section 2.2: The results of the study by Roperch et al. (2006, Tectonics) could be useful to add to this description.

 \rightarrow We are not convinced this study is relevant at our time scale and for this local description.

AR1, p. 1133, l. 1-2: To be more precise, Schildgen et al. (2009) stated that the uplift that occurred between 14 and 2.2 Ma was accommodated by normal faulting and monoclinal warping, but that doesn't necessarily mean that normal faulting did not continue afterwards (or did not start earlier). Rewriting could be as simple as changing "the Tertiary episode" to "a Tertiary episode", the latter of which doesn't exclude the possibility of later faulting.

 \rightarrow 0k, we rewrite the sentence as suggested by the reviewer.

AR1, p. 1133, l. 2-3: After reading the whole paragraph, I understand that the authors are talking about a strike-slip system, but calling them "rather vertical faults acting as thrust with punctual normal apparent movements" here is confusing as a technical description... This implies that the kinematics have changed over time, but I don't think that is what the authors are trying to say (?). Perhaps there are near-vertical faults with both apparent thrust and normal offsets, which vary laterally according to the fault orientation (transpression versus transtension)?

 \rightarrow 0k with the reviewer, and we clarify the text by using the term "strike-slip fault".

AR2, p. 1133, l. 3: A comma is missing after 'However'

 \rightarrow 0k, we add the comma.

AR2, p. 1133, l. 25-26: The age of Huaylillas Ignimbrite is mentioned here. Why not before?

 \rightarrow 0k, taken in account, we add the age in a previous paragraph.

AR2, p. 1133, l. 22: It is not mentioned exactly where is located the megafan, although it is a very important feature of the study.

 \rightarrow 0k, we add the location of the megafan in the text.

AR2, p. 1134, l. 2: Here, the Huaylillas volcanic unit is mentioned. It is the ignimbrite? If so, I suggest using the same name to avoid confusions.

 \rightarrow Yes, it is the ignimbrite; we change the name to avoid confusions.

AR2, p. 1134, l. 3: The Megafan Acoy is again mentioned as if it was the first time. It should be all together before, to avoid confusions.

 \rightarrow Ok, we completely reorganize the geomorphological setting part, we gathered the Acoy megafan description in a single paragraph.

AR1, p. 1134, l. 9-12: This material would be better placed at the end of the previous section, because it describes results from another study, not the "Sampling strategy and methods" of this paper.

 \rightarrow 0k, we move this paragraph in the geomorphological setting.

AR2, p. 1134, l. 9-12: The Megafan Acoy is again mentioned as if it was the first time. It should be all together before, to avoid confusions.

 \rightarrow 0k, we gather the Acoy megafan description in the same single paragraph.

AR2, p. 1134, l. 18-24: The description of the specific research is scarce. Can you explain more?

 \rightarrow We describe here the classical sampling strategy for TCN surface exposure dating, we are not convinced that the text need to be more detailed.

AR1, p. 1134, l. 21-24: I guess that an explanation for how a quartz pebble can be used to estimate the erosion rate on rhyolite blocks that are higher in elevation and subject to higher rainfall will come later, but for now, I'm skeptical.

→ As regional erosion rates are scares (Hall et al., 2012; Kober et al. 2007), we sampled the closest available surface from the Chuquibamba landslide to provide another individual erosion rate. Additionally, this new erosion rate is in agreement with those calculated by Hall et al. (2012) for southern Peru (~0.3 mm/ka) or by Kober et al. (2007) for the western escarpment in northern Chile (rhyolitic Oxaya formation; 0-0.5 mm/ka).

AR1, p. 1135, l. 11-24: Why not include this basic description of the morphology in section 2.3, where there is already some description of the geomorphology? This way, the results section could move directly into the cosmogenic data, instead of first reviewing more aspects of the morphology.

 \rightarrow We change the organization of the text and we gather our geomorphological results in the geomorphological setting part as suggested by the reviewers 1 and 2.

AR1, p. 1136, l. 2-7: There should at least be a caveat here that one would expect a quartz pebble to yield a lower erosion rate compared to rhyolite blocks. I think the assumption of low erosion is reasonable, but the authors should be a bit more careful in this argument.

→ We definitely agree with this statement. Indeed, as we should expect a higher erosion rate in rhyolite block than in quartz pebble, we looked for this lower pediment surface in order to calculate the erosion rate and discuss any eventual difference. In such an arid environment, and for 100kyrs time scales, our erosion rate is compatible with those calculated by Hall et al. (2012) for the southern Peru and by Kober et al. (2007) for the Oxaya ignimbrite in the western escarpment of the northern Chile.

AR2, p. 1136, l. 8: Ages in the text do not match those of Figure 5a and 5b.

 \rightarrow Ok, we change the ages in the figures 5a and 5b (those ages were calculated with zero erosion whereas we displayed the ages calculated with an erosion rate of 0.21 mm/ka in the manuscript).

AR2, p. 1136, l. 20-21: There is a link between the climate event and the megafan. What kind of relationship?

 \rightarrow Ok, we change the sentence to detail and explain the causal link between the Ouki climatic event and the Acoy megafan (Steffen et al., 2010).

AR1, p. 1137, l. 1: It's a reduction of the normal stress (or a reduction in shear strength, but certainly not a reduction in shear stress) due to hydrostatic pressure that typically leads to landsliding; Strasser and Schlunegger referred to decreased friction along a basal shear plane. To quote Ritter, Kochel, and Miller: "A body of material on a slope will remain in equilibrium (stable) as long as the sum of the applied shear stresses does not exceed the sum of the shear strength of the slope materials."

 \rightarrow 0k, we modify the text as suggested.

AR2, p. 1137, l. 7-9: It is suggested that an increase of pore pressure in the Huaylillas Ignimbrite by climate event triggered the megafan, but this is not well argued. What are the mechanical and lithological characteristics of the ignimbrite that let you make this proposal? Moreover, it is not establish directly that the age of the ignimbrite is correlated with the age of the climate event.

 \rightarrow The Huaylillas ignimbrite sheets are welded to strongly welded (Thouret et al., 2007) and interlayered with tuff. It suggests strong internal porosity variations in this unit but no geotechnical study of the Huaylillas ignimbrite is available.

AR1, p. 1137, l. 7-9: Sounds reasonable, but then why would the later wet events not have triggered additional landslides?

 \rightarrow Ok, we add a sentence to explain this point: for us the Ouki event is the last event strong enough (in duration or/and intensity) to trigger landslides in the arid western Andean flank.

AR1, p. 1137, l. 15: Can this generalization really be applied to all landslides? More evidence is needed to back that up, although it seems to be reasonably demonstrated in this case.

 \rightarrow Ok, we refocused the discussion on the Chuquibamba landslide and then we add some references to suggest the important role of the tectonic fracturing in preconditioning landsliding in the arid Western Cordillera.

AR2, p. 1137, l. 11-16: Similarity is suggested in the proposal on the flow direction of sliding Chuquibamba over other in northern Chile. In the case of Chile landslide it flowed towards the SW, perpendicular to structures; but here appears instead (maybe I'm

wrong) that megafan flowed parallel to the structures probably favored by the incision of the river Majes. Should put flow lines of the megafan in Fig.3a to clarify this point?

 \rightarrow We clarify in the text and discuss this point in the landscape evolution paragraph, pointing out the source of the megafan and detailing the succession of events in such a context. However we don't add any additional information on figure 3a as the Acoy megafan is highlighted in red.

AR1, p. 1137, l. 19-20: Not clear what the authors are trying to say here; both subduction earthquakes and El Nino events were linked to debris flows? The "succession in time" phrase is confusing. The authors might consider referencing McPhillips et al. (2014, Nature Geoscience) here.

 \rightarrow 0k, we change the sentence to clarify the text and we quote McPhillips et al. (2014).

AR1, p. 1137, l. 20-24: Not being able to exclude a seismic triggering (Lines 22-23) is quite far from "a likely link" (Lines 20-21). The authors should decide what interpretation they would like to make; I suggest deleting the sentence on lines 20-21, because they have not established any link between the recurrence of seismicity and landslide triggering.

 \rightarrow 0k, we delete the sentence lines 20-21 as suggested.

AR2, p. 1138, l. 4: Now you are talking about a Huaylillas paleosurface. Which one? It was never before been described. What is it? It is the roof of the Huaylillas Ignimbrite? It is the roof of the Huaylillas Formation?

 \rightarrow Yes, it is the weathered roof of the Huaylillas ignimbrite. We change the sentence and add a description of this surface in the geological context.

AR1 / AR2, p. 1138, l. 19: - Not clear what is meant by "permits new destabilizations"; are more recent mass movements sourced from that region? Is there a reason for that? / - The phrase 'This last debris-flow permits new destabilizations that enlarge the system' seems very ambiguous. Destabilization of what? Which system? hillsides, valleys, slopes?

 \rightarrow 0k, we change the sentence.

AR1, p. 1138, l. 19-24: This should be a separate paragraph, but "perfectly preserved" should be replaced with "well preserved". It seems dangerous to assume that over the 100 kyr history of exposure, there has been no rainfall, particularly considering that the deposit is interpreted to have been mobilized during a past wet phase. Overall, I'm not convinced that these final lines add anything useful to the discussion; the impact of aridity on surface preservation has already been stated a couple times previously.

 \rightarrow Ok, we remove this part.

AR1 / AR2, p. 1139, l. 2-7: - This is not a very effective paragraph for a conclusions section; it represents a fairly standard set of statements that can be applied to any study that measures TCNs in quartz from arid environments (of which there are many). / - I think that this paragraph does not add much to the conclusion.

 \rightarrow 0k, we remove this paragraph.

AR1, p. 1139, l. 10-11: Text needs to be tightened: the triggering of the debris flow is favored, not the debris flow itself. Also, what part of the results presented favors the climatic triggering? If only the age (i.e., a coincidence in timing), then that should be stated directly.

 \rightarrow Ok, we improve this sentence. Both the age of the debris-flow deposit and the Acoy megafan emplacement (with other debris-flow deposits in its stratigraphy) during the Ouki event favors the climatic triggering, we clarify this point in the text.

AR2, p. 1139, l. 11-12: You mentioned again that the triggering by seismicity but the arguments are lacking.

→ We improve the discussion on a potential seismic triggering of the debris-flow. But considering our results and the absence of historical earthquake catalog for this period (~100 ka) we can't conclude on this point.

AR2, p. 1139, l. 15: Now you talk about the hillslope processes and fluvial erosion, but before the discussion about these topics was ambiguous. Discuss earlier properly.

 \rightarrow We are suggesting further potential implications and opening the discussion rather than developing arguments about the hillslope processes. This is part of the conclusion and as R1 did not argue on that specific point, we would like to keep this in place.

AR1 / AR2, p. 1139, l. 19-23: - These sentences would be more appropriate in the discussion, particularly as they provide some of the broader regional context that is currently missing from the discussion. This argument in general though would benefit from references to any studies that may have attempted to quantify the amount of material mobilized in landslides from recent large earthquakes. Are there none from southern Peru or northern Chile? Without these, it is difficult to have much confidence in the dominance of climatic triggering, as the authors themselves point out that a seismic triggering cannot be ruled out. / - Here there is a discussion not mentioned before, place it on the discussion item.

 \rightarrow Ok, we place it in the discussion as suggested. There is no study, which quantify the amount of material mobilized in landslides triggered from a recent large earthquake but Lacroix et al.

(2013) demonstrated that in Peru during and after Pisco earthquake (Mw8), no important landslide were triggered along the Andean front. Similarly, no study reported major impact on landsliding of the Arequipa Mw8.5 event, and thus we base our statement on the fact that in the arid western Andean Cordillera none of the recent large earthquake triggered an important landslide.

AR1, p. 1139, l. 24-25: This statement needs to be supported, but I don't recall that it was brought up earlier in the discussion.

 \rightarrow 0k, we add a reference and add this point in the discussion.

AR2, p. 1139, l. 24-26: You talk about a strong tectonic control / climate on landslide. I think a better discussion about the tectonic control is lacking. There are still uncertainties (at least for me) in the Huaylillas Ignimbrite regard to its geomorphological features... and apparently is very important. You should have a good figure (map/geomorphologic image?) of the ignimbrite location. In addition, there are inaccuracies in the flow direction of Megafan Acoy to support the affirmation in these lines. Improving Fig. 3a is essential at this point.

 \rightarrow The map of the figure 1 clearly shows the ignimbrite location (in yellow). We improve the Acoy megafan description and location in the geomorphological setting.

AR1, p. 1139, l. 27-28: This sentence should be deleted, not only because grammatically it's incorrect, but also because it appears to add nothing to the conclusions.

 \rightarrow 0k, we delete the sentence as suggested.

EDITORIAL COMMENTS/CORRECTIONS

 \rightarrow We took in account, agree and correct all the typos and spelling mistakes the reviewer 1 point out. We thank AR1 for his help on editing the manuscript.

→ We only keep the notation ¹⁰Be-ka for the ¹⁰Be ages because this notation indicates that the ages can be posteriorly refined and re calculated in case of a refinement of the production rate. This convention is known and used by at least a part of the ¹⁰Be community (e.g. Gosse et al., 1995; Bigot-Cormier et al., 2005; Delunel et al., 2010).

TABLE AND FIGURES

AR2, fig. 1: What is the rectangle at the top of the figure?

 \rightarrow We remove it.

AR2, fig. 2c: The symbology of structures is lacking.

 \rightarrow Ok, we add the faults symbology in the legend.

AR2, fig. 3a: It is not well understood what are the Megafan, T1 and T2, there is a bad relationship with the legend. Improve the legend, the lineaments are not indicated. This figure is fundamental to understanding the manuscript, data presentation and discussion. However, it is unclear for a reader not familiar with the area.

 \rightarrow Ok, we change the legend (for the Acoy megafan, T1 and T2) to improve the readability. We indicate the lineaments in the legend.

AR2, fig. 3b: The image resolution is not good enough to review the morphological features.

 \rightarrow It's probably linked with the downsizing of the Figures for editorial purposes. We provide the full resolution for this figure.

AR2, fig. 3c: I think this image is underexploited in the text.

 \rightarrow 0k, we referenced this figure more carefully in the text (now 3 citations).

AR2, fig. 4: Verify that the ages in this figure match the text and table.

 \rightarrow The ages in this figure match those from the text and table.

AR2, fig. 5: The ages are used in the text are different from those used in Fig. 5.

Make coherent.

 \rightarrow 0k, we change the ages according to the text and table 1 (erosion = 0.21 mm/ka).

REFERENCES

Bigot Cormier, F., Braucher, R., Bourles, D., Guglielmi, Y., Dubar, M., and Stephan, J.: Chronological constraints on processes leading to large active landslides, Earth and Planetary Science Letters, 235, 141–150, 2005.

Delunel, R., Hantz, D., Braucher, R., Bourlès, D.L., Schoeneich, P., and Deparis, J.: Surface exposure dating and geophysical prospecting of the Holocene Lauvitel rock slide (French Alps), Landslides, 7, 393-400, 2010.

Gosse, J. C., Evenson, E.B., Klein, J., Lawn, B., and Middleton, R.: Precise cosmogenic 10Be measurements in western North America: support for a global Younger Dryas cooling event, Geology, 23, 877-880, 1995.

Hall, S. R., Farber, D. L., Audin, L., and Finkel, R. C.: Recently active contractile deformation in the forearc of southern Peru. Earth and Planetary Science Letters, 337-338, 85-92, 2012.

Kober, F., Ivy-Ochs, S., Schlunegger, F., Baur, H., Kubik, P. W., and Wieler, R.: Denudation rates and a topography-driven rainfall threshold in northern Chile: Multiple cosmogenic nuclide data and sediment yield budgets, Geomorphology, 83, 97-120, 2007.

Lacroix, P., Zavala, B., Berthier, E., and Audin, L.: Supervised Method of Landslide Inventory Using Panchromatic SPOT5 Images and Application to the Earthquake-Triggered Landslides of Pisco (Peru, 2007, Mw8.0), Remote Sensing, 5, 2590-2616, 2013.

McPhillips, D., Bierman, P. R., and Rood, D.H.: Millennial-scale record of landslides in the Andes consistent with earthquake trigger, Nature Geoscience, 2014.

Placzek, C. J., Quade, J., and Patchett, P. J.: A 130ka reconstruction of rainfall on the Bolivian Altiplano, Earth and Planetary Science Letters, 363, 97-108, 2013.

Placzek, C., Quade, J., and Patchett, P. J.: Geochronology and stratigraphy of late Pleistocene lake cycles on the southern Bolivian Altiplano: Implications for causes of tropical climate change, Geological Society of America Bulletin, 118, 515-532, 2006.

Steffen, D., Schlunegger, F., and Preusser, F.: Late Pleistocene fans and terraces in the Majes valley, southern Peru, and their relation to climatic variations, International Journal of Earth Sciences, 99, 1975-1989, 2010.

Thouret, J.C., Wörner, G., Gunnell, Y., Singer, B., Zhang, X., and Souriot, T.: Geochronologic and stratigraphic constraints on canyon incision and Miocene uplift of the Central Andes in Peru, Earth and Planetary Science Letters, 263, 151-166, 2007.

- 1 Tectonic and climatic controls on the Chuquibamba
- 2 landslide (Western Andes, southern Peru)
- 3

4 A. Margirier¹, L. Audin^{1, 2}, J. Carcaillet¹, S. Schwartz¹ and C. Benavente³

5 [1] {ISTerre, Université Grenoble I, CNRS, 1381 rue de la piscine, 38400 Grenoble Cedex 09,

6 France}

7 [2] {IRD, ISTerre, 1381 rue de la piscine, 38400 Grenoble Cedex 09, France}

8 [3] {INGEMMET, Av. Canadá 1470, San Borja, Lima, Peru}

9 Correspondence to: A. Margirier (audrey.margirier@ujf-grenoble.fr)

10

11 Abstract

12 The contribution of landslides to the Quaternary evolution of relief is poorly documented in arid contexts. In southern Peru and northern Chile, several massive landslides disrupt the arid 13 Western Andean front. The Chuquibamba landslide, located in southern Peru, belongs to this 14 set of large landslides. In this area, the Incapuquio fault system captures the intermittent 15 drainage network and localizes rotational landslides. Seismic activity is significant in this 16 region with recurrent Mw9 subduction earthquakes; however, none of the latest seismic 17 events have triggered a major landslide. New terrestrial cosmogenic dating of the 18 19 Chuquibamba landslide provides evidence that the last major gravitational mobilization of these rotational landslide deposits occurred at ~102 ka, during the Ouki wet climatic event 20 identified on the Altiplano between 120 and 98 ka. Our results suggest that wet events in the 21 arid and fractured context of the Andean forearc induced these giant debris-flows. Finally, our 22 study highlights the role of tectonics and climate on (i) the localization of large Andean 23 landslides in the Western Cordillera and on (ii) the long-term mass transfer to the trench along 24 25 the arid Andean front.

26

A. Margirier 6/3/15 16:43 Supprimé: and

A. Margirier 6/3/15 16:43 Supprimé: reliefs

A. Margirier 6/3/15 16:43 Supprimé:



1 1 Introduction

In active mountain ranges, landslides are an important process in long-term erosion and thus 2 contribute to the geomorphologic evolution of relief (Korup et al., 2007). Despite their 3 importance in terms of hazards, landslide maps remain rare (Guzzetti et al., 2012) and 4 information on the type, age or distribution of individual landslides is often lacking. Only a 5 few publications deal with landslide triggering and/or evolution in arid contexts such as the 6 7 western Andean flank, where several gigantic scarps disrupt the forearc piedmont (Audin and Bechir, 2006; Pinto et al., 2008; Strasser and Schlunegger, 2005; Wörner et al., 2002; Mather 8 et al., 2014; Crosta et al., 2015). In contrast, because of the potential seismotectonic trigger 9 10 (Keefer, 1984 and 2002; McPhillips et al., 2014), landslide triggering along subduction active margins has been studied for a number of years, but most previous studies focused on humid 11 climatic settings (Taiwan, New Zealand, Papua New Guinea, Japan; Meunier et al., 2008; 12 Hovius et al., 2011). In southern Peru, the topographic gradient (average slope of 4% between 13 the coast and the Western Cordillera), the crustal seismotectonic activity and the aridity of the 14 forearc region has been directly linked to the Andean uplift and the subduction of the Nazca 15 16 plate for the last 25 Myrs at least (Devlin et al., 2012; Alpers and Brimhall, 1988; Dunai et al., 2005). However, the Quaternary tectonic crustal activity and its role in the localization of 17 landslides along the Western Andean Escarpment has never been explored in southern Peru 18 19 and northern Chile. As a consequence, numerous questions remain concerning the importance of giant landslides in slope erosion, relative to other nonseismic agents of erosion such as 20 climatic forcing. 21

22 The "Chuquibamba landslide" is a large complex zone of imbricated landslides (about 80 km²) affecting the Western Andean Cordillera in southern Peru. It belongs to the Andean arid 23 piedmont, where most of the geomorphic markers are well preserved from erosion/transport 24 25 processes (Hall et al., 2008). This study of the Chuquibamba landslide aims to explore the 26 links between seismotectonic activity and abrupt climatic changes on the triggering and development of large landslides. In this paper, we map out the area and characterize the 27 tectonic and geomorphological settings, use Terrestrial Cosmogenic Nuclides (TCN) to date 28 pertinent markers of the last debris-flow event and document the evolution of the landslide 29 area. 30

31

A. Margirier 6/3/15 16:43

Supprimé: reliefs

A. Margirier 6/3/15 16:43 Supprimé: landslide

A. Margirier 6/3/15 16:43 Supprimé:

A. Margirier 6/3/15 16:43 Supprimé: such a

1 2 Context

2 2.1 Geologic and climatic setting of Chuquibamba region

The Andean Pacific arid front comprises different morphological units: the Coastal Cordillera 3 0-1000 m above sea level (asl), the Pacific Piedmont 1000-1500 m asl and the Western 4 Cordillera 1500-6000 m asl. In southern Peru, the Western Cordillera corresponds to a 5 Jurassic to Cretaceous volcanic arc (Atherton et al., 1985) (Fig. 1). These magmatic and 6 7 volcano-sedimentary rocks are emplaced within a Precambrian to Paleozoic basement. During the Eocene to Neogene, volcano-sedimentary deposits of the Moquegua group were overlain 8 onto the Western Cordillera. This group is partly covered by the Huaylillas ignimbrite, 9 produced during the Neogene volcanism and dated at 14.3-12.7 (⁴⁰Ar/³⁹Ar; Thouret *et al.*, 10 2007). The top of this ignimbrite forms a weathering surface. The Moquegua group and 11 Huaylillas ignimbrite reach a maximum of 1500 m thick (Thouret et al., 2007; Gunnell et al., 12 2010). Some volcano-sedimentary rocks were deposited on the coastal plains during the late 13 Neogene. 14

The Western Cordillera and Pacific Piedmont are incised, perpendicular to the NW-SE 15 striking Andean range, by 1 to 3 km-deep canyons (Tosdal et al., 1984; Gunnell et al., 2010; 16 Hartley and Evenstar, 2010). This area is affected by Quaternary tectonic deformation mainly 17 expressed in the Western Cordillera by an active fault system striking parallel to the range 18 (Sébrier et al., 1985; Hall et al., 2008, 2012) (Fig. 2). The piedmont region is exposed to 19 extremely low denudation rates ranging from 0.1-1 mm ka⁻¹ in the coastal desert and Pacific 20 piedmont to 1-46 mm ka⁻¹ in the Western Cordillera (values obtained for Quaternary time-21 scale by TCN dating; Hall et al., 2008; Kober et al., 2007) due to the arid climatic conditions 22 since at least the Neogene (Mortimer, 1980; Alpers and Brimhall, 1988; Dunai et al., 2005; 23 Rech et al., 2006). Along the Andes, the active subduction and crustal seismic activity largely 24 control the geomorphologic evolution of the forearc (Keefer, 1994; Keefer and Moseley, 25 2004; Audin et al., 2006; Tavera et al., 2007; Perfettini et al., 2010). Moreover, the Altiplano 26 has been the target of many paleoclimatic investigations (Thompson, et al., 1998, 2000; Cross 27 et al., 2001; Baker et al., 2001; Placzek et al., 2006, 2013) that highlighted climate 28 29 fluctuations for the last 130 ka. Steffen et al. (2009, 2010) as well as Carretier et al. (2013) highlighted the contribution of these wet climatic events to mass transport in major canyons 30 from the Altiplano to the Pacific coast. In contrast, McPhillips et al. (2014) questioned the 31 impact of the climatic fluctuations on landsliding and thus on erosion rate. 32

A. Margirier 6/3/15 16:43 Supprimé: on A. Margirier 6/3/15 16:43 Supprimé: Then during A. Margirier 6/3/15 16:43 Supprimé: caped A. Margirier 6/3/15 16:43 Supprimé: .

A. Margirier 6/3/15 16:43 Supprimé: perpendicularly

A. Margirier 6/3/15 16:43 Supprimé: climate lasting A. Margirier 6/3/15 16:43 Supprimé: for

A. Margirier 6/3/15 16:43 Supprimé: Placzeck A. Margirier 6/3/15 16:43 Mis en forme: Police :Non Italique A. Margirier 6/3/15 16:43 Supprimé: . (A. Margirier 6/3/15 16:43 Supprimé: evidenced A. Margirier 6/3/15 16:43 Supprimé: on the Altiplano

1 2.2 Fault geometry and kinematics

In southern Peru, the Incapuquio fault zone has a strong geomorphic imprint on the Andean 2 range (Huaman, 1985, Sébrier et al., 1985). Based on microtectonic studies Sébrier et al. 3 (1985, 1988) and Schildgen et al. (2009) identified different kinematic episodes. Sébrier et al. 4 (1985, 1988) defined major Tertiary and Early Quaternary compressional phases and a minor 5 Late Quaternary episode of normal faulting. However, Schildgen et al. (2009) proposed that a 6 7 Tertiary episode of strike slip and normal faulting occurred between 14 and 2.2 Ma, However relocated microseismicity (Grange et al., 1984) and teleseismic data (Devlin et al., 2012) 8 demonstrate the present-day reverse and strike slip components on the Lluta fault segment 9 10 and more regionally for the Incapuquio fault system in the Arequipa region (Fig. 2). The Neogene surface and Quaternary drainage network are also affected by these fault segments 11 (Fig. 3). Channel orientations and captures evidence a sinistral strike-slip component along an 12 extrado graben (Fig. 3c). These Quaternary kinematics are coherent with the vertical 13 geometry of the fault plane imaged at depth (20 km) by the distribution of crustal earthquakes 14 (Fig. 2b) and focal mechanisms (Fig. 2c) (Mw<5; <20 km, Grange et al., 1984; Devlin et al., 15 16 2012). Local reverse and normal apparent surface movements are known to occur along major strike-slip fault or flower structure as for the forearc of northern Chile (Victor et al., 2004) 17 and southern Peru (Hall et al., 2012). The present-day main sinistral strike-slip kinematics of 18 19 the Incapuquio fault system demonstrated by the seismicity is compatible with normal apparent displacements as observed by Schildgen et al. (2009) and Sébrier et al. (1985, 1988). 20

21 2.3 Geomorphological setting

22 The Chuquibamba landslide is located in the Majes river catchment, along the Western flank of the Andes, between 1000 and 4000 m asl (Fig. 1). The Chuquibamba landslide comprises 23 an imbricated set of rotational landslides, a debris-flow deposit and, in the lower area, a 24 megafan and alluvial terraces (Fig. 3a). Upstream, the rotational landslides remobilize the 25 thick Huaylillas ignimbrite formation fractured by the long lasting tectonic activity of the 26 Incapuquio fault system. The total volume mobilized for Chuquibamba landslide is estimated 27 to reach $\sim 40 \text{ km}^3$. These imbricated rotational landslides correspond to a succession of 3 head 28 29 scarps (Fig. 3b). The scar of the rotational landslides, cutting the Huaylillas weathering surface, delimits the landslide area (Fig. 3b) and form an elongated amphitheater trending in 30 the NW-SE direction. This direction does not fit with the overall southwest dipping 31 32 topographic slope, but rather corresponds to the structural trend of active faults (Fig. 3a).

A. Margirier 6/3/15 16:43 Supprimé: structures

A. Margirier 6/3/15 16:43
Supprimé: the
A. Margirier 6/3/15 16:43
Supprimé: At a crustal scale, these observations are mostly compatible with rather vertical faults acting as thrust with punctual normal apparent movements.
A. Margirier 6/3/15 16:43
Supprimé: senestral
A. Margirier 6/3/15 16:43
Supprimé: distribution
A. Margirier 6/3/15 16:43
Supprimé: senestral
A. Margirier 6/3/15 16:43
Supprimé: kinematic
A. Margirier 6/3/15 16:43
Supprimé: amphitheater-shaped scar of the rotational landslides delimits the landslide area (Fig.
A. Margirier 6/3/15 16:43
Déplacé vers le bas [1]: 3b).
A. Margirier 6/3/15 16:43
Déplacé vers le bas [2]: The total volume mobilized for Chuquibamba landslide is estimated to reach ~40 km ³ .
A. Margirier 6/3/15 16:43
Supprimé: The
A. Margirier 6/3/15 16:43
Supprimé: emplaced between 14 and 12 Ma (Thouret <i>et al.</i> , 2007) and
A. Margirier 6/3/15 16:43
Déplacé (insertion) [2]
A. Margirier 6/3/15 16:43
Déplacé vers le bas [3]: The debris-flow remobilizes the rotational landslides deposits.
A. Margirier 6/3/15 16:43
Supprimé: The debris-flow deposit is composed of mixed angular clasts, breccias and numerous meter-size boulders (Fig. 4a) entrapped in
A. Margirier 6/3/15 16:43
Déplacé (insertion) [1]
A. Margirier 6/3/15 16:43
Supprimé: thin volcanic matrix, reworked from

Supprimé: volcanic formation.

Moreover, several fault planes control the shape of the poly-lobed rotational landslides (Fig. 2 <u>3b).</u>

3 The base of the cirque (2900 m asl) is formed by smooth, sub-horizontal surfaces (Fig. <u>3b)</u>.

4 These surfaces likely correspond to former lateral landslide deposits re-incised by the river

5 <u>after the initiation of the Chuquibamba landslide.</u>

- 6 <u>The debris-flow remobilizes the rotational landslides deposits.</u> It consists of mixed angular
- 7 clasts, breccias and numerous meter-size boulders embedded in a thin volcanic matrix (Fig.
- 8 4a), reworked from the Huaylillas ignimbrite. The debris-flow deposit displays a smooth and
- 9 100 meters-scale ondulated surface (Fig. 4a). In its upper part, the debris-flow rests directly
- 10 on the basement (Fig. 4a); but 30 km downstream of the head scarp, in the vicinity of the
- 11 Majes River (900 m asl), the debris-flow overlays the Acoy megafan (Fig. 3a).
- 12 The megafan, located downstream of the Chuquibamba village at the outlet of the Rio
- 13 Grande, is about 100 meters thick and 8 kilometers long (Fig. 3a). Its stratigraphy indicates
- 14 <u>both a response to multiple phases of sediment production and surface erosion in the Rio</u>
- 15 Grande and fluvial incision in the Majes Canyon (Steffen et al., 2010), Steffen et al. (2010)
- 16 described an alternation of sheet flood units and debris-flow deposits, and a typical clast- to
- **17** <u>matrix-supported fabric.</u> OSL dating places the Acoy megafan formation between 107.0 ± 5.0
- 18 <u>ka</u> (base) and 101.6 ± 4.9 ka (top) (Steffen *et al.*, 2010). Two levels of alluvial terraces (T1,
- 19 T2) have been identified in the Majes canyon in the vicinity of the megafan. The Acoy
- 20 megafan and alluvial terraces development has been linked to wetter episodes, identified in
- sediments of some Altiplano paleolakes (Placzek et al., 2006, 2013), driving the fluctuations
- 22 of Colca/Majes hydrologic regime (Steffen et al., 2010).
- 23

24 3 Sampling strategy and methods

The last debris-flow deposit is perfectly preserved all along the Chuquibamba valley (Fig. 3a). Six meter-scale boulders of rhyolite entrapped in the Acoy megafan and debris-flow surfaces (Figs 4a and 4c) have been sampled. The preserved surface of the boulders (evidenced by desert varnish) indicates minimal post-abandonment erosion; they are well-anchored and sufficiently elevated on the debris-flow surface to minimize the possibility of postdepositional movement and potential covering by surficial material. In order to provide additional constrain on the regional erosion rate and consequently improve the exposure age A. Margirier 6/3/15 16:43 Déplacé (insertion) [4]

A. Margirier 6/3/15 16:43 Déplacé (insertion) [3]

A. Margirier 6/3/15 16:43 **Supprimé:** Two levels of alluvial terraces (T1, T2) and a megafan (Acoy megafan) have been identified downstream of Chuquibamba village in the Majes canyon (Fig. 3a). Their

A. Margirier 6/3/15 16:43 Déplacé (insertion) [5]

. Margirier 6/3/15 16:43

Supprimé: The Chuquibamba debris-flow deposit overlays the Acoy megafan in Majes valley (Figs 3a and 4b). The megafan is about 100 meters thick and 8 kilometers long (Fig. 3a). OSL dating situates the Acoy megafan formation between 107.0 ± 5.0 ka (base) and 101.6 ± 4.9 ka (top) (Steffen *et al.*, 2010). Its stratigraphy indicates both a response to multiple phases of sediment production and surface erosion in the Rio Grande and Majes catchments. A. Margirier 6/3/15 16:43 Déplacé vers le haut [5]: Steffen et al. (2010) described an alternation of sheet flood units and debris-flow deposits, and a typical clast- to matrix-supported fabric.

A. Margirier 6/3/15 16:43

Supprimé: minimizes

A. Margirier 6/3/15 16:43 **Supprimé:** avoid the post-deposition movements and

A. Margirier 6/3/15 16:43

Supprimé: materials. To

1 determination, we also sampled the closest surface from the landslide: the Pampa Jahuay

2 (quartzite pebble) in the pediplains 60 km southeast of Chuquibamba (Fig. 1 and Table 1).

Sample preparation was conducted at the Institut des Sciences de la Terre (ISTerre, 3 Grenoble). As the rhyolitic samples contained enough quartz, we extracted *in-situ* produced 4 beryllium-10 (¹⁰Be) using the chemical procedures developed by Brown et al. (1991) and 5 Merchel and Herpers (1999). The AMS ¹⁰Be measurements were performed at the ASTER 6 AMS French national facility (CEREGE, Aix en Provence). Analytical uncertainties include 7 uncertainties associated with AMS counting statistics, AMS external error (1%), standard 8 reproducibility and chemical blank measurements (${}^{10}\text{Be}/{}^{9}\text{Be}$ blank = 1.60 ± 0.72 × 10⁻¹⁵). 9 External uncertainties include 6% uncertainty in the production rate and 8% uncertainty in the 10 ¹⁰Be decay constant. Exposure ages were calculated using the online Cronus Calculator 11 (Balco et al., 2008). Results are computed using the time dependent scaling scheme of Lal 12 (1991) modified by Stone (2000). 13

14

29

15 4 TCN dating

In the debris-flow deposit and Acoy megafan samples, the ¹⁰Be concentrations are relatively 16 consistent and range from $6.67 \pm 0.28 \times 10^5$ to $1.38 \pm 0.08 \times 10^6$ atoms per gram of quartz (at 17 g⁻¹ qtz) (Tab. 1). The high ¹⁰Be content of the sample (1.33 $\pm 0.02 \times 10^7$ at g⁻¹ qtz, 11A28) 18 collected from the Pampa Jahuay suggests an extremely low erosion rate lasting at least for 19 the last 2 million years in the southern Peruvian forearc $(1.9 \pm 0.3 \text{ My})$. The computed erosion 20 rate $(0.21 \pm 0.05 \text{ mm ka}^{-1})$ agrees with rates published by Hall *et al.* (2012) for the South Peru 21 and, by Kober et al. (2007) for Chile (Oxaya formation). These results support the hypothesis 22 23 of an insignificant erosion of the sampled boulders of the debris-flow deposit. TCN exposure ages deduced from debris-flow boulders (with an erosion rate of 0.21 mm ka⁻¹) range from 24 96.1 \pm 8.9 to 108.5 \pm 10.2 ¹⁰Be-ka (Tab. 1, Fig. 5). Considering the uncertainties, exposure 25 ages are consistent and suggest a single remobilization event with a weighted average age of 26 101.9 ± 5.5 ka. The age of the large boulder sampled on the Acoy megafan (Fig. 4) is $105.3 \pm$ 27 10.2¹⁰Be-ka (Tab. 1). 28

A. Margirier 6/3/15 16:43 Mis en forme: Police :Arial A. Margirier 6/3/15 16:43 Supprimé: Results[1] A. Margirier 6/3/15 16:43 Déplacé vers le haut [4]: 3b) A. Margirier 6/3/15 16:43 Supprimé: This would correspond to former lateral landslide deposits re-incised by the river after the initiation of the Chuquibamba landslide. The debris-flow deposit consists of large boulders embedded in a thin matrix and displays a smooth and 100 meters-scale ondulated surface (Fig. 4a). In its upper part, this debris-flow rests directly on the basement (Fig. 4a): but 30 km downstream of the head scarp, in the vicinity of the Majes River (900 m asl), the debris-flow rests on the Acoy megafan (Figs. 3a and 4). The A. Margirier 6/3/15 16:43

A. Margirier 6/3/15 16:43 Déplacé (insertion) [6]

Supprimé: It is important to note that the associated analytical uncertainties are remarkably low (3-6%), probably due to the purity of the quartz mineral, allowing low interferences during AMS measurement and subsequently excellent analytical statistics (Tab. 1).

A. Margirier 6/3/15 16:43 Supprimé: in A. Margirier 6/3/15 16:43 Supprimé: for Chile,

- A. Margirier 6/3/15 16:43
- Supprimé:). This supports
- A. Margirier 6/3/15 16:43
- Supprimé: a non-significant

5 Discussion 1

5.1 Tectonic and climatic forcing on Chuquibamba landslide evolution 2

The weighted average age of debris-flow boulders indicates a last major debris-flow at 101.9 3 \pm 5.5 ka. The abandonment age of the megafan surface (105.3 \pm 10.2 ¹⁰Be-ka; Figs 4 and 5) 4 agrees with the OSL ages published by Steffen et al. (2010) (i.e. 107.0 ± 5.0 ka at the base of 5 the megafan and 101.6 ± 4.9 ka near the top). The Ouki wet event has been evidenced from 6 7 sediments collected in the eponym paleolake located in the higher Bolivian Altiplano (Placzek et al., 2006, 2013). The chronological framework deduced from U-Th dating on carbonates 8 indicates the Ouki deep lake cycle between 120 to 98 ka (Placzek et al., 2006, 2013). Steffen 9 et al. (2010) already suggested a correlation between wet time intervals on the Altiplano and 10 sediment aggradation in the Majes Valley, According to Steffen et al. (2010), the Acoy 11 megafan recorded two phases of Rio Grande catchment development characterized by 12 landsliding during the Ouki wet climatic event. It indicates that during the megafan 13 emplacement, between 112.0 and 96.7 ka, the Rio Grande permits the sediment transport 14 downstream. Similarly to the Lluta catchment in Chile, landsliding might have been initiated 15 by enhanced precipitation and a reduced friction along a basal shear plane due to increasing 16 hydrostatic pressure in the groundwater (Hoke et al., 2004; Strasser and Schlunegger, 2005). 17 The weighted average age of the last debris-flow deposit (101.9 ± 5.5 ka) also correlates with 18 the Ouki event (120-98 ka). In the Western Andes, other landslides have been associated with 19 20 wetter climatic conditions such as the older Lluta collapse (North Chile; 18°S; Wörner et al., 2002), yielding a minimum age of 2.5 Ma (Strasser and Schlunegger, 2005), the Tomasiri 21 landslide (South Peru; 17°30'S), dated at 400 ka (Blard et al., 2009) or younger Argentinian 22 landslides which have been associated with the Minchin event (40-25 ka; Trauth et al., 2003; 23 Hermanns and Schellenberger, 2008). We suggest that the debris-flow, which remobilized 24 rotational landslide deposits, was triggered by an increase in the pore water pressure in the 25 Huaylillas Ignimbrite during the Ouki wet climatic event. After this event none of the latest 26 wet climatic event identified on the Altiplano (e.g., Michin, 48-36 ka and Tauca, 26-15 ka) 27 triggered a large landslide or remobilized the Chuquibamba debris-flow. Similarly, in 28 29 Southern Peru, Keefer et al. (2003) identified only cm-scale debris-flows for the last 38 ka. Theses observations are consistent with the preservation of the Chuquibamba debris-flow 30 morphology since its emplacement. We suggest that the Ouki event was the latest event, 31 32 which duration and intensity permitted to trigger a large landslide in the arid western Andean

A. Margirier 6/3/15 16:43
Supprimé: ; they linked the Acoy megafan formation
A. Margirier 6/3/15 16:43
Supprimé: the Ouki wet climatic event. After
A. Margirier 6/3/15 16:43
Supprimé:
A. Margirier 6/3/15 16:43
Supprimé: stress

A. Margirier 6/3/15 16:43

A.

A.

7

Supprimé: wet events such as the older

flank (Wörner et al., 2002; Strasser and Schlunegger, 2005; Blard et al., 2009). Moreover, 1 Carretier et al. (2013) proposed that the contribution of rare and strong erosive events to the 2 long-term erosion of the Andean range is more than 90% in an arid climatic context. We 3 suggest climatic fluctuations favor landsliding and thus have a great impact on sediment 4 transport and on the western Andean flank morphology. 5 The link between the tectonic framework, localization, and flow direction of a mega-landslide, 6 has already been suggested for the Andean forearc domain (Pinto et al., 2008; Antinao and 7 Gosse, 2009; Mather et al., 2014). As the Chuquibamba landslide is elongated in an NW-SE 8 trend guided by the Incapuquio fault system, we suggest that the localization and geometry of 9 the landslide is mainly controlled by preferential fracturing orientations (Fig. 3b). More 10 broadly, in southern Peru and northern Chile most of the large landslides are located in 11 12 tectonically fractured regions (Audin and Bechir, 2006; Pinto et al., 2008; Strasser and Schlunegger, 2005; Wörner et al., 2002; Crosta et al., 2015). It suggests that the 13 prefracturation plays an important role in preconditioning landsliding in this arid area. 14 No relationship between a Mw 8.0 subduction earthquake and a giant landslide has been 15 previously documented in southern Peru (Lacroix et al., 2013) but Keefer et al. (2003) point 16 out that the succession in time of subduction earthquakes and El Nino events produces debris-17 flows in the coastal region (southern Peru). Here, even if the Ouki wet phase seems to 18 19 predispose the triggering, we can't exclude a contemporaneous seismic triggering (subduction or crustal earthquake) for the Chuquibamba debris-flows. More generally, robust arguments 20 indicating such a correlation for giant-landslides are scarce along the South American margin; 21 other triggering factors (increase of pore pressure, climate change, glacial debuttressing) are 22 usually invoked in addition to a seismic triggering (Keefer, 2002; Pinto et al., 2008). 23

24 5.2 Landscape evolution and tectono-climatic scenario

We propose a landscape evolution scenario based on geomorphic marker analysis and new TCN ages (Fig. 6). The initial drainage network flowed toward the coast (i.e. in a southerly direction) and incised the <u>roof of the weathered Huaylillas jgnimbrite</u> (Fig. 6a). These parallel incisions are still preserved (Fig. 3b, 3c) and sometimes even abandoned. Afterwards, the development of a graben structured by the Incapuquio fault system captured the drainage network along a 110°N trending direction (Fig. 3c) creating a new tributary of the Majes River. Rotational landslides were initiated by slope instability or pore pressure increase at the

A. Margirier 6/3/15 16:43 Déplacé (insertion) [7]

A. Margirier 6/3/15 16:43 **Supprimé:** at the base of the Huaylillas Ignimbrites during the Ouki wet climatic

A. Margirier 6/3/15 16:43 Supprimé: , A. Margirier 6/3/15 16:43 Supprimé: in

A. Margirier 6/3/15 16:43 **Supprimé:** landslides are A. Margirier 6/3/15 16:43 **Supprimé:** fig.

A. Margirier 6/3/15 16:43
Supprimé: relation
A. Margirier 6/3/15 16:43
Supprimé: -
A. Margirier 6/3/15 16:43
Supprimé: evidenced
A. Margirier 6/3/15 16:43
Supprimé: a likely link seems to be established between the recurrence of seismicity and landslide triggering. Even
A. Margirier 6/3/15 16:43
Supprimé: ,

A. Margirier 6/3/15 16:43 Supprimé: paleosurface base of the Huaylillas ignimbrite during wet climatic events, as proposed by Hoke *et al.* (2004) and Strasser and Schlunegger (2005) for northern Chile (Fig. 6b). Sliding surfaces of
 rotational landslides are localized on fault planes and progressively enlarge the valley, as
 evidenced by the successive rotational landslides scarps (Fig. 3).

The rotational landslides deposits <u>accumulated upstream in the valley (Fig. 6b) and were</u> remobilized by debris-flows during the Ouki event (120-98 ka; Fig. 6c). The accumulation of successive debris-flow and mudflow deposits at the outlet of the Rio Grande during the Ouki event formed the Acoy megafan (Steffen *et al.*, 2010). Finally, <u>at 101.9 ± 5.5 ka</u>, the last major debris-flow sealed the erosion/transport system (Fig. 6c). This pattern, alternating upper slopes destabilizations and its remobilization by debris-flow permits to enlarge the valley (Fig. 6d).

At the present time, the arid climate on the Western flank of the Cordillera preserves
geomorphologic features from erosion; the water yield comes only from rivers,

14

15 6 Conclusions

This study raises the question of the control of local tectonic activity and Altiplano climatic 16 fluctuations on landslide processes in the arid Western Andean front. Indeed, for the 17 Chuquibamba landslide region, the sequence of debris-flows (identified in the Acoy megafan 18 stratigraphy and overlaying its roof) seems to have been favored by the occurrence of a wet 19 climatic event (Ouki event), even if a contemporaneous seismic triggering cannot be 20 excluded. Our results suggest sediment accumulation in the valley during wet periods and 21 22 incision during dry periods. Wet climatic events then appear to control the growth of the drainage network, participating in regressive erosion and in the creation of new tributaries on 23 the Western Andean front. Our results also show that during wet events in a tectonically 24 fractured region, hillslope processes, rather than fluvial erosion, dictate the evolution of the 25 landscape at the channel head in the arid and high relief area. 26 As rare and strong erosive events represent the most important contribution to the long-term 27 erosion of the arid Western Cordillera (Carretier et al., 2013), we propose that massive 28 landslides disrupting the arid Western Andean front contribute significantly to long-term 29

- 30 erosion of the western escarpment of the Andes. This study also proposes that the Western
- 31 <u>Cordillera large landslides have been triggered during wet periods. Moreover, in southern</u>

A. Margirier 6/3/15 16:43

Supprimé: accumulate A. Margirier 6/3/15 16:43

- Supprimé: ago
- A. Margirier 6/3/15 16:4
- Supprimé: This last
- A. Margirier 6/3/15 16:43
- Supprimé: new destabilizations that
- A. Margirier 6/3/15 16:43

Supprimé: system

A. Margirier 6/3/15 16:43

Supprimé: ,

A. Margirier 6/3/15 16:43

Supprimé: . The debris-flow deposit surface is perfectly preserved; the Rio Grande just incises the debris-flow deposit (~100 m) due to the lack of a significant input of water issuing from the Western Cordillera

A. Margirier 6/3/15 16:43

Supprimé: In an arid context, Terrestrial Cosmogenic Nuclides are the perfect tool to date the exposure of surface older than ~100 ka. In the Peruvian arid piedmont, the

A. Margirier 6/3/15 16:43

Déplacé vers le haut [6]: TCN dating A. Margirier 6/3/15 16:43

Supprimé: of landslides is theoretically applicable over several hundreds of thousands of years, particularly in this case of gigantic boulder sampling. Additionally our data exemplify that quartz crystals extracted from rhyolites are particularly adapted to TCN dating and open new perspectives for quantitative geomorphologic applications in volcanic areas.

A. Margirier 6/3/15 16:43

Supprimé: last

A. Margirier 6/3/15 16:43 Supprimé: flow (101.9 ka

A. Margirier 6/3/15 16:43

Supprimé: be

A. Margirier 6/3/15 16:43

Déplacé (insertion) [8]

A. Margirier 6/3/15 16:43

Supprimé: erosive events

A. Margirier 6/3/15 16:43

Déplacé vers le haut [8]: Our results suggest sediment accumulation in the valley during wet periods and incision during dry periods.

A. Margirier 6/3/15 16:43

Supprimé: More broadly

- 1 Peru and northern Chile, most of the large landslides are located in tectonically fractured
- 2 regions. We thus suggest a strong regional tectonic and climatic control of the long-term
- 3 erosion of the arid western Andean front.

.

A. Margirier 6/3/15 16:43

Supprimé: In these arid regions, climatic fluctuations have a greater impact on landslide triggering and on sediment transport.

A. Margirier 6/3/15 16:43

Déplacé vers le haut [7]: Carretier *et al.* (2013) proposed that the contribution of rare and strong erosive events to the long-term erosion of the Andean range is more than 90% in an arid climatic context.

A. Margirier 6/3/15 16:43

Supprimé: As massive landslides disrupting the arid Western Andean front represent an important contribution to long-term Andean range erosion, we

A. Margirier 6/3/15 16:43

Supprimé: An important perspective of this study would be the systematic dating of the massive landslides disrupting the Western Andean front to quantify the long-term erosion rates. -

1 Acknowledgments

- 2 This project was supported by the Institut de Recherche pour le Développement (IRD). We
- 3 acknowledge additional support from Instituto Geologico Mineralogico y Metallurgico del
- 4 Peru (INGEMMET, Peru). We thank all the staff from the ASTER AMS facility at CEREGE.
- 5 This work has been carried out in the frame of the Master Thesis of A. Margirier in ISTerre.

1 References

- 2 Alpers, C.N., and Brimhall, G.H.: Middle Miocene climatic change in the Atacama Desert,
- northern Chile: Evidence from supergene mineralization at La Escondida, Geological Society
 of America Bulletin, 100, 1640-1656, 1988.
- 5 Antinao, J.L. and Gosse, J.: Large rockslides in the Southern Central Andes of Chile (32-
- 6 34.5°S): Tectonic control and significance for Quaternary landscape evolution,
- 7 Geomorphology, 104, 117-133, 2009.
- 8 Atherton, M.P., Warden, V. and Sanderson, L.M.: The Mesozoic marginal basin of Central
- 9 Peru: A geochemical study of within-plate-edge volcanism, in: Magmatism at a Plate Edge:
- the Peruvian Andes, Blackie Halsted Press, M.P. Pitcher, E.J. Cobbing and R.D. Beckinsale
- 11 (Eds), London, 47-58, 1985.
- 12 Audin, L., Bechir, A.: Active tectonics as determinant factor in landslides along the Western
- 13 Cordillera? Congreso Peruano de Geologia, Lima, Peru, 17-20 October, 237-239, 2006.
- 14 Audin L., David C., Hall S., Farber D., Hérail G.: Geomorphic evidences of recent tectonic
- activity in the forearc, southern Peru, Special Volume on Neotectonics in South America,
 Revista de la Asociación Geológica Argentina, 61, 545-554, 2006.
- 17 Baker, P.A., Seltzer, G.O., Fritz, S.C., Dunbar, R.B., Grove, M.J., Tapia, P.M., Cross, S.L.,
- 18 <u>Rowe, H.D., and Broda, J.P.: The history of South American tropical precipitation for the past</u>
 19 <u>25,000 years, Science, 291, 640-643, 2001.</u>
- 20 Balco, G., Stone, J.O., Lifton, N.A. and Dunai, T.J.: A complete and easily accessible means
- of calculating surface exposure ages or erosion rates from ¹⁰Be and ²⁶Al measurements,
 Quaternary Geochronology, 3, 174-195, 2008.
- Blard, P.H., Lavé, J., Farley, K.A., Fornari, M., Jiménez, N. and Ramirez, V.: Late local
 glacial maximum in the Central Altiplano triggered by cold and locally-wet conditions during
 the paleolake Tauca episode (17-15ka, Heinrich 1), Quaternary Science Reviews, 28, 34143427, 2009.
- Brown, E.T., Edmond, J.M., Raisbeck, G.M., Yiou, F., Kurz, M.D. and Brook, E.J.:
 Examination of surface exposure ages of Antartic moraines using in situ produced ¹⁰Be and
 ²⁶Al, Geochimica et Cosmochimica Acta, 55, 2269-2283, 1991.
 - 12

- 1 Carretier, S., Regard, V., Vassallo, R., Aguilar, G., Martinod, J., Riquelme, R., Pepin, E.,
- 2 Charrier, R., Hérail, G., Farias, M., Guyot, J.L., Vargas, G., and Lagane, C.: Slope and
- climate variability control of erosion in the Andes of central Chile, Geology, 41, 195-198,
- 4 2013.
- 5 Chmeleff J, von Blanckenburg F, Kossert K, Jacob D.: Determination of the 10Be half-life by
- 6 multicollector ICP-MS and liquid scintillation counting, Nuclear Instruments and Methods in
- 7 Physics Research, 268, 192-199, 2010.
- 8 Cross, S.L., Baker, P.A., Seltzer, G.O., Fritz, S.C., and Dunbar, R.B.: Late Quaternary
- 9 <u>Climate and Hydrology of Tropical South America Inferred from an Isotopic and Chemical</u>
 10 Model of Lake Titicaca, Bolivia and Peru, Quaternary Research, 56, 1-9, 2001.
- 11 Devlin, S., Isacks, B.L., Pritchard, M.E., Barnhart, W.D. and Lohman, R.B.: Depths and focal
- mechanisms of crustal earthquakes in the central Andes determined from teleseismicwaveform analysis and InSAR, Tectonics, 31, 2012.
- 14 Crosta, G.B., Paolo, F., Elena, V. and Hermanns, R. L.: The Cerro Caquilluco-Cerrillos
- Negros Giant Rock Avalanches (Tacna, Peru), Engineering Geology for Society and
 Territory, 2, 921-924, doi:10.1007/978-3-319-09057-3 159, 2015.
- 17 Dunai, T.J., González López, G.A. and Juez-Larré, J.: Oligocene-Miocene age of aridity in
- the Atacama Desert revealed by exposure dating of erosion-sensitive landforms, Geology, 33,
 321-324, 2005.
- 20 Grange, F., Hatzfeld, D., Cunningham, P., Molnar, P., Roecker, S.W., Suarez, G., Rodrigues,
- A. and Ocola, L.: Tectonic implications of the microearthquake seismicity and fault plane
 solutions in southern Peru, Journal of Geophysical Research, 89, 6139-6152, 1984.
- Gunnell, Y., Thouret, J.C., Brichau, S., Carter, A. and Gallagher, K.: Low-temperature
 thermochronology in the Peruvian Central Andes: implications for long-term continental
 denudation, timing of plateau uplift, canyon incision and lithosphere dynamics, Journal of the
 Geological Society, 167, 803-815, 2010.
- 27 Guzzetti, F., Mondini, A. C., Cardinali, M., Fiorucci, F., Santangelo, M., and Chang, K.-T.:
- Landslide inventory maps: New tools for an old problem, Earth Science Reviews, 112, 1-25,2012.
 - 13

- 1 Hall, S.R., Farber, D.L., Audin, L., Finkel, R.C. and Mériaux, A.S.: Geochronology of
- 2 pediment surfaces in southern Peru: Implications for Quaternary deformation of the Andean
- 3 forearc, Tectonophysics, 459, 186-205, 2008.
- Hall, S.R., Farber, D.L., Audin, L. and Finkel, R.C.: Recently active contractile deformation
 in the forearc of southern Peru, Earth and Planetary Science Letters, 337-338, 85-92, 2012.
- 6 Hartley, A.J. and Evenstar, L.: Cenozoic stratigraphic development in the north Chilean
- 7 forearc: Implications for basin development and uplift history of the Central Andean margin,
- 8 Tectonophysics, 495, 67-77, 2010.
- 9 Hermanns, R.L. and Schellenberger, A.: Quaternary tephrochronology helps define
 10 conditioning factors and triggering mechanisms of rock avalanches in NW Argentina,
 11 Quaternary International, 178, 261-275, 2008.
- 12 Hovius, N., Meunier, P., Ching-Weei, L., Hongey, C., Yue-Gau, C., Dadson, S., Ming-Jame,
- H. and Lines, M.: Prolonged seismically induced erosion and the mass balance of a large
 earthquake, Earth and Planetary Science Letters, 304, 347-355, 2011.
- Hoke, G. D., Isacks, B. L., Jordan, T. E. and Yu, J. S.: Groundwater-sapping origin for the
 giant quebradas of northern Chile, Geology, 32, 605-608, 2004.
- 17 Huaman, R.: Evolution tectonique Cenozoïque et néotectonique du piémont Pacifique dans la
- région d'Arequipa (Andes du Sud Pérou), Ph.D. Thesis, Université Paris Sud, Centre d'Orsay,
 220 pp., 1985.
- 20 INGEMMET: Mapa Geológico del cuadrángulo de Caravelí, Ocoña, La Yesera, y
 21 Chuquibamba, scale 1:100 000, Lima, 2001.
- Keefer, D. K.: Landslides caused by earthquakes, Geological Society of America Bulletin, 95,
 406-421, 1984.
- Keefer, D.K.: The importance of earthquake-induced landslides to long-term slope erosion
 and slope-failure hazards in seismically active regions, Geomorphology, 10, 265-284, 1994.
- Keefer, D.K.: Investigating landslides caused by earthquakes a historical review, Surveys in
 Geophysics, 23, 473-510, 2002.
- Keefer, D.K., and Moseley, M.E.: Southern Peru desert shattered by the great 2001
 earthquake: implications for paleoseismic and paleo-El Niño-Southern Oscillation records,

- 1 Proceedings of the National Academy of Sciences of the United States, 101, 10878-10883,
- **2** 2004.
- 3 Keefer, D. K., Moseley, M. E., and deFrance, S. D.: A 38 000-year record of floods and
- 4 debris flows in the Ilo region of southern Peru and its relation to El Niño events and great
- 5 earthquakes, Palaeogeography, Palaeoclimatology, Palaeoecology, 194, 41-77, 2003.
- 6 Kelly, M.A., Lowell, T.V., Applegate, P.J., Phillips, F.M., Schaefer, J.M., Smith, C.A., Kim,
- 7 H., Leonard, K.C., Hudson, A.M.: A locally calibrated, late glacial ¹⁰Be production rate from
- a low-latitude, high-altitude site in the Peruvian Andes, Quaternary Geochronology, 1-16, inpress.
- 10 Kober, F., Ivy-Ochs, S., Schlunegger, F., Baur, H., Kubik, P.W. and Wieler, R.: Denudation
- rates and a topography-driven rainfall threshold in northern Chile: Multiple cosmogenic
 nuclide data and sediment yield budgets, Geomorphology, 83, 97-120, 2007.
- 13 Korschinek, G., Bergmaier, A., Faestermann, T., Gerstmann, U. C., Knie, K., Rugel, G.,
- 14 Wallner A., Dillmann I., Dollinger G., Gostomski von C. L., Kossert K., Maiti M., Poutivtsev M. and
- 15 Remmert A.: A new value for the half-life of ¹⁰Be by Heavy-Ion Elastic Recoil Detection and
- 16 liquid scintillation counting, Nuclear Inst. and Methods in Physics Research, 268, 187-191,
- 17 2010.
- 18 Korup, O., Clague, J. J., Hermanns, R. L., Hewitt, K., Strom, A. L. and Weidinger, J. T.:
- Giant landslides, topography, and erosion, Earth and Planetary Science Letters, 261, 578-589,
 2007.
- 21 Lacroix, P., Zavala, B., Berthier, E. and Audin, L.: Supervised Method of Landslide Inventory
- Using Panchromatic SPOT5 Images and Application to the Earthquake-Triggered Landslides
 of Pisco (Peru, 2007, Mw8.0), Remote Sensing, 5, 2590-2616, 2013.
- Lal, D.: Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion
 models, Earth and Planetary Science Letters, 104, 24-439, 1991.
- Mather, A.E., Hartley, A.J. and Griffiths, J.S.: The giant landslides of northern Chile:
 Tectonic and climate interactions on a classic convergent plate margin, Earth and Planetary
- 28 Science Letters, 388, 249-256, 2014.
- 29 McPhillips, D., Bierman, P.R., and Rood, D.H.: Millennial-scale record of landslides in the
- 30 Andes consistent with earthquake trigger, Nature Geoscience, 1-6, 2014.



- Meunier, P., Hovius, N., and Haines, J.A.: Topographic site effects on the location of
 earthquake induced landslides, Earth and Planetary Science Letters, 275, 221-232, 2008.
- 3 Merchel, S. and Herpers, U.: An update on radiochemical separation techniques for the
- 4 determination of long-lived radionuclides via accelerator mass spectrometry, Radiochimica
- 5 Acta, 84, 215-219, 1999.
- Mortimer, C.: Drainage evolution of the Atacama Desert of northernmost Chile, Revista
 Geológica de Chile, 11, 3-28, 1980.
- 8 Perfettini, H., Avouac, J.-P., Tavera, H., Kositsky, A., Nocquet, J.-M., Bondoux, F., Chlieh,
- M., Sladen, A., Audin, L., and Farber, D.L.: Seismic and aseismic slip on the central Peru
 megathrust, Nature, 465, 7294, 78-81, 2010.
- 11 Pinto, L., Hérail, G., Sepúlveda, S.A. and Krop, P.: A Neogene giant landslide in Tarapacá,
- 12 northern Chile: A signal of instability of the westernmost Altiplano and palaeoseismicity
- 13 effects, Geomorphology, 102, 532-541, 2008.
- 14 Placzek, C., Quade, J. and Patchett, P.J.: Geochronology and stratigraphy of late Pleistocene
- 15 lake cycles on the southern Bolivian Altiplano: Implications for causes of tropical climate
- change, Geological Society of America Bulletin, 118, 515-532, 2006.
- Placzek, C.J., Quade, J., and Patchett, P.J.: A 130ka reconstruction of rainfall on the Bolivian
 Altiplano: Implications for the causes of tropical climate change, Earth and Planetary Science
- 19 Letters, 363, 97-108, 2013.
- 20 <u>Rech, J.A., Currie, B.S., Michalski, G., and Cowan, A.M.: Neogene climate change and uplift</u>
 21 in the Atacama Desert, Chile, Geology, 34, 761-764, 2006.
- 22 Roperch, P., Sempere, T., Macedo, O., Arriagada, C., Fornari, M., Tapia, C., Garcia, M., and
- 23 Laj, C.: Counterclockwise rotation of late Eocene-Oligocene fore-arc deposits in southern
- Peru and its significance for oroclinal bending in the central Andes, Tectonics, 25, TC3010,
 2006.
- 26 Schildgen T.F., Hodges K.V., Whipple K.X., Pringle M.S., van Soest M. and Cornell K.: Late
- 27 Cenozoic structural and tectonic development of the western margin of the central Andean
- 28 Plateau in southwest Peru, Tectonics, 28, TC4007, 2009.

A. Margirier 6/3/15 16:43 Mis en forme: Police :(Intl) Calibri



- 1 Sébrier, M., Mercier, J.L., Mégard, F., Laubacher, G. and Carey-Gailhardis, E.: Quaternary
- 2 normal and reverse faulting and the state of stress in the central Andes of south Peru,
- 3 Tectonics, 4, 739-780, 1985.
- Sébrier M., Lavenu A., Fornari M. and Soulas J.P.: Tectonics and uplift in Central Andes
 (Peru, Bolivia and Northern Chile) from Eocene to present, Géodynamique, 3, 85-106, 1988.
- 6 Steffen, D., Schlunegger, F., and Preusser, F.: Drainage basin response to climate change in
 7 the Pisco valley, Peru, Geology, 37, 491-494, 2009.
- 8 Steffen, D., Schlunegger, F. and Preusser, F.: Late Pleistocene fans and terraces in the Majes
- 9 valley (southern Peru) and their relation to climatic variations, International Journal of Earth
 10 Sciences, 99, 1975-1989, 2010.
- Stone, J.O.: Air pressure and cosmogenic isotope production, Journal of Geophysical
 Research, 105, 23753-23759, 2000.
- Strasser, M. and Schlunegger, F.: Erosional processes, topographic length-scales and
 geomorphic evolution in arid climatic environments: the "Lluta collapse," northern Chile,
- 15 International Journal of Earth Sciences, 94, 433-446, 2005.
- Tavera, H., Audin, L., and Bernal, I.: Source parameters of Sama (Tacna) earthquake (5.4
 Mw) of 20 November 2006, Bol. Soc. geol. Perù, 102, 109-115, 2007.
- 18 Thompson, L.G., Davis, M.E., Mosley-Thompson, E., Sowers, T.A., Henderson, K.A.,
- <u>Zagorodnov, V.S., Lin P.N., Mikhalenko V.N., Campen R.K., and Bolzan J.F.: A 25,000-year</u>
 <u>tropical climate history from Bolivian ice cores, Science, 282, 1858-1864, 1998.</u>
- Thompson, L.G., Mosley-Thompson, E., and Henderson, K.A.: Ice-core palaeoclimate
 records in tropical South America since the Last Glacial Maximum, Journal of Quaternary
 Science, 15, 377-394, 2000.
- Thouret, J.C., Wörner, G., Gunnell, Y., Singer, B., Zhang, X. and Souriot, T.: Geochronologic
 and stratigraphic constraints on canyon incision and Miocene uplift of the Central Andes in
 Peru, Earth and Planetary Science Letters, 263, 151-166, 2007.
- 27 Tosdal, R.M., Clark, A.H. and Farrar, E.: Cenozoic polyphase landscape and tectonic
- evolution of the Cordillera Occidental, southernmost Peru, Geological Society of America
 Bulletin, 95, 1318-1332, 1984.
 - 17

- 1 Trauth, M.H., Bookhagen, B., Marwan, N. and Strecker, M.R.: Multiple landslide clusters
- 2 record Quaternary climate changes in the northwestern Argentine Andes, Palaeogeography,
- 3 Palaeoclimatology, Palaeoecology, 194, 109-121, 2003.
- 4 Victor, P., Oncken, O. and Glodny, J.: Uplift of the western Altiplano plateau: Evidence from
- 5 the Precordillera between 20° and 21°S (northern Chile), Tectonics, 23, 2004.
- 6 Wörner, G., Uhlig, D., Kohler, I. and Seyfried, H.: Evolution of the West Andean Escarpment*
- 7 at 18°S (N. Chile) during the last 25 Ma: uplift, erosion and collapse through time,

A. Margirier 6/3/15 16:43 **Mis en forme:** Normal, Retrait : Première ligne : 0,05 cm

8 Tectonophysics, 345, 183-198, 2002.

Age ¹⁰ Be external	(ka) (f)	,			8.86	9.52	10.2	9.15	11.46	10.19		,
Age ¹⁸ Be internal moettaintv	(ka) (c)	,			3.68	4.09	4.87	4.19	8.34	5.57		,
Age ¹⁸ Be (ka) e=0.21 -1	mMa (d)		,		96.11	102.86	108.46	97.88	106.87	105.26		,
Age ¹⁰ Be external moettainty	(ka) (f)		,		8.83	9.49	10.17	9.10	11.42	10.15		271.34
Age "Be internal	(ka) (c)				3.67	4.07	4.85	4.17	8.30	5.55		94.88
Age ¹⁸ Be (ka) e = 0 u (d)					95.99	102.72	108.29	97.51	106.7	105.1		1894.08
Erosion rate external	(m Ma ⁻¹)		,		,	,	,	,	,			0.05
Erosion rate (m Ma ³) ¹					,	,	,	,	,			0.21
"Be concentration uncertainty	(× 10° atoms g ¹ Qez)	14.026	0.0076-1		25.748	793.97	34.974	34.175	84.507	27.905		227.531
¹⁸ Be concentration	(z) ammin g	31.274			\$28.574	938.088	992.821	992.144	1375.656	666.607		13318.936
Incertainty "Ber [®] Be	(%)	12.17	ţ		3.10	3.16	3.52	3.44	6.14	100 100 100		1.71
¹¹ Be [/] Be	8	1 604 10 ¹⁵	AT +000T		3.998 10 ¹⁵	3.257 10 ¹⁵	2.56610^{13}	5.468 10 ⁻¹⁵	6.981 10 ¹⁵	3.196 10 ¹⁵		1.32010^{11}
Be Jacob	ē.	A96	001-0		0.291	0.254	0.290	0.292	0.291	0.294		0.295
Quantz (g)					9.54	6.02	5.10	10.98	10.44	77.6		19.96
Sample thickness factor	(q)				0.98	0.99	0.97	0.98	0.98	66.0		0.97
Geomorphic scaling factor	(a)		,		0.99	66.0	0.98	0.99	96.0	1.00		1.00
ion rate $g^{\rm d} y r^{\rm d})$	Muons				0.27	0.28	0.28	67.0	0.33	0.24		0.30
Product (atoms	Spallation				7.15	7.43	7.39	8.40	10.54	5.13		8.76
(m asl)			,		1315	1356	1391	1543	1899	518		1592
.ongitude E (°W)					-72.5306	-72.9267	-72.9245	-72.5487	-72.5927	-72.4892		-72.0840
Latitude 1 (°N)			,	nsit	-15.9294	-15.5327	-15-5343	-15.9145	-15.8784	1600'91-		-16.3408
Sample		Blank DerCDEET	/nonunwo	Debris-flow dept	11A39	11A40	11A41	11A42	11A43	deor megafaw 11A28	Pantpa Jahuay	11A62

Table 1. TCN results of the Chuquibamba Valley and the Pampa Jahuay. (a) The topographic 1 scaling factor has been calculated following the method of Dunne et al. (1999). (b) 2 The sample thickness correction has been calculated with a 2.7 density factor. (c) AMS 3 4 (Accelerator Mass Spectrometry) analyses have been carried out at the French AMS facility ASTER. Calibration of ¹⁰Be concentrations were done with NIST Standard Reference 5 Material 4325 which use ${}^{10}\text{Be}/{}^9\text{Be}$ ratio of 2.79 × 10⁻¹¹ and a ${}^{10}\text{Be}$ half-life of 1.387 ± 0.012 × 6 10⁶ years (Chmeleff et al., 2010; Korschinek et al., 2010). Results have been corrected from 7 the chemical blank (10 Be/ 9 Be blank = 1.60 ± 0.72 × 10⁻¹⁵). Internal uncertainties consider the 8 analytical uncertainties including counting statistics, the instrumental variability (1%), the 9 standard deviation and chemical blank. External uncertainties include 6% uncertainty in the 10 production rate and 8% uncertainty in the ¹⁰Be decay constant. (d) Ages have been computed 11 with Cronus Calculator (Balco et al., 2008) using the time-dependent production rate of 12 Lal (1991) modified by Stone (2000). The production rate calibrate by Kelly et al. (in press) 13 on recent time scale in Peruvian Andes is not relevant for this study. For our range of ages 14 (100 ka) geomagnetic variations have to be considered. Ages are presented with the internal 15 (e) and the external uncertainties (f). 16

A. Margirier 6/3/15 16:43 **Mis en forme:** Motif : Transparente (Blanc)





3 Figure 1. Simplified geological map of South Peru (modified from Roperch et al., 2006;

4 INGEMMET, 2001), showing the Chuquibamba village (red star), the destabilization zone

5 and the Pampa Jahuay sampling site (J). Coordinates are given in WGS 84 longitude and

- 6 latitude (degrees). Inset shows the study area location within Peru and a part of South
- 7 America.

A. Margirier 6/3/15 16:43 **Mis en forme:** Police :(Intl) Times New Roman



3 Figure 2. Regional crustal seismicity and focal mechanisms. (a) Crustal seismicity (depth < 20 km) and faults are represented on the geological map of South Peru (modified from Roperch 4 et al., 2006; INGEMMET, 2001). (b) Cross section showing the vertical cluster of 5 6 earthquakes. (c) Focal mechanisms from Grange et al. (1984) and Devlin et al. (2012) studies, 7 the kinematic of the Incapuquio fault system is indicated by black arrows.





Figure 3. SRTM numerical elevation model overlay by Landsat image of the Chuquibamba 3 destabilization zone. Coordinates are given in WGS 84 longitude and latitude (degrees). (a) 4 Global view of the Chuquibamba area pointing major detachment scarps, debris-flow 5

- 1 deposits, Acoy megafan, alluvial terraces and faults. (b) Zoom on the amphitheater-shaped
- 2 scar of the rotational landslides showing the different head scarps. (c) Drainage network and

3

faults.



A. Margirier 6/3/15 16:43 **Mis en forme:** Police :(Intl) Times New Roman

5 6

Figure 4. Field photographs showing the debris-flow deposit morphology and the Acoy
megafan. (a) Characteristic block sampled on the debris-flow deposit. The arrow indicates
flowing direction of the debris-flow. (b) General view of the Acoy megafan (Acoy MF), red
lines highlight the top of the terrace. The OSL ages obtain by Steffen *et al.* (2010) for the

11 Acoy megafan are indicated with an asterisk (c) Block sampled on the megafan.





2

Figure 5. Samples location. (a) Digital elevation model (GeoMapApp, SRTM data, 90 m
resolution) of the study area, showing samples location and TCN ages obtained for the debrisflow deposit and the Acoy megafan surface. OSL ages obtained by (Steffen *et al.*, 2010) for
the Acoy megafan are noted in grey. (b) Elevation profile locating debris-flow and the Acoy

7 megafan ages.



Figure 6. Map and block diagrams of the evolution of the Chuquibamba landslide. (a) SRTM numerical elevation model, the white rectangle localizes the block diagrams. (b) Block diagram showing rotational landslide initiation on the flank of the Chuquibamba Valley during the Ouki wet event. The landslide enlarges the valley and accumulates material at the bottom. (c) The accumulated materials are remobilized by debris-flows during the wetter phases. (d) This remobilization allows new rotational landslides that enlarge the amphitheater shape valley.

10