

This discussion paper is/has been under review for the journal Earth Surface Dynamics (ESurFD).
Please refer to the corresponding final paper in ESurFD if available.

Modelling sediment clasts transport during landscape evolution

S. Carretier^{1,2,3,4}, P. Martinod^{1,2,3,4}, M. Reich^{4,5}, and Y. Godderis^{3,2,1}

¹IRD, UR 234, GET, 14 avenue Edouard Belin, 31400, Toulouse, France

²Université de Toulouse, UPS, GET, 14 avenue Edouard Belin, 31400 Toulouse, France

³CNRS, GET, 14 avenue Edouard Belin, 31400, Toulouse, France

⁴Department of Geology, FCFM, Universidad de Chile, Santiago, Chile

⁵Andean Geothermal Center of Excellence (CEGA), FCFM, Universidad de Chile, Santiago, Chile

Received: 27 September 2015 – Accepted: 28 September 2015 – Published: 14 October 2015

Correspondence to: S. Carretier (sebastien.carretier@get.omp.eu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Over thousands to millions of years, the landscape evolution is predicted by models based on fluxes of eroded, transported and deposited material. The laws describing these fluxes, corresponding to averages over many years, are difficult to prove with the available data. On the other hand, sediment dynamics are often tackled by studying the distribution of certain grain properties in the field (e.g. heavy metals, detrital zircons, ^{10}Be in gravel, magnetic tracers, etc.). There is a gap between landscape evolution models based on fluxes and these field data on individual clasts, which prevent the latter from being used to calibrate the former. Here we propose an algorithm coupling the landscape evolution with mobile clasts. Our landscape evolution model predicts local erosion, deposition and transfer fluxes resulting from hillslope and river processes. Clasts of any size are initially spread in the basement and are detached, moved and deposited according to probabilities using these fluxes. Several river and hillslope laws are studied. Although the resulting mean transport rate of the clasts does not depend on the time step or the model cell size, our approach is limited by the fact that their scattering rate is cell-size dependent. Nevertheless, both their mean transport rate and the shape of the scattering-time curves fit the predictions. Different erosion-transport laws generate different clast movements. These differences show that studying the tracers in the field may provide a way to establish these laws on the hillslopes and in the rivers. Possible applications include the interpretation of cosmogenic nuclides in individual gravel deposits, provenance analyses, placers, sediment coarsening or fining, the relationship between magnetic tracers in rivers and the river planform, and the tracing of weathered sediment.

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Numerical models of landscape evolution have significantly improved our understanding of relief dynamics by recasting competing theories within a general framework (e.g. Kooi and Beaumont, 1994). Within this framework, the relief dynamics are determined by the balance between detached, deposited and transferred sediment fluxes under the influence of tectonics and climate (e.g. Tucker and Hancock, 2010). These fluxes are set by constitutive laws, such as the diffusion law for hillslope creep (Culling, 1960) or the stream-power law for sediment river transport (Howard and Kerby, 1983). Nevertheless, it remains a challenge to justify these laws in geomorphology. Different laws produce different topographic evolutions, which have been widely studied (e.g. Howard, 1997; Willgoose et al., 1991; Kooi and Beaumont, 1994, 1996; Whipple and Tucker, 2002; Tucker and Whipple, 2002; Carretier et al., 2009; Davy and Lague, 2009). These laws have been justified or calibrated by mechanics (Whipple et al., 2000), experiments (Sklar and Dietrich, 2001) or by comparing their predictions with natural river profiles (e.g. Stock and Montgomery, 1999; Lague and Davy, 2003; Carretier et al., 2006; Logget et al., 2006), which has not always resulted in a good agreement (e.g. Tomkin et al., 2003; van der Beek and Bishop, 2003). The erosion-transport laws in landscape evolution models necessarily apply over long time steps ($\gg 1$ yr) and relatively large model cells ($\gg 10$ m). The uncertainty on these laws comes from a lack of methods to directly quantify sediment fluxes over such time and spatial scales.

At the same time, there are many techniques to trace the provenance and transport of rock fragments (clasts) and minerals. For example, detrital zircons, heavy minerals or trace elements in sedimentary rocks and river streams are routinely used to determine sedimentary provenance and/or constrain the exhumation history of orogenic highlands (e.g. Roddaz et al., 2005; Rodríguez et al., 2012). Other approaches use magnetic mineral tracers (e.g. Hassan et al., 1991; Haschenburger, 2011) or experimental devices (Lajeunesse et al., 2010; Kasprak et al., 2015) to better understand the dynamics of sediment transport in rivers. Furthermore, U-series disequilibrium stud-

ESURFD

3, 1221–1254, 2015

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Landscape with
clasts**

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ies in sediment fractions and cosmogenic nuclide measurements in individual clasts provide relevant information about erosion processes and rates (Chabaux et al., 2006; Codilean et al., 2008; Gayer et al., 2008; McPhillips et al., 2014). Many of these methods provide information about individual clasts that are part of bulk material fluxes over geological time scales. Therefore, we hypothesise that this information may be used to derive erosion-transport laws used in landscape evolution models. The link between erosion-transport laws and the spatial evolution of a clast population during landscape evolution requires a model that couples fluxes, topography and clasts.

In order to develop such a model, we couple a landscape evolution model with a clast dispersion model. The landscape evolution model is a modified version of Cidre (e.g. Carretier et al., 2009), which belongs to the family of complexity-reduced models (Murray, 2007). The clasts move according to probabilities depending on the erosion, deposition and transport fluxes calculated in Cidre. Our main goal here is to show that this algorithm yields a clast population movement that is consistent with the predicted sediment flux in some simple hillslope and river cases.

After briefly reporting previous modelling approaches based on flux-particle duality, we present Cidre and the probabilities used to move clasts. Then we analyse clast movement in the restricted cases of hillslope diffusion and river transport. Finally, we discuss the potential applications of this model. They include the 3-D tracing of weathered material which initially motivated this modelling approach.

2 Previous models coupling fluxes and particles

Models coupling fluxes and particles have been developed in other scientific fields, in particular in fluid mechanics and are known as Smoothed particle hydrodynamics models (SPH) (Gingold and Monaghan, 1977). The philosophy of these models is to simplify fluids to a limited set of discrete fluid particles moving under a force field. The distribution of the particles at the next time step is then used to estimate the new field of quantities such as forces, temperatures, densities or water flow. In geomor-

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



phology, the original landscape evolution model introduced by Mitasova et al. (2004) in the GIS software GRASS belongs to this class of models. In addition, authors used moving grains to predict topographical variations on hillslopes and in rivers. For example, Tucker and Bradley (2010) modelled the morphological evolution of 2-D scarps using moving grains, while Malmaeus and Hassan (2002) and Lajeunesse et al. (2013) among others used grains to model bed load transport. Grain transport modelling has also been carried out to predict the statistical distribution of certain geochemical properties. Within that scope, Repka et al. (1997), Gayer et al. (2008), Codilean et al. (2008), Carretier et al. (2009) and Carretier and Regard (2011) developed approaches in which grains move over a static landscape in order to trace their cosmogenic nuclide concentrations.

The modelling approach presented in this paper is different from these published works in the sense that: (1) particles are not used to estimate the water or erosion fields, (2) the topography evolves over time in our model, and (3) our modelling is 3-D instead of 2-D. Nevertheless, our model is inspired by the coupling of the landscape evolution model Eros and sediment particles (Davy and Lague, 2012).

3 Model

3.1 Erosion-sedimentation in Cidre

Cidre is a c^{++} code modelling the topography dynamics on a regular grid of square cells. At the beginning of a time step, a specified volume of rain falls. Cells are sorted by decreasing elevations. The propagation of water and sediment is proceeded in cascade starting from the highest cell to ensure mass conservation. A multiple flow algorithm propagates the water flux $Q [L^3 T^{-1}]$ toward all the downstream cells proportionally to the slope in each direction.

3.1.1 Mass balance

The elevation z (river bed or hillslope surface) changes on each cell according to the two following equations (Fig. 1):

$$\frac{\partial z}{\partial t} = -\epsilon + D + U, \quad (1)$$

and we define

$$D = \frac{q_s}{L}, \quad (2)$$

where ϵ is a local erosion (detachment or entrainment) rate [LT^{-1}], D is a local deposition rate [LT^{-1}], U is the uplift or subsidence rate [LT^{-1}], q_s is the incoming sediment flux per unit width [L^2T^{-1}] and L is transport distance. $L \geq dx$ to keep the deposition flux Ddx smaller than the incoming sediment flux q_s . For a complete justification of these equations in the case of river transport, see Charru (2006), Davy and Lague (2009) or Lajeunesse et al. (2015). We recall some of the elements in the Appendix A. Here, we generalise this approach for both hillslope and river sediment transport processes by specifying ϵ and L in both cases.

The length L determines the proportion of incoming sediment flux which is deposited. A large L means that the deposition is small, which is favoured in a real-case scenario by steep slope or high water discharge. The cell outflux per unit width q_s results from the sum of the sediment detached from this cell and the sediment eroded from upstream and which then crossed the cell without depositing. The sediment flux q_s is thus non-local (e.g. Tucker and Bradley, 2010; Lajeunesse et al., 2015).

3.1.2 Hillslope

In the following we establish equations for ϵ and L first for hillslope processes and then for rivers. As the hillslope model is new, we begin by recalling how long-term ($\gg 1$ ka)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



then most of the sediment entering a cell is deposited on this cell. This is the pure diffusion case. The sediment flux q_s does not include sediment eroded from above, it is thus “local” (Furbish and Haff, 2010). When $S \sim S_c$, L becomes infinity and there is no redeposition on the cell. This behaviour corresponds to mass wasting where a grain eroded from a place is able to go down a large distance before stopping. In that case, the flux q_s is “non-local” as it incorporates not only sediment that has been detached locally but also sediment in transit that has eroded from above. For an intermediate S , there is a progressive transition between pure creep and “ballistic” transport of material through the hillslope, which seems consistent with experiments (Roering et al., 2001).

Despite these conceptual differences, Eqs. (5) and (6) predict similar topographic evolutions as the “non-linear” diffusion equations for $\kappa' = \kappa dx$. This similarity is illustrated by Fig. 2, which displays the evolution of a transverse hill profile during uplift and relief decline. In order to solve Eqs. (3) and (4), we use an explicit finite difference approach along a topographic profile. In order to avoid numerical instability when S is close to S_c , we approximate Eq. (4) with its linear approximation when $S > 0.99S_c$ (e.g. Carretier et al., 2009). Equations (5) and (6) do not have this stability problem because L is simply set at a huge number when $S > S_c$. Figure 2 shows that both sets of equations (Eqs. 5, 6 and 3, 4) lead to the same evolution.

It would be difficult to experimentally verify Eqs. (5) and (6), because this would require separately observing the erosion ϵ and deposition D rates (Furbish and Haff, 2010; Furbish and Roering, 2013). This may explain why the non-linear diffusion model has been presented in the form of a local sediment flux q_s . Nevertheless, Eqs. (5) and (6) present several conceptual and numerical advantages. They may reconcile the different views concerning the “diffusive” or “non-local” nature of erosion on hillslopes as they predict a progressive transition between local and linear to non-local and non-linear sediment flux when the slope increases (e.g. Roering et al., 2001; van Milligen and Bons, 2002; Roering et al., 2002; Furbish and Haff, 2010; Tucker and Bradley, 2010; Fofoula-Georgiou et al., 2010; Falcini et al., 2013; Furbish and Roering, 2013). These equations are simple to implement in a 3-D model, particularly when there are

different sediment and bedrock layers because the erosion of these layers is separated from the sediment passing through the cell or deposited on it. We have not carried out an extensive comparison between the different resolution schemes of Eqs. (3), (4) (for example, see the solutions provided in Perron, 2011) and Eqs. (5), and (6).

5 Nevertheless, the solution of Eqs. (5) and (6) that we propose remains stable for time steps that are one order of magnitude larger than with Eqs. (3) and (4) in the examples given in Fig. 2.

3.1.3 Rivers

10 In the case of river processes, we describe here a simplified version of material detachment (sediment or bedrock), although the detachment threshold and the explicit expression of bed shear stress in particular can be included (e.g. Tucker, 2004):

$$\epsilon = Kq^m S^n, \quad (7)$$

$$L = \xi q, \quad (8)$$

15 where K is an erodibility coefficient, $q [L^2 T^{-1}]$ the water discharge per unit flow width on the cell, S is the steepest slope and the exponents are positive. The transport length L comes from the derivation of Davy and Lague (2009) for saltation and $\xi [T L^{-1}]$ is a factor depending on particle size and density. This law implies that the deposition rate decreases when the water discharge per unit width q increases. As demonstrated by Charru (2006), Davy and Lague (2009) or Lajeunesse et al. (2015), Eqs. (1) and (2) are mathematically equivalent to the “under-capacity” or “saturation” transport model (e.g. Beaumont et al., 1992; Whipple and Tucker, 2002; Andreotti et al., 2010), although L has a different physical meaning in both cases. It is a transport length in the Charru (2006) or Davy and Lague (2009) formulation (e.g. a characteristic transport distance of a population of grains entering a river locally by landsliding), and a saturation length to reach a maximum or “equilibrium” sediment flux, also called transport capacity in the Beaumont et al. (1992) formulation (e.g. the distance required for the suspended

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sediment flux to reach a maximum value downstream a brutal transition between the bedrock and alluvial river bed). As detailed by Davy and Lague (2009), the notion of transport capacity emerges by rewriting Eqs. (1) and (2) as $\frac{\partial z}{\partial t} = \frac{q_s - \epsilon L}{L}$, which is the form of the under-capacity model with ϵL defining a transport capacity. Thus, the formulation adopted in Cidre is equivalent to the under-capacity model.

For river processes, the flow width w can be set to the cell width dx or a river width. In the latter case, the flow width w is:

$$w = k_w Q^{0.5}, \quad (9)$$

where k_w is a coefficient depending on the lithology and $Q [L^3 T^{-1}]$ is the total water discharge at a river section. In this case, cell erosion ϵ in Eq. (1) is recast as $\epsilon \frac{w}{dx}$ (finite volume approach).

Erosion for sediment is different from that of bedrock (Eqs. 5 and 7), and within bedrock, different layers can be defined by their erodibility and detachment or slope thresholds (κ and S_c for hillslope processes and K for river processes). On one cell and during a time step dt , different layers are potentially eroded successively. The erosion of each layer consumes part of dt so that less time remains to erode the underlying layer. This time reduction is taken into account by multiplying the time step dt by $\left(1 - \frac{\text{volume layer}}{w dx \epsilon dt}\right)$ between the layers. If the surface layer is made of sediment, their erosion takes time and limits the erosion of the bedrock below. This process is known as the “cover” effect for river (e.g. Whipple and Tucker, 2002; Lague, 2010), and thus is taken into account in Cidre.

3.1.4 Bank or lateral erosion

Flowing water can erode lateral cells (Fig. 1a). These cells are those which are topographically above the considered cell and in a lateral direction perpendicular to each downstream direction. The lateral sediment volumetric flux $Q_{sl} [L^3 T^{-1}]$ is simply a fraction of the sediment flux in the considered direction (e.g. Murray and Paola, 1997;

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nicholas and Quine, 2007):

$$Q_{sl} = \alpha Q_s, \quad (10)$$

where α is a bank erodibility coefficient. α is specified for sediment and is implicitly determined for bedrock layers proportionally to their “fluvial” erodibility such that $\alpha_{\text{sediment}}/\alpha_{\text{bedrock}} = K_{\text{sediment}}/K_{\text{bedrock}}$ (K from Eq. 7). If sediment covers the bedrock of a lateral cell, α is weighted by its respective thickness above the target cell. Equation (10) is debatable and alternative formulations have been proposed (e.g. Davy and Lague, 2009). The main point is that bank erosion is introduced here to allow clasts from the sides of rivers to be incorporated into the river by lateral transport, as we expect this to control clast spreading.

3.1.5 Sediment spreading

The sediment leaving a cell is spread in the same way as water (Fig. 1a). This rule is justified by the probability of moving proportionally to the slope in the case of diffusive hillslope processes. As water is spread according to this rule, sediment in suspension is spread in the same way. Flume experiments may also support this rule for bed load (Seizilles et al., 2014).

Compared to previous published versions of Cidre (Carretier et al., 2009; Pepin et al., 2010; Carretier et al., 2014), significant modifications have been made: (1) the introduction of the transport length scale L , (2) the new hillslope erosion law and (3) the calculation of erosion using the steepest slope and then the spreading of sediment toward all downstream slopes. In the previous version, a transport capacity was calculated in each downstream direction and then summed to determine the cell erosion. The two approaches are not equivalent, in particular if the river erosion law depends non-linearly on water discharge. The advantage of the new version is that it is more stable and rapid while still allowing flow divergence and spreading. One motivation of these changes was to explicitly separate erosion, deposition and transfer rates on each cell, so that the treatment of clast movement will be straightforward.

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Adding and following Clasts

The philosophy of our approach is to use the local ϵ and D rates calculated by Cidre to determine a probability for a clast to move, cross or deposit onto a cell (Fig. 1b).

A clast has a specified radius R , with no particular limitation, between a small mineral to a large cobble. Its initial position and depth z_{clast} (of its base) are specified. When it is set initially in the bedrock, it corresponds to a piece of rock which will have this size once detached.

A clast is detached (eroded) if its depth is lower than the erosion calculated over the time step on that cell (if $z_{\text{clast}} \leq \epsilon dt$). Its probability to go in one of the downstream directions is simply the ratio of the local slope and the sum of the downstream slope (Fig. 1b).

For a moving clast entering a cell, the probability that it will be deposited is simply the ratio between the volumetric deposition flux and the volumetric incoming flux $Dwdx/Q_s$. Therefore, its probability to cross the cell is $1 - Dwdx/Q_s$ (Fig. 1b).

A clast may be detached but not leave the cell. This may occur if the clast was at depth. Removing material above the clasts takes part of the time step, so that the remaining time may prevent the clast from leaving the cell. Furthermore, a big clast should have a lower probability to leave the cell than a small one. In order to take these realities into account, a probability to leave the cell is determined by

$$1.25 \left(\frac{\epsilon dt}{2R} \right) \left(1 - \frac{z_{\text{clast}}}{\epsilon dt} \right) \delta, \quad (11)$$

where $\delta = 1$ if the direction of movement is parallel to rows or columns, and $\delta = 1/\sqrt{2}$ along diagonals (a longer distance decreases the probability to leave the cell). The value of probability law (Eq. 11) is set to 1 if it exceeds unity, which may occur if the clast is at the surface and the erosion is larger than the clast diameter. The first term accounts for the inverse relationship between entrainment probability and clast size (e.g. Malmaeus and Hassan, 2002). The numerator ϵdt renders the travel distance

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



independent of the time step. The second term decreases the probability to move if the clasts were at depth. The coefficient 1.25 is justified a posteriori to adjust the mean travel rate of a population of clasts to theoretical predictions (see below). If a clast is detached but does not leave the cell, it remains at the cell surface.

Note that the probability of deposition could also depend on clast size. This is not implemented here but we will return to this point in the discussion.

4 Results

4.1 Linear diffusion

One way to validate the above model is to demonstrate that the displacement of a clast population follows predictions in simple cases. We begin with the case of linear diffusion. In Cidre, linear diffusion is obtained by using Eqs. (5) and (6) only, where $\epsilon = \kappa S$ and $L = dx$. The resulting linear and local sediment flux per unit width is $q'_s = \kappa dx S$, and the product κdx defines the diffusivity $\kappa' [L^2 T^{-1}]$.

We consider an inclined plane of slope S . Clasts initially set at the top surface of the plan will travel at a mean velocity \dot{x} given by the ratio of the sediment flux per unit width q_s and a depth corresponding to one clast diameter: $\dot{x} = \frac{q_s}{2R} = \frac{\kappa' S}{2R}$. Thus, the mean travel distance of the clast population $\bar{x} = \frac{\kappa' S}{2R} t$. Furthermore, within the framework of Fickian's diffusion theory, the standard divergence of the distances between particles is predicted to vary as $\sigma = \sqrt{2\kappa' t}$ (Einstein, 1905).

These predictions are tested on a bedrock plan of $S = 0.1$. The same κ is used for the bedrock and sediment (product of erosion) to simplify. Sediment and clasts can only go out on the lowest boundary condition, the elevation of which is fixed at 0 m. Different values of κ , dx and dt are tested in order to evaluate the effect of cell size and time step on the results. The diffusivity is $\kappa' = 0.01 \text{ m}^2 \text{ yr}^{-1}$ in all cases. Initially, 1000 clasts are set at the top of the plan and then followed during their descent. Their radius $R = 0.005 \text{ m}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 3 illustrates the progressive spreading of the clast population during the plane's erosion. \bar{x} well fits the predicted evolution and is not sensitive to the cell size or the time step for a given κ' . This non-dependence comes from the chosen form of the probability law (Eq. 11). In this law, the factor 1.25 was added to better adjust the calculated \bar{x} to $\frac{\kappa'S}{2R}t$. This coefficient works for other κ , dx , dt values. It may correspond to the numerical diffusion or to the c^{++} random function used to move the clasts.

As predicted by the diffusion theory, the scattering of the clasts $\sigma \propto \sqrt{t}$. It does not depend on dt for a large range of time steps (5–50 years in Fig. 3), but σ is significantly lower for larger dt values (250 years). For this time step, the eroded thickness at each step is larger than the diameter of the clasts, so that the “leave cell” probability (Eq. 11) is increased. Hence, at each time step, there is less chance that a clast will stay while the other ones move. The clast population moves more homogeneously, and consequently, σ is lower than with a smaller time step. Note that the effect of this time step on σ is consistent with natural processes where erosion occurs by discrete events. A thick erosion event necessarily entrains clasts that have a smaller diameter. For large time steps, the mean travel rate of the clast population remains equal to $\frac{\kappa'S}{2R}$, the effect of the time step only affects the scattering.

In all cases, σ strongly overestimates $\sqrt{2\kappa dx t}$. However, σ tends towards this predicted value when dx decreases or dt increases. When a clast moves, it travels a distance which is at least dx . Thus, the distance between immobile and moving clasts necessarily depends on dx . This is a limitation of this modelling approach. It is possible to adjust σ to the correct value, independent of dx , by replacing $\left(\frac{edt}{2R}\right)$ by $\left(\frac{\kappa dt}{dx}\right)$ in Eq. (11). However, the mean travel distance becomes strongly dependent on dx in this case.

In this version, we prefer probability law (Eq. 11) which allows the mean travel distance to be respected, for three reasons. \bar{x} is directly linked to the sediment mass balance of a portion of the landscape, not σ . In practice, in a realistic topography, convergence zones and canals naturally limit the scattering of the clasts. Finally, many

potential applications will be more sensitive to \bar{x} , which determines the mean residence time, rather than σ .

4.2 “Non-linear” diffusion with a slope threshold

Here we illustrate the scattering of the clasts in the case of “non-linear” diffusion. We use Eqs. (5) and (6) with $S_c = 0.105$, a value just above the plan slope $S = 0.1$. This S_c value is much lower than published values ~ 0.6 (e.g. Roering et al., 2001), but our goal here is to compare this with the previous linear diffusion case. We also use coarser clasts with $R = 0.05$ m to obtain similar mean transport rates as in the previous case. The other parameter values are the same as in the linear diffusion case.

Figure 4 shows the pattern of clasts at different times. Compared to the linear diffusion case, once detached, clasts move a longer distance because $L \gg dx$ when $S \sim S_c$. Consequently, the clast population presents a much more elongated distribution along the hillslope. This pattern seems consistent with colluvial sediment covering steep hillslopes in the real world. The mean travel distance \bar{x} does not depend on dx nor dt , while σ does depend on dx . Note that σ varies as \sqrt{t} but with much larger values representing the downstream spreading of the clasts. The different patterns obtained in the linear and non-linear diffusion cases illustrate the progressive transition from diffusion for $S \ll S_c$ to ballistic movements for $S \sim S_c$.

4.3 River transport

River transport usually implies the formation of incisions, local depositions and lateral movements of sediment by bank erosion. The movement of clasts associated with this dynamic is an active research field (e.g. Seizilles et al., 2014). In order to verify that \bar{x} is consistent with the description of the river process using erosion and deposition rates, we simplify the problem by using a planar and stable portion of a river bed. This bed corresponds to the same bedrock plan with the slope of 0.1 used above. Detachment is determined by $\epsilon = KqS$. The same K is used for both bedrock and

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sediment. The flow width w is set at cell size dx , Thus, ϵ corresponds to the “stream power law” (e.g. Whipple and Tucker, 1999). The transport length $L = \xi q$ is close to dx , by setting ξ to a low value 10^{-6} (L cannot be smaller than dx). Consequently, the sediment flux per unit width q_s is close to $\epsilon dx \propto q$. The linear relationship between q_s and q prevents incisions from forming (Smith and Bretherton, 1972), which is what we want here. A precipitation rate of $P = 1 \text{ m yr}^{-1}$ is imposed on the grid. Thus, the water discharge per unit width $q = Px$ grows linearly downstream, as does the probability of clast entrainment.

The mean travel velocity of a clast \dot{x} can be written as $\dot{x} = \frac{q_s}{2R} = \frac{Kdx}{2R} SPx$. A solution to this equation is $x(t) = x_o \exp(\frac{Kdx}{2R} PSt)$ where x_o is the initial distance. For a clast population, \bar{x} must follow this solution.

We use larger clasts with a radius of 0.05 m in order to have a transport rate of the same order of magnitude as in the diffusion case. Figure 5 shows that clasts are spread from the injection point, forming a downstream plume. This pattern results from the downstream increase in the entrainment probabilities. \bar{x} fits the above solution acceptably well and does not depend on dx or dt . The scattering σ increases more than linearly because the water discharge increases downstream. Note that despite similar plan view patterns, σ evolves very differently in the non-linear diffusion and river cases. This difference illustrates the interest of coupling clasts and flux in order to identify erosion-transport processes.

4.4 A more general case

We present here an illustration of clasts moving in a mountain-foreland context. Our goal is not to precisely analyse the clast dynamics in that case, but to qualitatively describe a possible situation in a real case scenario.

The mountain-foreland system consists of an uplifting block (the mountain) and a stable domain where sediment eroded from the block is deposited or in transit. The grid is 200×200 cells, the cell size is 500 m, and the mountain is 37.5 km wide, which

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



could correspond to the Pyrenees, or certain Andean foothills in Argentina for example. A constant uplift rate is fixed at 0.5 mm yr^{-1} . The precipitation rate is homogeneous and constant at 1 m yr^{-1} . Sediment, water and clasts can escape the model grid on the southern boundary, where the elevation is fixed at 0 m. The east and west boundaries are periodic, so that sediment, water and clasts can go out from one side and re-enter on the other side. No sediment, water or clasts can leave the northern side. Erosion parameters are chosen so that the maximum elevation is 3000 m when the topography reaches a rough steady-state at around 3 My (see Fig. 6 caption).

The experiment begins from a flat topography with small random elevations ($< 1 \text{ m}$) (Fig. 6). The sediment eroded from the uplifting block is propagated through the foreland, forming alluvial fans and rivers. An upstream drainage develops in the mountain, and a divergent river network progrades over the alluvial apron. Initially, 2000 clasts were seeded at two different places in the mountain at depths of 500–600 m (green clasts) and 400–500 m (red clasts), as if they were belonging to two intrusive bodies. When erosion reaches this depth, clasts begin to exhume, move and deposit in the foreland. The red clasts exhume first and are spread within an alluvial fan. Deposition of the green clasts ends 0.5 Myr after the deposition of the red clasts. The complete deposition of both populations lasts 1 Myr. Both populations are mainly included in a sediment layer that is thinner than 100 m. They are mixed at similar depths at the transition zone between the two alluvial fans. This situation illustrates a lateral diachronism between close alluvial aprons which may have been erroneously correlated in the field.

If these grains were detrital zircons of different ages, the analysis of their age distribution at different places would allow the mixed zone to be mapped, and thus, the dynamics of the lateral alluvial fan to be reconstructed.

5 Perspectives

The modelling approach described in this paper and its developments may have different applications which we propose in the following.

5.1 River-erosion law

River dynamics involve processes acting on a large range of time periods, from hours in the case of catastrophic flooding to thousands of years to transport huge volumes of glacier sediment, for example. Determining simplified laws to predict this complexity remains a challenge (Murray, 2007). Recently, Kasprak et al. (2015) showed a tight relationship between the inter-bar distance and travel distance of marked grains in experiments on braided rivers. At the same time, Davy and Lague (2009) evidenced a relationship between this inter-bar distance and L , a fundamental transport-length parameter used in the Eros landscape evolution model and which has been also introduced in Cidre. The link between both observations can be made using the modelling approach proposed in this paper. By using a try-and-guess approach, investigating different erosion and deposition laws of different complexities, it may be possible to identify which laws predict the distribution of the clasts and the river bed forms observed in experiments (Kasprak et al., 2015) or rivers (Bradley et al., 2010). As far as we know, the study that was carried out by introducing moving clasts in Eros is the only study of this kind (Davy and Lague, 2012).

5.2 Grain sorting, downstream fining, coarsening

The origin of grain size patterns along fluvial systems is still being debated. Explanations include size-selective entrainment or deposition, and attrition (Paola et al., 1992a, b). By using grains of different sizes in Cidre, it may be possible to rank these processes along the whole fluvial system. Selective deposition could be simply included by allowing the deposition probability to depend on clast size. Physical attrition could also be included by decreasing the clast size according to the travel distance. Local surface coarsening could be also studied, as clast entrainment depends on their size. This model may represent an alternative to models requiring a depth discretisation of sediment layers containing different grain sizes (Cohen et al., 2009).

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.3 Grain-size dependent landscape evolution

In the real world, material erosion and transport depend on the clast size and the grain size evolves spatially. The feedback of grain size on the landscape dynamics has been little explored (Sklar et al., 2008). In Cidre, entrainment and deposition parameters may vary according to the mean radius of the grains on each cell. The number of model parameters would be kept small (K , L specified initially for the median size, for example), while accounting for complex feedbacks. This could be an alternative to other approaches which incorporate explicitly different classes of grain sizes to simulate fluvial dynamics (e.g. Coulthard et al., 2002).

5.4 Provenance problems and alluvial or placer-type deposits

Provenance studies on detrital grains help constrain the chronology of the exhumation of the sediment source (e.g. Roddaz et al., 2005; Rodríguez et al., 2012). Grains of a particular lithology can be used to identify the provenance of certain material of economic interest, like distant gravels eroded from a hidden porphyry copper deposit (Sil-litoe, 2010). These provenance studies come up against uncertainties regarding the landscape evolution. The coupled modelling of landscape evolution and clast movement may help constrain this link in specific cases.

Placer-type deposits are secondary ores that can contain free particles with very fine gold and other native metals (e.g., platinum-group elements, PGE). The occurrence of gold grains in supergene environments, such as soil, sediments and placers in rivers, is controlled by physical (as well as bio-geochemical) processes of redistribution from a distant gold-quartz vein (Reich and Vasconcelos, 2015). Gold grains between 0.1 and 4 mm in diameter constitute most of alluvial and eluvial gold, and represent economically important ores, such as the Witwatersrand paleoplacer gold deposits in South Africa (Mossman et al., 1999; Fairbrother et al., 2013). Measurements of geometric grain parameters such as roundness, degree of bending, and grain flatness have been used to constrain the source-type of the gold and transportation distances (e.g. Townley

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2003; Mudaliar et al., 2007). Therefore, an approach that involves the modeling of gold grain transport during the landscape evolution might provide an independent test to check the validity of transport distance estimations based on grain morphology.

5.5 Cosmogenic nuclides

5 The analysis and modelling of cosmogenic nuclide concentration in individual clasts give quantitative information about the erosion-transport processes at the landscape scale (Repka et al., 1997; Codilean et al., 2008; Gayer et al., 2008; McPhillips et al., 2014; Carretier et al., 2009; Carretier and Regard, 2011; Vassallo et al., 2011). It would be straightforward to compute the cosmogenic nuclide concentration evolution in clasts
10 in Cidre, using for example the model of Carretier and Regard (2011). This may be particularly useful to link cosmogenic nuclides from buried sediment in basins to quantify the erosion-sediment history of mountain-foreland systems.

5.6 Tracing weathering material

15 In the 1990's, Raymo and Ruddiman (1992) proposed that the uplift of Tibet has driven the post-Eocene global atmospheric cooling by accelerating the chemical weathering of silicate rocks and the associated consumption of atmospheric CO₂. However, recent studies at the foot of the Himalayas and Andes suggest that the weathering flux may be larger in the foreland basin than in the mountain range itself (Lupker et al., 2012). These studies call for for the development of 3-D models able to trace the weathered
20 material from source to sink (Anderson et al., 2012). The clast-flux modelling approach may be developed within that scope, by allowing clasts to weather. This will be the aim of a future paper.

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



6 Conclusions

The algorithm predicts a consistent clast velocity and surface erosion rate in simple cases. The mean travel distance of the clasts does not depend on the model cell size or time step. The scattering of the clasts depends on the cell size and is overestimated.

5 Nevertheless, decreasing the cell size decreases the overestimation. This model has numerous potential applications allowing field data on distinct grains to be linked to a large-scale landscape evolution. The differences between the simple river and hillslope cases illustrated here (e.g. Figs. 4 and 5) show that the movement of tracers in the field or in experiments may provide a way to establish local erosion-transport laws at spatial
10 and temporal scales adapted to landscape evolution models.

Appendix A

In order to derive the variation in the soil or river bed surface elevation z two downstream length dx systems are considered above and below the surface. In the first subsurface system, the mass balance is:

$$15 \quad \frac{\partial z}{\partial t} = -\frac{\partial q_s}{\partial x}, \quad (\text{A1})$$

where q_s [$L^2 T^{-1}$] is the sediment flux per unit width. In the second fluid system, from a Lagrangian point of view the transported mass of the sediment M varies as

$$\frac{DM}{Dt} = \frac{\partial M}{\partial t} + u \frac{\partial M}{\partial x} = \rho w dx \epsilon - \rho w dx D, \quad (\text{A2})$$

where u is the flow velocity, w is the flow width, ρ is the sediment density and ϵ and D
20 are the erosion-detachment and deposition rates, respectively.

Assuming the mass of the sediment in movement is locally constant in the water (steady state – $\frac{\partial M}{\partial t} = 0$) and recognising that $u \frac{\partial M}{\partial x} = \rho w dx \frac{\partial q_s}{\partial x}$, the mass balance in the

first system is

$$\frac{\partial z}{\partial t} = -\epsilon + D. \quad (\text{A3})$$

Acknowledgements. This paper is a contribution to the LMI COPEDIM (funding from IRD). S. Carretier thanks the Departamento de Geología of the Universidad de Chile for its welcome. M. Reich acknowledges funding by the MSI grant “Millennium Nucleus for Metal Tracing Along Subduction (NC130065)”. Cidre sources are available upon request. The initial Cidre core was developed by B. Poisson at BRGM. This paper is dedicated to her memory.

References

- Anderson, S., Anderson, R., and Tucker, G.: Landscape scale linkages in critical zone evolution, *C. R. Geosci.*, 344, 586–596, 2012. 1240
- Andreotti, B., Claudin, P., and Pouliquen, O.: Measurements of the aeolian sand transport saturation length, *Geomorphology*, 123, 343–348, 2010. 1229
- Andrews, D. J. and Hanks, T. C.: Scarp degraded by linear diffusion: Inverse solution for age, *J. Geophys. Res.*, 90, 10193–10208, 1985. 1227
- Beaumont, C., Fullsack, P., and Hamilton, J.: Erosional control of active compressional orogens, in: *Thrust Tectonics*, edited by: Mc Clay, K. R., Chapman and Hall, New York, USA, 1992. 1229
- Binnie, S. A., Phillips, W. M., Summerfield, M. A., and Fifield, L. K.: Tectonic uplift, threshold hillslopes, and denudation rates in a developing mountain range, *Geology*, 35, 743–746, 2007. 1227
- Bradley, D. N., Tucker, G. E., and Benson, D. A.: Fractional dispersion in a sand bed river, *J. Geophys. Res.-Earth*, 115, F00A09, doi:10.1029/2009JF001268, 2010. 1238
- Carretier, S. and Regard, V.: Is it possible to quantify pebble abrasion and velocity in rivers using terrestrial cosmogenic nuclides?, *J. Geophys. Res.-Earth*, 116, F04003, doi:10.1029/2011JF001968, 2011. 1225, 1240
- Carretier, S., Niviere, B., Giamboni, M., and Winter, T.: Do river profiles record along-stream variations of low uplift rate?, *J. Geophys. Res.-Earth*, 111, F02024, doi:10.1029/2005JF000419, 2006. 1223

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Carretier, S., Regard, V., and Soual, C.: Theoretical cosmogenic nuclide concentration in river bedload clasts: Does it depend on clast size?, *Quat. Geochronol.*, 4, 108–123, 2009. 1223, 1224, 1225, 1227, 1228, 1231, 1240
- Carretier, S., Godderis, Y., Delannoy, T., and Rouby, D.: Mean bedrock-to-saprolite conversion and erosion rates during mountain growth and decline, *Geomorphology*, 209, 29–52, 2014. 1227, 1231
- Chabaux, F., Granet, M., Pelt, E., France-Lanord, C., and Galy, V.: 238U-234U-230Th disequilibrium and timescale of sedimentary transfers in rivers: clues from the Gangetic plain rivers, *J. Geochem. Explor.*, 88, 373–375, 2006. 1224
- Charru, F.: Selection of the ripple length on a granular bed sheared by a liquid flow, *Phys. Fluids*, 18, 121508-1–121508-9, 2006. 1226, 1229
- Codilean, A., Bishop, P., Stuart, F., Hoey, T., Fabel, D., and Freeman, S. P. H. T.: Single-grain cosmogenic ²¹Ne concentrations in fluvial sediment reveal spatially variable erosion rates, *Geology*, 36, 159–162, 2008. 1224, 1225, 1240
- Cohen, S., Willgoose, G., and Hancock, G.: The mARM spatially distributed soil evolution model: A computationally efficient modeling framework and analysis of hillslope soil surface organization, *J. Geophys. Res.*, 114, F03001, doi:10.1029/2008JF001214, 2009. 1238
- Coulthard, T. J., Macklin, M. G., and Kirkby, M. J.: Simulating upland river catchment and alluvial fan evolution, *Earth Surf. Proc. Land.*, 27, 269–288, 2002. 1239
- Culling, W.: Analytical theory of erosion, *J. Geol.*, 68, 336–344, 1960. 1223
- Davy, P. and Lague, D.: The erosion/transport equation of landscape evolution models revisited, *J. Geophys. Res.*, 114, F03007, doi:10.1029/2008JF001146, 2009. 1223, 1226, 1229, 1230, 1231, 1238
- Davy, P. and Lague, D.: Sediment transfer in rivers: linking sediment grain motions with geomorphic dynamics with a focus on process stochasticity, in: *Colloque de l'Académie des Sciences, Paris, France, 26–27 March 2012, 24–25, 2012.* 1225, 1238
- Einstein, A.: The motion of elements suspended in static liquids as claimed in the molecular kinetic theory of heat, *Annalen Der Physik*, 17, 549–560, 1905. 1233
- Fairbrother, L., Brugger, J., Shapter, J., Laird, J., Southam, G., and Reith, F.: Supergene gold transformation: Biogenic secondary and nano-particulate gold from arid Australia, *Chem. Geol.*, 320–321, 17–31, 2013. 1239

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Falcini, F., Foufoula-Georgiou, E., Ganti, V., Paola, C., Voller, V. R.: A combined nonlinear and nonlocal model for topographic evolution in channelized depositional systems, *J. Geophys. Res.-Earth*, 118, 1617–1627, 2013. 1228
- Foufoula-Georgiou, E., Ganti, V., and Dietrich, W. E.: A nonlocal theory of sediment transport on hillslopes, *J. Geophys. Res.-Earth*, 115, F00A16, doi:10.1002/jgrf.20108, 2010. 1228
- Furbish, D. J. and Haff, P. K.: From divots to swales: Hillslope sediment transport across divers length scales, *J. Geophys. Res.-Earth*, 115, F03001, doi:10.1029/2009JF001576, 2010. 1228
- Furbish, D. J. and Roering, J. J.: Sediment disentrainment and the concept of local versus nonlocal transport on hillslopes, *J. Geophys. Res.-Earth*, 118, 937–952, 2013. 1228
- Gayer, E., Mukhopadhyay, S., and Meade, B.: Spatial variability of erosion rates inferred from the frequency distribution of cosmogenic he-3 in olivines from hawaiian river sediments, *Earth Planet. Sci. Lett.*, 266, 303–315, 2008. 1224, 1225, 1240
- Gingold, R. and Monaghan, J.: Smoothed particle hydrodynamics: theory and application to non-spherical stars, *Mon. Not. R. Astron. Soc.*, 181, 375–389, 1977. 1224
- Hanks, T. C.: The age of scarp-like landforms from diffusion-equation analysis, in: *Quaternary geochronology: methods and applications*, edited by: Noller, J. S., Sowers, J. M., and Lettis, W. R., AGU Washington DC, USA, 313–338, 1999. 1227
- Haschenburger, J. K.: The rate of fluvial gravel dispersion, *Geophys. Res. Lett.*, 38, L24403, doi:10.1029/2011GL049928, 2011. 1223
- Hassan, M., Church, M., and Schick, A.: Distance of movement of coarse particles in gravel bed streams, *Wat. Resour. Res.*, 27, 503–511, 1991. 1223
- Howard, A. D.: Badland morphology and evolution: interpretation using a simulation model, *Earth Surf. Proc. Land.*, 22, 211–227, 1997. 1223
- Howard, A. D. and Kerby, G.: Channel changes in badlands, *Geol. Soc. Am. Bull.*, 94, 739–752, 1983. 1223
- Kasprak, A., Wheaton, J. M., Ashmore, P. E., Hensleigh, J. W., and Peirce, S.: The relationship between particle travel distance and channel morphology: Results from physical models of braided rivers, *J. Geophys. Res.-Earth*, 120, 55–74, 2015. 1223, 1238
- Kooi, H. and Beaumont, C.: Escarpment evolution on high-elevation rifted margins: Insights derived from a surface processes model that combines diffusion, advection, and reaction, *J. Geophys. Res.*, 99, 12191–12209, 1994. 1223

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kooi, H. and Beaumont, C.: Large-scale geomorphology: Classical concepts reconciled and integrated with contemporary ideas via surface processes model, *J. Geophys. Res.*, 101, 3361–3386, 1996. 1223

Lague, D.: Reduction of long-term bedrock incision efficiency by short-term alluvial cover intermittency, *J. Geophys. Res.*, 115, F02011, doi:10.1029/2008JF001210, 2010. 1230

Lague, D. and Davy, P.: Constraints on the long-term colluvial erosion law by analyzing slope-area relationships at various tectonic uplift rates in the Siwaliks (Nepal), *J. Geophys. Res.* 108, 2129, doi:10.1029/2002JB001893, 2003. 1223

Lajeunesse, E., Malverti, L., and Charru, F.: Bed load transport in turbulent flow at the grain scale: Experiments and modeling, *J. Geophys. Res.-Earth*, 115, F04001, doi:10.1029/2009JF001628, 2010. 1223

Lajeunesse, E., Devauchelle, O., Houssais, M., and Seizilles, G.: Tracer dispersion in bedload transport, *Adv. Geosci.*, 37, 1–6, doi:10.5194/adgeo-37-1-2013, 2013. 1225

Lajeunesse, E., Devauchelle, O., Lachaussée, F., Claudin, P., 2015. Bedload Transport in Laboratory Rivers: the Erosion-Deposition Model, in: *ravel-bed Rivers: Gravel Bed Rivers and Disasters*, <http://www.gbr8.dpri.kyoto-u.ac.jp/paper/Lajeunesseetal..pdf>, 2015. 1226, 1229

Loget, N., Davy, P., and van den Driessche, J.: Mesoscale fluvial erosion parameters deduced from modeling the Mediterranean sea level drop during the Messinian (late Miocene), *J. Geophys. Res.*, 11, F03005, doi:10.1029/2005JF000387, 2006. 1223

Lupker, M., France-Lanord, C., Galy, V., Lave, J., Gaillardet, J., Gajurel, A. P., Guilmette, C., Rahman, M., Singh, S. K., and Sinha, R.: Predominant floodplain over mountain weathering of Himalayan sediments (Ganga basin), *Geochemical et Cosmochemica Acta*, 84, 410–432, 2012. 1240

Malmaeus, J. and Hassan, M.: Simulation of individual particule movement in a gravel streambed, *Earth Surf. Process. Landforms*, 27, 81–97, 2002. 1225, 1232

McPhillips, D., Bierman, P. R., and Rood, D. H.: Millennial-scale record of landslides in the Andes consistent with earthquake trigger, *Nat. Geosci.*, 7, 925–930, 2014. 1224, 1240

Mitasova, H., Thaxton, C., Hofierka, J., McLaughlin, R., Moore, A., Mitas, L.: Path sampling method for modeling overland water flow, sediment transport and short term terrain evolution in Open Source GIS, in: *Proceedings of the XVth International Conference on Computational Methods in Water Resources (CMWR XV)*, edited by: Miller, C., Farthing, M., Gray, V., and Pinder, G., 1479–1490, 2004. 1225

**Landscape with
clasts**

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Mossman, D., Reimer, T., and Durstling, H.: Microbial processes in gold migration and deposition: modern analogues to ancient deposits, *Geosci. Canada*, 26, 131–140, 1999. 1239
- Mudaliar, G., Richards, J., and Eccles, D.: Gold, Platinum and Diamond Placer Deposits in Alluvial Gravels, Whitecourt, Alberta, Alberta Geological Survey Report, 89, 22 pp., 2007. 1240
- 5 Murray, A. B.: Reducing model complexity for explanation and prediction, *Geomorphology*, 90, 178–191, 2007. 1224, 1238
- Murray, A. B. and Paola, C.: Properties of a cellular braided-stream model, *Earth Surf. Proc. Land.*, 22, 1001–1025, 1997. 1230
- 10 Nicholas, A. and Quine, T.: Crossing the divide: Representation of channels and processes in reduced-complexity river models at reach and landscape scales, *Geomorphology*, 90, 38–339, 2007. 1231
- Paola, C., Heller, P. L., and Angevine, C. L.: The large-scale dynamics of grain-size variation in alluvial basins – 1: Theory, *Basin Res.*, 4, 73–90, 1992a. 1238
- 15 Paola, C., Parker, G., Seal, R., Sinha, S., Southard, J., and Wilcock, R.: Downstream fining by selective deposition in a laboratory flume, *Science*, 258, 1757–1760, 1992b. 1238
- Pepin, E., Carretier, S., and Hérail, G.: Erosion dynamics modelling in a coupled catchment-fan system with constant external forcing, *Geomorphology*, 122, 78–90, 2010. 1231
- Perron, J. T.: Numerical methods for nonlinear hillslope transport laws, *J. Geophys. Res.-Earth*, 20 116, F02021, doi:10.1029/2010JF001801, 2011. 1227, 1229
- Raymo, M. and Ruddiman, W.: Tectonic forcing of late Cenozoic climate, *Nature*, 359, 117–122, 1992. 1240
- Reich, M. and Vasconcelos, P.: Geological and economic significance of supergene metal deposits, *Elements*, 11, 305–310, 2015. 1239
- 25 Repka, J. L., Anderson, R. S., and Finkel, R. C.: Cosmogenic dating of fluvial terraces, Fremont River, Utah, *Earth Planet. Sci. Lett.*, 152, 59–73, 1997. 1225, 1240
- Roddaz, M., Viers, J., Brusset, S., Baby, P., and Hérail, G.: Sediment provenances and drainage evolution of the Neogene Amazonian foreland basin, *Earth Planet. Sci. Lett.*, 239, 57–78, 2005. 1223, 1239
- 30 Rodríguez, M., Pinto, L., and Encinas, A.: Cenozoic erosion in the Andean forearc in Central Chile (33–34° S): Sediment provenance inferred by heavy mineral studies, *GSA Spec. Papers*, 487, 141–162, 2012. 1223, 1239

Landscape with clasts

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Roering, J. J., Kirchner, J. W., and Dietrich, W. E.: Evidence for nonlinear, diffusive sediment transport on hillslopes and implications for landscape morphology, *Wat. Resour. Res.*, 35, 853–870, 1999. 1227, 1250

Roering, J. J., Kirchner, J. W., Sklar, L. S., and Dietrich, W. E.: Hillslope evolution by nonlinear creep and landsliding: An experimental study, *Geology*, 29, 143–146, 2001. 1227, 1228, 1235

Roering, J. J., Kirchner, J., Sklar, L., and Dietrich, W.: Hillslope evolution by nonlinear creep and landsliding: An experimental study: Reply, *Geology*, 30, 482–482, 2002. 1228

Seizilles, G., Lajeunesse, E., Devauchelle, O., and Bak, M.: Cross-stream diffusion in bedload transport, *Phys. Fluids*, 26, 013302, doi:10.1063/1.4861001, 2014. 1231, 1235

Sillitoe, R. H.: Porphyry Copper Systems, *Economic Geology*, 105, 3–41, 2010. 1239

Sklar, L. S. and Dietrich, W. E.: Sediment and rock strength controls on river incision into bedrock, *Geology*, 29, 1087–1090, 2001. 1223

Sklar, L., Dietrich, B., and William, E.: Implications of the saltation-abrasion bedrock incision model for steady-state river longitudinal profile relief and concavity, *Earth Surf. Proc. Land.*, 33, 1129–1151, 2008. 1239

Smith, T. R. and Bretherton, F. P.: Stability and the conservation of mass in drainage basin evolution, *Wat. Resour. Res.*, 8, 1506–1529, 1972. 1236

Stock, J. and Montgomery, D. R.: Geological constraints on bedrock river incision using the stream power law, *J. Geophys. Res.*, 104, 4983–4993, 1999. 1223

Tomkin, J., Brandon, M. T., Pazzaglia, F. J., Barbour, J. R., and Willet, S. D.: Quantitative testing of bedrock incision models for the Clearwater River, NW Washington State, *J. Geophys. Res.*, 108, 2308, doi:10.1029/2001JB000862, 2003. 1223

Townley, B., Herail, G., Maksaev, V., Palacios, C., de Parseval, P., Sepulveda, F., Orellana, R., Rivas, P., and Ulloa, C.: Gold grain morphology and composition as an exploration tool: application to gold exploration in covered areas, *Geol. Soc. London*, 3, 29–38, 2003. 1239

Tucker, G. E.: Drainage basin sensitivity to tectonic and climatic forcing: Implications of a stochastic model for the role of entrainment and erosion thresholds, *Earth Surf. Proc. Land.*, 29, 185–205, 2004. 1229

Tucker, G. E. and Bradley, D. N.: Trouble with diffusion: Reassessing hillslope erosion laws with a particle-based model, *J. Geophys. Res. Earth*, 115, F00A10, doi:10.1029/2009JF001264, 2010. 1225, 1226, 1228

- Tucker, G. E. and Hancock, G. R.: Modelling landscape evolution, *Earth Surf. Proc. Land.*, 35, 28–50, 2010. 1223
- Tucker, G. E. and Whipple, K.: Topographic outcomes predicted by stream erosion models: Sensitivity analysis and intermodel comparison, *J. Geophys. Res.*, 107, 2179, doi:10.1029/2001JB000162, 2002. 1223
- 5 van der Beek, P. and Bishop, P.: Cenozoic river profile development in the Upper Lachlan catchment (SE Australia) as a test of quantitative fluvial incision models, *J. Geophys. Res.*, 108, 2309, doi:10.1029/2002JB002125, 2003. 1223
- van Milligen, B. and Bons, P.: Hillslope evolution by nonlinear creep and landsliding: An experimental study: Comment, *Geology*, 30, 481–482, 2002. 1228
- 10 Vassallo, R., Ritz, J.-F., and Carretier, S.: Control of geomorphic processes on ^{10}Be concentrations in individual clasts: Complexity of the exposure history in Gobi-Altay range (Mongolia), *Geomorphology*, 135, 35–47, 2011. 1240
- Whipple, K. X. and Tucker, G. E.: Dynamics of the stream-power incision model: implication for height limits of mountain ranges, landscape response timescales, and research needs, *J. Geophys. Res.*, 104, 17661–17674, 1999. 1236
- 15 Whipple, K. X. and Tucker, G. E.: Implication of sediment-flux-dependent river incision models for landscape evolution, *J. Geophys. Res.*, 107, 2039, doi:10.1029/2000JB000044, 2002. 1223, 1229, 1230
- 20 Whipple, K. X., Hancock, G. S., and Anderson, R. S.: River incision into bedrock: Mechanics and relative efficacy of plucking, abrasion and cavitation, *Geol. Soc. Am. Bull.*, 112, 490–503, 2000. 1223
- Willgoose, G., Bras, R. L., and Rodriguez-Iturbe, I.: Result from a new model of river basin evolution, *Earth Surf. Proc. Land.*, 16, 237–254, 1991. 1223

**Landscape with
clasts**

S. Carretier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 1. Illustration of erosion-deposition processes in Cidre. **(a)** Multiple flow and the different calculated fluxes: q_s [$L^2 T^{-1}$] incoming sediment flux per unit width. T_i [$L T^{-1}$] transferred sediment rate in each downward direction. ϵ [$L T^{-1}$] detachment rate of material (sediment or bedrock) from the cell calculated in the steepest direction. D [$L T^{-1}$] deposited sediment rate on the cell. Q_{sl} [$L^3 T^{-1}$] lateral volumetric sediment flux deposited on the cell. S_i slope in each downward direction. w = flow width. dx = cell width. The incoming water flow is propagated towards lower cells proportionally to the local slope in each direction (not illustrated). **(b)** Probabilities that clasts will sediment, transfer or detach using fluxes calculated by Cidre.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Landscape with
clasts

S. Carretier et al.




Figure 2. Comparison between the profile evolutions of the hill predicted by Eqs. (5) and (6) (the model proposed in this paper – in red) and Eqs. (3) and (4) (the “non-linear” diffusion model proposed for example by Roering et al., 1999 – in black). The “non-linear” diffusion model is solved by explicit finite differences along a 2-D profile. The distance between points $dx = 10$ m, $\kappa' = 0.01 \text{ m}^2 \text{ yr}^{-1}$, $S_c = 0.6$, $dt = 1000 \text{ yr}$. For the model presented in this paper, the diffusivity $\kappa' = \kappa dx = 0.01 \text{ m}^2 \text{ yr}^{-1}$ and $S_c = 0.57$ ($= \tan 30^\circ$). The red profiles represent 2-D averaged elevations across 3-D topographies. Note that S_c is slightly larger in the “non-linear” model because of stability reasons, the non-linear flux in Eq. (4) is approximated by the tangent to the curve $q_s = f(S)$ when S is close to S_c , resulting in a small underestimation of q_s . **(a)** Relaxation from a triangular hill with a slope of 0.5. Profiles every 2.5 Myr. **(b)** Uplift (0.1 mm yr^{-1}) from an initial horizontal surface. Profiles every 0.5 Myr. The elevation is fixed at 0 m at the two boundary conditions in **(a)** and **(b)**.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. Model test in the linear diffusion case ($e = \kappa S$ and $L = dx$). The initial plan slope is $S = 0.1$. 1000 clasts are initially set at the surface and top of this plan. The diffusivity $\kappa' = \kappa dx = 0.01 \text{ m}^2 \text{ yr}^{-1}$ in all the cases. **(a)** Example of clasts spreading in the case $\kappa = 0.0001$, $dx = 100 \text{ m}$ and $dt = 50 \text{ yr}$. **(b)** Evolution of the mean travel distances of the clasts for the different dx , dt and κ values. Note that the mean transport distance fits well with the prediction and does not significantly depend on dt or dx . **(c)** Evolution of the standard deviation of the travel distances of the clasts σ . Note that $\sigma \propto \sqrt{t}$ as predicted by the diffusion theory, although it overestimates by more than one order of magnitude the predicted value of 0.036 km at $t = 50 \text{ kyr}$ for the green and red curves.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Landscape with
clasts

S. Carretier et al.




Figure 4. Model test in the non linear diffusion case ($\epsilon = \kappa S$ and $L = \frac{dx}{1-(S/S_c)^2}$). The initial plan slope is $S = 0.1$ and S_c is set close at 0.105. The mean sediment flux per unit width $q_s = \epsilon dx$ is the same in all cases (same $\kappa' = \kappa dx$). 1000 clasts are initially set at the surface and top of this plan. **(a)** Example of clasts spreading in the case $\kappa = 0.0001$, $dx = 100$ m and $dt = 50$ yr. **(b)** Evolution of the mean travel distances of the clasts for the different dx , dt and κ values. Note that the mean transport distance fits well with the prediction and does not depend significantly on dt or dx . **(c)** Evolution of the standard deviation of the travel distances of the clasts σ . Note that $\sigma \propto \sqrt{t}$ as predicted by the diffusion theory.

Landscape with
clasts

S. Carretier et al.




Figure 5. Model test in the river case ($\epsilon = KqS$ and $L = dx$). The initial plan slope is $S = 0.1$ and the precipitation rate is 1 m yr^{-1} . Initially, 1000 clasts are set at the surface and top of this plan. The mean sediment flux per unit width $q_s = \epsilon dx$ is the same in all cases by keeping $Kdx = 310^{-5}$ constant. **(a)** Example of clasts spreading in the case $K = 310^{-7} \text{ m}^{-1}$, $dx = 100$ m and $dt = 50$ m. **(b)** Evolution of the mean travel distances of the clasts for the different dx , dt and K . The blue color indicates an increase in the downstream water discharge. Note that the mean transport distance fits well with the prediction and does not significantly depend on dt or dx . **(c)** Evolution of the standard deviation for the travel distances of the clasts σ . Note that σ increases exponentially downstream, differently from the diffusion cases.

Landscape with
clasts

S. Carretier et al.




Figure 6. Example of clasts exhumation from two locations (intrusive body) located initially at a depth of 0.5 km (green clasts) and 0.4 km (red clasts). The final maximum elevation for the mountain is 3000 m. The domain size is 100×100 km. There is constant rainfall over the whole grid (1 m yr^{-1}). The northern block is continuously uplifting at 0.5 mm yr^{-1} . The elevation of the clasts is increased to see them. Note that the green grains stop depositing 1 Myr after the red ones. Nevertheless, both populations are mixed between the two alluvial fans at similar depth, showing that 1 Myr of difference may separate two adjacent sediment layers that are apparently synchronous. $K_{\text{sediment}} = 0.6 \times 10^{-3} \text{ m yr}^{-1}$, $K_{\text{bedrock}} = 0.3 \times 10^{-3} \text{ m yr}^{-1}$, $m = 0.5$, $n = 1$, $\xi = 0.1 \text{ yr m}^{-1}$, $\alpha_{\text{sediment}} = 0.01$, $\kappa_{\text{sediment}} = \kappa_{\text{bedrock}} = 10^{-4} \text{ m yr}^{-1}$, $S_{\text{sediment}} = S_{\text{bedrock}} = 0.8$, $dx = 500 \text{ m}$, $dt = 100 \text{ yr}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

