

Dr. Louis Andreani

TU Bergakademie Freiberg, Institut für Geologie,
Bernhard-von-Cotta Str. 2, 09599 Freiberg, Germany.



**TECHNISCHE UNIVERSITÄT
BERGAKADEMIE FREIBERG**

Die Ressourcenuniversität. Seit 1765.

**Faculty of Geosciences, Geoengineering and Mining
Institute of Geology**

**To: Dr. Giulia Sofia
Associate Editor, Earth Surface Dynamics.**

Freiberg, December 22, 2015

Dear Editor,

Please find attached the revised version of our manuscript. We added the suggested references to the size of the moving window and modified our manuscript accordingly. We provide with this letter the revised version of our manuscript (modified sections are highlighted).

Thank you again for the editorial handling of this manuscript.

Sincerely,

Louis Andreani,

On behalf of all co-authors.

Es bestehen Unterschiede zwischen den Dokumenten.

Neues Dokument:

[Manuscript_Esurf_v3](#)

55 pages (32.28 MB)

22.12.2015 19:09:34

[Zum Anzeigen der Ergebnisse verwendet.](#)

Altes Dokument:

[Manuscript_Esurf](#)

55 Seiten (32.28 MB)

22.12.2015 19:09:32

[Beginn: Erste Änderung auf Seite 7.](#)

Es wurden keine Seiten gelöscht

Wie ist dieser Bericht zu lesen

Highlight zeigt eine Änderung an.

Gelöscht zeigt gelöschten Inhalt an.

 zeigt an, dass Seiten geändert wurden.

 zeigt an, dass Seiten verschoben wurden.

Geomorphic analysis of transient landscapes in the Sierra Madre de Chiapas and Maya Mountains (northern Central America): implications for the North American – Caribbean – Cocos plate boundary

L. Andreani^{1,2} and R. Gloaguen^{1,2}

¹TU Bergakademie Freiberg, Institut für Geologie, Freiberg, Germany.

²Helmholtz Zentrum Dresden Rossendorf, Helmholtz Institute Freiberg for Resource Technology, Division "Exploration Technology", Freiberg, Germany.

Correspondence to: L. Andreani (louis.andreani@googlemail.com), R. Gloaguen (r.gloaguen@hzdr.de)

Abstract. We use a geomorphic approach in order to unravel the recent evolution of the diffuse triple junction between the North American, Caribbean, and Cocos plates in northern Central America. We intend to characterize and understand the complex tectonic setting that produced an intricate pattern of landscapes using tectonic geomorphology, and available geological and geophysical data. We classify regions with specific relief characteristics and highlight uplifted relict landscapes in northern Central America. We also analyze the drainage network from the Sierra Madre de Chiapas and Maya Mountains in order to extract information about potential vertical displacements.

Our results suggest that most of the landscapes of the Sierra Madre de Chiapas and Maya Mountains are in transient stage. Topographic profiles and morphometric maps highlight elevated relict surfaces that are characterized by a low amplitude relief. The river longitudinal profiles display upper reaches witnessing these relict landscapes while lower segments characterized by multiple knickpoints, that adjust to new base-level conditions.

These results backed by published GPS and seismotectonic data allow us to refine and extend existing geodynamic models of the triple junction. Relict landscapes are delimited by faults and thus result from a tectonic control. The topography of the Sierra Madre de Chiapas evolved as the result of (1) the inland migration of deformation related to the coupling between the Chiapas Massif and the Cocos fore-arc sliver, and (2) the compression along the northern tip of the Central America Volcanic Arc. Although most of the shortening between the Cocos fore-arc sliver and the North American plate is accommodated within the Sierra de Chiapas and Sierra de los Cuchumatanes, a small part may be still transmitted to the Maya Mountains and the Belize margin through a “rigid” Petén basin.

1 Introduction

The aim of this work is to examine geomorphic features along two key areas (the Sierra Madre de Chiapas and the Maya Mountains, Fig. 1) in order to test existing models of the North American-
25 Caribbean-Cocos plate boundary (e.g., Malfait and Dinkelman, 1972; Burkart, 1983; Guzmán-Speziale et al., 1989; Lyon–Caen et al., 2006; Ratschbacher et al., 2009; Authemayou et al., 2011). The oceanic Cocos plate is subducted beneath the North American and Caribbean plates along the Middle America Trench while the North American–Caribbean plate boundary is a sinistral transform system which accommodates the eastward escape of the Caribbean plate (e.g., Lyon–Caen et al.,
30 2006; Manea and Manea, 2006; Andreani et al., 2008a; Authemayou et al., 2011).

The complexity of the plate boundary comes from the fact that both the Caribbean and North American plates are limited to the west by a forearc sliver (Fig. 1) which is coupled to the Cocos slab (e.g., Turner et al., 2007; Phipps-Morgan et al., 2008). This sliver rotates counterclockwise as it is pulled by the eastward escape of the Caribbean plate (e.g., Andreani et al., 2008a; Authemayou et al., 2011). Over the past decades, numerous models regarding the structure and the evolution of the
35 triple junction have been proposed (Authemayou et al., 2011, and references therein). Recent models agree on the fact that the dextral Jalpatagua fault (Fig. 2) coupled with extension along the grabens of Guatemala represents the limit between the forearc sliver and the Caribbean plate (e.g., Lyon–Caen et al., 2006; Andreani et al., 2008a; Authemayou et al., 2011; Franco et al., 2012). However, there is
40 no clear consensus on the boundary between the forearc sliver and the North American plate. This is due to the fact that this boundary is highly diffuse and most of the active deformation is distributed over the Sierra Madre de Chiapas orogenic belt. On top of that, none of the revisited tectonic models of the North American–Caribbean–Cocos plate boundary attempted to include the Maya Mountains (Fig. 1 and 2). In spite of evidence for recent (Late Neogene or Pliocene) tectonics (e.g., Weidie,
45 1985; Lara, 1993; Purdy et al., 2003; Bauer-Gottwein et al., 2011), this region was seldom studied due to a dense vegetation cover and scarce roads.

Using geomorphic indices, we attempt to understand the link between tectonics and landscape evolution within these two regions. We discuss the potential effects of varying uplift, climatic regimes and lithology on geomorphic markers. In active orogens, landscapes result from a competition be-
50 tween vertical uplift, which modifies the base-level of rivers, and erosional processes, which result in the progressive rejuvenation of topographic features through time (e.g., Mather, 2000; Snyder et al., 2000). Geomorphic indices are commonly used to detect the response of landscapes to recent deformation processes (Burbank and Anderson, 2001; Keller and Pinter, 1996, and references therein) and several algorithms and toolboxes allow their extraction from digital elevation models
55 (e.g., Schwanghart and Kuhn, 2010; Shahzad and Gloaguen, 2011a). We use topographic profiles and morphometric maps to classify landscapes. The characterization of elevated surfaces allow us to discriminate uplifted relict landscapes and the propagating front of river incision. We also analyze river longitudinal profiles in order to estimate base-level changes which affect the drainage network.

Finally, we combine the results from the geomorphic analyses with available geophysical data (GPS and seismicity, Fig. 3) in order to propose a model for the North American–Caribbean–Cocos triple junction.

2 Tectonic setting

2.1 Models of the triple junction

Early geodynamic models of the junction between the North American, Caribbean and Cocos plates consisted in a simple transform-trench boundary. For instance, Burkart (1983) proposed a model in which the Polochic fault (and perhaps also the Motagua fault) connected to the Middle America trench. He proposed that the 300 km Neogene motion between the North American and Caribbean plates was accommodated by left-lateral slip along the Polochic and Motagua faults and by extension within the depressions of Honduras and Guatemala. The slip between these two plates was compensated by a left-lateral offset of the trench.

Later works challenged the idea of a junction between the Polochic-Motagua fault system and the trench because the Polochic fault trace terminates within the Chiapas Massif and the Motagua fault trace is lost within the Central America volcanic arc (e.g., Guzmán-Speziale et al., 1989; Guzmán-Speziale and Meneses-Rocha, 2000). As a result, more recent models (Fig. 4) argue that the motion between the North American and Caribbean plates does not result in an offset of the trench but is rather absorbed onshore within a complex zone of deformation. However, there is no clear consensus on the geometry of the plate boundary and on how the deformation is distributed inland.

Using seismotectonic data (Fig. 3a), Guzmán-Speziale and Meneses-Rocha (2000) proposed a model in which the Polochic-Motagua fault system is unable to propagate across the Chiapas Massif (Fig. 4a). As a result, a part of the motion between the North American and Caribbean plates is absorbed by the reverse and strike-slip faults of Chiapas (Fig. 4a). In this model, the reverse faults act as a restraining bend between the strike-slip faults and the Polochic-Motagua fault system. Later on, Guzmán-Speziale (2001) suggested that part of the plate motion is also absorbed by the grabens of Central America, south of the Motagua fault (Fig. 4a).

A second category of models (Fig. 4b) mainly focused on the extensional province located south of the Motagua fault. Gordon and Muehlberger (1994) proposed a model in which the Chortis block is stretched and rotating counterclockwise along the dextral Jalpatagua fault, which affects the volcanic arc (Fig. 2), and the Guayape fault in Honduras. However, more recent works suggest that dextral slip along the Central America volcanic arc is in fact related to a slip partitioning along the trench (e.g., DeMets, 2001; Turner et al., 2007; Correa-Mora et al., 2009). Using GPS data (Fig. 3b), Lyon-Caen et al. (2006) proposed a model of the plate boundary which took into account both the extensional province and the forearc sliver. They suggested that the eastward escape of the Caribbean plate was accommodated by dextral slip along the Jalpatagua fault, sinistral slip along the Polochic-

Motagua fault system and by extension within the grabens of Central America (Fig. 4b). This model
95 was refined by Franco et al. (2012) which proposed a difference in coupling along the subduction
interface (Fig. 4b) in order to explain the GPS velocity field in southern Mexico (Fig. 3b). However,
these two models did not address the tectonic evolution of the Sierra Madre de Chiapas.

The latest models (Fig. 4c and 4d) attempted to explain both the eastward escape of the Caribbean
plate as described by Lyon–Caen et al. (2006) and tectonics of the Sierra Madre de Chiapas. Andreani
100 et al. (2008b) suggested a connection between the strike-slip faults of Chiapas, the Veracruz shear
zone which affects the Veracruz basin further north, and the transtension of the Mexican volcanic
arc (Fig. 4c). They proposed that these three zones represent the boundary of the so-called ‘Southern
Mexico block’ Andreani et al. (2008a). In their model (Fig. 4c), the dynamic of the plate boundary
is driven by the escape of the Caribbean plate and by the counterclockwise rotation of the Southern
105 Mexico block. However, this model does not solve the interactions between the Southern Mexico
block and the Central America forearc sliver. An alternative model was proposed by Authemayou
et al. (2011), in which the Central America forearc sliver extends offshore southern Mexico (Fig. 4d).
In their model, the eastward escape of the Caribbean plate induces a counterclockwise rotation of the
forearc sliver and a ‘zipper’ process (i.e., progressive suturing) between the Jalpatagua and Motagua
110 fault as the space between the two faults is left ‘empty’ by the moving Caribbean plate. According to
their model, the Tonalá Shear Zone is a suture resulting from this ‘zipper’ process. However, both the
Motagua and Jalpatagua fault traces are lost west of 91°W. The western termination of both faults
seems to be associated with extension along the Guatemala City graben (Lyon–Caen et al., 2006).
Furthermore, the Tonalá Shear Zone clearly connects to the Polochic Fault (Fig. 2, Guzmán-Speziale
115 et al., 1989; Guzmán-Speziale and Meneses-Rocha, 2000).

In summary, current views on the plate boundary agree on the fact that the North American and
Caribbean plates are limited by a forearc sliver to the west. The counterclockwise rotation of the
forearc sliver results in transpression north of the Polochic-Motagua fault system while the east-
ward escape of the Caribbean plate is accommodated by the strike slip motions of the Motagua and
120 Jalpatagua faults and by extension along the grabens of Ipala and Guatemala city. However, the
connections between the western corner of the Caribbean plate and the transpressional tectonics of
the Sierra Madre de Chiapas are still unclear. In addition, none of these recent models attempted to
integrate the recent tectonics of the Maya Mountains and Yucatán peninsula into the dynamics of the
plate boundary.

125 **2.2 North-America–Caribbean plate boundary**

The present-day North American–Caribbean plate boundary consists of three main sinistral faults
which delimitates two crustal-scale slivers: the Motagua (Malfait and Dinkelman, 1972; Plafker,
1976), Polochic (Burkart, 1978; Deaton and Burkart, 1984; Sánchez–Barreda, 1981) and Ixcán
(Guzmán-Speziale, 2010) faults (Fig. 2). These faults are seismically active (Fig. 3a; White, 1984;

130 Singh et al., 1984; Ambraseys and Adams, 1996; Guzmán-Speziale, 2010) and Quaternary displace-
ments are documented (e.g., Burkart, 1978, 1983; Authemayou et al., 2012). Brocard et al. (2011)
described a relict Middle Miocene planation surface which formed at low elevation and covered most
of the Polochic-Motagua sliver (the so-called ‘Mayan paleosurface’). This surface was subsequently
uplifted and deformed. Using this surface as a marker, Authemayou et al. (2011) estimated the crustal
135 shortening between 17 and 35 km along the western segment of the Polochic fault (Fig. 2), where the
13 to 7 Ma old Mayan paleo-surface rests on top of the Sierra de los Cuchumatanes at a maximum
elevation of 3800 m.

A part of the motion between the North American and Caribbean plates is also accommodated
along at least thirteen N-trending grabens (e.g., Mann and Burke, 1984; Guzmán-Speziale, 2001)
140 located south of the Motagua fault. The main structures form the Guatemala City, Ipala, Esquipulas,
Sula and Comayagua depressions (Fig. 2). Rogers et al. (2002) related these grabens to the uplift
of the Central America plateau following a slab break-off underneath Central America. The onset
of extension has been determined at 11-8 Ma (Gordon and Muehlberger, 1994; Ratschbacher et al.,
2009). However, most of the present-day extension (11-12 mm.yr⁻¹) seems to be accommodated
145 along the westernmost grabens (Lyon-Caen et al., 2006; Rodriguez et al., 2009; Franco et al., 2012).

2.3 Sierra Madre de Chiapas

The Sierra Madre de Chiapas is constituted by four main structural domains: the Chiapas Massif,
the Central Valley, the Sierra de Chiapas and the frontal fold-and-thrust belt (Guzmán-Speziale and
Meneses-Rocha, 2000; Andreani et al., 2008a; Witt et al., 2012b). The Chiapas Massif mostly con-
150 sists of Permian igneous rocks affected by medium to high grade metamorphism and Middle to late
Miocene intrusions related to the subduction of the Cocos plate (Damon and Montesinos, 1978; We-
ber et al., 2007; Molina-Garza et al., 2015). The western flank of the Chiapas Massif is bounded
by the left-lateral Tonalá Shear Zone which connects to the south to the western termination of the
Polochic fault (Wawrzyniec et al., 2005; Weber et al., 2005). The displacement along the Tonalá
155 Shear Zone is synchronous with Miocene magmatic intrusions and could have reached 100 km
(Molina-Garza et al., 2015). The Central Valley is a ~170 km long and ~30 km wide depression
corresponding to a NW-trending synclinalorium with superimposed smaller folds (Witt et al., 2012b).
The Sierra de Chiapas has a roughly sigmoidal shape and is bounded to the west by the promi-
nent NW-trending left-lateral Tuxtla fault. It is constituted by blocks culminating at heights between
160 2000 m and 2400 m and delimited by E-trending left-lateral faults defined as the High Sierra Fault
System by Witt et al. (2012b). These faults probably connect to the Malpaso fault to the west while
strike-slip motion is absorbed at their eastern terminations by thrusting and folding. The Chiapas
frontal fold-and-thrust belt is located between the Sierra de Chiapas and the western border of the
Yucatán platform. It is constituted by closely spaced faulted folds which are rooted in shallow levels
165 (Witt et al., 2012b).

Thermochronological, tectonic and stratigraphic evidence suggest that renewed exhumation and topographic growth occurred along the Chiapas region during the middle Miocene (16-10 Ma) and late Miocene-Pliocene (6-5 Ma), following a phase of rapid exhumation to upper crustal levels at ~ 30 Ma (Ratschbacher et al., 2009; Witt et al., 2012a). The Tonalá shear zone may have accommodated significant deformation since 16 Ma while the displacement along the transpressive Tuxtla and Malpas faults occurred during the last 6-5 Ma and could have reached 50 to 70 km, involving 0.5 to 0.8 cm.yr^{-1} of left-lateral motion (Meneses-Rocha, 2001; Witt et al., 2012b).

2.4 Maya Mountains and Belize margin

The Maya mountains and the Belize margin are located north to the Polochic-Motagua fault system and east to the Yucatán platform. Due to a dense vegetation cover and scarce roads, this region was seldom studied in comparison with other areas related to the North American–Caribbean plate boundary.

The eastern border of the Yucatán peninsula is crosscut by a series of NNE-trending normal faults (e.g., Weidie, 1982, 1985), which are referred as East Yucatán fault zone (EYFZ) in Fig. 2. The fault zone is ~ 80 km wide and extends over 500 km between the NE tip of the Yucatán peninsula and the Maya Mountains. The surface expression of this fault zone is seen in the alignment of topographic scarps, hydrological features (cenotes, lakes and rivers) and coastal bays (Weidie, 1985; Lesser and Weidie, 1988; Bauer-Gottwein et al., 2011). The East Yucatán fault zone represents the onshore continuation of an extensive horst and graben system which affects the western margin of the Yucatán Basin (Weidie, 1985; Rosencrantz, 1990).

According to Rao and Ramanathan (1988) and Purdy et al. (2003), the Maya Mountains correspond to a roughly NE-trending structural high where the Paleozoic basement is uplifted. Paleozoic rocks constitute the highest elevations of the Maya Mountains. The orogen mainly consists in metamorphosed Late Carboniferous to Middle Permian volcano-sedimentary rocks overlying Late Silurian granites and are bounded by the Northern and Southern Boundary faults (Kesler et al., 1974; Bateson and Hall, 1977; Steiner and Walker, 1996). The Maya Mountains are delimited by fault-bounded E- to NE-trending depressions: the Corozal Basin located to the north and the Belize Basin that borders the Maya block offshore to the east and onshore to the south. According to Purdy et al. (2003), unroofing of the Cretaceous carbonate cap of the Maya Mountains siliciclastic sediment source did not occur until late Neogene, perhaps no earlier than late Pliocene. Faulting along the Belize margin is not accurately documented. The most recent tectonic event is Pliocene or younger and the resulting structures affected Quaternary carbonate deposition. However, Lara (1993) related this event to transtensional faulting while Purdy et al. (2003) rather interpreted this event as the result of a transpression.

200 3 Geomorphic analyses: methods and tools

3.1 Swath topographic profiles

Swath topographic profiles condense elevation data of a complex region into a single profile (e.g., Isacks, 1992; Masek et al., 1994; Telbisz et al., 2013; Hergarten et al., 2014). Topography is extracted from a rectangular swath using a series of parallel profiles, rather than using a single line as in
205 conventional topographic profiles. Elevation data are then projected onto a vertical plane parallel to the longitudinal axis of the swath rectangle, and statistical parameters (the maximum, minimum and mean elevations) are calculated. The curve for maximum elevation corresponds to the ridgelines and helps to identify topographic features, such as relicts of paleo-surfaces. The curve for minimum elevation corresponds to the valley floors or river beds. A quick estimate of the incision is given
210 by the arithmetic difference between the maximum and minimum elevations in a given longitudinal distance (window).

Swath topographic profiles were extracted from 3 arc-seconds SRTM data from CIAT (Jarvis et al., 2008) using a MATLAB script. The swath width was fixed to 20 km. This width is large enough to contain both elevated surfaces and major rivers and small enough to avoid topographic
215 features that are too oblique with respect to the swath axis. Elevation data were sampled using ~220 parallel profiles separated by 90 m. Elevations along each individual profile were also sampled using a 1-pixel (90 m) interval.

3.2 Surface analyses

The combined use different morphometric indices proves to be an efficient way to classify landscapes
220 according to their state of dynamic equilibrium (e.g., Andreani et al., 2014; Domínguez–González et al., 2015). Hypsometric integral (HI) efficiently highlights topographic scarps, surface roughness (SR) substantially increases with incision and relief anomaly (RA) and surface index (SI) highlight elevated low relief landscapes.

The hypsometric integral shows the distribution of landmass volume remaining beneath or above
225 a basal reference plane (Strahler, 1952; Schumm, 1956). This index proved to be efficient in evaluating the response of landscapes to active tectonics (e.g., Pérez-Peña et al., 2009; Mahmood and Gloaguen, 2012; Siddiqui and Soldati, 2014; Andreani et al., 2014; Domínguez–González et al., 2015). According to Pike and Wilson (1971), the hypsometric integral (HI) for a given area can be approximated with Eq. 1:

$$230 \quad HI = \frac{h_{mean} - h_{min}}{h_{max} - h_{min}} \quad (1)$$

with h_{mean} , h_{min} and h_{max} being the mean, minimum and maximum elevations of the analyzed area.

Surface roughness can be described by several parameters (Smith, 2014, and references therein). In this work we used the area ratio approach which evaluate the similarities between a topographic surface with a given area and a flat surface with the same geographic extend (e.g., Hobson, 1972; Grohmann, 2004; Grohmann et al., 2009; Shahzad and Gloaguen, 2011b). The ratio is close to 1 for flat areas and increases rapidly as the real surface becomes irregular. The method used to obtain the ‘real’ and flat surfaces is adapted from the GRASS-R algorithm of Grohmann (2004). First, a slope map is produced using the neighborhood algorithm included in ArcGIS (Burrough and Mcdonell, 1998). The ‘real’ surface S_R is then obtained for each pixel using Eq. 2:

$$S_R = res \times \sqrt{res^2 + (\tan(\alpha) \times res)^2} \quad (2)$$

where res is the DEM resolution in meters and α is the pixel slope in degrees. The flat area S_F is defined for each pixel by $S_F = res \times res$. The surface roughness is then obtained by summing the pixel values of S_R and S_F within a moving window and by dividing the sum of S_R pixels by the sum of S_F pixels.

We computed the hypsometric integral and surface roughness from 90 m resolution SRTM data (CIAT, Jarvis et al., 2008) using TecDEM, a MATLAB-based toolbox (Shahzad and Gloaguen, 2011b). Each pixel of the output raster represents the hypsometric integral and surface roughness values for a 15×15 km moving window. The choice of the window size depends on the objectives. The size of the window will affect the amount of smoothing on the relief (low-pass filtering). To get rid of high-frequency signal such as erosion patterns, the window must encompass a sufficient portion of the analyzed landscape and thus be larger than the wavelength of the undesired signal. Otherwise, computed values will reflect local-scale variations in relief such as hillslopes. **Smallest window scales, aside from being noisy due to high-frequency signals, can also be more influenced by errors (e.g., Albani et al., 2004; Tarolli et al., 2012; Sofia et al., 2013, 2014).** In the present case, a 15 km window allows to filter out the effect of the drainage system as the window size represents an area that will include several valleys and ridges. (Andreani et al., 2014).

Finally, we used the relief anomaly (Scotti et al., 2014) and the surface index (Andreani et al., 2014) to highlight elevated and low relief landscapes. The relief anomaly (RA) represents the elevations normalized by the local relief. We computed this index for a given area using Eq. 3:

$$RA = \frac{h_{mean}}{h_{max} - h_{min}} \quad (3)$$

with h_{mean} , h_{min} and h_{max} being the mean, minimum and maximum elevations of the analyzed area. Highest values are obtained for flat and elevated surfaces.

The surface index (SI) combines the elevations from the DEM with the computed maps of hypsometric integral and surface roughness. It allows to discriminate areas with low local relief landscapes from areas with a more rugged topography. To compute this index, rasters of elevations, hypsometric integral and surface roughness are normalized by their respective minimum and maximum values in

order to obtain pixels values between 0 and 1. We then combine the newly created rasters using Eq. 4:

$$270 \quad SI = (N_h \times N_{HI}) - N_{SR} \tag{4}$$

with N_h , N_{HI} and N_{SR} being the normalized elevations, hypsometric integral and surface roughness values, respectively. Positive surface index values correspond to elevated surfaces with low local relief while negative values highlight rugged landscapes.

3.3 Modeling of the Drainage Network

275 We extracted the drainage network from 30 m resolution SRTM data. The extraction was done using TecDEM (Shahzad and Gloaguen, 2011a) by calculating flow directions and contributing area for each pixels using the D8 algorithm (O’Callaghan and Mark, 1984; Fairfield and Leymarie, 1991; Jones, 2002). Streams were identified using a minimum contributing area of 1 km². This threshold is introduced in order to take into account the transition from colluvial to stream-flow (fluvial) dominated channel, which is usually observed at ~ 1 km² in temperate humid regions (e.g., Montgomery and Foufoula-Georgiou, 1993; Wobus et al., 2006). It is assumed that a theoretical flow accumulation that drains an area larger than 1 km² can be associated to a real water channel. Extracted streams were then organized hierarchically using Strahler (1957) order.

A DEM-based procedure allows to easily extract and analyze a regional-scale drainage network. 285 However, there are several uncertainties related to the DEM and methods. Extracted drainage network and contributing areas are affected by the quality and sampling of the DEM. Free and commonly used data include the 1 arc-second (ca. 30 m) and the 3 arc-second (ca. 90 m) resolution SRTM. For both datasets the absolute vertical error is reported to be less than 20 m. In some areas, the available 1 arc-second SRTM data presented voids (i.e. pixels with no data) mainly located in 290 ridges. We filled these voids with 3 arc-second SRTM data.

The flow-routing method we use (D8 algorithm) may introduce bias in flow path orientation as it discompose flow directions into units of 45° (Fairfield and Leymarie, 1991). This is especially the case in flat areas. Additional known bias are related to methodological aspects. In our study area the commonly encountered problem concerned nested depressions which are related to DEM 295 imperfections. These pits need to be filled to create flow directions. Other artifacts are mainly found along entrenched rivers and canyons, where bad pixels values ‘block’ the path of the extracted rivers. In two cases (Sumidero and La Venta canyons) the DEM filling resulted is a significant deviation of the modeled rivers with respect to the actual rivers. We thus manually carved the bad pixels blocking these two canyons in order to obtain a more accurate flow path. The original topography is also 300 affected by human artifacts such as dammed rivers. This is especially the case for the Grijalva river in southern Mexico. Artificial flats related to DEM filling or dammed rivers introduces errors in

extracted river paths. These errors are easily detected in river longitudinal profiles and were taken into account when interpreting the knickzones in extracted rivers.

3.4 Analysis of River Longitudinal Profiles

305 Deviations from the typical concave-up shape of stream longitudinal profiles, such as knickpoints or convex segments, indicate a disequilibrium state resulting from tectonic, base-level or lithological perturbations (Kirby and Whipple, 2001; Chen et al., 2003; Troiani and Della Seta, 2008; Pedrera et al., 2009; Font et al., 2010). The normalized steepness index (k_{sn}) is widely used to investigate tectonically-induced perturbations in river longitudinal profiles and has been used to propose patterns of uplift (Kirby and Whipple, 2001; Wobus et al., 2006; Whittaker et al., 2008). The relationships between slope and catchment area which define the equilibrium state channel gradient are given by Eq. 5 and 6 (Flint, 1974; Kirby and Whipple, 2001; Wobus et al., 2006):

$$S = k_s \times A^{-\theta} \quad (5)$$

with

315 $k_s = \left(\frac{U}{K} \right)^{\frac{1}{n}} \quad (6)$

where S is the local channel slope, θ is the channel concavity, k_s is the steepness index, A is the upstream drainage area, U is the rock uplift rate and K is the dimensional coefficient of erosion. As suggested by Wobus et al. (2006, and the references therein) a normalized steepness index k_{sn} is used, since k_s and θ are strongly correlated.

320 In some cases, the upstream portions located above prominent knickpoints are associated with an upper-relict landscape. Landscapes tend towards an equilibrium in which rivers are graded to sea level or local base-level. Tectonic or climatically-induced base-level falls modify the equilibrium of the drainage. The result is an erosion wave propagating upstream and the areas not yet affected by the erosion wave form an upper-relict landscape (e.g., Clark et al., 2005; Reinhardt et al., 2007; Pérez-Peña et al., 2015). The reconstruction of the original stream profile downstream of the convexity provides an estimate for the amount of base-level change and subsequent incision (Schoenbohm et al., 2004; Clark et al., 2005; Gallen et al., 2013). To reconstruct the original stream profile, we used the power law between slope and distance defined by Eq. 7 (Hack, 1957):

$$S = i \times D^j \quad (7)$$

330 S is the local channel slope and D is the distance from the drainage divide. Parameters i and j are obtained by regressing the upper segment of the stream profile in a logarithmic plot of slope against distance.

We analyzed river longitudinal profiles using TecDEM (Shahzad and Gloaguen, 2011a). Normalized steepness indices were computed from Eq. 5 using regressions of slope versus catchment area

335 in logarithmic plots with a given reference concavity $\theta_{ref} = 0.45$ (Kirby and Whipple, 2001; Wobus
et al., 2006). Prominent knickpoints or convex anomalies can be observed directly on river longi-
tudinal profiles. However, logarithmic plots of slope against catchment area allow a more detailed
analysis, as minor anomalies in the gradient of rivers can be easily detected. For each longitudinal
profile, we selected and regressed several segments delimited by changes in the gradient of the river.
340 We then plotted regressed segments and their assigned k_{sn} values on a map. The reconstruction of
stream profiles was done by regressing the segments located upstream of knickzones in logarith-
mic plots of the slope against the distance (Eq. 7). We used a Matlab-based script implemented in
TecDEM by Andreani et al. (2014).

The integer format of DEMs locally produces multiple flats with zero slope, which cannot be
345 handled in logarithmic plots (e.g., Kirby and Whipple, 2001; Wobus et al., 2006). This issue is com-
monly solved by smoothing extracted river profiles (Wobus et al., 2006). However, this may induce
bias in extracted geomorphic indices, especially for segments located close to major knickpoints.
In this study we smoothed extracted river profiles using a 20 pixels (ca. 600 m) moving window.
We used a bootstrapping approach in order to address uncertainties related to both DEM artifacts
350 and methods (Andreani et al., 2014). Linear regressions used to compute both normalized steepness
indices and reconstruction of stream profiles are based on subsamples. Typically, we randomly se-
lected from logarithmic plots 75 % of the points corresponding to the analyzed segment in order to
obtain 500 subsets. We then performed a regression on each subset. We obtain the mean values as
well as a range for both normalized steepness indices and reconstructed profiles.

355 3.5 Isobase Maps from Rivers

The spatial distribution of stream height (isobase map) is a useful proxy for investigating geologic
or tectonic processes (e.g., Dury, 1952; Filosofov, 1960; Golts and Rosenthal, 1993; Grohmann
et al., 2007, 2011). Drainage networks are commonly organized according to Strahler (1957) stream
order. According to Golts and Rosenthal (1993) streams of similar orders are of similar geological
360 age and are related to similar geological events. Hence the interpolation of isobase lines, which
connect streams with a similar order, produce a surface resulting from the same erosional events. As
suggested by Grohmann et al. (2011), isobase maps can be seen as a smoothed version of the original
topographic surface, from which was removed the ‘noise’ of the 1st Strahler order streams erosion.
Sharp topographic changes affecting an isobase surface were used to infer potential structures or
365 uplifted blocks associated to tectonic movements (e.g., Golts and Rosenthal, 1993; Grohmann et al.,
2007; Domínguez-González et al., 2015).

Isobase maps were computed using the drainage network extracted from the 30 m resolution
SRTM data (see section 3.3). Using SAGA GIS software we selected streams with a Strahler or-
der ≥ 2 and we interpolated the elevations from extracted rivers using a natural neighbor method.

4.1 Swath topographic profiles

We analyzed the topography of the Sierra Madre de Chiapas and Maya Mountains along five swath profiles (location in Fig. 5a). For each profile (Fig. 6) we extracted the minimum, mean and maximum elevation along the swath. We estimated the local incision by subtracting the maximum and minimum elevation. We also plotted the annual averaged precipitation derived from the Tropical Rainfall Measuring Mission (TRMM, Fig. 5b). Processed TRMM data for the 1998-2009 period are provided by Bookhagen (2009). Swath profiles 1 to 3 (Fig. 6) are perpendicular to major mountain belts and faults in the Sierra Madre de Chiapas. The fourth one (profile 4 in Fig. 6) intersects the volcanic arc, the Sierra de los Cuchumatanes and the Petén basin. The last profile (Fig. 6) displays main topographic trends across the Maya Mountains and the Yucatán platform.

4.1.1 Sierra Madre de Chiapas

The Chiapas Massif (profiles 1 to 3, Fig. 6) dominates the pacific coastal plain. Its SW boundary is abrupt and corresponds to the linear trace of the Tonalá shear zone. The topography of the Chiapas Massif increases from 1500 m to 3000 m toward the SE. River incision (green curves in Fig. 6) increases significantly southward (from ~ 700 m to ~ 1500 m). The range appears strongly asymmetric in profiles 1 and 2. The drainage divide (indicated by a peak in minimum elevations) is located close to the trace of the Tonalá shear zone. The SW flank facing the Pacific coastal plain is steep while the opposite side of the range is gently dipping towards the NE. The curve for minimum elevations also indicates that rivers flowing towards the inner part of the Sierra Madre de Chiapas have a gentle gradient in comparison with those flowing into the Pacific.

The Central Valley (profiles 2 and 3, Fig. 6) corresponds to a ~ 40 km wide depression. Despite a mean elevation of ~ 600 m the local incision appears to be less than 100 m. The topographic scarp associated to the Concordia fault is barely marked. In contrast, La Venta fault and the Tuxtla fault delimitate a ~ 1500 m topographic high (referred as “La Venta block”, profile 1 in Fig. 6) which acts as a barrier for rivers coming from the Chiapas Massif.

In profile 1, the Tuxtla and Malpas faults represent the boundary between the La Venta block and the Chiapas fold-and-thrust-belt. In profiles 2 and 3 The Tuxtla fault represents the limit between the Central Valley and two topographic highs: the Sierra de Chiapas and the Comitán High. The Sierra de Chiapas forms an impressive bulge which dominates both the Central Valley and the Chiapas fold-and-thrust-belt. Its central area is relatively flat and culminates at ~ 2500 m. The topography of the Comitán High is flat with very low incision by the drainage network and the mean elevation is ~ 1600 m. This flat surface is limited to the east by a topographic high (referred as “Leyva Velázquez block”, profile 3 in Fig. 6) which belongs to the Chiapas fold-and-thrust-belt. While in profile 1 the

elevation of the fold-and-thrust-belt is constant, in profiles 2 and 3 the topography of the belt is
405 asymmetric and elevations gently decrease toward the flat Tabasco coastal plain.

Peaks in precipitation indicate orographic effects along the Pacific side of the Chiapas Massif
(profiles 1 to 3 in Fig. 6) and along the northern part of the Chiapas fold-and-thrust belt (profiles
1 and 2 in Fig. 6). However, the shadowing effect of these ranges appear limited as we observe no
410 significant drop in the amount of precipitation between the inner parts of the Sierra Madre de Chiapas
and surrounding areas (Pacific coast and Tabasco coastal plain).

4.1.2 Volcanic arc and Sierra de los Cuchumatanes

In profile 4 (Fig. 6) the volcanic arc domain appears as an uplifted plateau with elevations reaching
3500 m. The curve for minimum elevations indicate that the main drainage divide is located close to
the NE limit of the volcanic arc. Rivers flowing towards the Pacific deeply incised the volcanic arc as
415 the local incision reaches up to 2500 m. We also observe an orographic effect on precipitation. The
region beyond the volcanic arc consists in a relatively flat depression which may in fact represent
a back-arc plateau (profile 4 in Fig. 6). The shadowing effect of the surrounding ranges does not
result in significantly dryer conditions. Precipitation within the depression range between 1000 and
1500 mm/yr while they range between 1500 and 2000 mm/yr along the Pacific coast and Petén
420 basin. This area shows a moderate incision by the drainage network in comparison with the volcanic
arc and the Sierra de los Cuchumatanes. The curve for minimum elevations suggests a significant
base-level drop between rivers flowing within the inner part of the back-arc plateau and the drainage
associated to the Polochic fault trace.

The Sierra de los Cuchumatanes (profile 4 in Fig. 6) is bounded by two impressive scarps as-
425 sociated to the Polochic and Ixcán faults. The Sierra de los Cuchumatanes is also deeply incised
with local incision values up to 1500 m. The general topographic trend of the range, with higher
elevations along the Polochic fault, suggests a tilt towards the NE. The asymmetry of the drainage is
possibly related to the tilt of the range or to an orographic effect (or to a combination of both factors).
However, the peak in precipitation is located along the Ixcán fault. In spite of a SW gradient (from
430 2500 to 1500 mm/yr), most of the Sierra de los Cuchumatanes receives as much as rain as the Petén
basin (2000 mm/yr). Finally, the flat and regular topography of the Petén basin contrasts with the
elevated and incised topography of the others areas.

4.1.3 Maya Mountains

In topographic profile (profile 5 in Fig. 6) the Maya Mountains are delimited by two major scarps
435 related to the NE-trending Southern Boundary fault and the two NE- and E-trending Northern bound-
ary faults. The topography of the Maya Mountains is asymmetric and the range appears slightly tilted
towards the NW. The main drainage divide as well as highest elevations (up to 1000 m) are located
close to the Southern Boundary fault. As indicated by the curve for local incision (profile 4 in Fig. 6),

the areas located East of the main drainage divide are more eroded (up to 750 m of incision) in comparison with the areas located on the other side of the drainage divide where the main river incised the initial topography by 300 to 400 m. We also observe a drop in the amount of precipitation (from 2500 to 1500 mm/yr) along the southern side of the Maya Mountains.

The topography of the Yucatán platform (profile 4 in Fig. 6) appears to be controlled by the NE-trending Eastern Yucatán Fault Zone. The faults are located within a ~80 km wide corridor and delimitate tilted blocks. Highest elevations and incisions are located close to the fault traces and decrease towards the NW. The inner part of the Yucatán platform (located west of the fault zone) is characterized by a well defined and continuous surface. This surface is poorly incised (local incision is less than 100 m) and appears to be tilted towards the NW.

4.2 Morphometric maps

The hypsometric integral values (Fig. 7a) appear to be essentially controlled by tectonic features. Highest values are mainly found along fault-related topographic scarps while lowest values are located within depressions and flat areas. In Chiapas region areas such as the coastal plain, the Central Valley and the inner part of the Comitán High are associated to low hypsometric integral values ($HI \leq 0.35$). These are flat areas as seen in swath topographic profiles. Both the Tonalá shear zone and the Tuxtla fault scarps are highlighted by higher values ($HI > 0.35$). However hypsometric integral values are significantly greater for the Tuxtla fault scarp ($HI > 0.55$).

The structural control is also obvious in some other regions. The mountain belt north to the Polochic fault displays rather high hypsometric values. Within the extensional province of Central America, structures such as the Ipala, Comayagua and Sula grabens are associated to hypsometric lows. The pattern of hypsometric values also highlights the N- to NE-trending lineaments of the Eastern Yucatán Fault Zone (e.g., Rio Hondo fault) which are located along the Belize margin.

The surface roughness map (Fig. 7b) reveals the location of strongly incised areas as well as flat surfaces. Peak values (> 1.07 in Fig. 7b) are essentially found along the central and southern part of the Tonalá shear zone, the NW tip of the Central America volcanic arc, the mountain belt located north to the Polochic fault (e.g., Sierra de los Cuchumatanes) and Las Minas range (located between the Motagua and Polochic faults). These areas are both elevated (altitudes comprised between 2500 and 4000 m) and strongly eroded. As seen in our swath topographic profiles (Fig. 5), the difference in elevation between ridges and valley floors (local incision) is comprised between 1500 and 2500 m. Intermediate surface roughness values (1.03 to 1.07 in Fig. 7b) are mainly found within the inner part of the Chiapas fold-and-thrust-belt, along the eastern part of the Maya mountains and within the grabens of Central America.

As for hypsometry, the distribution of surface roughness values in the extensional province is structurally controlled. Lowest values highlights graben floors while highest values correspond to horsts. Lowest surface roughness values (< 1.03 in Fig. 7b) emphasize flat (i.e. low amplitude) land-

475 scapes. Some of these areas (e.g., Central Valley of Chiapas, Comitán High) correspond to elevated surfaces and plateaus observed in swath topographic profiles (Fig. 5).

Surface roughness and hypsometric integral do not allow to discriminate low relief landscapes according to their topographic elevations. For this purpose we use two additional geomorphic indices: the relief anomaly (Fig. 8a) and the surface index (Fig. 8b). Both maps give similar results
480 regarding the distribution of elevated and low relief landscapes, which are identified by the highest relief anomaly values (> 16) in Fig. 8a and by positive (0 to 0.5) surface index values in Fig. 8b. Within the Sierra Madre de Chiapas, elevated and low relief landscapes are mainly located within the Central Valley of Chiapas and the areas located east of the Tuxtla fault (Sierra de Chiapas and Comitán High). The drainage system of the Chiapas Massif (consisting of large and flat valleys) is
485 also highlighted. The two maps also show the extend of the elevated plateaus located at the northern tip of the Central America volcanic arc as well as the relatively flat landscapes of the Petén basin and Yucatán platform. In other areas, the distribution of the low relief landscapes appears more fragmented. This is the case for the areas located between the Ixcán and Motagua faults, where our maps highlight the remnants of the Mayan paleosurface described by Brocard et al. (2011).

490 We propose to compare the distribution of geomorphic indices (Fig. 7 and 8), precipitation data (Fig. 5a) and geological contours (Fig. 1b) within the Sierra Madre de Chiapas. Our aim is to detect possible effects of climate or lithology on the location of incised areas and low relief surfaces. At first glance, peaks in precipitation (Fig. 5a) seem to coincide with more rugged topography (along the Pacific coast, the northern Sierra de Chiapas and the eastern part of the Sierra de los Cuchumatanes). However, peaks in precipitation are located at the limits of the orogenic belt while the
495 rugged topography extends inward the Chiapas orogen. On top of that, some areas display a rugged topography in spite of relatively low precipitation (e.g., west of the Sierra de los Cuchumatanes) and vice-versa (e.g., eastern part of the Chiapas fold-and-thrust belt). To compare the distribution of precipitation and geomorphic indices, we randomly selected 20000 points within the Sierra Madre de
500 Chiapas. For each point we extracted the corresponding pixel values from surface roughness, surface index, hypsometric integral, local relief and precipitation maps. We then compared the distribution of the values using scatterplots (Fig. 9). The co-linearity of the scatterplots was estimated using the Pearson correlation coefficient (PCC). PCC values vary between +1 and -1 (1 is total positive correlation, 0 is no correlation, and -1 is total negative correlation). PCC values for plots of precipitation
505 against several geomorphic indices (Fig. 9) suggest no significant correlation between the amount of precipitation and the ruggedness of the landscape.

We also compared the distribution of geomorphic indices for different geological units. We made three sets of points located within the Sierra Madre de Chiapas. Each set of points corresponds to one of the main geological units of the Sierra Madre de Chiapas: the Paleogene terrigenous deposits,
510 the Cretaceous limestones and the Permian granodiorite of the Chiapas Massif (Fig. 2). For each point we extracted the corresponding surface roughness, surface index, hypsometric integral and

local relief value. Histograms in Fig. 10 reveals slight changes in the distribution of the geomorphic indices which may be related to an effect of lithology. Cretaceous carbonates show lower surface roughness values when compared to Paleogene deposits or Permian rocks. Most of the Permian granodiorite display negative surface index values. Cretaceous carbonates show slightly lower Local relief and higher hypsometric integral values.

4.3 River longitudinal profiles

Topographic uplift, subsidence or climatically-induced sea-level changes, modify the base-level of rivers and result in the progressive erosion or rejuvenation of pre-existing topographic features through time. Thus, the analysis of deviations from the typical concave-up shape of stream longitudinal profiles (i.e., knickpoints or convex segments) is a suitable approach to explore these phenomena.

4.3.1 Tonalá Shear Zone and volcanic arc

We analyzed 36 rivers draining the western flank of the Chiapas Massif and the Central America volcanic arc (Fig. 11). We focused our analysis on the areas located between the main drainage divide and the alluvial Pacific coastal plain. Longitudinal profiles were extracted from the modeled drainage network (Fig. 11 and 12).

Figure 12 displays characteristic stream profiles. Longitudinal profiles 6 and 15 (Fig. 12) are representative of main trunk rivers located upstream of the Tonalá Shear Zone (streams and catchments 1 to 17 in Fig. 11). The shear zone marks the boundary between the Permian batholith of the Chiapas Massif to the northeast and the Pacific coastal plain to the southwest. Two third of the analyzed rivers display a rather smooth and graded longitudinal profile with a single concave segment (such as profile 6 in Fig. 12). The remaining rivers display a gentle knickzone which separates two main segments (such as profile 15 in Fig. 12). Profiles 23 to 36 (Fig. 11) drain the Central America volcanic arc domain. Half of the analyzed rivers display a prominent knickzone (such as profile 24 in Fig. 12) which separates a gently concave upper reach from a steeper lower reach. Profiles 18 to 22 (Fig. 11) are located within a transition zone between the two previous regions, where the NW-trending Tonalá Shear Zone connects to the E-trending Polochic fault system along a WNW-tending fault zone. The longitudinal profiles of streams 21 and 22 display two main knickzones (see profile 21 in Fig. 12). The upper knickzone extends a few kilometers upstream of the fault which makes the junction between the Tonalá Shear Zone and the Polochic fault system. However, the lower knickzone appears more prominent and thus seems to be associated to a larger base-level change.

We extracted two parameters from analyzed longitudinal profiles in order to illustrate the along-strike changes in the geometry of rivers. The first one is the steepness index (k_{sn}) extracted from logarithmic plots of slope against area (see examples in Fig. 12). Graded profiles show a single k_{sn} value while non-equilibrated profiles are characterized by changing k_{sn} values associated to several segments. In this latest case we considered the k_{sn} value of the lowermost segment as it is the one

along which rivers adjust to new base-level conditions (e.g., Kirby and Whipple, 2001; Wobus et al., 2006; Whittaker et al., 2008). The second parameter is the estimated base-level change. If the profile displayed a well defined upper reach, we projected the segment located above the knickzone using a logarithmic plot of slope against distance (see equation 7). We then estimated the base-level change according to the difference in elevations between the projected and the actual profiles at the outlet (see examples in Fig. 12).

Both k_{sn} values and estimated base-level changes were plotted along a profile passing through the outlets of the analyzed catchments (Fig. 11). The results suggest a substantial alongstrike increase in k_{sn} values, from ~ 100 along the northern tip of the Chiapas Massif to >200 along the volcanic arc. Estimated base-level changes seem to be low for the Chiapas Massif (<200 m) but increase significantly along the southern tip of the Tonalá Shear Zone (where it connects to the E-trending Polochic fault) and are higher within the volcanic arc (1000 to 2000 m).

4.3.2 Grijalva river

The Grijalva river (profile 1 in Fig. 13) flows from the northern tip of the volcanic arc towards the Tabasco coastal plain. Two main knickzones are located upstream of the Necta fault. They mark the transition between the elevated volcanic arc plateau, the back-arc domain and the local base-level of the central depression of Chiapas. The reconstructed profile made using the segment located above the lowest knickzone suggests a ~ 500 m base-level fall between the back-arc domain and the central depression of Chiapas. Segments located within the central depression of Chiapas (between the Necta and Tuxtla faults) and the Tabasco coastal plain display a lower gradient in comparison with those located upstream of the Necta fault. Despite numerous artifacts related to hydroelectric dams (e.g., La Angostura, Chicoasén and Malpaso dams) and canyons (e.g., Sumidero), we can infer a major knickzone located between the Tuxtla fault and the Chicoasén dam. We attempted to infer the geometry of the Grijalva river profile using the segments upstream and downstream of the flooded areas. The segments of the river upstream and downstream of La Angostura dammed lake could be fit by the same regression in a logarithmic plot of slope vs area (see profile 1 in Fig. 13). We are confident that no knickpoints are hidden below this dammed lake otherwise it would have been witnessed by a change in gradient of the river). In contrast, the gradient of the river upstream and downstream of the Chicoasén dammed lake differ and can not be fit by a single regression. The gradient of the Grijalva river is 200 m between the Tuxtla fault and the Chicoasén dam (~ 30 km apart) while it is less than 50 m between La Angostura dam and the Tuxtla fault (~ 60 km apart). We thus suspect that the Chicoasén dammed lake hides a major flooded knickzone which marks a ~ 350 m base-level fall between the central depression of Chiapas and the Tabasco coastal plain. For the Malpaso and Peñitas dam the gradient of the river became too low and the scatter in slope-area data too high to make such tests.

The Seleguá river (profile 2 in Fig. 13) is a right bank tributary of the Grijalva river which flows from the southern tip of the Sierra de los Cuchumatanes into the central depression of Chiapas. A prominent knickzone is observed upstream of the Polochic fault. This knickzone separates an upper segment with a gentle gradient ($k_{sn} < 50$) from a lower and much steeper ($k_{sn} > 300$) segment. The upper segment is associated to the relicts of the Mid-Miocene Mayan paleosurface which is preserved on top of the Sierra de los Cuchumatanes (Authemayou et al., 2011). Downstream of the Polochic fault the profile displays an overall convex shape related to numerous knickzones. The lowermost segment is located within the central depression of Chiapas and shows a gentle gradient.

585

590

595

600

605

Longitudinal profiles 3 to 5 (Fig. 13) are from left-bank tributaries located within the Chiapas Massif. Longitudinal profile 3 (southern tip of the Chiapas Massif) display two prominent knickzones. The upper knickzone is located close to the drainage divide and separates an upper segment with a gentle gradient ($k_{sn} < 75$) from a lower and much steeper ($k_{sn} > 150$) segment. The lower knickzone is more gentle and located downstream of the Necta fault trace. Profiles 4 and 5 intersect the lineament made by La Venta and Concordia faults. In both profiles the upper reach (60 to 80 km from the drainage divide) is gently concave (k_{sn} values around 30). The upper segment are limited downstream by a knickzone which correspond both to the boundary between the Chiapas Permian batholith and the Cretaceous/Tertiary sedimentary cover and to La Venta–Concordia lineament. In both profiles the knickzone is associated to canyons as the river incised through the sedimentary cover (mainly Cretaceous limestones). In profile 4 this knickzone marks a minor base-level fall (approximately 50 m) between the upper reach and the central depression of Chiapas. In profile 5 the knickzone is associated to a major base-level fall as the lower reach connects to the portion of the Grijalva river which is in equilibrium with the base-level of the Tabasco coastal plain. The reconstruction of the upper reach suggests approximately 200 m base-level fall. However, the lower part of the tributary is flooded by the Malpaso dammed lake and the real base-level fall could be higher.

4.3.3 Sierra de Chiapas

Figure 14 displays river longitudinal profiles from the northern part of the Sierra de Chiapas. Profiles 1 to 3 are characteristic of the tributaries of the Grijalva river flowing across the Tuxtla fault. Almost all longitudinal profiles display prominent knickzones located upstream of the fault scarp. Upper reaches display either a smooth concave shape (as in profiles 1 and 3, Fig. 14) or a convex shape (as in profile 2). Convex upper reaches are mainly observed in the areas where the fault trace intersect volcano-sedimentary deposits related to Plio-Quaternary volcanoes or in the area where the fault trend change from NW to NNW. The lower reaches are strongly concave and are associated to higher k_{sn} values. Reconstructions of upper reaches (profiles 1 and 3 in Fig. 14) suggest a 500 to 800 m base-level change along the Tuxtla fault scarp.

615

Longitudinal profiles 4 to 6 are located within the northern parts of the Sierra de Chiapas. They are complex, as rivers flow through the E-trending left-lateral faults and then through the NW to NNW-

trending structures of the Chiapas fold-and-thrust belt. As a result, these profiles display several knickzones (e.g., profile 4 in Fig. 14). However, the upper knickzones are more prominent as they mark a sharp transition between the uppermost segments characterized by gentle gradients and low k_{sn} values (<100) and the lower segments which show steeper gradients and higher k_{sn} values (between 150 and 250). Reconstructions of upper reaches suggest a 700 to 1100 m base-level change with respect to the Grijalva river (profiles 5 and 6 in Fig. 14) or the Tabasco coastal plain (profile 4 in Fig. 14).

625 4.3.4 Steepness index map

The analysis of river longitudinal profiles shows that major base-level changes (i.e., higher than 100 m) are associated to prominent knickzones and to a substantial increase in steepness index (k_{sn}) values. We propose thus to map the variations of k_{sn} values within the Sierra Madre de Chiapas in order to assess the locations of main drainage network perturbations. We computed k_{sn} values from extracted river profiles using a 5 km moving window. We then interpolated obtained values using a natural neighbor (NN) method. The result is shown in Figure 15.

The overall distribution of k_{sn} appears to be tectonically controlled, as highest values are found close to major tectonic features. In Chiapas highest k_{sn} values (>150 in Fig. 15) are mainly distributed along the Tuxtla fault, the E-trending strike slip faults of the Sierra de Chiapas and the portion of the fold-and-thrust belt located north of the Sierra de Chiapas. Areas located along the La Venta fault and the Tonalá Shear Zone display intermediate k_{sn} values (90 to 120). Uniformly low k_{sn} values suggests that most of the tributaries of the Grijalva river flowing in the northern part of the Chiapas Massif and the Central Valley of Chiapas share a common base-level.

To the south, the Polochic fault exerts a strong control over the drainage network. The Sierra de los Cuchumatanes and the SE tip of the Chiapas Massif (limited by the Polochic fault to the south and the Ixcán and Necta faults to the north) display especially high values (>180). By contrast, most of the areas located south of the Polochic fault are associated to lower k_{sn} values (<150). Lowest k_{sn} values (<90) are found within the back-arc region and the graben of Guatemala City. The last area displaying high k_{sn} values (>180 in Fig. 15) corresponds to the SW flank of the Central America volcanic arc where rivers are affected by a very steep topographic gradient.

4.3.5 Maya mountains

We analyzed the drainage network of the Maya Mountains using a map of isobases from rivers and several longitudinal profiles (Fig. 16). Longitudinal profiles 1 to 5 (Fig. 16) are part of the Belize river watershed, which drain the northern and western side of the Maya Mountains. All these profiles display prominent knickpoints which mark the transition between the upper reaches which are usually gently concave and the lower segments which are either concave with a steeper gradient or convex. The knickpoints are located upstream of the ENE- to E-trending Northern Boundary fault

for profiles 1 to 3 (Fig. 16) and upstream of a NE-trending fault zone in profiles 4 and 5 (Fig. 16). The reconstructed profiles made using the upper reaches suggest that the drainage network faced a
655 300 to 500 m base level change.

Profiles 6 to 9 (Fig. 16) are located in the southern and eastern parts of the Maya Mountains and flow directly into the Caribbean sea. Only southernmost profiles display prominent knickpoints which delimitate a well defined upper reach (e.g., profile 6 in Fig. 16). These knickpoints are located upstream of the NE-trending Southern Boundary fault and a ENE-trending fault zone. By contrast,
660 most of profiles located along the eastern side of the Maya Mountains display an almost smooth and concave shape such as profile 7 (Fig. 16). In particular, there is no significant change in the gradient of the rivers along the Southern Boundary fault. However, in some cases a knickpoint associated to a relict upper reach is found close to the drainage divide (profiles 8 and 9 in Fig. 16). The reconstructed upper reaches in profiles 5 and 8 (Fig. 16) suggest a 500 m base level change.

665 Isobases from rivers (Fig. 16) allows to summarize the geometry of the drainage network. Within the Maya Mountains, the isobase surface forms a roughly sigmoidal-shaped plateau. Spaced contour lines indicate gentle gradient in rivers flowing in the central part of the range. The top of the isobase surface is more elevated (>800 m) in the southern part of the Maya Mountains (between 16.30°N and 16.60°N) where it forms a flat structural high delimited by two parallel ENE-trending fault
670 zones. North of 16.60°N , the top of the isobase surface rather forms a tilted panel, with elevations decreasing towards the NW (from ca. 700 m to ca. 400 m). The isobase surface of this tilted panel is almost flat but the drainage network of the Macal river (stream 3 in Fig. 16) appears entrenched with elevations ca. 200 m below the average elevation of the isobase surface. This can also be observed in swath profiles (Fig. 6) crossing this drainage basin. The western and northern sides of the tilted
675 northern panel are controlled by major structures, as indicated by sharp isobase gradients associated to a NE-trending fault zone and to the ENE- to E-trending Northern Boundary fault. Areas located along the northern segment of the Southern Boundary Fault appear significantly incised and the top of the isobase surface is located ca. 25 km East of this fault.

5 Discussion

680 5.1 Equilibrium of landscapes

The evolution of landscapes results from complex interactions between climatic and tectonic processes (e.g., Hack, 1960; Willett and Brandon, 2002; Matmon et al., 2003; Pazzaglia, 2003; Ferrater et al., 2015, and the references therein). Recent tectonic or climatically-induced base-level fall are associated to the propagation of an erosion wave, which represents the limit between an upper-relict
685 landscape and a lower adjusting zone (e.g., Reinhardt et al., 2007; Pérez-Peña et al., 2015). As a result, relict portions of landscapes (i.e., where uplift is not yet counterbalanced by fluvial erosion)

may persist though time (e.g., Burbank and Anderson, 2001; Clark et al., 2005; Legrain et al., 2014; Giletycz et al., 2015).

690 Previous works indicate the occurrence of such relict landscapes along the Polochic-Motagua
fault system and within the Chortis block (western corner of the Caribbean plate). Rogers et al.
(2002) interpreted the topography of the northern Central America highlands (east of the modern
volcanic arc and south of the Motagua fault) as an uplifted ignimbritic plateau (Fig. 17). More
recently, Brocard et al. (2011) and Authemayou et al. (2011) described a relict Middle Miocene
695 planation surface (the so-called ‘Mayan paleosurface’) which covered most of the areas between
the Ixcán and Motagua faults (Fig. 17) and was subsequently uplifted and deformed. Our results
suggest that relict landscapes are also preserved within the Sierra Madre de Chiapas and Maya
Mountains. Topographic profiles (Fig. 5 and 6) and both relief anomaly and surface index maps
(Fig. 8) highlight elevated areas which are characterized by a low amplitude relief. These areas are
surrounded by more dissected landforms which are characterized by high surface roughness values
700 (Fig. 7). The surface index map (Fig. 8) shows very well this dual distribution of elevated and low
relief (positive values in Fig. 8) and dissected (negative values in Fig. 8) landforms. This is a clear
indication that landscapes within the Sierra Madre de Chiapas and Maya Mountains are in transient
stage, and morphometric maps allows to discriminate between the areas forming part of an upper-
relict landscape and the propagating front of river incision. Moreover, river longitudinal profiles
705 display an upper reach associated to these relict landscapes while lower segments represent one or
several knickzones along which channels adjust to new base-level conditions.

Our results suggest a limited effect of climate. Within the Sierra Madre de Chiapas we found no
clear correlation between peaks in precipitation and the distribution of rugged topography. There is
no scaling between the amount of precipitation and the ruggedness of the topography or the amount
710 of incision (Fig. 9). We observe a clear orographic effect on precipitation (swath profiles in Fig 6).
This effect is confined to the borders of the orogen and may locally be superimposed to a rugged
topography. However the incision by the drainage network extends more inward where precipitation
are no significantly different from the surrounding coastal plains. The erosion is related to the uplift
of the Sierra Madre de Chiapas rather than to a change in climatic conditions. Indeed, recent studies
715 suggest a dominance of tectonics over climate in the denudation of active mountain ranges (e.g.,
Godard et al., 2014; Fuchs et al., 2015). There is a possible effects of lithology on the location
of incised areas and low relief surfaces. The distribution of morphometric indices suggests slight
changes related to lithology (Fig. 10). Lower surface roughness and local relief values observed in
Cretaceous limestones are probably related to karst processes. The fact that a part of the drainage is
720 subterranean may have contributed to the extensive preservation of La Venta and Sierra de Chiapas
elevated plateaus. By contrast in other areas where bedrock erosion prevail (for instance within the
Paleogene terrigenous deposits of the northern Sierra de Chiapas) elevated relict landscapes have
been deeply incised.

We attempted to map the distribution of elevated relict landscapes from northern Central America (Fig. 17). The surface index and relief anomaly maps (Fig. 8) highlights well areas where elevated and low relief surfaces are extensively preserved and provide thus a good basis for mapping the extend of relict landscapes. In some areas where elevated surfaces are incised by the drainage network and for the Chiapas Massif where surfaces are nested between ridges, we refined our map using slopes and topographic contours derived from the SRTM data.

730 Relict landscapes from the Sierra Madre de Chiapas can be divided in four main domains. The westernmost domain consist in a monadnock-type landscape which developed over the Chiapas Massif (Fig. 17). In morphometric maps, this area is characterized by low hypsometric integral and surface roughness values (Fig. 7). In fact, these geomorphic indices respond to an extensive system of large and relatively flat valleys. The drainage system appears in steady-state as rivers display smooth concave profiles and low k_{sn} values (upper segments of profiles 4 and 5 in Fig. 13). The isobase surface of the rivers is almost flat (see curves for minimum elevations in swath profiles 1 and 2, Fig. 5). Such low gradients are characteristic of lowland alluvial rivers. We interpret the topography of the Chiapas Massif as the result of a sustained erosion which resulted in the development of a low relief surface and a system of alluvial valleys. In other words, the Permian batholith of the Chiapas Massif and its sedimentary cover were incised until rivers reached an almost flat and steady-state base-level. 740 However, this isobase surface is now located ~ 400 m above the present-day sea level. We suspect thus that the topography of the Chiapas Massif was much lower than it is today and that it has been subsequently uplifted.

The large valleys of the Chiapas Massif connects to the east to two morphological domain: the Central Valley of Chiapas and La Venta plateau (Fig. 17). The eastern boundary of these two domains corresponds to the Tuxtla and Malpaso faults. The Central Valley of Chiapas consists in gently folded tertiary sediments. The topography of this area is smooth as suggested by high relief anomaly (Fig. 8) and low surface roughness (Fig. 7) values. However, high hypsometric integral values along the Grijalva river reflect an entrenchment of the drainage network. The canyons are filled by the water of La Angostura dammed lake but their depth probably did not exceeded 150 m. The Mesozoic carbonates of La Venta plateau were uplifted and gently tilted along La Venta and Tuxtla faults. This area is also associated to an entrenchment of the drainage network. The analysis of river longitudinal profiles (Fig. 13) suggests a ~ 350 m base level change between the inner part of the Sierra Madre de Chiapas and the Tabasco coastal plain. We relate the entrenchment of the drainage network within the Central Valley of Chiapas and La Venta plateau to the uplift of the areas located between the Tuxtla fault and the Tonalá shear zone. This uplift resulted in an erosional wave propagating upstream within the drainage of the Grijalva river. 750

The last domain encompasses the elevated plateaus east of the Tuxtla fault (Comitán High and Sierra de Chiapas, Fig. 17). The northwest part of this plateau is heavily incised as shown by high surface roughness values (Fig. 7). By contrast, the areas south of the Tectapan fault appear to be less 760

dissected. In spite of elevations up to 2400 m, surface roughness values (Fig. 7) remains low. Indeed, the top of the plateau is associated to a low relief topography which is disrupted by incisions along the E-trending faults. These differences are probably related to several factors, including lithology (the southern areas consist mainly in Mesozoic carbonates with subterranean drainage limiting surface erosion), climate (the northern areas may receive more precipitation from the Gulf of Mexico), and local base-level changes. The analysis of river longitudinal profiles (Fig. 13) suggests a ~500-700 m base level change along the Tuxtla fault and ~700-1100 m in the northern part of the Sierra de Chiapas.

The age of the relict landscapes in Sierra Madre de Chiapas is unknown. Thermochronological and stratigraphic evidence suggest two main periods of topographic growth. The first one occurred during the middle Miocene (16-10 Ma) and the latest one started during the late Miocene-Pliocene and is still ongoing (Guzmán-Speziale and Meneses-Rocha, 2000; Meneses-Rocha, 2001; Ratschbacher et al., 2009; Witt et al., 2012a). We assume that the relict landscapes of the Sierra Madre de Chiapas have developed following the middle Miocene phase of topographic growth. The development of large alluvial valleys within the Chiapas Massif suggests that the Permian batholith and its sedimentary cover were uplifted and significantly eroded until rivers reached an almost flat base-level. Indeed, Witt et al. (2012a) show that the middle to late Miocene period is associated to marked increase in erosion of the Chiapas Massif and sedimentation within the basins of the Tabasco region. The distribution of morphometric indices (Fig. 7) and the location of knickzones upstream of major faults such as the Tuxtla fault (profiles 1 to 3 in Fig. 14) indicate that this late Miocene erosional topography was subsequently uplifted and fragmented by tectonics. This probably happened during the latest period of deformation (~5 Ma to present). A middle Miocene age for the erosional topography of the Sierra Madre de Chiapas would be consistent with the timing of similar topographic features, such as the so-called "Mayan paleosurface" which covered most of the Polochic-Motagua sliver (Authemayou et al., 2011; Brocard et al., 2011). According to Brocard et al. (2011), this widespread planation surface formed at low elevation during the middle Miocene (between 13 and 7 Ma).

Our results show that most of the Maya Mountains consist in an elevated relict landscape (Fig. 17). Geomorphic maps (Fig. 7 and 8) as well as isobases from rivers (Fig. 16) indicate that low relief surfaces have been extensively preserved in the central and western parts of the Maya Mountains, while the eastern part of the range has been more eroded. Most of the analyzed river longitudinal profiles (Fig. 16) display an upper reach associated to these low relief surfaces. Here again, the age of this relict landscape is unknown. According to Purdy et al. (2003), uplift in the Maya Mountains did not occur until late Neogene and perhaps no earlier than late Pliocene. The low relief surfaces we observe in the Maya Mountains may have formed during the Middle Miocene as the Mayan paleosurface. In the southern part of the range, the low relief surface of the Maya Mountains gently dip towards the west and seems to connect to the flat topography of the Petén basin.

5.2 Morpho-tectonic interpretations

Landscapes from the Sierra de Chiapas and the Maya Mountains were fragmented and locally rejuvenated by tectonics. We propose to combine our topographic profiles (Fig. 5 and 6), geomorphic maps (Fig. 7 and 8) and analyzed river profiles (Fig. 11 to 16) with published GPS and seismotectonic data in order to locate main active deformation zones. We attempted to define coherent tectonic blocks (Fig. 18) in order to produce a comprehensive map of the plate boundary.

The Tonalá Shear Zone has been depicted as a relict left-lateral structure (Wawrzyniec et al., 2005; Weber et al., 2005). Indeed, it shows no signs of seismic activity (Fig. 3b). However, the differential motion between GPS stations ESPO and CONC (Fig. 18) suggests that the fault zone currently accommodates a $\sim 2.5 \text{ mm.yr}^{-1}$ ENE extension. Our geomorphic analyzes suggest that the landscapes of the Chiapas Massif are mostly in steady-state. However, the southern tip of the massif displays evidence of topographic rejuvenation. River profiles crossing the northern segment of the Tonalá Shear Zone (profile 6 in Fig. 12) show no major perturbations and indicate thus an equilibrium between erosion and uplift. Towards the SE, river profiles display gentle knickzone associated to a 50-150 m base-level fall (profile 15 in Fig. 12). In the same area, Authemayou et al. (2011) observed that alluvial fans across the fault zone are affected by vertical offsets ranging between 15 and 60 m. The base-level drop we observe in river profiles is thus related to a gentle uplift along the central portion of the Tonalá Shear Zone. Finally, river profiles along the southern termination of the fault zone (profile 21 in Fig. 12) exhibit prominent knickpoints and convex segments which indicate a major disequilibrium. The k_{sn} values increase southward along the Tonalá Shear Zone but in fact all the areas located between the Polochic fault and the Necta fault display high k_{sn} values (Fig. 15). Under constant climate and lithologies, the steepness index (k_{sn}) correlates with uplift rates (e.g., Kirby and Whipple, 2001; Wobus et al., 2006; Kirby and Whipple, 2012). We suspect thus higher uplift rates along the southern border of the Chiapas Massif (north of the Polochic fault and south of the Necta fault). The Necta fault separates the northern domain, where the topography is almost in steady-state, from the southern domain, where landscapes are being rejuvenated due to an increase in uplift rates. We propose to define a rigid block (referred as 'Chiapas Massif sliver' in Fig. 18b) delimited by the Tonalá shear zone to the west, by the La Venta and Tuxtla faults to the east and by the Necta fault to the south. The Tonalá shear zone seems to accommodate a 2.5 mm.yr^{-1} extension while the Tuxtla fault instead accommodates a 2.5 mm.yr^{-1} shortening (Fig. 18b).

The northern tip of the Central America volcanic arc forms an uplifted plateau with a mean elevation of $\sim 3000 \text{ m}$. The drainage network developed prominent knickzones as it responded to high-amplitude base level changes (Fig. 11 and 12 and k_{sn} map in Fig. 15). The differential motion between GPS stations TPCH and SOL (Fig. 18) indicates a $\sim 5.8 \text{ mm.yr}^{-1}$ NNE shortening between the Pacific coastal plain and the Sierra de los Cuchumatanes. We also noted that the motion of GPS stations located along the volcanic arc (MAZ, CHL and SSIA in Fig. 18) is intermediate between the Cocos forearc sliver and the Caribbean plate. From the motion of these GPS stations

835 we infer a dextral fault zone bounding the southern edge of the volcanic arc. We suspect that the volcanic arc acts as a tectonic sliver which accommodates a significant part of the motion between the Cocos forearc sliver and the overriding plates. East of the Graben of Guatemala City, the sliver is possibly delimited by two right-lateral fault zones which accommodates each half of the motion ($\sim 13 \text{ mm.yr}^{-1}$) between the Cocos forearc sliver and the Caribbean plate. West of the Graben of
840 Guatemala City, the volcanic arc sliver is pinned between the Cocos forearc sliver to the SW, the Polochic sliver to the NE and the Chiapas Massif to the north (Fig. 18b). Though a 6.3 mm.yr^{-1} shortening is expected between GPS stations MAZ and JOY, we could not find a clear limit between this volcanic arc sliver and the Motagua sliver.

Within the Sierra Madre de Chiapas orogenic belt, most of the present day seismicity (Fig. 3b)
845 is located along the strike-slip faults of Chiapas, which delimitate the La Venta block (profile 1 in Fig. 5) and the northern part of the Sierra de Chiapas, and along the inner part of the Chiapas fold-and-thrust belt, east of the Sierra de Chiapas and Comitán High uplifted domain. In addition, two historical events were documented within the Central valley of Chiapas: the 14 March 1591 (García-Acosta and Suárez-Reynoso, 1996; Peraldo and Montero, 1999; Guzmán-Speziale, 2010)
850 and the 23 September 1902 (Böse, 1903; Figueroa, 1973) earthquakes. Historical earthquakes of this area were attributed either to the Concordia (Guzmán-Speziale, 2010) or to the Tuxtla (Andreani et al., 2008a) faults. The differential motion between GPS stations CONC and SOLE suggests a $\sim 2.5 \text{ mm.yr}^{-1}$ NE-trending compression which is perpendicular to the azimuth of mapped faults. Our geomorphic data suggest that the Tuxtla fault is more likely to have produced these earthquakes.
855 The Concordia fault trace is not associated to any major fault scarps and rivers display a rather low degree of perturbation. By contrast the Tuxtla fault scarp displays a very young and prominent morphology (high hypsometry and low surface roughness) and the drainage network responded to a significant base-level fall ($> 500 \text{ m}$, profiles 1 to 3 in Fig. 14). Moreover, major topographic uplift took place east of the Tuxtla fault (Comitán High and the Sierra de Chiapas) where the elevation of
860 relict landscapes is between 1600 and 2400 m.

Finally, our results bring new evidence for topographic uplift within the Maya Mountains. Most of the Maya Mountains consist in an uplifted low relief landscape. The isobase surface from rivers is more elevated in the southern part of the Maya Mountains. It suggests thus stronger uplift in the south. The analysis of rivers longitudinal profiles is consistent with this interpretation. The reconstruction of profiles made using the upper reaches (associated to the low relict surface) show that
865 rivers faced a $\sim 500 \text{ m}$ base level fall in the south (profiles 6, 4 and 8 in Fig. 16). Northern rivers (profiles 1 to 3 in Fig. 16) were affected by a lower (ca. 300 m) base level change. The topographic profile across the Maya Mountains and the Yucatán peninsula (Fig. 5) is consistent with a geometry of blocks tilted towards the NW along NE-trending normal faults.

870 5.3 Proposed model of the plate boundary

The proposed model for the North American–Caribbean–Cocos plate boundary is displayed in Fig. 19. This model incorporate main ideas from previous works. East of 91°W the dynamic of the plate boundary is guided by the eastward escape of the Caribbean plate. This motion results in sinistral shear along the Polochic and Motagua slivers, dextral shear along the Jalpatagua fault and extension mainly concentrated along the grabens of Guatemala City and Ipala (Fig. 18b and 19; e.g., 875 Lyon–Caen et al., 2006; Authemayou et al., 2011). Recent models of the triple junction also agree on the fact that both North American and Caribbean plates are limited by a rigid forearc sliver to the west (Turner et al., 2007; Phipps-Morgan et al., 2008; Authemayou et al., 2011). The eastward escape of the Caribbean plate results in a counterclockwise rotation of the rigid forearc sliver 880 (Phipps-Morgan et al., 2008; Authemayou et al., 2011).

The interactions between the forearc sliver and the North American plate were summarized as distributed compression north of the Polochic fault (Guzmán-Speziale and Meneses-Rocha, 2000; Andreani et al., 2008a; Witt et al., 2012b). Our interpretation, based on the characterization of elevated relict landscapes and published geophysical data, is that the compression between the forearc 885 sliver and the North American is mainly absorbed at the boundary of small tectonic blocks which act as ‘buffers’ (Fig. 18b). As a result, significant topographic uplift occurred along two narrow domains. The first one encompasses the Sierra de Chiapas and the western tip of the Polochic sliver (Sierra de los Cuchumatanes) and the second one correspond to the northern tip of the volcanic arc (Fig. 19).

890 Within the Sierra Madre de Chiapas, the distribution of highest relict landscapes (>1500 m) indicates that significant topographic uplift occurred between the Tuxtla fault and reverse structures located east of the Sierra de Chiapas and Comitán High. Our results and those of Authemayou et al. (2011) also provide evidence for vertical motions along the Tonalá Shear Zone. There are evidence for a Miocene sinistral motions along the Tonalá Shear Zone and for Tuxtla and Malpaso faults 895 (Guzmán-Speziale and Meneses-Rocha, 2000; Witt et al., 2012b; Molina-Garza et al., 2015). However, the GPS data suggest a present-day $\sim 2.5 \text{ mm.yr}^{-1}$ extension along the Tonalá Shear Zone and a similar amount of compression along the Tuxtla fault (Fig. 18). These two opposite movements are possibly related to a counterclockwise rotation of the Chiapas Massif (Fig. 19) following a reorganization of the subduction interface. The Middle Miocene to present evolution of the Sierra Madre 900 de Chiapas is associated to an inland migration of the Chiapanecan volcanic arc (Damon and Montesinos, 1978) resulting from a flattening of the subduction beneath southern Mexico (Manea and Manea, 2006; Manea et al., 2013). This flattening would have resulted in higher coupling along the subduction interface, as suggested by Franco et al. (2012). As a result, the areas west of the Tuxtla fault (which is located along most of the present day volcanic centers) became partially incorporated 905 into the forearc domain. This would explain why significant Plio-Quaternary topographic growth (> 1000 m) mainly occurred east of the Tuxtla fault rather than along the Tonalá Shear Zone.

The compression between the rotating forearc sliver and the North American plate resulted in topographic uplift along the Sierra de los Cuchumatanes and the northern tip of the Central America volcanic arc (Authemayou et al., 2011, this study). In existing models, the westernmost corner of the Caribbean plate is stretched along NE-trending normal faults and the counterclockwise rotation of the forearc sliver is associated to a zipping (i.e. a suture zone) between the inferred prolongation of the Jalpatagua and Motagua faults (Authemayou et al., 2011; Franco et al., 2012). However, a suture zone between the Jalpatagua and Motagua faults may not fully explain why the volcanic arc is uplifted, mainly because most of the elevated Volcanic arc plateau is located west of the area where both faults would be zipped. To explain the atypical motions of GPS stations along the volcanic arc and the topographic elevation of the plateau (MAZ, CHL and SSIA in Fig. 18a) we infer a fault zone bounding the western edge of the volcanic arc (Fig. 18 and 19) and along which the volcanic arc plateau is uplifted. The indentation of the volcanic arc to the north possibly resulted in the topographic rejuvenation of the southern Chiapas Massif as well.

The interaction between the Maya Mountains and the rest of the plate boundary are unclear. Our hypothesis is that a part of the compression associated to the uplift of the volcanic arc and the Sierra de Cuchumatanes is transmitted towards the Yucatán platform and the Maya Mountains through the Petén basin. The Petén basin is a surprisingly flat and low area with a mean elevation of ~ 150 m (swath profile 4 in Fig. 5). It contrast greatly with the topography of the Sierra de Chuchumatanes which instead culminates at ~ 3800 m and the scarp of the Ixcán fault is almost 1000 m high (profile 4 in Fig. 5). We suspect that the Petén basin is underlain by a rigid basement which acts as a rigid crustal block. The extreme topography of the Sierra de Chuchumatanes can be explained by an abutment along this more rigid block. A rigid Petén block would also explain how a residual part of the motion along the triple junction may have been transmitted to the Yucatán peninsula and Maya Mountains. This residual motion may have reactivated NE-trending faults inherited from the Eocene opening of the Yucatán basin (Rosencrantz, 1990; Leroy et al., 2000).

6 Conclusions

We demonstrate that a geomorphic analysis allows to classify zones of similar relief patterns that we assume witnessed different tectonic and erosive histories. Using DEM-based geomorphic indices, we examined the topography of the Sierra Madre de Chiapas and the Maya Mountains. We used topographic profiles and morphometric maps in order to understand the spatial distribution of landscapes and we also analyzed in detail the disequilibrium of drainage network in order to map potential vertical displacements. Finally, we combined our results with existing GPS and seismological data in order to better understand the interactions between tectonics and landscapes within the highly diffuse North American–Caribbean–Cocos plate boundary.

Our analysis indicates that the topography of the Sierra Madre de Chiapas and the Maya Mountains is in transient stage. Topographic profiles and morphometric maps highlight elevated relict landscapes which are characterized by a low amplitude relief. These relict landscapes are surrounded by areas being actively eroded. River longitudinal profiles display knickzones which separate an upper reach associated to these relict landscapes from lower and steeper segments along which channels adjust to new base-level conditions. The relict landscapes from the Sierra Madre de Chiapas and the Maya Mountains probably evolved from an initially low topography which was then uplifted and fragmented by tectonics. This initial low relief topography is possibly a northern equivalent of the so-called "Mayan paleosurface" (Authemayou et al., 2011; Brocard et al., 2011) which covered most of the Polochic-Motagua sliver and formed at low elevation during the Middle Miocene (between 13 and 7 Ma).

East of 91°W the dynamic of the triple junction is mainly related to the eastward escape of the Caribbean plate (Lyon–Caen et al., 2006; Authemayou et al., 2011), which resulted in sinistral shear along the Polochic and Motagua slivers, dextral shear along the Jalpatagua fault and extension mainly concentrated along the grabens of Guatemala City and Ipala. West of 91°W we mainly observe compression resulting from the counterclockwise rotation of the Cocos forearc sliver. Within the Sierra Madre de Chiapas, this compression resulted in a major topographic uplift east of the Tuxtla and La Venta faults (Sierra de Chiapas and La Venta slivers in Fig. 18b). Within the Sierra de Chiapas and Comitán High, relict landscapes are found at elevations between 1600 and 2400 m. The areas between the Tuxtla and La Venta faults to the east and the Tonalá shear zone to the west were also uplifted but to a lesser extent (400 to 600 m). Major uplift is also observed along the western tip of the Polochic sliver (Sierra de los Cuchumatanes) and the northern tip of the volcanic arc. Using GPS data and morphotectonic analyses, we propose that the northern tip of the Central America volcanic arc is part of a tectonic sliver which has been pinned between the Cocos forearc sliver and the North American plate (Fig. 18b). Our study also brings new evidence for a 250 to 500 m topographic uplift within the Maya Mountains. Our hypothesis is that a residual part of the compression along the plate boundary is transmitted towards the Yucatán platform and the Maya Mountains through a rigid basement underlying the Petén basin.

Acknowledgements. The authors thank John Armitage, José Vicente Pérez-Peña and Giulia Sofia for their constructive reviews and comments.

References

- Albani, M., Klinkerberg, B., Andison, D., and P., K. J.: The choice of window size in approximating topographic surfaces from Digital Elevation Models, *Int. J. Geogr. Inf. Sci.*, 18, 577–593, 2004.
- Ambraseys, N. N. and Adams, R. D.: Large-magnitude Central American earthquakes, 1898–1994, *Geophysical Journal International*, 127, 665–692, 1996.
- 975 Andreani, L., Le Pichon, X., Rangin, C., and Martínez-Reyes, J.: The Southern Mexico Block : main boundaries and new estimation for its Quaternary motion, *Bulletin de la Société Géologique de France*, 179, 209–223, 2008a.
- Andreani, L., Rangin, C., Martínez-Reyes, J., Le Roy, C., Aranda-García, M., Le Pichon, X., and Peterson-Rodriguez, R.: The Neogene Veracruz fault: evidences for left-lateral slip along the southern Mexico block, *Bulletin de la Société Géologique de France*, 179, 195–208, 2008b.
- 980 Andreani, L., Stanek, K. P., Gloaguen, R., Krentz, O., and Domínguez-González, L.: DEM-based analysis of interactions between tectonics and landscapes in the Ore Mountains and Eger Rift (East Germany and NW Czech Republic), *Remote Sensing*, 6, 7971–8001, doi:10.3390/rs6097971, 2014.
- 985 Authemayou, C., Brocard, G., Teyssier, C., Simon-Labric, T., Gutiérrez, A., Chiquín, E. N., and Morán, S.: The Caribbean–North America–Cocos triple junction and the dynamics of the Polochic–Motagua fault systems: pull-up and zipper models, *Tectonics*, 30, 1–23, doi:10.1029/2010TC002814, 2011.
- Authemayou, C., Brocard, G., Teyssier, C., Suski, B., Cosenza, B., Morán-Ical, S., González-Véliz, C. W., Aguilar-Hengstenberg, M. A., and Holliger, K.: Quaternary seismo-tectonic activity of the Polochic Fault, *Guatemala, J. Geophys. Res.*, 117, B07403, doi:10.1029/2012JB009444, 2012.
- 990 Bateson, J. H. and Hall, I. S. H.: The geology of the Maya Mountains, Belize, Institute of Geological Sciences (Great Britain), Natural Environment Research Council, Overseas Memoir, 3, 1–44, 1977.
- Bauer-Gottwein, P., Gondwe, B. R. N., Charvet, G., Marín, L. E., Rebolledo-Vieyra, M., and Merediz-Alonso, G.: Review: The Yucatán Peninsula karst aquifer, Mexico, *Hydrogeology Journal*, 19, 507–524, 2011.
- 995 Bookhagen, B.: High resolution spatiotemporal distribution of rainfall seasonality and extreme events based on a 12-year TRMM time series, available at: <http://www.geog.ucsb.edu/~bodo/TRMM/#tif> (last access: 20 November 2015), 2009.
- Böse, E.: Informe sobre los temblores de Zanatepec a fines de septiembre de 1902 y sobre el estado actual del volcán de Tacaná, vol. 1, *Parergones del Instituto Geológico, Imprenta y Fototipía de la Secretaría de Fomento, Ciudad de México, México*, 1903.
- 1000 Brocard, G., Teyssier, C., Dunlap, W. J., Authemayou, C., Simon-Labric, T., Cacao-Chiquín, E. N., Gutiérrez-Orrego, A., and Morán-Ical, S.: Reorganization of a deeply incised drainage: role of deformation, sedimentation and groundwater flow, *Basin Research*, 23, 631–651, doi:10.1111/j.1365-2117.2011.00510.x, 2011.
- Burbank, D. W. and Anderson, R. S.: *Tectonic Geomorphology*, Blackwell Science, Cambridge, 2001.
- 1005 Burkart, B.: Offset across the Polochic fault of Guatemala and Chiapas, Mexico, *Geology*, 6, 328–332, 1978.
- Burkart, B.: Neogene North American-Caribbean plate boundary across Northern Central America: offset along the Polochic fault, *Tectonophysics*, 99, 251–270, 1983.
- Burrough, P. A. and Mcdonell, R.: *Principles of Geographical Information Systems*, Oxford University Press, New York, US, 1998.

- 1010 Chen, Y.-C., Sung, Q., and Cheng, K.-Y.: Along-strike variations of morphotectonic features in the Western Foothills of Taiwan: tectonic implications based on stream-gradient and hypsometric analysis, *Geomorphology*, 56, 109–137, 2003.
- Clark, M. K., Maheo, G., Saleeby, J., and Farley, K. A.: The non-equilibrium landscape of the southern Sierra Nevada, California, *GSA Today*, 15, 4–10, 2005.
- 1015 Correa-Mora, F., DeMets, C., Alvarado, D., Turner, H. L., Mattioli, G., Hernandez, D., Pullinger, C., Rodriguez, M., and Tenorio, C.: GPS-derived coupling estimates for the Central America subduction zone and volcanic arc faults: El Salvador, Honduras and Nicaragua, *Geophys. J. Int.*, 179, 1279–1291, 2009.
- Damon, P. E. and Montesinos, E.: Late Cenozoic volcanism and metallogenesis over an active Benioff zone in Chiapas, Mexico, *Arizona Geological Society Digest*, X1, 155–168, 1978.
- 1020 Deaton, B. C. and Burkart, B.: Time of sinistral slip along the Polochic fault of Guatemala, *Tectonophysics*, 102, 297–313, 1984.
- DeMets, C.: A new estimate for present-day Cocos-Caribbean plate motion: implications for slip along the Central American volcanic arc, *Geophysical Research Letters*, 28, 4043–4046, 2001.
- Domínguez-González, L., Andreani, L., Stanek, K. P., and Gloaguen, R.: Geomorpho-tectonic evolution of the Jamaican restraining bend, *Geomorphology*, 228, 320–334, 2015.
- 1025 Dury, G. H.: *Map Interpretation*, Sir Isaac Pitman & Sons, Ltd, London, U.K., 1952.
- Fairfield, J. and Leymarie, P.: Drainage networks from grid digital elevation models, *Water Resources Research*, 27, 709–717, 1991.
- Ferrater, M., Booth-Rea, G., Pérez-Peña, J. V., Azañón, J. M., Giaconia, F., and Masana, E.: From extension to transpression: Quaternary reorganization of an extensional-related drainage network by the Alhama de Murcia strike-slip fault (eastern Betics), *Tectonophysics*, 663, 33–47, doi:10.1016/j.tecto.2015.06.011, 2015.
- 1030 Figueroa, J.: *Sismicidad en Chiapas*, Instituto de Ingeniería de la UNAM, Ciudad de México, México, 1973.
- Filosofov, V. P.: *Brief Guide to Morphometric Methods in Search of Tectonic Structures (in russian)*, Saratov University Publishing House, Saratov, Russia, 1960.
- 1035 Flint, J. J.: Stream gradient as a function of order, magnitude and discharge, *Water Resources Research*, 10, 969–973, 1974.
- Font, M., Amorese, D., and Lagarde, J. L.: DEM and GIS analysis of the stream gradient index to evaluate effects of tectonics: the Normandy intraplate area (NW France), *Geomorphology*, 119, 172–180, 2010.
- Franco, A., Lasserre, C., Lyon-Caen, H., Kostoglodov, V., Molina, E., Guzmán-Speziale, M., Monterosso, D., Robles, V., Figueroa, C., Amaya, W., Barrier, E., Chiquin, L., Moran, S., Flores, O., Romero, J., Santiago, J. A., Manea, M., and Manea, V. C.: Fault kinematics in northern Central America and coupling along the subduction interface of the Cocos Plate, from GPS data in Chiapas (Mexico), Guatemala and El Salvador, *Geophysical Journal International*, 189, 1223–1236, 2012.
- Fuchs, M. C., Gloaguen, R., Merchel, S., Pohl, E., Sulaymonova, V. A., Andermann, C., and Rugel, G.: Denudation rates across the Pamir based on ^{10}Be concentrations in fluvial sediments: dominance of topographic over climatic factors, *Earth Surf. Dyn.*, 3, 423–439, 2015.
- 1045 Gallen, S. F., Wegmann, K. W., and Bohnenstiehl, D. R.: Miocene rejuvenation of topographic relief in the southern Appalachians, *GSA Today*, 23, 4–10, doi:10.1130/GSATG163A.1, 2013.


- García-Acosta, V. and Suárez-Reynoso, G.: Los sismos en la historia de México. Tomo I, Fondo de Cultura Económica, Ciudad de México, México, 1996.
- 1050 Garrity, C. and Soller, D.: Database of the Geologic Map of North America; adapted from the map by J.C. Reed, Jr. and others (2005), U.S. Geological Survey Data Series 424, 2009.
- 1055 Giletycz, S., Loget, N., Chang, C. P., and Mouthereau, F.: Transient fluvial landscape and preservation of low-relief terrains in an emerging orogen: example from Hengchun Peninsula, Taiwan, *Geomorphology*, 231, 169–181, 2015.
- Godard, V., Bourlès, D. L., Spinabella, F., Burbank, D. W., Bookhagen, B., Burch Fisher, G., Moulin, A., and Léanni, L.: Dominance of tectonics over climate in Himalayan denudation, *Geology*, 42, 243–246, 2014.
- Golts, S. and Rosenthal, E.: A morphotectonic map of the northern Arava in Israel, derived from isobase lines, *Geomorphology*, 7, 305–315, 1993.
- 1060 Gordon, M. and Muehlberger, W. R.: Rotation of the Chortis block causes dextral slip on the Guayapé fault, *Tectonics*, 13, 858–872, 1994.
- Grohmann, C. H.: Morphometric analysis in Geographic Information Systems: applications of free software GRASS and R, *Comput. Geosci.*, 30, 1055–1067, 2004.
- Grohmann, C. H., Riccomini, C., and Alves, F. M.: SRTM-based morphotectonic analysis of the Pocos de 1065 Caldas Alkaline Massif, southeastern Brazil, *Comput. Geosci.*, 33, 10–19, 2007.
- Grohmann, C. H., Smith, M. J., and Riccomini, C.: Surface roughness of topography: a multi-scale analysis of landform elements in Midland Valley, Scotland, *Proceedings of Geomorphometry 2009, Zurich, Switzerland*, 31 August - 2 September 2009, pp. 140–148, 2009.
- Grohmann, C. H., Riccomini, C., and Chamani, M. A. C.: Regional scale analysis of landform configuration 1070 with base-level (isobase) maps, *Earth Syst. Sci.*, 15, 1493–1504, 2011.
- Guzmán-Speziale, M.: Beyond the Motagua and Polochic faults: active strike-slip faulting along the western North America–Caribbean plate boundary zone, *Tectonophysics*, 496, 17–27, 2010.
- Guzmán-Speziale, M. and Meneses-Rocha, J. J.: The North America–Caribbean plate boundary west of the Motagua-Polochic fault system: a fault jog in southeastern Mexico, *Journal of South American Earth Sci-* 1075 *ences*, 13, 459–468, 2000.
- Guzmán-Speziale, M., Pennington, W. D., and Matumoto, T.: The triple junction of the North America, Cocos, and Caribbean plates: seismicity and tectonics, *Tectonics*, 8, 981–997, 1989.
- Guzmán-Speziale, M.: Active seismic deformation in the grabens of northern Central America and its relationship to the relative motion of the North America–Caribbean plate boundary, *Tectonophysics*, 337, 39–51, 1080 2001.
- Hack, J. T.: Studies of longitudinal stream profiles in Virginia and Maryland, U.S. Geological Survey Professional Paper, 294, 45–97, 1957.
- Hack, J. T.: Interpretation of erosional topography in humid temperate regions, *Am. J. Sci.*, 258, 80–97, 1960.
- Hergarten, S., Robl, J., and Stüwe, K.: Extracting topographic swath profiles across curved geomorphic features, 1085 *Earth Surf. Dynam.*, 2, 97–104, doi:10.5194/esurf-2-97-2014, 2014.
- Hobson, R. D.: Surface roughness in topography: quantitative approach, in: *Spatial analysis in geomorphology*, edited by Chorley, R. J., pp. 225–245, Methuen, London, UK, 1972.

- Isacks, B. L.: Long term land surface processes: erosion, tectonics and climate history in mountain belts, in: TERRA-1: Understanding the Terrestrial Environment, edited by Mather, P. M., pp. 21–36, Taylor and Francis, London, UK, 1992.
- 1090 Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled seamless SRTM data V4, International Centre for Tropical Agriculture (CIAT), available at: <http://srtm.csi.cgiar.org> (last access: 25 November 2014), 2008.
- Jones, R.: Algorithms for using a DEM for mapping catchment areas of stream sediment samples, *Computers & Geosciences*, 28, 1051–1060, 2002.
- 1095 Keller, E. A. and Pinter, N.: *Active tectonics : earthquakes, uplift, and landscape*, Prentice Hall, Englewood Cliffs, N.J., 1996.
- Kesler, S. E., Kienle, C. F., and Bateson, J. H.: Tectonic Significance of Intrusive Rocks in the Maya Mountains, British Honduras, *Geological Society of America Bulletin*, 85, 549–552, 1974.
- Kirby, E. and Whipple, K. X.: Quantifying differential rock–uplift rates via stream profile analysis, *Geology*, 1100 29, 415–418, 2001.
- Kirby, E. and Whipple, K. X.: Expression of active tectonics in erosional landscapes, *J. Struct. Geol.*, 44, 54–75, 2012.
- Lara, M. E.: Divergent wrench faulting in the Belize southern lagoon: implications for Tertiary Caribbean plate movements and Quaternary reef distribution, *American Association of Petroleum Geologists Bulletin*, 77, 1105 1041–1063, 1993.
- Legrain, N., Stüwe, K., and Wölfler, A.: Incised relict landscapes in the eastern Alps, *Geomorphology*, 221, 124–138, 2014.
- Leroy, S., Mauffret, A., Patriat, P., and Mercier de Lépinay, B.: An alternative interpretation of the Cayman trough evolution from a reidentification of magnetic anomalies, *Geophysical Journal International*, 141, 539–1110 557, 2000.
- Lesser, J. M. and Weidie, A. E.: Region 25, Yucatan Peninsula, in: *Hydrogeology*, edited by Back, W., Rosenhein, J. S., and Seaber, P. R., pp. 237–242, Geological Society of America, Boulder, Colorado, 1988.
- Lyon–Caen, H., Barrier, E., Lasserre, C., Franco, A., Arzu, I., Chiquin, L., Chiquin, M., Duquesnoy, T., Flores, O., Galicia, O., Luna, J., Molina, E., Porrás, O., Requena, J., Robles, V., Romero, J., and Wolf, R.: Kinematics 1115 of the North American–Caribbean–Cocos plates in Central America from new GPS measurements across the Polochic–Motagua fault system, *Geophysical Research Letters*, 33, L19 309, doi:10.1029/2006GL027694, 2006.
- Mahmood, S. A. and Gloaguen, R.: Appraisal of active tectonics in Hindu Kush: Insights from DEM derived geomorphic indices and drainage analysis, *Geosci. Front.*, 3, 407–428, doi:10.1016/j.gsf.2011.12.002, 2012.
- 1120 Malfait, B. T. and Dinkelman, M. G.: Circum–Caribbean tectonic and igneous activity and evolution of the Caribbean plate, *Geological Society of America Bulletin*, 83, 251–272, 1972.
- Manea, V. C. and Manea, M.: Origin of the modern Chiapanecan volcanic arc in southern Mexico inferred from thermal models, *Geological Society of America Special Paper*, 411, 27–38, 2006.
- Manea, V. C., Manea, M., and Ferrari, L.: A geodynamical perspective on the subduction of Cocos and Rivera 1125 plates beneath Mexico and Central America, *Tectonophysics*, 609, 56 – 81, 2013.
- Mann, P. and Burke, K.: Cenozoic rift formation in the Northern Caribbean, *Geology*, 12, 732–736, 1984.

- Masek, J. G., Isacks, B. L., Gubbels, T. L., and Fielding, E. J.: Erosion and tectonics at the margins of continental plateaus, *J. Geophys. Res.*, 99, 13 941–13 956, doi:10.1029/94JB00461, 1994.
- 1130 Mather, A. E.: Adjustment of a drainage network to capture induced base-level change, *Geomorphology*, 34, 271–289, 2000.
- Matmon, A., Bierman, P., Larsen, J., Southworth, S., Pavich, M., and Caffee, M.: Temporally and spatially uniform rates of erosion in the southern Appalachian Great Smoky Mountains, *Geology*, 31, 155–158, 2003.
- Meneses-Rocha, J. J.: Tectonic evolution of the Ixtapa Graben, an example of a strike-slip basin of southeastern Mexico: implications for regional petroleum systems, in: *The Western Gulf of Mexico Basin: Tectonics, Sedimentary Basins and Petroleum Systems*, AAPG Memoir 75, edited by Bartolini, C., Buffler, R. T., and Cantú-Chapa, A., pp. 183–216, The American Association of Petroleum Geologists, Tulsa, Oklahoma, 2001.
- 1135 Molina-Garza, R. S., Geissman, J. W., Wawrzyniec, T. F., Peña Alonso, T. A., Iriando, A., Weber, B., and Aranda-Gómez, J.: Geology of the coastal Chiapas (Mexico) Miocene plutons and the Tonalá shear zone; syntectonic emplacement and rapid exhumation during sinistral transpression, *Lithosphere*, 7, 257–274, 2015.
- 1140 Montgomery, D. R. and Foufoula-Georgiou, E.: Channel network source representation using digital elevation models, *Water Resour. Res.*, 29, 3925–3934, 1993.
- O’Callaghan, J. F. and Mark, D. M.: The extraction of drainage networks from digital elevation data, *Comput. Vision Graph.*, 28, 323–344, 1984.
- 1145 Pazzaglia, F. J.: Landscape evolution models, in: *The Quaternary Period in the United States*, edited by Gillespie, A., Porter, S., and Atwater, B., pp. 247–274, Elsevier Science Ltd.: Oxford, UK, 2003.
- Pedrerá, A., Pérez-Peña, J. V., Galindo-Zaldívar, J., Azañón, J. M., and Azor, A.: Testing the sensitivity of geomorphic indices in areas of low-rate active folding (eastern Betic Cordillera, Spain), *Geomorphology*, 105, 218–231, 2009.
- 1150 Peraldo, G. and Montero, W.: *Sismología Histórica de América Central*, Instituto Panamericano de Geografía e Historia, Ciudad de México, México, 1999.
- Pérez-Peña, J. V., Azañón, J. M., Booth-Rea, G., Azor, A., and Delgado, J.: Differentiating geology and tectonics using a spatial autocorrelation technique for the hypsometric integral, *J. Geophys. Res. Earth Surface*, 114, F02 018, doi:10.1029/2008JF001092, 2009.
- 1155 Pérez-Peña, J. V., Azañón, J. M., Azor, A., Booth-Rea, G., Galve, J. P., Roldán, F. J., Mancilla, F., Giaconia, F., Morales, J., and Al-Awabdeh, M.: Quaternary landscape evolution driven by slab-pull mechanisms in the Granada Basin (Central Betics), *Tectonophysics*, 663, 33–47, doi:10.1016/j.tecto.2015.07.035, 2015.
- Phipps-Morgan, J., Ranero, C. R., and Vannucchi, P.: Intra-arc extension in Central America: links between plate motions, tectonics, volcanism, and geochemistry, *Earth and Planetary Science Letters*, 272, 365–371, 2008.
- 1160 Pike, R. J. and Wilson, S. E.: Elevation relief ratio, hypsometric integral, and geomorphic area-altitude analysis, *Geological Society of America Bulletin*, 82, 1079–1084, 1971.
- Plafker, G.: Tectonic aspects of the Guatemalan earthquake of 4 February 1976, *Science*, 193, 1201–1208, 1976.
- Purdy, E. G., Gischler, E., and Lomando, A. J.: The Belize margin revisited. 2. Origin of Holocene antecedent topography, *International Journal of Earth Sciences*, 92, 552–572, 2003.
- 1165

- Rao, R. P. and Ramanathan, R.: Belize 1988-89 petroleum activity keyed to prices, *Oil and Gas Journal*, 86, 81-91, 1988.
- Ratschbacher, L., Franz, L., Min, M., Bachmann, R., Martens, U., Stanek, K., Stübner, K., Nelson, B. K., Herrmann, U., Weber, B., López-Martínez, M., Jonckheere, R., Sperner, B., Tichomirowa, M., McWilliams, M. O., Gordon, M., Meschede, M., and Bock, P.: The North American-Caribbean Plate boundary in Mexico-Guatemala-Honduras, *Geological Society, London, Special Publications*, 328, 219-293, 2009.
- 1170 Reinhardt, L. J., Bishop, P., Hoey, T. B., Dempster, T. J., and Sanderson, D. C. W.: Quantification of the transient response to base-level fall in a small mountain catchment: Sierra Nevada, southern Spain, *J. Geophys. Res.*, 112, F03S05, doi:10.1029/2006JF000524, 2007.
- 1175 Rodriguez, M., DeMets, C., Rogers, R., Tenorio, C., and Hernandez, D.: A GPS and modelling study of deformation in northern Central America, *Geophysical Journal International*, 178, 1733-1754, 2009.
- Rogers, R. D., Kárasón, H., and van der Hilst, R. D.: Epeirogenic uplift above a detached slab in northern Central America, *Geology*, 30, 1031-1034, 2002.
- Rosencrantz, E.: Structure and tectonics of the Yucatan basin, Caribbean Sea, as determined from seismic-reflection studies, *Tectonics*, 9, 1037-1059, 1990.
- 1180 Sánchez-Barreda, L. A.: Geologic evolution of the continental margin of the Gulf of Tehuantepec in southern Mexico, Ph.D. thesis, University of Texas, Austin, Texas, 1981.
- Schoenbohm, L. M., Whipple, K. X., Burchfiel, B. C., and Chen, L.: Geomorphic constraints on surface uplift, exhumation, and plateau growth in the Red River region, Yunnan Province, China, *Geol. Soc. Am. Bull.*, 116, 895-909, 2004.
- 1185 Schumm, S. A.: Evolution of drainage systems and slopes in badlands at Perth Amboy, New Jersey, *Geological Society of America Bulletin*, 67, 597-646, 1956.
- Schwanghart, W. and Kuhn, N. J.: TopoToolbox: A set of Matlab functions for topographic analysis, *Environmental Modelling & Software*, 25, 770-781, 2010.
- 1190 Scotti, V., Molin, P., Faccenna, C., Soligo, M., and Casas-Sainz, A.: The influence of surface and tectonic processes on landscape evolution of the Iberian Chain (Spain): quantitative geomorphological analysis and geochronology, *Geomorphology*, 206, 37-57, 2014.
- Shahzad, F. and Gloaguen, R.: TecDEM: A MATLAB based toolbox for tectonic geomorphology, Part 1: Drainage network preprocessing and stream profile analysis, *Computers & Geosciences*, 37, 250-260, 2011a.
- 1195 Shahzad, F. and Gloaguen, R.: TecDEM: A MATLAB based toolbox for tectonic geomorphology, Part 2: Surface dynamics and basin analysis, *Computers & Geosciences*, 37, 261-271, 2011b.
- Siddiqui, S. and Soldati, M.: Appraisal of active tectonics using DEM-based hypsometric integral and trend surface analysis in Emilia-Romagna Apennines, northern Italy, *Turk. J. Earth Sci.*, 23, 277-292, doi:10.3906/yer-1306-12, 2014.
- 1200 Singh, S. K., Rodríguez, M., and Espindola, J. M.: A catalog of shallow earthquakes of Mexico from 1900 to 1981, *Bulletin of the Seismological Society of America*, 74, 267-279, 1984.
- Smith, M. W.: Roughness in the Earth Sciences, *Earth Sci. Rev.*, 136, 202-225, doi:10.1016/j.earscirev.2014.05.016, 2014.

- 1205 Snyder, N. P., Whipple, K. X., Tucker, G. E., and Merritts, D. J.: Landscape response to tectonic forcing: digital elevation model analysis of stream profiles in the Mendocino triple junction region, northern California, *Geological Society of America Bulletin*, 112, 1250–1263, 2000.
- Sofia, G., Pirotti, F., and Tarolli, P.: Variations in multiscale curvature distribution and signatures of LiDAR DTMs errors, *Earth Surf. Proc. Land.*, 38, 1116–1134, 2013.
- 1210 Sofia, G., Dalla Fontana, G., and Tarolli, P.: High-resolution topography and anthropogenic feature extraction: testing geomorphometric parameters in floodplains, *Hydrol. Process.*, 28, 2046–2061, 2014.
- Steiner, M. B. and Walker, J. D.: Late Silurian plutons in Yucatan, *Journal of Geophysical Research*, 101, 17 727–17 735, doi:10.1029/96JB00174, 1996.
- Strahler, A. N.: Hypsometric (area-altitude) analysis of erosional topography, *Geological Society of America Bulletin*, 63, 1117–1142, 1952.
- 1215 Strahler, A. N.: Quantitative analysis of watershed geomorphology, *Transactions of the American Geophysical Union*, 8, 913–920, 1957.
- Tarolli, P., Sofia, G., and Dalla Fontana, G.: Geomorphic features extraction from high-resolution topography: landslide crowns and bank erosion, *Natural Hazards*, 61, 65–83, 2012.
- Telbisz, T., Kovács, G., Székely, B., and Szabó, J.: Topographic swath profile analysis: a generalization and sensitivity evaluation of a digital terrain analysis tool, *Z. Geomorph.*, NF 57, 485–513, doi:10.1127/0372-8854/2013/0110, 2013.
- Troiani, F. and Della Seta, M.: The use of the Stream Length–Gradient index in morphotectonic analysis of small catchments: a case study from Central Italy, *Geomorphology*, 102, 159–168, doi:10.1016/j.geomorph.2007.06.020, 2008.
- 1225 Turner, H. L., LaFemina, P., Saballos, A., Mattioli, G. S., Jansma, P. E., and Dixon, T.: Kinematics of the Nicaraguan forearc from GPS geodesy, *Geophysical Research Letters*, 34, L02 302, doi:10.1029/2006GL027586, 2007.
- Wawrzyniec, T., Molina-Garza, R. S., Geissman, J., and Iriondo, A.: A newly discovered, relic, transcurrent plate boundary: The Tonalá shear zone and paleomagnetic evaluation of the western Maya block, SW Mexico, *Geological Society of America, Salt Lake City Annual Meeting*, October 16–19, Abstracts with Programs, 37, p. 68, 2005.
- 1230 Weber, B., Cameron, K. L., Osorio, M., and Schaaf, P.: A Late Permian tectonothermal event in Grenville crust of the southern Maya Terrane: U–Pb zircon ages from the Chiapas Massif, southeastern Mexico, *International Geology Review*, 47, 509–529, 2005.
- 1235 Weber, B., Iriondo, A., Premo, W., Hecht, L., and Schaaf, P.: New insights into the history and origin of the southern Maya block, SE Mexico: U–Pb–SHRIMP zircon geochronology from metamorphic rocks of the Chiapas massif, *International Journal of Earth Sciences*, 96, 253–269, 2007.
- Weidie, A. E.: Lineaments of the Yucatan Peninsula and fractures of the Central Quintana Roo coast, in: Field trip no. 10 – Yucatan, road log and supplement to 1978 guidebook, pp. 21–25, GSA annual meeting, New Orleans, Louisiana, October 18–21, 1982.
- 1240 Weidie, A. E.: Geology of Yucatan platform, in: *Geology and Hydrogeology of the Yucatan and Quaternary Geology of Northeastern Yucatan Peninsula*, edited by Ward, W. C., Weidie, A. E., and Back, W., pp. 1–19, New Orleans Geol. Soc., New Orleans, Louisiana, 1985.

- White, R. A.: Catalog of historic seismicity in the vicinity of the Chixoy-Polochic and Motagua faults, **1245** Guatemala, Open-File Report 84–88, U. S. Geological Survey, Office of Earthquakes, Volcanoes and Engineering, Menlo Park, California, pp. 1–26, 1984.
- Whittaker, A. C., Attal, M., Cowie, P. A., Tucker, G. E., and Roberts, G.: Decoding temporal and spatial patterns of fault uplift using transient river long profiles, *Geomorphology*, 100, 506–526, 2008.
- Willett, S. and Brandon, M.: On steady states in mountain belts, *Geology*, 30, 175–178, 2002.
- 1250** Witt, C., Bricchau, S., and Carter, A.: New constraints on the origin of the Sierra Madre de **Chiapas** (south Mexico) from sediment provenance and apatite thermochronometry, *Tectonics*, 31, TC6001, doi:10.1029/2012TC003141, 2012a.
- Witt, C., Rangin, C., Andreani, L., Olaz, N., and Martínez, J.: The transpressive left-lateral Sierra Madre de Chiapas and its buried front in the Tabasco plain (Southern Mexico), *Journal of the Geological Society of London*, 169, 143–155, 2012b. 
- Wobus, C., Whipple, K. X., Kirby, E., Snyder, N., Johnson, J., Spyropoulou, K., Crosby, B., and Sheehan, D.: Tectonics from topography: procedures, promises and pitfalls, *Geological Society of America Special Paper*, 398, 55–74, 2006.




Figure 1. Main plate boundaries in Central America (black lines) and location of the Sierra Madre de Chiapas (SMC) and Maya Mountains (MM) in red. The red box shows the extend of Fig. 2. Abbreviations: CB – Chortis block, CFS – Central America forearc sliver, MB – Maya block, SMB – Southern Mexico block. Topography and bathymetry from the General Bathymetric Chart of the Oceans (GEBCO, <http://www.gebco.net/>)




Figure 2. Topography (A) and simplified geological map (B) for northern Central America. Geological contours from Garrity and Soller (2009). Structures compiled from Guzmán-Speziale (2001); Meneses-Rocha (2001); Rogers et al. (2002); Purdy et al. (2003); Ratschbacher et al. (2009); Authemayou et al. (2011) and Witt et al. (2012b). Morpho-tectonic domains: CAVA – Central America volcanic arc, CFTB – Chiapas fold-and-thrust belt, CH – Comitán High, CM – Chiapas Massif, CR – Chuacus range, CVC – Central Valley of Chiapas, EYFZ – East Yucatán fault zone, GCA – Grabens of Central America, LMR – Las Minas range, MM – Maya Mountains, PB – Petén basin, SCH – Sierra de Chiapas, SdC – Sierra de los Cuchumatanes, TCP – Tabasco coastal plain, YP – Yucatán platform. Main structures: CoF – Concordia fault, CoG – Comayagua graben, GuG – Guatemala City graben, HSFS – High Sierra fault system, IpG – Ipala graben, IxF – Ixcán fault, JaF – Jalpatagua fault, LVF – La Venta fault, MaF – Malpas Fault, MoF – Motagua fault, NaF – Necta fault, NBF – Northern boundary fault, PoF – Polochic fault, RHF – Rio Hondo fault, SBF – Southern boundary fault, SuG – Sula graben, TSZ – Tonalá shear zone, TuF – Tuxtla fault.




Figure 3. Superficial (depth <40 km) seismicity in northern Central America (A) and GPS velocities with respect to the fixed North American plate (B). Focal mechanisms from Guzmán-Speziale et al. (1989); Guzmán-Speziale (2010) (in greys) and Global CMT Catalog (in black, <http://www.globalcmt.org>). Historical earthquakes (red circles) from White (1984); Singh et al. (1984) and Guzmán-Speziale (2010). GPS velocities are from Franco et al. (2012). See Fig. 2 for abbreviations.




Figure 4. Recent models for the North American–Caribbean–Cocos plate boundary in northern Central America. A: ‘fault-jog’ model from Guzmán-Speziale and Meneses-Rocha (2000) and Guzmán-Speziale (2001). B: model from Lyon–Caen et al. (2006) and Franco et al. (2012). C: model from Andreani et al. (2008a). D: ‘zipper’ model from Authemayou et al. (2011). Main plates and blocks: CAFS – Central America forearc sliver, CB – Chortis block (part of the Caribbean plate), NAP – North American plate, SMB – Southern Mexico block. Main structures: JF – Jalpatagua fault, GCA – grabens of Central America, MAT – Middle America trench, PMFS – Polochic-Motagua fault system, SSF – strike-slip faults of Chiapas, TSZ – Tonalá shear zone.




Figure 5. Location of swath topographic profiles (A) and annual averaged precipitation derived from the Tropical Rainfall Measuring Mission (TRMM) for the 1998 to 2009 period (B). Processed TRMM data are provided by Bookhagen (2009). See Fig. 2 for abbreviations.




Figure 6. Swath profiles across the Sierra Madre de Chiapas, volcanic arc, and Maya Mountains. Upper plots: topographic profiles (swath width is 20 km). Vertical exaggeration is 10 for profiles 1 to 4 and 25 for profile 5. Red, black and blue curves represent the maximum, mean and minimum elevations, respectively. Lower plots: Green lines represent the local incision in km (subtraction of minimum and maximum elevations). Blue dashed lines represent annual averaged precipitation (derived from TRMM data) in m/yr. See Fig. 2 for abbreviations.




Figure 7. Hypsometric integral (A) and surface roughness (B) for northern Central America. See Fig. 2 for abbreviations.




Figure 8. Relief anomaly (A) and surface index (B) for northern Central America. See Fig. 2 for abbreviations.




Figure 9. Comparison between precipitation(derived from TRMM data) and morphometric indices within the Sierra Madre de Chiapas. Points represents pixel values extracted from the precipitation (Fig. 5b) and morphometric maps (Fig. 7 and 8). The linear correlation between the variables was estimated using the Pearson correlation coefficient (PCC).




Figure 10. Distribution of morphometric indices for the main lithological groups of the Sierra Madre de Chiapas. Histograms are based on three sets of 5000 pixel values extracted from the morphometric maps (Fig. 7 and 8). Red line represents the median.




Figure 11. Analyzed streams in Chiapas Massif and Central America volcanic arc. Top: location map. Blue lines represent channels extracted from the modeled drainage network. White dots represent major knickzones. Red lines show limits of catchments. Black lines and red triangles represent main faults and active volcanoes, respectively. Bottom: plot of steepness index (k_{sn}) values and estimated base-level changes along a profile passing through the outlet of each catchment. Abbreviations: MoF – Motagua fault. NaF – Nectá fault. PoF – Polochic fault. TSZ – Tonalá shear zone.




Figure 12. Examples of stream profiles extracted from the modeled drainage network in Chiapas Massif and volcanic arc. Upper plots: blue lines represent longitudinal profiles, green lines show the contributing area (flow accumulation), red lines show reconstructed profiles (interpolation of the slope-distance relation from segments above the upper knickzones, see equation 7) and pink polygons represent the uncertainties (based on a bootstrapping method). Central plots: black lines represent extracted steepness index (k_{sn}) values for a given segment and grey polygons show the uncertainties (based on a bootstrapping method). Lower plots: logarithmic plots of slope vs area (grey crosses) and envelopes of regressions (black lines) used to estimate the k_{sn} values.




Figure 13. Longitudinal profiles for the Grijalva river and the tributaries from the Sierra de los Cuchumatanes and Chiapas Massif. Profiles were extracted from the modeled drainage network. See Fig. 2 for faults abbreviations and Fig. 12 for plot description.




Figure 14. Longitudinal profiles for the Sierra de Chiapas. Channels and profiles were extracted from the modeled drainage network. See Fig. 12 for plot description. Abbreviations: LVF – La Venta fault, MaF – Malpaso Fault, SCF – San Cristobal fault, TecF – Tectapan fault, TenF – Tenejapa fault, TuF – Tuxtla fault.




Figure 15. Interpolated map of steepness index (k_{sn}) for the Sierra Madre de Chiapas and the northern segment of the Central America volcanic arc. Values were extracted from river longitudinal profiles using 5 km long segments and interpolated using a natural neighbor (NN) method. See Fig. 2 for abbreviations.




Figure 16. Analysis of drainage network for the Maya Mountains. Top: map of isobases (i.e., interpolated elevations) from rivers with a Strahler order ≥ 2 . Main faults (black lines), analyzed streams (blue lines) and main knickpoints (blue dots) are also reported. Bottom: longitudinal stream profiles. Blue dots are main knickpoints (location on the map). Red lines show profiles reconstructed using segments located above the knickpoints (see equation 7). Black arrows show the intersection with main faults. Abbreviations: NBF – Northern Boundary fault, SBF – Southern Boundary fault, RHF – Rio Hondo fault, F – other faults.




Figure 17. Proposed extension of elevated surfaces and relict landscapes in northern Central America (interpretative map based on morphometric maps, Fig. 7 and 8). Legend: 1 – Remnants of the Miocene ignimbritic plateau (Rogers et al., 2002), 2 – Plio-Quaternary volcanic arc plateau, 3 – Middle Miocene Mayan paleosurface (Authemayou et al., 2011; Brocard et al., 2011), 4 – relict landscape of the Maya Mountains, 5 – tilted surfaces of the Yucatán platform, 6 – Petén basin, 7 – elevated plateaus of La Venta and northern Sierra Madre de Chiapas, 8 – inferred extension of the plateau within the fold-and-thrust belt, 9 – monadnock landscape developed over the Chiapas batholith, 10 – Central valley of Chiapas. See Fig. 2 for faults and morpho-tectonic domains abbreviations.




Figure 18. Proposed limits of tectonic slivers forming the plate boundary. **A:** GPS velocities with respect to the fixed North American plate (Franco et al., 2012). Colors emphasize vectors with similar azimuths and velocities. The diagram of northern and eastern components (bottom left corner) allows to deduce motions between paired GPS stations. **B:** Main tectonic blocks defined using known major faults (Fig. 2), our geomorphic maps (Fig. 7 and 8) and published GPS data (Fig. 18a) and seismicity (Fig. 3a). Black arrows show the motion along blocks boundaries deduced from paired GPS stations (values are in mm.yr⁻¹). See Fig. 2 for faults abbreviations.




Figure 19. Proposed model for the North American–Caribbean–Cocos plate boundary in northern Central America. Abbreviations: EYFZ – East Yucatán fault zone, GCA – Grabens of Central America, HSFS – High Sierra fault system, IxF – Ixcán fault, JaF – Jalpatagua fault, MAT – Middle America Trench, MoF – Motagua fault, PoF – Polochic fault, TSZ – Tonalá shear zone, TuF – Tuxtla fault.