

Response to comments by Anonymous Referee #2

We are grateful for the constructive comments by an anonymous reviewer (hereafter: reviewer) on our manuscript. Below we respond to all points raised in the review and how we changed our manuscript accordingly.

General:

The manuscript deals with the formation of theater-head hydrological channels by analogical sandy box experiments, testing different water supply processes (groundwater or association of precipitation and groundwater) and different locations of water sources (local water table or distant water table). My general point of view concerning this part of manuscript is quite positive, even if some clarifications must be done. Indeed, authors used "seepage" word in the title and used "sapping" word in the text. These are two different processes. At the manuscript reading, authors seem themselves unclear on what process(es) they modeled. It is quite confusing. Therefore, it is necessary to add the definition of these processes, list parameters that they differentiate and determine which process they model.

We can understand the view of the reviewer in this regard in terms of explaining more fully the definitions of the terminology to avoid confusion for those unfamiliar with the definitions of seepage and sapping. Seepage and sapping are of course linked: Seepage is the hydrological process of groundwater coming to the surface and groundwater sapping is a geomorphological process of erosion triggered by that seepage. We have extended the introduction to explain these differences more fully. We stick to using seepage in the title as seepage is the driving process, and other subsequent processes, other than sapping alone play a role in the formation of the resulting valleys.

Minor points must be added concerning the experimental design and parameters to help readers. In my point of view, they modeled seepage process in no cohesive material forming fluvial channels. They do not find channel bifurcations. How could they explain that?

We have added text to explain our thoughts on the absence of the formation of bifurcation in the discussion of the manuscript: the slope suppresses the formation of bifurcations.

Different Martian fluvial valley analogs have been explored in regards to the experiment results. Although this approach seems interesting and promising, the choice of examples is not relevant and has no direct relationships with parameters tested in analogical experiment. Indeed, they studied Martian valley networks (not channel networks) with theater-heads showing a lowly branching pattern. This 2D channel/valley organization is not observed in their experiments. Therefore, this part must be either removed from the part dealing with analogical experiment in other manuscript or try to find good Martian analogs without channel/valley bifurcations.

We disagree with the reviewer here. The formation of bifurcations was not observed in the experiment. The aim of our manuscript is to provide a framework of the range of possible landscapes and our experiments complement previous work. Our experiments are used to

improve our insight into seepage processes, which is required to complete the framework we present. The Martian examples we use are classic exemplary cases that we use to show how our framework maps onto these features. We think that some of the concerns raised here result from confusion on the terminology used, we think they are now addressed within the new introduction section where clearer definitions are outlined.

The second point concerns the valley formation in relation with the channel formation on Earth and Mars. How do they explain the formation processes in regards to their experiment? The third point is the origin of distant water table? How do they supply the distant water table, on Earth and on Mars? What are the climatic implications?

This is a good point. We have now extended the last section of the discussion (5.5) to incorporate sources of water for the groundwater system and relate this to previous work on Martian hydrology (Harrison and Grimm, 2004 / Clifford 1993).

The fourth point and the more critical is the lithology. Is it possible to compare a non cohesive material like sand to Martian lithology (expected to be volcanic rocks (from lava to volcanic ash) and locally sedimentary rocks not sediments)?

The reviewer has a fair point here. The lithology on Mars is largely unknown. It is likely to be very diverse in nature and in states of weathering or consolidation. This issue is unlikely to be resolved soon. In our manuscript, the unknown material properties are an important reason for focusing our research on entire landscapes rather than valley-head morphology alone (page 131, lines 16-24 of initial manuscript). In section 5.2 we do deal with limitations of the applicability of our experiment, and we direct the reader to Marra et al., 2014 / Icarus for further discussion on this point.

This point is not discussed. The manuscript is well enough written, but requires typewriting corrections. The revised manuscript is now extensively proof-read by the native speakers of our team.

Abstract

Rewrite abstract in regards to seepage vs sapping to be in line with the main manuscript outline or title. Line 5. "but the climate implications are quite different". Remove or explain why. Line 12. " We study sapping: : " seepage or sapping ? Line 14. Use another word than "piracy" or explain the physical process. Lines 19-21. You mention only two Martian examples whereas you present other examples in the part 4. Be coherent.

We have adjusted the abstract to avoid the use of the term sapping (we explain this term now later in the introduction).

We rephrased the statement on climate to: 'This morphological ambiguity hampers the interpretation of formative processes and responsible hydrology of such valleys on Mars'.

We also rephrased 'groundwater piracy' to the more common term 'groundwater flow-piracy' to make this part more clear. In our opinion, explaining the entire process is not appropriate for an abstract.

The two examples of Louros and Nirgal are used to exemplify our framework, the other 'examples' where the reviewer points at are mere illustrations to make a more appealing figure, which are not analyzed. We made this point clear in the abstract by rephrasing the linking sentence to: 'To exemplify our interpretive framework, we apply these results to two Martian cases.'

1. Introduction

p.2. Lines 23-25. It will be good that you describe geometrically the theater-head channels or valleys before listing terrestrial and Martian examples. In addition, it is necessary to discriminate formation processes between valleys and channels.

We now explain the difference between valleys and channels in the introduction. Furthermore, the addition of an extra figure (2) helps in identifying the shape of these valleys for the reader.

p.3. lines 5-15. Be clearer between processes and observed objects. For example, Laity and Malin (1985) explained the sapping process and not the seepage process.

We do not agree with the reviewer on their reading of the Laity and Malin paper. Just as our work, Laity and Malin describe the valley formation by groundwater sapping, i.e. undercutting and erosion by the seepage of groundwater. They also use both terms in the same way we do. We hope our reformulated introduction will address this and avoid mis-interpretation of these terms.

You mentioned that the "head-wall retreat is due to undercutting by water fall erosion of a hard cap rocks (Lamb et al. 2006) would be the best explanation for Snake river and Chilean valleys formation (Fig. 1 d and e). You ignore the contribution of runoff, overflow effect at local scale and infiltration forming temporary water table at regional scale. Lines 16-17. "the main argument against a groundwater origin of the Martian valleys is the limited erodibility by the low seepage discharge (Lamb et al., 2006). It is quite surprising, isn't it? How do you form valleys on Mars? Could you explain this point. p.3. Lines 19-21. No, there are Martian examples showing these morphological relationships (e.g., Mangold et al., 2004; Mangold et al., 2008, Ansan et al. 2008, Lucas et al., 2009).

These points by the reviewer highlighted that this part of our introduction needed rephrasing. We have clarified and restructured parts of the introduction based on the concerns of both reviewers.

p.3. Lines 22-24. I do not understand your arguments about climatic implications. How do you supply your groundwater table ?

We mean hydrological conditions, which we now explain explicitly in the text. Indeed, groundwater also requires a source, but this source could be far away in both space and time,

which puts less constraint on the current and local conditions. We reformulated the text to now read: "Waterfall erosion requires a large flood event or a long period with flowing water on the surface, which requires atmospheric conditions to sustain liquid water. Whereas seepage erosion could occur under surface conditions where liquid water is not stable, as the recharge of groundwater could be far in the past."

p.4. Line 1. Could you explain "ground water piracy" in terms of physical process.

This was explained in the previous sentence, we rephrased this to refer to that explanation better.

P.4 Lines 24-25. Could add references as Schumm (1986), Dunne et al. (1980)

These references are used in the introduction to set the stage and describe previous work and the current state of knowledge. In the lines referred to by the reviewer, we start describing the aims of our research, which does not seem an appropriate place to cite this work again.

p.4. Lines 5-23. Once again, sapping or seepage ? Flow piracy ?

The sapping vs seepage terminology issue has been clarified in response to the reviewer's comments, see comments above. Seepage is the process of outflow, sapping the undercutting by erosion. We avoid the use of sapping by using terminology as undercutting where appropriate. Flow piracy has now been more fully explained at the first occurrence in the introduction (4th paragraph)

p.4. Lines 24-30. Remove valley formation because you model only channel formation. Choose definitely what you model: Seepage or sapping ?

The seepage-sapping terminology issue has been resolved, see above. In this case, the channel / valley terminology is correct, since we do model valley formation.

p.5. Lines 1-7. "shows the arrangement of multiple valleys" Is it really your main objective? In your experiment you do not obtain multiple valleys with bifurcation.

We do show multiple valleys in our experiment. Indeed there are no bifurcations, but we combine previous results with novel experiments to make a complete interpretive framework. This is indeed the main objective.

Do you really constrain the climate conditions with your experiment? Give arguments.

This is a fair point of the reviewer, we do not constrain the climate conditions. Our research is limited to the hydrological conditions, which we now explain more clearly in both the introduction and discussion.

2. Methods

p.5. Lines 21-26. Could you justify your choices about sand granulometry, topographic slope of sandy surface (0.22 mm⁻¹) which is quite high for granular material. Why do you choose a partially sloping impermeable floor with a decreasing in slope value by a factor 2? This leads to a wedge of sandy layer from top to bottom of your experiment box, why this choice? How do justify the horizontal offset in slopes between the sandy surface and the impermeable surface? Which implications do you expect?

We used natural sand to mimic natural processes as far as possible and the grain size was such that the experiment progressed neither too quickly nor too slowly. We have added this detail on the choices made and clarifications in the manuscript.

We stated in the original manuscript that the wedging was done to ensure seepage at the surface and the fake floor was installed to reduce the amount of sediment required. This clarification comes directly after the description of the setup, and we feel this is in the right place. To guide the reader and give more focus to this part, we have also split this paragraph in the revised text.

p.6. Lines 9-11. Could you explain clearer how does work your head tank? At which depth does water flow: on entire tank height or at different heights? It is not clear in text and in Figure 2.

We have now clarified this point in the methods section.

p.6. Lines 20-28. Could you justify and explain your different mean discharges in local and distal experiments (cf. table1): one order of magnitude?

In the local experiment, there was more groundwater flow due to the higher groundwater table and a part of the precipitation fell directly into the valleys due to the uniform distribution of precipitation. We have extended the first paragraph of section 3.2 to explain this in the manuscript.

2.3 Valley development and eroded volume Change valley word by channel p.8. Line 25. Could you explain why you use SLc (shape index of channel cross-section) and 0.5 coefficient in your formula of channel volume. You obtain for each channel the DEM by stereogrammetry, so you can extract directly volume without this correction.

Full DEMs are only taken at a few points in time, whereas the length and width are captured at each time-lapse interval, so using the volumes of the DEM does not give the desired temporal resolution to show the cyclic behavior. Furthermore, we are not interested in the exact volume of erosion, but mainly in the patterns of erosion.

We use the volumes from the DEMS to check if the calculations using width, length and estimates of depth and valley shape are appropriate. Which they are, as shown in Figure 6. We added a short paragraph at the start of section 2.3 to clarify the purpose of these data.

In addition, this formula is not appropriate because you want to calculate channel V-shape whereas you obtain wide rectangular shape.

The formula is correct, we use the shape index to take into account the shape of the cross-section profile (V vs U shaped), the factor of 0.5 is to correct for the longitudinal shape, which is triangular (see Figure 5). We believe that the text is quite clear on this point and we have made no change concerning this point.

p.9. Line 5. Once again, I do not understand your calculation strategy in erosion rate. Why do use the density of sand? Why the high density value of your sand (2300 kg.m-3 and not 2300 kg.m-1)? How do you measure the porosity of sediment at the begin of experiment and during the experiment?

In line with the points discussed above, we are interested in the patterns of change and check calculated and measured erosion. Therefore we assume a default porosity and density for sand, which are values that are not subject to change more than a few percent; taking a measurement could introduce larger errors. Again, the short clarification we added at the start should help evaluating the meaning and required accuracy of these calculations.

Indeed, the dimensions for density were incorrectly typed into the manuscript, we have corrected this typographical error.

2.4 Martian landscape metrics In this part, change the word channel by valley because you analyze valleys at the surface of Mars.

Indeed, we corrected this (and in other places as indicated by the reviewers).

p.9. Line 24. How do you analyze Nirgal valley whereas you do not have accurate DEM?

As stated in our methods, we use planform (x-y) valley segments, which does not require elevation data. We rephrased the first paragraph of 2.4 to avoid the focus on the limited resolution.

Here we present the morphological analyze of valley bifurcations but you do not write about that in experiment methodology. It is incoherent.

We discussed this point already in this review: our experiments are designed to complete the set of knowledge and together with previous experiment we provide a framework of interpretation for these valleys. Our experiments do show details of groundwater-flow piracy that causes the valley length distribution that we show with the data presented for the Martian cases (this point is clarified in the introduction and discussed earlier in this response).

3.1. Distal groundwater source Here change the word "valley" by channel. p.11 Lines 13-15.

See discussion on valleys and channels elsewhere. Indeed, this instance should be channel, we corrected this line.

Could you explain the physical process of "groundwater piracy? Why is typical of groundwater sapping? Sapping, why? You model seepage.

We have modified the manuscript to explain this process in greater detail in the introduction and at the end of section 3.1 where we mention groundwater piracy.

p.11. Line 17. "the valleys did not bifurcate, which is the result of the high initial slope" where ? at the top or the bottom of sand layer? Do you test for small slope? In this case, do you observe bifurcations?

The discussion of headward valley bifurcations is based on previously published experiments available in the literature but this discussion should not be included in the results section. We have removed the comparison of our experiments to previous experiments from this section. The relevant discussion on this point was already included in the discussion section of the manuscript and was superfluous in any case.

p.11 Line 23. Change the word "mud flow". It is not appropriate here, it would correspond to concentrate flow but not mudflow. Idem p.12 line 13.

Indeed, the reviewer is correct. We rephrased these sentences.

p.11 Lines 5-14. I do not understand where you measured channel width (Fig. 6 and 7). Is it at the head? At outlet? A mean value? Could you clarify this.

We clarified this point in the methods section (2.3): the width is taken just downstream of the head where the walls are parallel to each other.

I suggest to permute §3.2 with §3.3 because it is easier to understand the influence of local topography on distal groundwater table.

Good suggestion, we moved these sections.

3.2. Effect of initial morphology on distal experiment. In this experiment you have channel widening and deepening without increase in headward erosion. It is interesting but you do not analyze what experiment parameters could influence this process. You remain evasive suggesting that only groundwater piracy would be the main parameters and the channel geometry develops independently from the initial topography. It is not really the case, because they do not crosscut previous channels. Could you investigate which parameters (geometry of water table, slopes etc. : :) in your experiment could explain this final morphology?

The valleys that develop in the Distant after local experiment are of similar size and spacing as the Distal source experiment (with an initial flat surface). Therefore we conclude, that the initial morphology does not have a significant effect on the valley development from a distal groundwater source. The reason for the absence of cross cutting is simply the result of the valleys developing in the same direction.

We do agree that the details mentioned by the reviewer influence the details of the morphological development, but it is the aim of our manuscript to show the differences between contrasting initial and boundary conditions (slope, no slope, local, distant). Looking at details and understanding these processes even better is an interesting challenge for further

research and requires experiments focusses on those details. For the interpretation of the valley patterns on the scale of landscapes, as we provide in this manuscript, we believe we have chosen the right scale and the correct level of detail.

3.3. Local groundwater source Why do you model a discharge with a higher factor of 5 than that in distal groundwater source? It could have to misinterpretation when we compare experiment results. This water comes from local precipitation in your experiment. What was the precipitation rate? In which terrestrial climatic conditions?

Discharge rates of the experiments are explained and quantified in the methods section, we incorporated the comments of the reviewer there. The higher factor between the constant-head tank and precipitation experiment is the result of the design of the precipitation experiment where we wanted infiltration in the flat area and nearby seepage. The resulting and reported discharges are merely the result of that design. We added a sentence to the end of section 2.1 to clarify. Secondly, the experiments were carried out in terrestrial conditions, which we now state in section 2.1.

p.13. Line 13. "valley heads with a v-shaped planform (Fig. 3b-ii)" It is not clear on your figure. However this paragraph is quite convincing. But you have no channel bifurcation.

We have indicated the v-shaped heads in the figure with the time-lapse images. Indeed there were no bifurcations, as expected for the steep slope.

3.4. Effect of initial morphology on local groundwater table. This part is interesting but authors do not analyze completely the experiment result. It is pity. Channels widen and stretch out by headward erosion up to flat surface where a feather morphology develops. You observe channel bifurcation ! Which is process at the origin of that? Which parameters control that because you have not overpression? Could you explain.

The points raised by the reviewer here are already part of the revised manuscript, we describe the widening of the valleys and the overflow of the flat upstream area. However, the bifurcations that develop here are purely the result of converging overland-flow, which is a completely different process than the bifurcation that develop due to the groundwater flow processes described in the discussion of the paper. Since this is a different process and this last stage of the experiment does not contribute to the goal (effect of initial morphology), and we have thus shortened this section. Furthermore, in many places we now try to be as clear as possible when it comes to 'headward bifurcations', to avoid confusion.

4. Examples of Martian valley system. This part is not convincing in relation with the experimental results. Valley vs channel? Could you explain? Examples are not well chosen. This part must be reworked.

The aim of our work is not to just recreate a few seepage valleys and find perfect analogues to them, the overarching aim is to create an interpretative framework for a range of possible

morphologies. We do this by combining previous to novel experiment. Therefore, to show the applicability of our work, we choose more complex systems with multiple valleys that split as this both shows the piracy effect and bifurcation dynamics as indicated in previous work. For our purpose, i.e. contrasting a more local vs a more distal scenario, we think our chosen case studies work well in showing those differences.

We expanded the introduction paragraph of section to explain our reasons for these examples to the reader more fully.

5. Discussion. Once again, you model channels not valleys !

We do not agree with the reviewer on this point. We have both in the experiment. There are fluvial channels conveying the water and there is valley formation by erosion and incision. Valley formation is the main thing we are after. We have added terminology in the introduction to avoid confusion.

p. 17. Line 16. It is very ambiguous "The general morphology of our experimental channels (by seepage) agrees well with studies on sapping" Could you clarify. You discuss the nature of material as major parameter in experiment result (sand). You do not discuss the geometry of your experimental model on channel formation (thickness of sand, variation in slope between the upper and bottom surface, etc. : :)

Scale-effects of the experiments are discussed in a later section, 5.2, and are not discussed here for clarity to the reader. That section (5.2) has been extended based on the concerns of both the reviewer.

5.4. Headward channel bifurcation You conclude that the lack of bifurcation is due to the high slope in your experiment. Is it the only parameter? Others experiments obtain bifurcation with a horizontal surface. It is the only geometric parameter (topographic surface?)?

The bifurcation behavior mainly depends on slope and is favored by groundwater from a local source, as explained in paragraph 2 of section 5.4, flat horizontal surfaces and a local source increase the tendency to bifurcate. Our experiments do not feature these bifurcations as there is a steep slope, i.e. we study one extreme end of the parameter space, which complements previous experimental work. We have not changed our manuscript in light of this point, since we think section 5.4 adequately discusses this point.

5.5. Origin of Martian examples This part must be reworked, examples are not in correlation with your experiment results. You wrote sapping and seepage, which is very confusing. You do not discuss the scaling between your channels and Martian valley, the lithology on Mars: : The analogy is quite limited.

Our experiments were not designed to be a direct scale model or analogue to these Martian cases. Instead, they are designed to provide insight into the processes and morphological development associated with seepage erosion. We added a paragraph to section 5.2 on scaling where we explain our view and the applicability of the experiments.

Table 1. Could you add the signification of each term in legend, for example d=day etc: : :

For consistency, we changed hr. to h. for hours and we added explanation to the legend as suggested by the reviewer.

Groundwater seepage landscapes from distant and local sources in experiments and on Mars

Wouter A. Marra¹, Stuart J. McLelland², Daniel R. Parsons², Brendan J. Murphy², Ernst Hauber³, and Maarten G. Kleinhans¹

¹Faculty of Geosciences, Universiteit Utrecht, Heidelberglaan 2, 3584 CS, Utrecht, the Netherlands

²Department of Geography, Environment and Earth Sciences, University of Hull, Cottingham Road, Hull, HU6 7RX, UK

³Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Planetenforschung, Rutherfordstraße 2, 12489 Berlin, Germany

Correspondence to: W. A. Marra (w.a.marra@uu.nl)

Abstract. ~~Theater-headed valleys can form by undercutting caused by the seepage of groundwater~~, but such valleys can also be the Valleys with theater-shaped heads can form due to the seepage of groundwater or as a result of knick-point (waterfall) erosion generated by overland flow. This ~~morphological ambiguity~~ ambiguity in the mechanism of formation hampers the interpretation of formative processes and responsible hydrology of such valleys on Mars, especially due to insufficient particularly since there is limited knowledge of material properties. ~~Instead of single-valley morphology, metrics of the entire landscape~~ Moreover, the hydrological implications of a groundwater or surface water origin are important for our understanding of the evolution of planet Mars and quantification of valley morphologies at the landscape scale may provide diagnostic ~~insight in~~ insights on the formative hydrological conditions. However, flow patterns and the resulting landscapes ~~are different for different~~ produced by different sources of groundwater ~~and are~~ poorly understood. We aim to ~~increase our~~ improve the understanding of the formation of ~~the entire landscapes by seepage from different sources of groundwater and to entire valley landscapes through seepage processes from different groundwater sources that will~~ provide a framework of landscape metrics ~~of such systems to aid for the~~ interpretation of such ~~landscapes systems~~. We study ~~seepage from local and distal sources of groundwater in~~ groundwater seepage from a distant source of groundwater and from infiltration of local precipitation in a series of sandbox experiments and combine our results with previous experiments and observations of the Martian surface. Key results are that groundwater ~~flow piracy acts on distally-fed valleys, which~~ flow piracy acts on valleys fed by a distant groundwater source and results in a sparsely dissected landscape of many small and a few large valleys ~~while locally-fed valleys~~. In contrast, valleys fed by a local groundwater source, i.e. nearby infiltration, result in a densely dis-

sected landscape. In addition, ~~distally-fed valleys grow into the direction of the groundwater source while locally-fed valleys~~ valleys fed by a distant groundwater source grow towards that source while valleys with a local source grow in a broad range of directions and have a strong tendency to bifurcate, particularly on ~~flat horizontal surfaces~~. ~~To exemplify our interpretive framework, we apply these results~~ flatter surfaces. We consider these results with respect to two Martian cases. ~~The valleys of Louros Valles seem to have formed by seepage erosion from~~: Louros Valles show properties of seepage by a local source of groundwater and Nirgal Vallis shows evidence of a ~~distal source of groundwater~~. distant source, which we interpret as groundwater flow from Tharsis.

30 1 Introduction

~~Add short section on a step broader: channels, valleys, v-shaped, seepage, sapping. Then elaborate a bit on theatre-shaped valley heads and link with next.~~

Valleys with theater-shaped heads exist in the landscapes of Earth and Mars. On Mars, examples of such valleys are ~~for example~~ Louros Valles (Fig. 1a) and Nirgal Vallis (Harrison and Grimm, 2005). Terrestrial examples can be found in the Atacama Desert in Chile (Fig. 1b), on the Canterbury Plain, New Zealand, on the Colorado plateau and on Hawaii (Schumm and Phillips, 1986; Howard and Kochel, 1988; Craddock et al., 2012). Furthermore, much smaller ~~;~~ but examples that are similar in shape, are valleys that emerge in eroding riverbanks (Fig. 1c) or those on the beach that develop during a receding tide (Higgins, 1982; Otvos, 1999; Fox and Wilson, 2010; Hagerty, 1991).

~~The small valleys on the beach and riverbanks form in several hours by seeping~~ Such theater-headed valleys can form by the seepage of groundwater in ~~unconsolidated sediment~~. ~~The governing processes and morphology are the same as in groundwater seepage experiments (e.g. Howard and McLane, 1988) and consist of undercutting by erosion by groundwater seepage and resulting head wall collapse. This process is also known to be responsible for the valleys in the Colorado plateau sandstone (e.g. Laity and Malin, 1985), and has long been thought to be the cause of the formation of valleys on Hawaii and on Mars (Kochel and Piper, 1986). However, head wall retreat due to undercutting by waterfall erosion of a hard cap rock produces similar valley morphology (Lamb et al., 2006) and is probably a better explanation for the erodible sediment (e.g. Howard and McLane, 1988). These valleys form due to headward erosion that is produced by mass-wasting processes where groundwater seeps to the surface (Fig. 2a). In this paper we define *seepage* as the hydrological process of groundwater emerging at the surface and groundwater *sapping* as the geomorphological process of erosion by undercutting which is triggered by seepage, although not all erosion by seepage of groundwater results in undercutting. We define a *valley* as an erosional geomorphological feature, where a *channel* is a body of flowing water. The aim of this study is to investigate valley formation that develops from groundwater seepage that is sufficient to maintain an active channel.~~

Valley formation by seepage (Fig. 2a) is different from valley formation by overland flow. In the former, headward progression is the result of knickpoint retreat and fluvial incision (Fig. 2b). However, overland flow can also produce similar theater-headed valleys when incising into a substrate with an erosion resistant top-layer (Lamb et al., 2006). This process is a likely candidate for the formation of the theater-headed valleys next to the Snake River in Idaho (Fig. 1d) and the Chilean valleys (Irwin et al., 2014).

The main argument against a groundwater origin of the valleys hampers the interpretation of their origin based on their theater-shaped morphology alone. This ambiguity is particularly problematic for the Martian valleys is the limited erodibility by the low seepage discharge (Lamb et al., 2006), but the material properties of Mars are largely unknown, which hinders unambiguous interpretation of the formative mechanism of the theater-headed valleys. Furthermore, small upstream channels feeding the main valleys are expected for waterfall erosion which are visible in terrestrial cases (e.g., Fig. 1e), but evidence for such a source is missing for many Martian cases. Most importantly, formation by either waterfall erosion or groundwater seepage has different implications for the responsible hydrological conditions. Waterfall erosion requires a large flood event or a long period with flowing water on the surface, which requires atmospheric conditions to sustain liquid water. Whereas seepage erosion could occur under surface conditions where liquid water is not stable, as the recharge of groundwater could be far in the past. explanation of theater-shaped valley-heads on Mars where direct field observations and material properties are lacking, and a long period of weathering obscures morphological details.

Besides the single-valley morphology, the landscape properties of the morphological properties of entire landscapes with multiple valleys may help in the interpretation of the Martian valleys these Martian valleys, when single entities have an ambiguous origin. An important feedback for mechanism for the landscape formation by groundwater seepage is flow piracy, groundwater flow piracy: since valleys are topographic depressions, hence they attract more groundwater from their surrounding, possibly resulting in the abandonment of other valleys (Howard and McLane, 1988). surroundings, resulting in a decrease of discharge to nearby valleys. (Howard and McLane, 1988). As a result, smaller valleys cease to develop in favor of larger valleys. Landscape metrics may provide information on show the presence of feedback mechanisms like groundwater piracy this feedback mechanism.

Furthermore, the angle of valley bifurcations, and the splitting of valleys during their headward development (headward bifurcation) produces typical angles between valley segments Devauchelle et al. (2012); Glines and Fassett (2012). In case of valley formation by seepage from uniform precipitation, the theoretical angle between valley segments becomes 72° (Devauchelle et al., 2012). Such properties as well as the orientation of valleys could indicate the (Jaumann and Reiss, 2002) can be extracted from the landscape and show the responsible hydrological system and the location of the groundwater source (Jaumann and Reiss, 2002; Devauchelle et al., 2012; Glines and Fassett, 2012).

Our knowledge of the groundwater sapping groundwater seepage processes and their relation to landscape evolution is rather limited. Particularly as systems with only groundwater processes are

absent on Earth and previous ~~experimental~~ studies were mostly limited to the same boundary conditions of groundwater from ~~a~~ an upstream constant-head tank (e.g. Howard and McLane, 1988; 95 Lobkovsky et al., 2004; Schorghofer et al., 2004; Pornprommin et al., 2010; Marra et al., 2014a). These experiments simulated a distant groundwater source and showed the basic morphology of ~~sapping valleys~~, revealing valleys formed by groundwater seepage and also revealed the importance of groundwater flow piracy as a key process for valley evolution. However, alternate hydrological systems exist where other processes ~~play a role, which results are significant and result~~ in a different 100 landscape evolution. ~~For example, in~~ In contrast to a distant groundwater source, ~~a local groundwater source can release water from a shallow aquifer, generated by local~~ groundwater seepage could come from nearby infiltration of precipitation. Such systems exist on Earth (Abrams et al., 2009; Fig. 1f) and have been studied in experiments to some extent (Berhanu et al., 2012), but require more attention in terms of their morphological impact on landscape dynamics. We hypothesize that a local 105 groundwater source, e.g. as result of locally infiltrated precipitation, results in less groundwater flow piracy compared to groundwater that first travels some distance ~~because a local~~ before seepage to the surface, because seepage from a nearby source is less influenced by the ~~regional topographic gradient~~ topographic gradient responsible for flow piracy. Since flow piracy is an important mechanism in the formation of ~~theater-headed~~ valleys by groundwater seepage, we expect ~~a different~~ 110 different and distinct morphological development for ~~locally-fed sapping valleys~~. valleys formed by seepage from a local source compared to those produced by distally fed systems. Specifically, distant sources of groundwater produces landscapes with abandoned valleys whereas in landscapes produced by local sources of groundwater, there are no or hardly any abandoned valleys.

In this paper, we aim to ~~increase~~ improve our understanding of groundwater seepage processes, 115 specifically on the resulting valley formation. ~~In order to expand our knowledge on the possible range of processes, we~~ at a landscape scale using morphological experiments. We specifically study the difference in morphological development of valleys that ~~emerge from distant and local sources of groundwater in sealed experiments~~ result from a distant groundwater source (simulated with an upstream constant-head tank), and a local groundwater source (simulated by infiltrating precipitation). 120 Furthermore, we combine our experimental insight with previous studies in order to ~~cover~~ show a complete range of ~~possible sapping systems~~ landscapes formed by groundwater seepage under different conditions. The objective is ~~essentially~~ to provide a framework that shows the arrangement of multiple valleys, i.e., the orientation and length distribution that ~~result~~ results from different hydrological boundary conditions. These properties will aid in the identification of the formative 125 processes when the single valley morphology is ambiguous, and will constrain the ~~hydrological and climate conditions~~ during formation. As an example, we apply our developed framework to two often-cited underlying hydrological conditions. To demonstrate the application of this framework for landscape interpretation, we use two frequently cited cases of groundwater seepage on Mars, Nirgal Vallis and Louros Valles, and relate their landscape metrics to the possible sources of groundwater.

130 2 Methods

2.1 Experimental design

~~We conducted several~~ We conducted a series of flume experiments in the Total Environmental Simulator (TES) facility at the University of Hull to ~~study~~ investigate the morphological development of theater-headed valleys by ~~seeping groundwater~~ groundwater seepage. Moreover, with these experiments we ~~studied the effect of groundwater seepage from a local and from a distal source~~. ~~We simulated groundwater seeping from a distal source by~~ simulated the difference between groundwater seepage generated by a distant source of groundwater, using a ~~fixed-head constant~~ groundwater level at the upstream end of the ~~setup~~. ~~As a local source of groundwater, we applied precipitation on the entire experiment domain, using an intensity~~ experimental setup, and groundwater seepage produced
140 by infiltration of precipitation applied over the entire experimental domain. The distant groundwater source was maintained using a constant-head reservoir at a level below the sediment surface and the precipitation was generated using an array of nozzles spraying water at a rate lower than the infiltration capacity of the sediment. We repeated both experiments with an idealized initial morphology and with a heterogeneous initial morphology that was the result of a previous experiment.
145 In this section, we present the setup, the initial topography and applied boundary conditions across the experimental set.

The initial idealized morphology consisted of a volume of sand with a median grain size of 0.7 mm, that comprised (1) a flat area of 1.7 m upstream and (2) a slope of 0.22 m m^{-1} for 3.5 m (Fig. 3), which was ~~equal uniform~~ over a width of 4 m. ~~The~~ We used natural, moderately angular sand
150 to mimic natural groundwater and surface flow processes. The grain size was such that groundwater flow was neither too rapid nor too slow for the formation of valleys within a reasonable period of time. The sloping section ensured seepage of groundwater from a hydrostatic groundwater level, ~~is not pressurized~~. ~~that is, without applying extra pressure to the groundwater.~~

The sediment was placed on top of a partially sloping impermeable floor to increase groundwater flow in the downstream half of the setup and to reduce the amount of sediment required. This
155 floor was flat for 2.6 m and the slope was 0.11 m m^{-1} for 2.6 m. Pond liner ensured the impermeability of the floor and walls. A rough cloth on top of the pond liner prevented the entire block of sediment from sliding down the smooth pond liner surface. The total sediment depth was 0.5 m in the upstream flat part, sloping towards the downstream end. At the downstream end, a row of 6 cm
160 high bricks truncated the wedging slope to prevent the sediment from sliding down. In addition, the small spaces between these 10 cm wide bricks acted as initial surface perturbations. This ensured ~~evenly distributed channel initiation~~ valley initiation was evenly distributed over the entire width of the sediment surface.

~~To simulate a groundwater system from a distal source, we used a constant head tank at the upstream end of the setup. This constant head tank was constructed over~~ The constant head tank
165

was designed to simulate a distant groundwater source and was constructed opposite of the sloping wedge of sediment. It spanned the entire width ~~of the setup using and~~ depth of the sediment fill with a 0.6 mm mesh fabric, braced with chicken wire and steel gratings ~~. To simulate a local groundwater system at the water-side to retain the sediment and avoid collapse into the reservoir under the weight of the sand. This setup enabled water from the constant-head tank to enter the body of sediment over the entire width and depth at the upstream side of the flat section of the sediment fill.~~

For the experiment with precipitation as the water source, we used ~~a series~~ an array of spray nozzles above the setup to supply water over the sediment surface. These nozzles were ~~set to spray at such discharge fed with a discharge such~~ that the water infiltrated, ~~which corresponds with a discharge spray infiltrated in the flat area and seeped-out on the sloping wedge. The discharge feed was slightly~~ lower than the infiltration capacity of the sediment. A rising groundwater table induced seepage, but, in contrast to the constant head tank, the seepage areas were fed by ~~locally-nearby~~ infiltrated groundwater. Twelve spray nozzles with square spray patterns were used ~~, which were pressurized to ensure uniform spray distribution and pressurized water was fed~~ from a ring main to ensure equal spray rates for all nozzles.

We carried out five experiments with the above described boundary condition combinations (Table 1) ~~. Experiment “Distal” under terrestrial conditions. The experiment labelled *Distant source* was carried out entirely with the constant head tank as boundary condition with the constant-head tank throughout as the boundary condition using the initial topography described above. The final morphology of this surface morphology from the *Distant source* experiment was used as initial morphology of experiment “Local after Distal” the experiment labelled *Local after distant*, which was run with water input from the spray nozzles. Experiments “Local” The experiment labelled *Local precipitation* was run with water input from the spray nozzles on the above-described initial topography. Experiment “Local 2” was only The experiment labelled *Local precipitation 2* was run to generate an initial morphology with the same conditions as “Local”, but *Local precipitation and* this experiment was ceased early. The final morphology of “Local 2” *Local precipitation 2* was the initial morphology of “Distal after Local” *Distant after local*, which was subject to groundwater flow from the constant head generated by the constant-head tank.~~

2.2 Experimental imagery and elevation models

We captured the morphological development of the experiments using time-lapse photography. These images ~~allowed~~ enabled us to study the experiments in detail from different angles. Moreover, we derived ~~channel-valley~~ dimensions from orthorectified time-lapse images. The time-lapse setup consisted of six cameras (Canon PowerShot A640), mounted around the experimental setup (see C1–C6 in Fig. 3), which were ~~synchronously triggered~~ triggered *synchronously* at set intervals. These intervals ~~differed~~ ranged from 30 s to 5 min, based on the ~~pace of the morphological development in rate of morphological development during~~ the ongoing experiment (values in Table 1).

For each experiment, time-lapse imagery from four ~~angles were processed to a single orthorectified image. Camera's cameras~~ (C1, C2, C4 and C5) ~~were processed to construct a single orthorectified photograph~~ (Fig. 3) ~~were used to construct orthorectified images~~. Orthorectification was ~~conducted on the initial surface elevation model. Using~~ implemented using the 'Image Processing' and the 'Camera Calibration' Toolboxes in the MATLAB software suite. Orthorectification was performed using the initial surface elevation ~~, the orthorectified images were distorted in morphologically altered areas. However, the dimensions of erosional features, such as the valleys in our experiments, remain correct~~ model, due to absence of elevation data at each time-step. This method resulted in warped imagery in areas with elevation change, i.e. within the valleys. However, these images were used to extract valley lengths and widths which are calculated from the distances between non-eroded valley walls. The outside edge of these walls form in the original surface and are therefore correctly represented in the orthorectified photograph using this method. ~~We performed time-lapse orthorectification in MATLAB, using the Image Processing Toolbox and the Camera Calibration Toolbox for MATLAB.~~

215

For detailed morphological analysis ~~and morphometrical analysis, we collected~~, we generated digital elevation models and orthorectified images at the end of each experiment using a large set of images and a structure-from-motion (SfM) algorithm (Forsyth and Ponce, 2011). In addition, we also acquired these data ~~several times during the Distal and Distal-after-Local~~ every day during the ~~Distant source and Distant after local~~ experiment at irregular intervals (Table 1).

220

~~Elevation models were acquired at the end of each experiment and during the experiments Distal and Distal-after-Local every day of the experiment (see Table 1) using SfM. For each~~ Each digital elevation model and associated orthorectified image ~~, we took a set of about 70–100~~ were derived from 70 to 100 digital images with about 80 % overlap. We took these images by hand, allowing us to capture the area of interest. Twenty-four targets with known coordinates within the experimental setup enabled referencing of the images. Camera positions and orientations were solved using these ~~known~~ target coordinates and matched features between images. To ~~increase~~ improve the quality of the result, we removed features in wet areas ~~in order~~ to eliminate faulty matches in reflective areas due to different lightning angles between images. Elevation models were generated for each set of images, which were processed to a gridded elevation model with 1 mm cell size and a 0.5 mm orthorectified images. We used a Canon 550D DSLR camera with an 18–55 mm f/2.8 lens to take the photos, which we processed in RAW format to 16 bit TIF images to eliminate compression artifacts. We used Agisoft PhotoScan for SFM processing (Agisoft, 2014).

230

~~In addition, we used a~~ A laser scanner ~~to obtain~~ was also used to obtain digital elevation models at the end of each experiment. Point-cloud elevation data of the final morphology were ~~obtained using a laser scanner from two~~ scanned from two different angles in order to eliminate data shadows outside the line-of-sight of the ~~laser~~ scanner (see Fig. 3). These point-clouds were oriented ~~and merged~~ using fixed targets in the experimental setup ~~. The combined scans were gridded to to produce~~

235

a combined scan gridded onto a digital elevation model with a 2 mm ~~elevation models~~ cell size.

240 We used a Leica ScanStation 2 laser scanner for data acquisition, CloudCompare for point-cloud orientation and ArcGIS for gridding of the point-cloud data.

2.3 Valley development and ~~eroded volume~~ erosion rates

To quantify the morphological development, we plotted measured valley widths, lengths and depths during the experiments. Based on these data, we calculated valley shapes, and erosion rates and
245 compared the latter to measured sediment output.

The length L (m) and width W (m) of each valley that formed during the experiments were determined for each time-lapse interval from the orthorectified time-lapse images. The valley width was taken just downstream of the valley-head where the valley walls were parallel. Valley depth D (m) was defined as the deepest point of each valley, i.e. the largest elevation difference between
250 the original surface and the eroded surface. Valley depth was measured at each SfM interval. The final valley floor slope S_f (m m^{-1}) was extracted from the final DEM. We defined the valley floor as the lowest point in each valley cross section. We estimated the erosion rates of each valley during the experiments by combining the valley dimensions and valley shapes.

First, the eroded volume V (m^3) was estimated as:

$$255 \quad V = W \cdot L \cdot D \cdot \text{SI}_c \cdot 0.5, \quad (1)$$

for which W and L were taken at each time-lapse interval, D (m) is the ~~channel-valley~~ depth that was interpolated between ~~measurements at SfM measurements for~~ each time-lapse interval. SI_c is the shape-index of the valley cross-section, which is the average ratio of the actual valley cross section to the square cross-section of $W \times D$. The factor 0.5 corrects for the longitudinal profile of
260 the valley, which is in all cases ~~an approximate triangle~~ approximately triangular. Valley volume was transformed ~~to~~ into an erosion rate E (g s^{-1}):

$$E = \frac{\Delta V \rho_s (1 - n) \cdot 10^{-3}}{\Delta T}, \quad (2)$$

where ΔV is the change in volume, ρ_s is the density of sand (2300 kg m^{-3}), n the porosity of the sediment (0.3) and ΔT (s) is the time over which the change in volume occurred. Cumulative
265 erosion was compared to sediment output measurements ~~using buckets~~ collected from bucket traps.

2.4 Martian landscape metrics

We constructed elevation profiles and extracted the orientation in degrees from North (i.e. azimuth) of valley segments, the ~~bifurcation angles~~ angles between converging valleys and valley lengths of Louros Valles and Nirgal Vallis. ~~Channel segments were digitized based on~~ Elevation profiles were

270 ~~extracted from HRSC image H0380_0001 (125 m resolution DEM) for Louros Valles, valleys of~~
~~Nirgal Vallis were too small to produce elevation profiles. Valley segments for both systems were~~
~~digitized from~~ THEMIS Daytime Infrared Mosaic (Ferguson et al., 2013) and HRSC (Jaumann et al.,
2007) imagery. We distinguished different stream-orders, based on the Hack stream ordering number
(Hack, 1957). In this system, the first-order is the main, downstream, valley; the first tributary is
275 the second order and so on. We choose this system since it represents the chronology of valley
formation by headward erosion. ~~Elevation profiles were extracted from HRSC image H0380_0001~~
~~(125 resolution DEM) for Louros Valles, valleys of Nirgal Vallis were too small to produce useful~~
~~profiles.~~

The dataset of ~~channel-valley~~ segments was transformed to a network-topology to distinguish
280 ~~between~~ upstream and downstream ~~direction~~directions, using logical operators based on the meth-
ods described in Marra et al. (2014d) using ArcGIS and MATLAB. In this dataset, we identi-
fied ~~bifureations~~converging valley segments, valley heads and outlets based on the ~~topology of the~~
~~network.~~ ~~From this dataset~~network topology. Building upon the work of Jaumann and Reiss (2002),
the orientation relative to north of each valley segment was extracted for each stream order (~~building~~
285 ~~upon the work of Jaumann and Reiss, 2002~~)identified in the dataset. Orientation distributions were
normalized per stream order to ~~better show the difference and reveal the effect of bifureations of~~
~~valley orientation~~clearly show the differences between valley orientations of different stream-orders.
At each ~~bifureation~~node of converging valleys, we calculated the ~~bifureation~~angle between the up-
stream ~~channel-valley~~ segments (following Glines and Fassett, 2013). ~~Furthermore, for each valley~~
290 ~~head in the network, we calculated the distance to the first lower-order valley segment.~~

~~What we here refer to as bifureation is a valley that splits in upstream direction. This~~ ~~Such~~
~~converging valleys is referred to as headward bifurcation, this~~ definition relates to the chronological
order of events in valley formation. In active rivers, the term bifurcation is used for a ~~channel that~~
~~splits into two channels in~~ fluvial channel that splits into two in the downstream direction, which re-
295 lates to the ~~motion~~direction of water ~~movement.~~ ~~Furthermore, for each valley head in the network,~~
~~we calculated the distance to the first lower-order valley segment.~~

3 Experimental results

In the following section, we first describe the observed morphological development during the ex-
periments, and then we link this morphological development to the acting processes. Time-lapse
300 imagery and elevation models support these observations (time-lapse movies are available in the
~~Supplement~~online supplement).

3.1 ~~Distal groundwater~~ ~~Distant~~ source

~~The distal groundwater experiment~~ The experiment with seepage from a distant constant-head tank

was characterized by slowly developing valleys. This experiment took over three days to complete
305 and was carried out with a constant discharge of 2.4 liter per minute (Table 1).

The sediment saturated in the first hours of the experiment. During this stage, a visible wetting
front at the surface progressed from the upstream constant head tank in downstream direction. The
sediment became fully saturated at the foot of the slope where seepage occurred after 2.5 h over
the full width of the sediment surface (Fig. 4a-i). The initial seepage pattern remained roughly the
310 same, though the seepage area extended upslope to about 1 m from the foot of the slope. Initially,
the seepage was too low for fluvial transport to occur. As the seepage rate increased, fluvial transport
started after 4 h and the first ~~valleys~~ channels started to form at the foot of the slope within the
seepage area.

The initial ~~valleys that formed~~ channels at the foot of the slope featured a combination of mass-
315 wasting and fluvial processes. Mass-wasting of saturated sediment at the head caused headward
erosion, and fluvial processes ~~removed the material from the valley in the channel~~ resulted in incision
and the formation of valleys (Fig. 4a-iii). As the valleys developed in upstream direction, the seepage
area retreated and seepage focused within the valleys as shown by drying of the sediment between the
valleys and a concurrent increase of discharge within the valleys (Fig. 4a-iii). Seepage was limited to
320 a declining number of valleys, as the valleys that reached ~~furthermost~~ most upstream progressively
attracted more groundwater. From the ten valleys that started to form in the initial stage of the
experiment, only six remained active after a few hours (Fig. 4a-iv), and only three remained active
for several days (Fig. 4a-v).

The decreasing number of ~~active valleys testifies to the presence of groundwater piracy, which~~
325 ~~is typical for the process of groundwater sapping. This groundwater piracy actively developing~~
valleys illustrates the process of groundwater flow piracy as the largest valleys attract most of the
groundwater flow since these are the deepest point in the landscape. As a result, more groundwater is
directed to those largest valleys, which are therefore more active and smaller valleys cease developing.
This feedback resulted in a final morphology ~~comprising with~~ a few large and several small valleys
330 (Fig. 5). ~~The valleys did not bifurcate, which is the result of the high initial slope (Pornprommin et al., 2010).~~

The remaining valleys grew and as they became deeper, the head- and side-walls gained strength
by cohesion as the sediment was moist. As a result, the head-wall retreat was governed by collapse
due to undercutting at the toe, in contrast to the slumping of the entire valley head before the devel-
335 opment of this cohesive top layer. In this process, the toe of the head wall was destabilized by fluvial
erosion, resulting in collapse of the head-wall. The collapsed material ~~transformed into a mudflow.~~
~~Therefore, the collapsed material~~ spread over a distance of 0.1 to 0.2 m into the valleys. Fluvial trans-
port removed this material from the upstream end to the downstream end of the experimental setup.
These processes showed a cyclic behavior: head collapse only occurred after a destabilization of the

340 head due to the removal of the sediment by fluvial transport. This cycle is essential for the continuation of the process as the ~~channel-valley~~ head would stabilize without such erosion and sediment transport. The final morphology shows the former presence of these various processes. In the three most developed valleys, the upstream end had a steeper slope than the downstream sections with a ~~clear knick-point break in slope~~ separating the two (Fig. 6a). This ~~corresponded to mudflow and~~
345 ~~fluvial flow dominating in these areas, respectively~~change in slope is the result from the transition of mass-wasting processes upstream to fluvial processes downstream.

The collapse of unsaturated material at the valley heads resulted in steep head walls (Fig. 5). The step-wise increase of valley width and length shows ~~this cyclic behavior~~distinct peaks of collapse (Fig. 7a and b) and ~~results in peaks in the estimated instantaneous~~ erosion rates (Fig. 7d). Steps in
350 width and depth of the valleys are not simultaneous, which shows that collapse of the head- and side-walls occur at different moments. Although erosion takes place in distinct peaks of activity, ~~the distally-fed~~these valleys show a linear trend in valley length and width during most part of their development. Not only the length and width ratio of a single valley remain the same, but this ratio is the same for the three main valleys (Fig. 8a).

355 3.2 ~~Effect of initial morphology on Distal experiment~~Local precipitation

~~We studied the effect of an initial morphology on the valley development in experiment Distal-after-Local by repeating the experiment on an initial morphology. This initial morphology was the result of experiment Local-2, which consists of multiple parallel shallow valleys created by overland flow (Fig. 9).~~

360 ~~Experiment Distal-after-Local showed the same general characteristics and development as the Distal experiment. The main difference is valley initiation processes. We consider other differences as natural variations, which could have occurred without the altered initial morphology as well. In the Distal-after-Local experiment, initial seepage at the downstream end focused within the valleys of the initial morphology. However, due to groundwater flow piracy, only a limited number of these valleys fully developed. Six valleys started to form, but only two valleys fully developed (Fig. 5e). Development in the two remaining valleys was the same as described above. In the early stages of valley development, the valleys followed the path of the existing valleys in the initial morphology. When they became larger, they still followed the path of the initial valleys, although these were straight at this stage and the new valleys were much wider than the initial valleys. In our view, these~~
365 ~~mature valleys seemed to develop independently from the initial morphology.~~

370 3.3 ~~Local groundwater source~~

~~The local groundwater experiment took one hour and~~The *Local precipitation* experiment took one hour and 50 min with an average discharge of
min with an average discharge of 11.9 L min⁻¹. This discharge is higher than in the *Distal source* experiment. A part of the precipitation fell directly into

375 the valleys since the precipitation was distributed evenly over the experimental domain. Furthermore,
the groundwater table in the *Local precipitation* experiment was close to the surface compared to a
relative deep groundwater table in the *Distal source* experiment. As a result this setup allowed for
more seepage due to the higher seepage area, explaining the higher discharge and shorter run of this
experiment. In this experiment we distinguished two stages in valley development. In roughly the
380 first half of the experiment, overland flow was the main source of water for valley development. In
the second half, groundwater fed the valleys at their heads.

In the first stage of the experiment, the sediment in the downstream part of the slope saturated
rapidly due to the limited sediment thickness (Fig. 4b-i). On this saturated sediment, precipitation
transformed directly into runoff, which in turn induced valley formation by fluvial processes. These
385 valleys started to form over the entire width of the sediment and showed valley heads with a v-
shaped planform (Fig. 4b-ii). During this stage, valleys developed in headward direction by fluvial
erosion and valley heads were within the area of saturated sediment. Seepage inherently occurred in
the valleys due to the setup of the experiment. However, the overland flow processes dominated the
seepage processes.

390 As the groundwater table rose during the experiment, the boundary of saturated sediment moved
upslope. This progression of the saturated area slowed down as it progressed. In the first stage, valley
development did not keep up with this moving front. However, the valleys caught up and developed
upstream of the saturated area as the experiment progressed (Fig. 4b-iii). This marks the second stage
in valley development wherein groundwater seepage rather than surface runoff fed the valley heads.
395 From the moment the valley heads were fed by groundwater, their planform changed from v-shaped
to theater-shaped (Fig. 4b-iv and b-v). This change indicates a change from fluvial flow to mass-
wasting processes at the ~~head-wall~~headwall. The headward growth showed similar characteristics as
in the ~~distal-seepage-experiment~~experiment with seepage from a distant source: growth governed
by failure of the headwall and fluvial transport that removed the failed material. Similar to the ~~distal~~
400 *Distal source* experiments, there was also a distinguishable difference in slope in the upstream and
downstream half of the valleys (Fig. 6), although this difference was less pronounced.

The valleys in the ~~local~~*Local precipitation* experiment were shallow compared to the valleys in
the ~~distal~~*Distal source* experiment. In both cases, the valleys developed around the groundwater
table, which was close to the surface in the ~~local~~*Local precipitation* experiments. The limited depth
405 was presumably the result of the high groundwater table, there was no zone of unsaturated sediment
resulting in valleys without steep walls (Figs. 5b and 6a).

The valleys in the ~~local~~*Local precipitation* experiment became longer and slightly wider by lateral
erosion during the experiment (Fig. 7f and g). An important difference with the ~~distal~~*Distal source*
case was that all valleys remained active and had similar sizes during the experiment (Fig. 7i).
410 This is due to the absence of groundwater flow piracy since each ~~channel~~valley was fed by locally
infiltrated groundwater (Fig. 5). In contrast to the ~~distally-fed-valleys~~valleys from a distant, the

relation between valley length and width in the locally-fed valleys is not linear and different valleys do not have the same ratio (Fig. 8b).

3.3 Effect of initial morphology on ~~Local experiment~~seepage from a distant source

415 We studied the effect of an initial morphology on the valley development in experiment *Distant after local* by repeating the experiment on an initial morphology. This initial morphology was the result of experiment *Local precipitation 2*, which consists of multiple parallel shallow valleys created by overland flow (Fig. 9).

~~In the Local-after-Distal experiment,~~ Experiment *Distant after local* showed the same general
420 characteristics and development as the *Distant source* experiment. The main difference is valley initiation processes. We consider other differences as natural variations, which could have occurred without the altered initial morphology. In the *Distant after local* experiment, initial seepage at the downstream end focused within the valleys of the initial morphology. However, due to groundwater flow piracy, only a limited number of these valleys fully developed. Six valleys started to form,
425 but only two valleys fully developed (Fig. 5c). Development in the two remaining valleys was the same as described for the distant experiment. In the early stages of valley development, the valleys followed the path of the existing valleys in the initial morphology. When they became larger, they still followed the path of the initial valleys, although these were straight at this stage and the new valleys were much wider than the initial valleys. In our view, these mature valleys seemed to develop
430 independently from the initial morphology.

3.4 Effect of initial morphology on local precipitation experiment

In the *Local after distant* experiment, we studied the effect of the initial morphology in the opposite way as in the ~~Distal-after-Local~~*Distant after local* experiment; the final morphology of ~~Distal~~the *Distant source* experiment acted as the initial morphology of this run (Fig. 5a). The same processes acted in this experiment, though the initial morphology had a much larger effect on the final morphology in this case.
435

In the first stage of the ~~Local-after-Distal~~*Local after distant* experiment, the present valleys reactivated as the sediment saturated. Due to the rising groundwater level, the steep side- and headwalls of the previous valleys became unstable and collapsed. This resulted in a decreasing valley depth and increasing width. The valleys that were abandoned in the ~~Distal~~*Distant source* experiment due to groundwater flow piracy reactivated as they were fed by local precipitation and subsequently infiltrated groundwater, resulting in a smaller difference in valley size (Fig. 5d), compared to the initial situation (Fig. 5a). Collapse of the head-wall caused headward erosion and lateral erosion caused widening of the valleys. These are the same processes as in the ~~Local-run~~*Local precipitation*
445 experiment with no initial morphology, but the final morphology showed much wider valleys. The

initially present valleys were relatively deep in comparison to the final valleys. The reduction in depth of these valleys corresponded with this widening.

At the end of the ~~Local-after-Distal~~ experiment, a body of ponding water at the flat area upstream of the valleys was released in the central valley. This resulted in the development of a feather-shaped drainage network upstream of the valley head. ~~Local after distant~~ experiment, water ponded at the upstream flat section of the experimental setup. This ponding seemed to be the result of the sediment becoming fully saturated towards the end of the experiment. The headward developing valleys tapped into this shallow reservoir, resulting in a final slightly catastrophic stage of erosion due to the breach of this reservoir (Fig. 5d). ~~We do not further consider this event in this paper, but a similar morphological feature develops after the overflow of a shallow lake as result from pressurized groundwater outflow, which is discussed in Marra et al. (2014e)~~ This stage is not representative for the main objectives of this paper and therefore not further considered here.

4 Examples of Martian valley systems

In this section, we show the morphology of Louros Valles and Nirgal Vallis (Figs. 10 and 11), two Martian valley systems that were previously attributed to a groundwater seepage origin (e.g. Jaumann and Reiss, 2002; Harrison and Grimm, 2005; Glines and Fassett, 2013). ~~These two system serve as an example on how to apply our experimental results and have received much attention in recent literature. Furthermore, these systems show branching valleys which also aids the interpretation of these systems. In this section, we describe Louros and Nirgal Valles, their interpretation is part of the discussion.~~

4.1 Louros Valles

Louros Valles are located at the north and south flanks of Valles Marineris. These valleys have circular valley heads cutting into flat plains. The valleys have a total relief of several kilometers and are between 10 and 100 km long (Fig. 10b and c). Upstream of the valley heads, there are no visible tributaries or depressions in the elevation data or imagery. Downstream of the valleys, in Valles Marineris, there are no clear deposits associated with these valleys. Sediment output in ~~the experiments was constantly removed, in the Martian case slow could have removed the produced sediment. Alternatively, deposits could this case could~~ be spread over a large area on the floor of Ius Chasma as a thin veneer ~~layer~~ and not recognizable as fluvial deposits. The valleys on the northern flank are shorter than the valleys at the south. Of the southern valleys, there are two larger valleys in the west; all other valleys are approximately equal in size. The valleys are closely spaced and several valleys touch or intersect, resulting in a relatively densely dissected plain (Fig. 10c and d).

As an example, we show an elevation profile of the largest valley of Louros Valles (Fig. 10e). These and other elevation data show a rough profile with many bumps, likely related to post-valley

480 formation wall collapse or tectonism. At the ~~distal~~-downstream end of the valley, the elevation quickly drops, which shows the onset of Valles Marineris. Based on a 10 km moving average, we show the valley has a change in slope about halfway to a lower slope than the upstream part (Fig. 10e). However, the irregular elevation data limits the interpretation of these observations.

All valleys show ~~headward-bifurcations~~headward bifurcations. For the northern valleys, the orientation of the first-order valleys varies from northwest to northeast with most valleys oriented to the north-northeast (Fig. 12). The first tributaries, or second order (Hack, 1957) valleys have a similar spread in orientation, but most valleys are oriented to the west-northwest. For the southern valleys, most first-order valleys are oriented towards the southwest while higher-order valleys are directed towards the south-southwest or towards the west-southwest (Fig. 12b). Interestingly, a few third- and fourth-order segments are oriented in opposite direction (180°) as the first-order valleys (e.g. Fig. 10d).

The lengths of the tributaries range between 5 and 15 km (the main valleys are longer, but most are outside the graph), with no specific trend in the distribution of valley length (Fig. 12d). Mean heaward bifurcation angles of the different stream-orders are between 70 and 90° (Fig. 12f).

495 4.2 Nirgal Vallis

Nirgal Vallis consists of one more than 500 km long main valley and several sparsely distributed side valleys of various sizes (Fig. 11). Valley depths range from several tens to several hundreds of meters. The valley cuts into the plateau through several north-to-south oriented wrinkle ridges, which are in places the highest points in the landscape. Several side-valleys align with these wrinkle ridges (Glines and Fassett, 2013).

The orientation of the main valley is dominantly west-northwest, the first-order tributaries have the same dominant orientation, but a large part is deflected north- and southward (Fig. 12c). This tendency of dominantly westward-oriented ~~channels~~-valleys is shown in the landscape (Figs. 11d and 12c). There a few larger side valleys, but most side-valleys are very short (Figs. 11d and 12e). This results in the sparsely dissected landscape. The mean heaward bifurcation angle between valleys is 70.7° with a standard deviation of 18.6 (Fig. 12g), similar but slightly less than the results of Glines and Fassett (2013) for the same valley network but with less measured junctions. ~~Bifurcation~~ Headward bifurcation angles are similar for different stream orders.

5 Discussion

510 5.1 Valley morphology related to groundwater source~~character~~

~~In this paper, we aim to find morphological properties related to different sources of groundwater in the formation of valleys by groundwater seepage. The general morphology of our experimental valleys agrees well with previous studies on sapping valleys (e.g. Howard and McLane, 1988; Hagerty, 1991; Dunne, 1980; Fox et al~~

~~These valleys consist of~~ The valleys in experiments with a distant source of groundwater have half-
515 round, theater-shaped ~~;~~ valley heads with a sharp transition to the upstream, ~~uneroded surface. They~~
~~develop in un-eroded surface.~~ These are similar to those found in previous studies on valleys formed
by groundwater seepage (e.g. Howard and McLane, 1988; Hagerty, 1991; Dunne, 1980; Fox et al., 2006; Pornprommin et al., 2010)
Valleys in our experiment fed from nearby infiltrated precipitation also featured round valley-heads
but lacked the steep theater-shaped head wall. The valleys from both boundary conditions developed
520 in a headward direction by destabilization of the ~~channel head, which results from valley head due~~
~~to either~~ undercutting or slumping. The eroded material is transported ~~though along~~ the valley by
fluvial processes. ~~These two main processes showed a cyclic behavior as the fluvial erosion in the~~
~~valley was the trigger for collapse at the valley head.~~ Furthermore, in both experiments, the slope
in the upstream section of the valley floor ~~is was~~ steeper than in the downstream ~~end valley floor~~
525 (Fig. 6), which ~~results from the transition of hyper-concentrated flow with high sediment availability~~
~~at the channel heads to regular fluvial flow further relates to the transition from mass-wasted material~~
~~released at the valley head to the fluvial transport of material~~ downstream.

The morphological similarity between theater-headed valleys in groundwater sapping features
(~~undercutting and failure by groundwater seepage erosion~~) at the beach (Higgins, 1982), in sand-
530 box experiments (Howard and McLane, 1988), ~~on~~ the Colorado plateau (Laity and Malin, 1985)
and ~~valleys~~ on Mars is often used as an argument for a groundwater origin of the Martian valleys
(e.g. Schumm and Phillips, 1986; Mangold and Ansan, 2006; Harrison et al., 2013). A compli-
cation in the study of such morphology on Mars is that different processes yield a similar mor-
phology, ~~for example.~~ ~~For example,~~ waterfall erosion (Lamb et al., 2006) or groundwater weather-
535 ing (Pelletier and Baker, 2011) ~~can also produce theatre-headed valleys.~~ We do not solve ~~that this~~
~~controversy~~ in this paper since the experiments here do not explore the morphological ~~difference~~
~~between different differences between all these possible~~ processes. Here, we focus on ~~groundwater~~
~~processes morphology related to groundwater flow processes and subsequent erosion~~ in further detail
~~and provide metrics of entire landscapes to aid the interpretation of Martian landscapes.~~

540 In the following discussion, we ~~start by considering the applicability of our experiments. Then, we~~
propose different end-member landscapes based on knowledge from our experiments combined with
previous experimental ~~and work,~~ modeling results and theoretical considerations. The main land-
scape properties are the distribution of valley lengths, valley order, valley ~~spacing and bifurcations.~~
~~We interpret the orientation and the angle between valley segments. We close the discussion with an~~
545 ~~interpretation of the~~ Martian valleys described ~~above in this framework and end with considerations~~
~~on the applicability of scaled experiments of such systems, using the proposed landscape metrics~~
~~framework.~~

5.2 Scalability of experimental results

The experiments described in this paper are not dimensionally scaled or direct analogues to the
550 ~~Martian case studies.~~ Instead, the experiments provide insights into the fundamental processes that
~~result from groundwater seepage and the resultant morphology.~~ These experiments were devised to
~~contrast distinct sources of groundwater and complement previous work.~~ The experimental setup
~~was designed to be simple in order to show clearly the effect of different hydrological boundary~~
~~conditions.~~ Different initial conditions will produce different landscapes, but again, our work is
555 ~~focused on the essential underlying processes and representative morphological features.~~

The experiments presented in this paper, and previous work on seepage erosion, applies to landscapes
formed by groundwater and thus landscapes that form in porous and erodible material. The overall
patterns are expected to be similar on different scales and for different materials that meet these
conditions, but details will differ. Our analyses are therefore limited to the large-scale patterns in
560 the landscapes and not expected to explain details. Below we point out the scale-effects in our
experiments and how we take these into account.

An important difference between the ~~distal and local groundwater~~ *Distal source* and *Local*
precipitation experiments was the steepness of the valley heads and sidewalls, which were much
steeper in the ~~distal groundwater~~ *Distal source* case. In the ~~Distal~~ *Distal source* experiment, the
565 groundwater table was deeper, resulting in an unsaturated (moist) top layer above the groundwater
table, which has more apparent cohesion than the saturated (wet) top layer in the ~~locally infiltrated~~
~~groundwater case~~ *Local precipitation* experiment. In natural systems, such ~~properties show contrasts~~
~~in material strength arise from~~ differences in soil ~~or substrate~~ properties rather than formative pro-
cesses. The depth of the unsaturated layer relates to capillary forces, which are scale-independent
570 and thus relatively small for large experimental valleys (see discussion in Marra et al., 2014a). Nev-
ertheless, theater head formation took place in both cases with and without an unsaturated top-layer,
which indicates that this process takes place under both conditions and is not the result of this scale-
effect.

Destabilization of the headwall is a necessary condition for the development of ~~sapping valleys~~
575 ~~valleys by seepage~~. This only takes place if sufficient sediment is removed from the toe of the head-
wall, which requires enough discharge for sediment ~~transport~~ *erosion and transport along the entire*
valley length. In previous ~~smaller sapping experiments~~ (Marra et al., 2014a), ~~there were problems~~
~~with sustaining the process without artificial removal of sediment.~~ In our current experiment, the
~~discharge was sufficient to allow continuous downstream sediment transport.~~ Marra et al. (2014a) ~~smaller-scale~~
580 ~~experiments,~~ (Marra et al., 2014a), report on experiments in a 1 × 3 m ~~setup flume~~ with sand where
the valleys clogged due to the absence of downstream sediment removal (~~these are not the main~~
~~experiments reported in that paper, they are mentioned in the discussion~~). Their solution to ~~keep~~
~~the sustain~~ upstream processes and valley formation ~~going~~ was to flush away sediment at the down-
stream end. In that same setup, ~~sapping valleys~~ *valleys from groundwater seepage* did develop when
585 lightweight plastic sediment was used, ~~resulting in which enabled~~ sufficient sediment transport ~~due~~

to the lower material density. In other words, sufficient downstream erosion by fluvial processes is essential to keep the cyclic formation characteristic for seepage valleys going. Similarly, the larger experiment by Howard and McLane (1988) featured a downstream gutter to remove the sediment and maintain the process of sediment evacuation from the growing valley. In the experiments described in this study paper, sediment was not flushed at the downstream end and the downstream elevation is kept constant by a row of bricks. In other words, the processes that occurred were not forced by the removal of downstream sediment, which shows that the scale-effect of having a too low insufficient discharge for sediment erosion and transport was overcome in our setup.

5.3 Groundwater piracy, valley spacing and length distribution

Additional work is required to understand the morphological details of valleys formed by groundwater seepage. In particular, we expect important effects on valley shapes to result from layered substrates with alternations in material erodibility. These effects can be studied experimentally, but to model erosion rates on larger scales than can be represented in the laboratory, numerical modelling will be more informative about the formative time-scales of such systems and may elucidate on terraces found in Martian valley systems. Furthermore, using such models, Martian scenarios with a thick layer of permafrost can be simulated which are unpractical to recreate in most laboratories.

5.3 Groundwater flow piracy, valley spacing and length distribution

The morphology of the entire landscape shows important differences that are related to subsurface groundwater flow processes. In the distally-fed *Distant source* experiment, a decreasing number of channels-valleys remained active when small-smaller valleys ceased to develop. This behavior is related to groundwater piracy of relates to groundwater flow piracy by the larger valleys and results in an increase of valley spacing. The valleys are depressions and therefore, since these depressions attract groundwater. Due to the travel distance and direction of the groundwater, areas downstream of large valleys receive less groundwater or even no groundwater as shown by surfacing drying. The resulting landscape consists of a few large and several small valleys in between, which ceased developing due to this cutoff of groundwater several small inactive valleys with no groundwater supply (Fig. 5a), and valleys are mainly in between a few large active valleys oriented towards the groundwater source (Fig. 13a). In contrast, the locally-fed channels-*Local precipitation* experiments did not feature groundwater flow piracy, which results in since the groundwater source is distributed everywhere and therefore cannot be captured by nearby valleys. The resulting landscape is densely dissected landscape with valleys of similar size in close proximity of to each other (Figs. 5b and 13b). The close proximity is a strong indication that the availability of water was not limiting and piracy can therefore not have been important.

An important parameter for groundwater flow piracy is the fraction of the groundwater flow that a valley captures. This is controlled by the fraction-ratio of cross-stream to downstream groundwater

flux (Pelletier, 2003; Schorghofer et al., 2004), which is proportional to the groundwater gradient in isotropic conditions. In the case of ~~sapping valleys~~ valleys formed by groundwater seepage, the emerging valley itself leads to a topographic low that introduces a cross-stream groundwater slope, which increases the flow towards that valley. This morphological feedback causes flow piracy when
625 a valley attracts enough flow to cease the development of another valley. This feedback and tendency for flow piracy, is stronger for ~~lower regional slopes as a perturbation~~ flat surfaces in contrast to valley formation on a slope, since a depression in a flat surface has a ~~relative large~~ larger effect on the convergence of groundwater flow (Pornprommin et al., 2010).

Our experiments show that the valley width-to-length ratio is similar for valleys formed by a ~~distal~~
630 distant source of groundwater (Fig. 8a), ~~which was but this is~~ not the case for ~~the locally-fed valleys~~ valleys generated by a local groundwater source (Fig. 8a). ~~Such~~ The similarity in the development of several distally-fed valleys is indicative for valley formation by the same source of groundwater. The size of the ~~valleys dominantly control~~ valley is the dominant control on the amount of water delivered to that valley since a larger ~~valley source yields more~~ and deeper valley yields more groundwater
635 seepage. In turn, the amount of erosion relates to the size of the valley and hence, the morphological development is similar for the different valleys. The amount of water delivered to the ~~locally-fed~~ channels ~~valleys fed by local precipitation~~ is only partly controlled by this mechanism. In this case, the amount of groundwater delivered to the valley head also depends on upstream area and local watersheds.

Initial conditions may affect the location ~~and thus of valley initiation and thereby the~~ spacing of
640 valleys in the final landscape. In our ~~distally-fed~~ Distant after local experiment with minor initial morphology (~~Distal-after-Local~~) and a distant groundwater source, the initiation of ~~sapping valleys~~ seemed ~~valleys appeared to be~~ related to the initial perturbation of the surface, but the resulting processes and the final landscape was similar to the experiment with no initial perturbations. This
645 shows that seepage is robustly driven by the subsurface flow pattern and agrees with the observations of Schumm and Phillips (1986) ~~÷that~~ valleys of a composite origin dominantly reflect the last process. In contrast, in the ~~locally-fed experiment with initial perturbations~~ (~~Local-after-Distal~~) Local after distant experiment with, there was a ~~major effect~~ significant effect of the initial morphology as old valleys reactivated ~~and dominated the final morphology~~.

~~The strong difference of local effects is the result of the groundwater flow patterns : in the locally-fed experiments, local perturbations affect the flow pattern, while groundwater from a distal source is driven dominantly by the large-scale morphology. Furthermore, flow piracy dampens the development of small perturbations. An important.~~ Importantly, the valley patterns in the Distant after local experiment are similar as in the Distant source experiment, an hardly influenced by the
650 initial morphology. An implication is that the location and orientation of ~~distally-fed sapping valleys~~ strongly ~~valleys fed from a distant source~~ relate to the hydrological system and ~~can thus be used as an indicator of~~ thereby provide an indicator for the possible source of groundwater.

5.4 Headward ~~channel~~ bifurcations

Our experimental valleys did not bifurcate at their ~~channel-valley~~ head, which is considered a typical
660 property for ~~sapping-valleys~~ valleys fed by groundwater seepage (e.g. Howard and McLane, 1988).
The absence of headward bifurcations in our experiments is the result of the steep slope in the
downstream half of the setup. Pornprommin et al. (2010) showed that ~~a low regional slope, and~~
~~thus lower morphological feedback, resulted~~ for valleys that formed in a flat surface, result
665 when growing in a headward direction. Our results show that seepage on a ~~relatively~~-steep slope
suppresses the tendency to form such headward bifurcations, compared with similar experiments
~~with a horizontal surface~~ on horizontal surfaces, which do show headward bifurcations (Fig. 13a and
c; e.g. Berhanu et al., 2012).

Besides the ~~regional~~-initial slope, Berhanu et al. (2012) showed that valleys fed from a local source
670 have a higher tendency to bifurcate in a headward direction. This tendency relates to the groundwater
flow that enters the ~~channel-valley~~ head from a wide range of directions and not mainly from up-
stream ~~of the valley. The typical flow field that results from local infiltration in a flat substrate results~~
~~in bifurcations with a typical angle of 72~~(Devauchelle et al., 2012). ~~In such system, the valleys would~~
~~have many or the direction of the groundwater source. As a result, valleys formed by seepage from~~
675 a local source on a flat topography have many headward bifurcations which results in a densely dis-
sected landscape (Fig. 13d). ~~This pattern is~~ similar to the Apalachicola Bluffs (Florida) which ~~were~~
have been shown to be formed by seepage of locally infiltrated precipitation (Abrams et al., 2009).
~~The valleys in this system could be subject to flow piracy, but as a result by other valleys cutting~~
~~each other off. This mechanism of piracy is different from the piracy by groundwater flow capture,~~
680 ~~which acts on a larger distance and is typical for distally-fed valleys-~~

The flow field that results from local infiltration into a flat substrate results in headward bifurcations
angles of 72° (Devauchelle et al., 2012). This value has been considered as evidence for a groundwater
origin of Nirgal Vallis by Glines and Fassett (2013), but this value is only characteristic for seepage
from uniform precipitation on a flat surface and is therefore not universally applicable. Furthermore,
685 structural controls from tectonics may also dictate the angles between valleys (Luo et al., 1997),
which is also likely the case for Nirgal Vallis (Glines and Fassett, 2013).

The combined occurrence of developing headward bifurcations and groundwater flow piracy re-
sults in the formation of typically stubby tributaries. When a valley bifurcates in headward direction,
there are two ~~channel-valley~~ heads close to each other, which will in ~~many cases~~-result in abandon-
690 ment of one of the two (Fig. 13c) due to the presence of groundwater flow piracy. This behavior and
the resulting morphology is indicative of ~~sapping-valley formation by seepage~~ from a ~~distal~~-distal
groundwater source.

5.5 Origin of Martian examples

~~In this paper, we~~ We analyzed the landscape of Louros and Nirgal Vallis, two valley systems often
695 attributed to groundwater seepage. Two additional examples are shown in Fig. 13a and b and these
examples are described below to serve as an illustration of how the morphology can appear, but are
not analyzed further. The valleys in Noctis Labyrinthus (Fig. 13a) show valleys with no headward
bifurcations and a few small valleys in between larger valleys. Although the plateau where these
valleys cut into is flat, there is a strong gradient between valley head and outflow point, which is
700 ~~what we mean with~~ illustrates our contention that a steep slope ~~and which is what~~ suppresses the
tendency ~~to bifurcate~~ for headward bifurcation to form. The small valleys in Gale Crater (Fig. 13b)
are similar in size, shallow and ~~the~~ there is a regional slope. ~~These examples serve as an illustration~~
~~on how the morphology can appear, but are not further analyzed here.~~

Alternate hypotheses for similar valley ~~formation~~ formations are bedrock erosion by catastrophic
705 release of surface water (Lamb et al., 2006) or bedrock weathering ~~rather and erosion~~ by groundwater
(Pelletier and Baker, 2011). ~~The latter would yield a similar morphology as sapping in unconsolidated~~
~~material, and the two processes would likely act~~, which would result in the same ~~systems as~~
~~the weathered bedrock material is further removed by seeping groundwater. However, weathering~~
~~in bedrock could result in steep slopes, which are not likely for large valleys in case of seepage~~
710 ~~erosion in unconsolidated material (Marra et al., 2014a)~~ combination of boundary conditions and morphology
as seepage in unconsolidated materials. Another hypothesis for the formation of Louros Valles is
focused erosion by meltwater in between patches protected from erosion by the presence of an
ice-cover (“glacial selective linear erosion”, Lee, 2000).

The evidence in favor of a groundwater origin ~~on both cases~~ over surface flow in both Nirgal
715 and Louros, is the absence of upstream feeder channels ~~to into~~ the main valleys. However, billions
of years of weathering and a dust cover could have obscured such small channels. The elevation
profile of one of the Louros Valles is bumpy (Fig. 10e), likely due to later activity. ~~Based on filtered~~
~~elevation profile, the valley shows a lower slope downstream, similar to the experiments, but the~~
~~implication of this is limited due to the bumpiness of the profile~~ Consequently, the interpretation of
720 such profile in comparison to elevation profiles of the experiments (Fig. 6) is limited. Therefore,
we use the properties of multiple valleys and valley segments in the entire landscape rather than
single-valley morphology for our interpretation.

The orientation of valley segments of Louros Valles is diverse ~~with wide bifurcation angles and~~
and has a broad range of valley lengths, resulting in a densely dissected landscape (Fig. 10). Such a
725 landscape is typical for a local groundwater source (Fig. 13d). ~~The bifurcation angles have a much~~
~~broader distribution than the theoretical value for such system. This could be the effect of valleys~~
~~aligning with the tectonic structure, which would result in 90 angles (Luo et al., 1997).~~ Furthermore,
in Louros Valles, some higher-order valley segments are oriented in ~~opposite direction~~ downstream
direction with respect of the main valleys (Fig. 10d), which can ~~only be explained by~~ be the result

730 ~~of~~ a local source of groundwater and not ~~by surface flow following the topography~~ groundwater
coming from greater distances. Additionally, the presence of valleys on both sides of Valles Marineris
suggests a local groundwater source and not groundwater coming from a great distance.

A possible local source of groundwater for Louros Valles is precipitation; melt of snow, ice or
permafrost; or upwelling groundwater from a cryosphere-confined aquifer. ~~Another source of water~~
735 ~~could be vertical upwelling groundwater from an aquifer. Such aquifer~~ (e.g. Clifford, 1993). The
presence of this type of aquifer in this region may ~~relate to the~~ also have been the source of water for
the outflow channels further northeast, ~~although the~~ which are likely to have formed by the release
of pressurized groundwater from a confined aquifer. The timing of events here is crucial. ~~The since~~
the presence of Valles Marineris, and the clear formation of Louros Valles after the opening of
740 Valles Marineris, suggest that this aquifer was at that point cut-off, ~~and in fact split between and~~
split between the north and south. Seepage at Louros Valles rather than the formation of an outflow
channel could represent a low aquifer pressure, which fits a trend of lower groundwater pressures
at higher elevations in Ophir and Lunae Plana (Marra et al., 2014b). ~~An alternate hypothesis for the~~
~~formation of Louros Valles is by focussed erosion of meltwater in between patches protected from~~
745 ~~erosion by the presence of an ice cover (“glacial selective linear erosion”, Lee, 2000) Furthermore,~~
the subsidence of Valles Marineris into the aquifer may have been a trigger for outflow or upwelling
of groundwater. This hypothesis could be further explored and the asymmetry between the valleys
on the North and South flank may provide additional insight into the nature of such aquifer.

~~Nirgal Vallis cuts through some ridges in the landscape where surface runoff would have flowed~~
750 ~~around. If this landscape is indeed formed by groundwater, the source is likely distal~~ The landscape
of Nirgal Vallis is an exemplar of valley formation by seepage from a distant source, given the
large number of small valleys typical for groundwater ~~piracy due to a distal source flow piracy~~
(Figs. 11, 5a, 13a and c). The groundwater source was likely to have been west of the valley due
to the orientation of most valleys towards that direction. ~~The average bifurcation angle is close to~~
755 ~~72, which is considered a typical angle for groundwater seepage valleys and mentioned as evidence~~
~~for a groundwater origin by Glines and Fassett (2013). However, this value is only valid for valleys~~
~~formed by a local source of groundwater (Devauchelle et al., 2011), which is not the case here, and~~
~~such value is heavily influenced by subsurface conditions like tectonic structures (e.g. Luo et al., 1997).~~

760 A possible source of groundwater ~~for Nirgal Vallis is groundwater~~ flow from the west, ~~which is~~
~~ultimately recharged from Tharsis~~ could be recharge in the Tharsis region (Harrison and Grimm, 2004).
Seepage of groundwater ~~could have taken likely took~~ place before the formation of a ~~confining~~
cryosphere. ~~Alternatively, if the cryosphere was already present, discontinuities or rupture could~~
~~have triggered outflow from a confined aquifer. However, in this case the groundwater should have~~
765 ~~flown laterally first to account for the morphology, which would be global~~ confining cryosphere,
which is considered a requirement for aquifer pressurization for the Martian outflow channels in

the Hesperian (Clifford, 1993; Marra et al., 2014c). Alternatively, a regional discontinuity may be the reason for seepage at Nirgal Vallis. In that case, seepage took place during an early stage in the case if the groundwater pressure was not sufficient to induce artesian seepage. Seepage would then occur further downstream at topographic depressions formation of the cryosphere. The climatic implications are the presence of precipitation in the source region which could be aqueous or icy (Wordsworth et al., 2013), but widespread precipitation is not required and a groundwater system as the dominant element of the hydrology shows that these valleys could form in the absence of a long-lived hydrological cycle at the surface.

Based on our and previous experiments, this study now provides a framework which links the landscape that links landscape properties to the groundwater source characteristics location. The two Martian examples shown, further illustrate this link. Although different processes could produce similar valley morphologies, the strong correspondence of the landscape metrics of these examples and those produced by sapping hints seepage points towards a groundwater origin. Particularly In particular the distant groundwater source of Nirgal Vallis implies a well-developed groundwater system. Perhaps most significantly, outflow and valley formation of such a system could have taken place regardless of climate conditions optimal for being optimal for the sustained presence of liquid water on the surface.

6 Conclusions

We studied groundwater seepage processes and subsequent valley formation using a series of large sandbox experiments. Our experiments focused on the difference between locally or distally fed valleys on a steep slope valleys fed from groundwater originating from a distant source or from infiltrated local precipitation. In both cases, valley heads developed in headward direction by mass-wasting processes triggered by steepening due to fluvial sediment transport out of the valley.

Combined with previous experimental work, we provide a framework of driving processes and resulting landscape metrics for sapping valleys fed by a distal and local source distant source and local precipitation, and for a steep and flat topography. Their main characteristics are as follows.

(1) Due to groundwater flow piracy, seepage erosion from a distant groundwater source results in a sparsely dissected landscape with a few large and many small valleys. Valleys fed by a local groundwater source from a local source of groundwater, e.g. precipitation, for example infiltrating precipitation or melt of ground ice, are not characterized by flow piracy and have a range sizes, resulting in a densely dissected landscape. (2) Valley formation in horizontal surfaces promote the development of headward bifurcations in contrast to steep surfaces where this tendency is suppressed. For distally fed sapping valleys fed by a distant source of groundwater, the combined occurrence of bifurcating valleys and flow piracy results in valley systems with stubby tributaries. Locally fed sapping valleys Valleys fed by locally infiltrated groundwater on horizontal surfaces grow in a wide range

of directions due to the ~~occurrence and prevalence of bifurcations~~ development of many headward bifurcations which remain morphologically active.

As an example, we applied these characteristics to two Martian systems. Firstly, Louros Valles
805 shows a densely dissected landscape with a broad range of valley orientations and valley lengths. This landscape is typical for a local groundwater source. Such local source could relate to an aquifer that fed the outflow channels, but is more likely related to local precipitation or melt of ice or snow. Secondly, Nirgal Vallis ~~whos~~ illustrates a sparsely dissected landscape with many small and only a few large valleys ~~that are oriented dominantly to the west~~ with a single dominant orientation. This
810 indicates a ~~distal~~ distant groundwater source in the west, which is likely produced from recharge at Tharsis. Further study of similar landscape properties as a result of overland flow is required to ~~further resolve~~ advance the ambiguous interpretation of these valleys.

Author contributions. Study conception: W. A. Marra, E. Hauber, M. G. Kleinans; Experimental design: W. A. Marra, S. J. McLelland, B. J. Murphy, D. R. Parsons, M. G. Kleinans; Analyses and interpretation:
815 W. A. Marra; Discussion of the results: W. A. Marra, E. Hauber and M. G. Kleinans with contribution from all other authors; W. A. Marra wrote the paper.

Acknowledgements. This work is part of the PhD research of W. A. Marra, financially supported by NWO grant ALW-GO-PL/10 01. The authors are grateful for the time-lapse camera triggering device build by Henk Markies, for the help by Tjalling de Haas, Steven de Jong, Leiping Ye, Chris Unsworth, Susan Conway and Rob
820 Thomas in running the experiments and Tjalling de Haas for reviewing a draft of this paper. Experimental data are available upon request from W. A. Marra (w.a.marra@uu.nl) or M. G. Kleinans (m.g.kleinans@uu.nl).

References

- Abrams, D. M., Lobkovsky, A. E., Petroff, A. P., Straub, K. M., McElroy, B., Mohrig, D. C., Kudrolli, A., and Rothman, D. H.: Growth laws for channel networks incised by groundwater flow, *Nat. Geosci.*, 2, 193–196, 825 doi:10.1038/ngeo432, 2009.
- Agisoft: Agisoft PhotoScan User Manual: Professional Edition, Version 1.0, available at: <http://www.agisoft.com/>, 2014.
- Berhanu, M., Petroff, A., Devauchelle, O., Kudrolli, A., and Rothman, D. H.: Shape and dynamics of seepage erosion in a horizontal granular bed, *Phys. Rev. E*, 86, 041304-1–041304-9, 830 doi:10.1103/PhysRevE.86.041304, 2012.
- [Clifford, S. M. \(1993\), A model for the hydrologic and climatic behavior of water on Mars, *J. Geophys. Res.*, 98\(E6\), 10,973, doi:10.1029/93JE00225.](#)
- Craddock, R. A., Howard, A. D., Irwin, R. P., Tooth, S., Williams, R. M. E., and Chu, P.-S.: Drainage network development in the Keanakākoʻi tephra, Kīlauea Volcano, Hawaiʻi: implications for fluvial erosion and valley 835 network formation on early Mars, *J. Geophys. Res.*, 117, 1–19, doi:10.1029/2012JE004074, 2012.
- Devauchelle, O., Petroff, A. P., Lobkovsky, A. E., and Rothman, D. H.: Longitudinal profile of channels cut by springs, *J. Fluid Mech.*, 667, 38–47, doi:10.1017/S0022112010005264, 2011.
- Devauchelle, O., Petroff, A. P., Seybold, H. F., and Rothman, D. H.: Ramification of stream networks., *P. Natl. Acad. Sci. USA*, 109, 20832–20836, doi:10.1073/pnas.1215218109, 2012.
- 840 Dunne, T.: Formation and controls of channel networks, *Prog. Phys. Geog.*, 4, 211–239, doi:10.1177/030913338000400204, 1980.
- Ferguson, R., Lee, E. M., and Weller, L.: Themis geodetically controlled mosaics of Mars, *Lunar Planet. Sci. Conf.*, XLIV, abstract 1642, The Woodlands, Texas, 18–22 March, 2013.
- Forsyth, D. A. and Ponce, J.: *Computer Vision: a Modern Approach*, 2nd Edn., Prentice Hall, Upper Saddle 845 River, New Jersey, 2011.
- Fox, G. A. and Wilson, G. V.: The role of subsurface flow in hillslope and stream bank erosion: a review, *Soil Sci. Soc. Am. J.*, 74, 717–733, doi:10.2136/sssaj2009.0319, 2010.
- Fox, G. A., Wilson, G. V., Periketi, R. K., and Cullum, R. F.: Sediment transport model for seepage erosion of streambank sediment, *J. Hydraul. Eng.-ASCE*, 11, 603–611, doi:10.1061/(ASCE)1084- 850 0699(2006)11:6(603), 2006.
- Glines, N. and Fassett, C. I.: Evidence for Groundwater Sapping on Mars from Junction Angles of Nirgal Vallis Tributaries, in: *Lunar Planet. Sci. Conf.*, Vol. XLIV, p. abstract 2011, The Woodlands, Texas, 18–22 March, 2013.
- Hack, J. T.: *Studies of Longitudinal Stream Profiles in Virginia and Maryland*, US Geol. Survey Prof. Pap., 855 294-B, Washington D.C., 1957.
- Hagerty, D. J.: Piping/sapping erosion. I: Basic considerations, *J. Hydraul. Eng.-ASCE*, 117, 991–1008, doi:10.1061/(ASCE)0733-9429(1991)117:8(991), 1991.
- [Harrison, K. P. and R. E. Grimm \(2004\), Tharsis recharge: A source of groundwater for Martian outflow channels, *Geophys. Res. Lett.*, 31\(14\), L14,703, doi:10.1029/2004GL020502.](#)
- 860 Harrison, K. P. and Grimm, R. E.: Groundwater-controlled valley networks and the decline of surface runoff on early Mars, *J. Geophys. Res.*, 110, E12S16, doi:10.1029/2005JE002455, 2005.

- Harrison, S., Balme, M., Hagermann, A., Murray, J., Muller, J.-P., and Wilson, A.: A branching, positive relief network in the middle member of the Medusae Fossae Formation, equatorial Mars-Evidence for sapping?, *Planet. Space Sci.*, 85, 142–163, doi:10.1016/j.pss.2013.06.004, 2013.
- 865 Higgins, C. G.: Drainage systems developed by sapping on Earth and Mars, *Geology*, 10, 147–152, doi:10.1130/0091-7613(1982)10<147:DSDBSO>2.0.CO;2, 1982.
- Howard, A. D. and Kochel, R. C.: Introduction to Cuesta landforms and sapping processes on the Colorado Plateau, in: *Sapping Featur. Color. Plateau*, edited by: Howard, A. D., Kochel, R. C., and Holt, H. R., chap. 2, 6–56, NASA Spec. Publ. 491, Washington D.C., 1988.
- 870 Howard, A. D. and McLane, C. F.: Erosion of cohesionless sediment by groundwater seepage, *Water Resour. Res.*, 24, 1659–1674, doi:10.1029/WR024i010p01659, 1988.
- Irwin, R. P., Tooth, S., Craddock, R. A., Howard, A. D., and de Latour, A. B.: Origin and development of theater-headed valleys in the Atacama Desert, northern Chile: morphological analogs to Martian valley networks, *Icarus*, 243, 296–310, doi:10.1016/j.icarus.2014.08.012, 2014.
- 875 Jaumann, R. and Reiss, D.: Nirgal Vallis: evidence for Extensive Sapping, in: *Lunar Planet. Sci. Conf.*, p. abstract 1579, The Woodlands, Texas, 11–15 March, 2002.
- Jaumann, R., Neukum, G., Behnke, T., Duxbury, T., Eichertopf, K., Flohrer, J., Gasselt, S., Giese, B., Gwinnner, K., Hauber, E., Hoffmann, H., Hoffmeister, a., Köhler, U., Matz, K.-D., McCord, T., Mertens, V., Oberst, J., Pischel, R., Reiss, D., Ress, E., Roatsch, T., Saiger, P., Scholten, F., Schwarz, G., Stephan, K.,
880 and Wählisch, M.: The high-resolution stereo camera (HRSC) experiment on Mars Express: instrument aspects and experiment conduct from interplanetary cruise through the nominal mission, *Planet. Space Sci.*, 55, 928–952, doi:10.1016/j.pss.2006.12.003, 2007.
- Kochel, R. C. and Piper, J. F.: Morphology of large valleys on Hawaii: evidence for groundwater sapping and comparisons with Martian valleys, *J. Geophys. Res.*, 91, E175, doi:10.1029/JB091iB13p0E175, 1986.
- 885 Laity, J. E. and Malin, M. C.: Sapping processes and the development of theater-headed valley networks on the Colorado Plateau, *Geol. Soc. Am. Bull.*, 96, 203–217, doi:10.1130/0016-7606(1985)96%3C203:SPATDO%3E2.0.CO;2, 1985.
- Lamb, M. P., Howard, A. D., Johnson, J., Whipple, K. X., Dietrich, W. E., and Perron, J. T.: Can springs cut canyons into rock, *J. Geophys. Res.*, 111, 7002, doi:10.1029/2005JE002663, 2006.
- 890 Lee, P.: Selective fluvial erosion on mars: glacial selective linear erosion on Devon island, Nunavut, arctic Canada, as a possible analog, *Lunar Planet. Sci. Conf.*, XXXI, abstract 2080, The Woodlands, Texas, 13–17 March, 2000.
- Lobkovsky, A. E., Jensen, B., Kudrolli, A., and Rothman, D. H.: Threshold phenomena in erosion driven by subsurface flow, *J. Geophys. Res.*, 109, 1–10, doi:10.1029/2004JF000172, 2004.
- 895 Luo, W., Arvidson, R. E., Sultan, M., Becker, R., Katherine Crombie, M., Sturchio, N., and El Alfy, Z.: Groundwater sapping processes, Western Desert, Egypt, *Geol. Soc. Am. Bull.*, 109, 43–62, doi:10.1130/0016-7606(1997)109<0043:GWSPWD>2.3.CO;2, 1997.
- Mangold, N. and Ansan, V.: Detailed study of an hydrological system of valleys, a delta and lakes in the Southwest Thaumasia region, Mars, *Icarus*, 180, 75–87, doi:10.1016/j.icarus.2005.08.017, 2006.

- 900 Marra, W. A., Braat, L., Baar, A. W., and Kleinhans, M. G.: Valley formation by groundwater seepage, pressurized groundwater outbursts and crater-lake overflow in flume experiments with implications for Mars, *Icarus*, 232, 97–117, doi:10.1016/j.icarus.2013.12.026, 2014a.
- Marra, W. A., Hauber, E., de Jong, S. M., and Kleinhans, M. G.: Pressurized groundwater systems in Lunae and Ophir Plana (Mars): insights from small-scale morphology and experiments, submitted, 2014b.
- 905 Marra, W. A., Hauber, E., McLelland, S. J., Murphy, B. J., Parsons, D. R., Conway, S. J., Roda, M., Govers, R., and Kleinhans, M. G.: Pressurized groundwater outflow experiments and numerical modeling for outflow channels on Mars, *J. Geophys. Res.-Planet.*, 119, 2668–2693, doi:10.1002/2014JE004701, 2014c.
- Marra, W. A., Kleinhans, M. G., and Addink, E. A.: Network concepts to describe channel importance and change in multichannel systems: test results for the Jamuna River, Bangladesh, *Earth Surf. Proc. Land.*, 39, 910 766–778, doi:10.1002/esp.3482, 2014d.
- Otvos, E. G.: Rain-induced beach processes; landforms of ground water sapping and surface runoff, *J. Coastal Res.*, 15, 1040–1054, 1999.
- Pelletier, J. D.: Drainage basin evolution in the Rainfall Erosion Facility: dependence on initial conditions, *Geomorphology*, 53, 183–196, doi:10.1016/S0169-555X(02)00353-7, 2003.
- 915 Pelletier, J. D. and Baker, V. R.: The role of weathering in the formation of bedrock valleys on Earth and Mars: a numerical modeling investigation, *J. Geophys. Res.*, 116, 1–13, doi:10.1029/2011JE003821, 2011.
- Pornprommin, A., Takei, Y., Wubneh, A. M., and Izumi, N.: Channel inception in cohesionless sediment by seepage erosion, *J. Hydro-Env. Res.*, 3, 232–238, doi:10.1016/j.jher.2009.10.011, 2010.
- Schorghofer, N., Jensen, B., Kudrolli, A., and Rothman, D. H.: Spontaneous channelization 920 in permeable ground: theory, experiment, and observation, *J. Fluid Mech.*, 503, 357–374, doi:10.1017/S0022112004007931, 2004.
- Schumm, S. A. and Phillips, L.: Composite channels of the Canterbury Plain, New Zealand: a Martian analog?, *Geology*, 14, 326, doi:10.1130/0091-7613(1986)14<326:CCOTCP>2.0.CO;2, 1986.
- Wordsworth, R., Forget, F., Millour, E., Head, J. W., Madeleine, J.-B. Charnay, B.: Global modelling of the 925 early martian climate under a denser CO₂ atmosphere: Water cycle and ice evolution, *Icarus*, 222(1), 1–19, doi:10.1016/j.icarus.2012.09.036, 2013.

~~Experimental runs, their duration, discharge setting and data acquisition intervals, video number corresponds with videos in the Supplement.~~




Figure 1. Examples of theater-headed valleys. (a) Louros Valles on Mars, (perspective view); (b) valleys on the coast of the Atacama Desert, Chile (oblique photo, human for scale); (c) a-riverbank (oblique photo); (d) Side valleys of Snake River, Idaho, USA (orthorectified image); (e) valleys on the coast of the Atacama Desert, star indicates position of viewpoint of panel b (orthorectified image); (f) Apalachicola bluffs near Bristol, Florida, USA (orthorectified image). Image credits: (a) Google Earth (NASA/USGS, ESA/DLR/FU Berlin), (b) Tjalling de Haas, (c) Wouter Marra, (d,f) Bing Maps Imagery, (d) Courtesy of GFZ Potsdam.




Figure 2. Fundamental processes at valley heads for overland flow and groundwater seepage. (a) Valleys formed by groundwater seepage extend in headward direction by mass-wasting processes. (b) Valleys by overland flow deepen by fluvial incision and extend in headward direction by knickpoint retreat.




Figure 3. Setup of the **sapping** experiments. **(a)** (a) Oblique photo from downstream end of the flume, showing the initial sediment surface, gutter and constant head tank in the back. The rain simulators and cameras are above the photographed area; their approximate locations (C1-C6) are shown. **(b)** (b) Cross section showing setup with impermeable floor, constant head tank (ch), gutter, approximate location of spray nozzles (sn) and brickwork. **(c)** (c) Plan view showing the locations of the rain simulators (x), camera locations (C1-C6) and positions of the laser scanner (LS1, LS2).




Figure 4. Stills from time-lapses of the experiments showing the main morphological development. Full time-lapse movie are available in the [Supplementonline supplementary materials](#).




Figure 5. Erosion maps / final morphology of the experiments showing valley letters used in subsequent figures. ~~(a)~~ (a) Distant source experiment, ~~(b)~~ (b) Local precipitation experiment ~~(c)~~ (c) Distant after Local local and ~~(d)~~ (d) Local after Distant distant.




Figure 6. Channel profiles and slopes. (a) Channel profiles of valleys C, D and E of the distant source experiment and valleys N and L of local precipitation experiment, displayed with factor 2 vertical exaggeration. Elevations are arbitrary and plotted with offset for clarity. The three distant source experiment profiles show three arrested stages of development also seen in larger valleys: incipient sapping-valley without a steep valley head (D), developing valley with moderate steep valley head (C) and developed valley with steep valley head and reduced valley floor gradient (E), arrows indicate break in slope at the valley floor. (b) Boxplot of slope in upstream and downstream part of all valleys in the distant source ($n = 9$) and local precipitation ($n = 14$) experiments. The horizontal dotted line shows the initial surface slope $S_i = 0.22$.




Figure 7. Valley development in the [distal-distant source](#) (left panel) and local [precipitation](#) (right panel) [experiment](#)[experiments](#). Main valleys indicated with colors, letters in legend correspond with letters in [Fig. Figure 5](#). [\(a, f\)](#) [\(a, f\)](#) Valley width and [\(b, g\)](#) [\(b, g\)](#) length derived from orthorectified time-lapse imagery, [\(c, h\)](#) [\(c, h\)](#) valley depth derived from SFM-dems, [\(d, i\)](#) [\(d, i\)](#) estimated erosion rate from these properties and [\(e, j\)](#) [\(e, j\)](#) total cumulative erosion estimate from valley volume compared to measured sediment output.




Figure 8. Development of valley width ~~vs.~~ versus length of the ~~(a) distal~~ (a) distant source and ~~(b) local precipitation~~ (b) local precipitation experiment. Colored symbols in panel a ~~represent~~ represent the three main valleys. Values are plotted for all time-lapse intervals (valley dimensions increase with time). The open symbols in panel a ~~represent~~ represent valley dimensions when the measured valley section flowed at the side wall which influenced the valley width. Dotted line indicates trend of the three persisting valleys when the flume wall did not influence their width.




Figure 9. Final morphology of experiment Local precipitation 2, which is the initial morphology of experiment Distal-Distant after Local local.




Figure 10. Maps and profile of Louros Valles. (a) Overview map showing location of (b) panel (b) (MOLA shaded relief). (b) THEMIS daytime infrared mosaic with color-coded MOLA DEM, showing location of (c) panels (c) and (d). (c) Valley centerlines, color-coded with stream-order on THEMIS day-IR mosaic. (d) Detail of the network showing a densely dissected landscape and bifurcating valleys. (e) Elevation profile based on HRSC data, moving average using a 10-km window (plotted with 1000-m vertical offset) and slopes of two segments. Location of this profile is the first order valley indicated in (c) panel (c).




Figure 11. Maps of Nirgal Vallis. (a) Overview map showing location of (b) panel b (MOLA shaded relief). (b) THEMIS daytime infrared mosaic with color-coded MOLA DEM, showing location of (c) panels c and (d). (c) Valley centerlines, color-coded with stream-order on THEMIS day-IR mosaic. (d) Detail of the network showing a sparsely dissected landscape with many small and a few large valleys.




Figure 12. Landscape metrics of Louros and Nirgal Vallis. (a-c) Valley orientation for valleys on the north (a) and south flank (b) of Louros Valles and Nirgal Vallis (c). (d and e) Valley length (distance to lower-order valley) distribution for different stream orders, most main valleys (order 1) plot far outside the shown window for (d) Louros and (e) Nirgal Vallis. (f and g) Distribution of bifurcation orientation and boxplots per stream order for (f) Louros and (g) Nirgal Vallis.




Figure 13. Landscape end-members formed by groundwater seepage as result from a ~~distal~~ distant source or local source precipitation, and steep or horizontal surface. Each panel shows a ~~schematic~~ schematic diagram (upper left), an Martian case showing a ~~similar~~ similar morphology (upper right) as an example and the expected valley length distribution and valley orientation (bottom). A ~~distal~~ distant source ~~(a, c)~~ (a, c) results in valley abandonment due to upstream capture of groundwater, ~~where~~ whereas a ~~local~~ local groundwater source ~~(b, d)~~ (b, d) is less prone to flow piracy. Horizontal surfaces ~~(c, d)~~ (c, d) have a ~~strong~~ strong tendency to form valley bifurcations in contrast to steep slopes ~~(a, b)~~ (a, b). ~~Distally fed valleys~~ Valleys emerging from a distant groundwater source result in an open landscape as no valleys develop downstream of large valleys. Similar Martian landscapes in ~~(a)~~ (a) Noctis Labyrinthus (THEMIS image), ~~(b)~~ (b) Gale Crater (CTX image), ~~(c)~~ (c) Nirgal Vallis (THEMIS image) and ~~(d)~~ (d) Louros Valles (THEMIS image).

Table 1. Experimental runs, their duration, discharge setting and data acquisition intervals, video number corresponds with videos in the online supplementary materials. Abbreviation used: d=days, h=hours, min=minutes.

<u>Experiment</u>	<u>Duration</u>	<u>Mean Q (l/min)</u>	<u>Cumm. Q (m³)</u>	<u>Time-lapse interval</u>	<u>SfM interval</u>	<u>video</u>
<u>Distant source</u>	<u>3 d. 3 h.</u>	<u>2.4</u>	<u>10</u>	<u>5 min.</u>	<u>1 d. 3 h.</u> <u>2 d. 2 h.</u> <u>2 d. 22 h.</u> <u>3 d. 3h.</u>	<u>video 1</u>
<u>Local precipitation</u>	<u>1 h. 50 min.</u>	<u>11.9</u>	<u>0.95</u>	<u>30 s.</u>	<u>end of exp.</u>	<u>video 2</u>
<u>Local precipitation 2</u>	<u>40 min.</u>					
<u>Local after distant</u>	<u>3 h. 10 min.</u>	<u>10.5</u>	<u>1.9</u>	<u>30 s.</u>	<u>end of exp.</u>	<u>video 3</u>
<u>Distant after local</u>	<u>3 d. 16 h.</u>	<u>2</u>	<u>10</u>	<u>5 min.</u>	<u>2 h.</u> <u>21 h.</u> <u>2 d. 0 h.</u> <u>2 d. 20 h.</u> <u>3 d. 16 h.</u>	<u>video 4</u>