

We wish to thank the reviewers for their insightful comments, which have led to a significantly improved paper. We previously responded to reviewer 1's comments. Here we take the opportunity to add some additional comments and make clearer how the manuscript has or has not been revised to address his/her concerns. In addition, we have included our responses to reviewer 2. Finally, a tracked-changes version of the revised manuscript is provided.

Reviewer 1:

Q: *“My large quibble is the authors use of a modern 30-year sediment yield data set (erosion rate data) as an input parameter into a model to predict equilibrium drainage density over millennial time-scales without little discussion of the appropriateness of applying a short-term data set to evaluate how differences in erosion rates might control variations in relief, drainage density and predicted differences from divides to valley bottoms. Given that previous work (e.g. Kirchner et al., 2001) point to a discrepancy between decadal-scale measurements of stream fluxes and erosion rates over longer time scales this is a point that needs clarification and discussion in the current study While the Kirchner et al. study was in mountainous terrain, their conclusion that the episodic nature of sediment delivery may limit the use of decadal-scale sediment yield studies in evaluating long-term landscape evolution is extremely pertinent to the Walnut Gulch Experimental Watersheds. I was disappointed in the minimal discussion in the paper of potential episodic events in the watersheds not captured by the data (e.g. extreme rain events, fire) or of the potential for skew in comparing the grassland (n=1) to shrubland (n=5) watershed erosion rate data if episodic events such as fire were non-evenly represented in the watersheds. Additionally, while the mean sediment yield data reflects a 10x difference, the standard deviation for the shrub sites is quite large ($2.03 \pm 2.2 \text{ t h}^{-1} \text{ yr}^{-1}$) which suggest perhaps exploring potential outcomes across a greater range of reasonable parameters.”*

A: Sediment yield cannot be negative, so the measured yields should not be reported as $2.03 \pm 2.2 \text{ t h}^{-1} \text{ yr}^{-1}$. The sediment yield is a function of drainage area, hence some of the variation in measured yields reflects a systematic increase in sediment yield with drainage area at WGEW, a dependence we accounted for. In our analysis we used the sediment yield data to quantify sediment transport coefficients that account for the drainage area dependence. For the shrubland sites the sediment transport coefficient is reported on lines 424-426 as follows: “ $k_{Q_{ss}} = 2 \times 10^{-6} \text{ m}^{1.56} \text{ yr}^{-1}$ (with a range of values including one standard error from 2×10^{-7} to 2×10^{-5}), $p = 1.44 \pm 0.2$, and $R^2 = 0.93$.” The value of $k_{Q_{ss}}$ one standard error below the mean is still more than 3 times larger than the value of $k_{Q_{sg}}$, which is $6 \times 10^{-8} \text{ m}^{1.56} \text{ yr}^{-1}$. We would certainly be willing to include additional potential outcomes (which would simply show a range of possible drainage densities depending on the ratio of $k_{Q_{ss}}$ to $k_{Q_{sg}}$), but we believe that our fundamental conclusion that sediment transport coefficients are ~10x greater in shrublands relative to grasslands is not undermined by the variability in the data.

We appreciate that sediment yields can differ among time scales. However, we believe that our decadal-scale sediment yields are an appropriate estimate (and certainly the best-available estimate given the likely impossibility of ^{10}Be cosmogenic erosion-rate determination in WGEW due to the Pleistocene age of the bedrock) for millennial-scale yields based on three lines of argument:

- 1) Published analyses have shown that fluvial sediment transport in WGEW, while highly episodic, is not dominated by a very small number of extreme events. The effective discharge (i.e. the discharge above which half of the total load is transported) occurs many times within a 30-year record. In a study of sediment transport in the 1995-2005

period, for example, Nearing et al. (2007) addressed this issue as follows: “For six of the seven watersheds, between 6 and 10 events produced 50% of the total sediment yields over the 11-year period.” That is, the effective discharge has a recurrence interval of between approximately 1 and 2 years. This is consistent with many other studies demonstrating that the effective discharge in low-gradient alluvial channels not subject to debris flows has a recurrence interval of approximately 1 to 2 years (e.g., Wolman and Miller, 1960; Andrews, 1980; Andrews and Nankervis, 1995).

References:

Andrews, E. (1980), Effective and bankfull discharge in the Yampa River Basin, Colorado and Wyoming, *J. Hydrol.*, 46, 311-330.

Andrews, E. D., and, J. M. Nankervis (1995), Effective discharge and the design of channel maintenance flows for gravel-bed rivers, *Natural and Anthropogenic Influences in Fluvial Geomorphology*, AGU Monograph Series 89, Washington, D.C., 151-164.

Wolman, M. G., and, J. P. Miller (1960), Magnitude and frequency of forces in geomorphic processes, *J. Geol.*, 68, 54-74.

- 2) Sediment yields calculated from 1995-2005 by Nearing et al. (2007) closely match yields measured over approximately 50 years using ^{137}Cs (Nearing et al., 2005), as we noted on lines 205-208. The 50 year time scale includes significant droughts at WGEW. We are not suggesting that this proves that sediment yields are invariant out to time scales of millennia, but it does establish constancy of rates out to “short” time scales that are significantly longer than most studies of short-term erosion rates, including Kirchner et al. (2001).
- 3) Kirchner et al. (2001) showed order-of-magnitude differences in sediment yields/erosion rates between interannual and millennial-scale erosion rates from a forested landscape subject to high-severity (i.e. stand-replacing) forest fires. The recurrence interval of high-severity forest fires in the western U.S. ranges from 150 to 400 yr based on frequency distributions of even-aged stands and fire-related sedimentation studies (Meyer et al., 1995; Veblen, et al., 1994; Kipfmüller and Baker, 2000; Sibold et al., 2006; Margolis et al., 2007; 2011; Fitch and Meyer, 2015). Erosion rates commonly increase by 2 to 4 orders of magnitude for one to several years following high-severity forest fires (e.g. Wagenbrenner and Robichaud, 2014; Orem and Pelletier, 2015). These numbers strongly suggest that, *in forests similar to those studied by Kirchner et al. (2001)*, millennial-scale erosion rates may be dominated by the erosion that occurs shortly following high-severity wildfires. This suggestion is consistent with the work of Fowler (1979). In his remarkable compilation of 90 sediment yield studies in forested areas of the U.S., Fowler (1979) found that approximately 20% of the 90 studies report short-term erosion rates of $<1 \mu\text{m}/\text{yr}$, i.e. 2 orders of magnitude below typical long-term erosion rates (i.e. $\sim 10\text{-}100 \text{ m}/\text{Myr}$ in mid-latitude areas of low to moderate relief). Of the remaining 80% of studies he compiled (i.e. those with short-term erosion rates $>1 \mu\text{m}/\text{yr}$), Fowler (1979) demonstrated that nearly all were associated with the occurrence of wildfire and/or landsliding/debris flows. As such, it is likely that some of the episodicity in erosion rates documented by Kirchner et al. (2001) is the result of high-severity forest fires and/or episodic mass movements. Neither of these processes occurs at WGEW in the late Holocene. The modest slopes of WGEW preclude debris flows. Wildfires are of very limited size and severity in shrublands. Grassland fires in Arizona typically result in modest (if any) increase in runoff and erosion rates (e.g. Stone et al., 2003).

References:

- Fitch, E.P., and Meyer, G. (2015), Temporal and spatial climatic controls on Holocene fire-related erosion and sedimentation, Jemez Mountains, New Mexico, *Quat. Res.*, doi:10.1016/j.yqres.2015.11.008.
- Fowler, J.M. (1979), The interface of forestry and agriculture as nonpoint sources of suspended sediment: A national modeling approach, Ph.D. dissertation, Iowa State Univ., Ames, Iowa, 244 p.
- Kipfmüller, K.F., and Baker, W.L. (2000), A fire history of a subalpine forest in southeastern Wyoming, USA, *J. Biogeogr.*, 27, 71-85, doi: 10.1046/j.1365-2699.2000.00364.x.
- Margolis, E. Q., Swetnam, T. W., and C. D. Allen (2007), A stand-replacing fire history in upper montane forests of the southern Rocky Mountains, *Can. J. Forest Res.*, 37(11), 2227-2241.
- Margolis, E. Q., Swetnam, T. W., and C. D. Allen (2011), Historical stand-replacing fire in upper montane forests of the Madrean Sky Islands and Mogollon Plateau, southwestern USA, *Fire Ecol.*, 7(3), 88-107.
- Meyer, G.A., Wells, S.G., and Jull, A.J.T. (1995), Fire and alluvial chronology in Yellowstone National Park: Climatic and intrinsic controls on Holocene geomorphic processes, *Geol. Soc. Am. Bull.*, 107(10), 1211-1230, doi: 10.1130/0016-7606(1995)107<1211:FAACIY>2.3.CO;2.
- Orem, C., and Pelletier, J.D. (2015), Quantifying the time scale of elevated geomorphic response following wildfires using multi-temporal LiDAR data: An example from the Las Conchas fire, Jemez Mountains, New Mexico, *Geomorphology*, 232, 224-238, doi: 10.1016/j.geomorph.2015.01.006.
- Sibold, J.S., Veblen, T.T., and Gonzalez, M.E. (2006), Spatial and temporal variation in historic fire regimes in subalpine forests across the Colorado Front Range in Rocky Mountain National Park, Colorado, USA, *J. Biogeogr.*, 32, 631-647, doi:10.1111/j.1365-2699.2005.01404.x.
- Stone, J.J. et al., (2003), Post-wildfire runoff and erosion response on grassland and oak woodlands in southeastern Arizona, available at https://www.firescience.gov/projects/03-2-3-11/project/03-2-3-11_03_2_3_11_Deliverable_03.pdf
- Veblen, T.T., Hadley, K.S., Nel, E.M., Kitzberger, T., Reid, M., and Villalba, R. (1994), Disturbance regime and disturbance interactions in a Rocky Mountain subalpine forest, *J. Ecol.*, 82(1), 125-135, doi:10.2307/2261392.
- Wagenbrenner, J.W., and Robichaud, P.R. (2014), Post-fire bedload sediment delivery across spatial scales in the interior western United States, *Earth Surf. Process. Landf.*, doi:10.1002/esp.3488.

We added the following text to the Discussion section on uncertainty to address this comment:

“ In particular, Kirchner et al. (2001) demonstrated the potential pitfall of applying sediment yields or erosion rates measured at one time scale to another time scale by demonstrating that rates over different time scales can differ by orders of magnitude. We agree that this is a significant potential concern. However, we believe that our decadal-scale sediment yields are an appropriate estimate for millennial-scale sediment yields based on three lines of argument. First, as noted above, published analyses have shown that fluvial sediment transport in WGEW, while highly episodic, is not dominated by just a few extreme events. The effective discharge (i.e. the discharge above which half of the total load is transported) occurs many times

within a 30-year record. In a study of sediment transport in the 1995-2005 period, for example, Nearing et al. (2007) addressed this issue as follows: “For six of the seven watersheds, between 6 and 10 events produced 50% of the total sediment yields over the 11-year period.” That is, the effective discharge has a recurrence interval of between approximately 1 and 2 years. Second, sediment yields calculated from 1995-2005 by Nearing et al. (2007) closely match yields measured over approximately 50 years using ¹³⁷Cs (Nearing et al., 2005). This 50 year time scale includes significant droughts at WGEW, i.e. time periods when vegetation cover would have been anomalously low and erosion rates high. Third, recent work suggests that the order-of-magnitude increase in sediment yields measured by Kirchner et al. (2001) between interannual and millennial time scales may partly reflect the influence of high-severity (i.e. stand-replacing) forest fires in their study area in Idaho. Orem and Pelletier (2016) measured wildfire-affected and non-wildfire-affected erosion rates measured over interannual time scales together with erosion rates measured over millennial and million-year time scales in the forested Valles Caldera, New Mexico. Orem and Pelletier (2016) observed a similar increase in erosion rates to that of Kirchner et al. (2001). They were also able to demonstrate that the increase they measured was due to the episodic effects of high-severity wildfire. In contrast to forested areas, wildfires are of limited size and severity in shrublands. Similarly, grassland fires in Arizona typically result in modest (if any) increases in runoff and erosion rates (e.g., Stone et al., 2003). As such, there is a basis for concluding that the large increase in erosion rates measured by Kirchner et al. (2001) between interannual and millennial time scales is unlikely to occur in WGEW.

Q: “While MAP is similar between sites, the authors do not discuss differences in rainfall intensity or aspect that may also contribute to different runoff patterns and drainage density. While I am in full agreement with the stated hypothesis regarding a climate-modulated vegetation shift controlling variations in topographic attributes, additional discussion (and perhaps rejection) of alternative controls would strengthen the paper. While the author’s do address some of this uncertainty in the discussion section, it seems appropriate to consider alternative scenarios earlier on.”

A: Certainly we can include these data, which demonstrate minimal differences between the shrubland and grassland sites. Nearing et al. (2015) (referenced in our paper) demonstrated that rainfall erosivity (which includes intensity at 30 min duration) is approximately 20% lower in Lucky Hills compared to Kendall (their Figure 1b). This is somewhat larger than the difference in MAP we noted on line 145, but still insignificant when compared to the 30-fold difference in erosion rates/sediment transport coefficients. Kendall drains to the west and has roughly equal areas of N- and S-facing hillslopes. The shrubland sites have a range of aspects and exhibit erosion rates approximately 30x higher than the grassland sites across all aspects. These points were included in the revision in the section introducing the study sites.

Q: “Also I found it puzzling that while Nearing et al. 2005 concluded that the differences in sediment yields between grass and shrub sites was controlled primarily by watershed morphology this study attributes sediment yield differences to the amount of cover. (Though admittedly there may be subsequent studies to the Nearing 2005 paper of which I am unaware).”

A: A key point of our paper is that watershed morphology and vegetation cover are related, so it is not inconsistent to invoke both watershed morphology and vegetation cover as controls, as the two are related. Also, Nearing et al. (2005) invoked vegetation cover as a cause of erosion rate differences. In order to emphasize this fact, we took the unusual step of directly quoting Nearing

et al. (2005) on this point in lines 214-220: “Nearing et al. (2005) interpreted the differences in erosion rates between Lucky Hills and Kendall to be primarily a function of vegetation cover, i.e. “hydrologic response differences as a function of vegetation differences are probably largely responsible for the differences in hillslope erosion rates between the two watersheds. If flows are more concentrated and vegetative cover is less, as on the Lucky Hills site, flow shear stresses and stream power will tend to be greater, resulting in a greater hydrologic potential for erosion. Also important is probably the higher litter cover and plant basal area cover on the grassland site that would have a direct protective effect against erosion.” No change made on this point.

Q: *“It would help to clarify if any of the topographic data was smoothed and if so what was the smoothing window for deriving hillslope gradient, curvature, and the stream profiles used for extracting slope-area plots and the associated power-law fits as values as the smoothing radius choices may influence interpretation.”*

A: As stated on line 233 an Optimal Weiner Filter was used to smooth the topography, following Pelletier (2013). No change made on this point.

Q: *“Also, as highlighted below in the line-specific comments, it would be helpful to report the shrubland standard error when comparing values between the grassland and shrubland sites.”*

A: For the shrubland sites this value is reported on lines 424-426 as follows: “ $k_{QSS} = 2 \times 10^{-6} \text{ m}^{1.56} \text{ yr}^{-1}$ (with a range of values including one standard error from 2×10^{-7} to 2×10^{-5}), $p = 1.44 \pm 0.2$, and $R^2 = 0.93$.” Note that the standard error is included. No change made on this point.

Q: *“Some additional justification on the use diffusive equations for colluvial sediment transport in non-soil mantled shrubland (bare earth) settings seems necessary as geomorphic transport laws do not yet exist for erosion driven by overland flow nor would I expect the hillslope erosion to be smooth but rather episodic and patchy.”*

A: WGEW is soil mantled almost everywhere (outcrops are extremely rare). Hence, we do not think a discussion of non-soil-mantled landscapes is relevant to our study area. It may be that the reviewer has interpreted our reference to “bare soil” on line 182 as referring to an absence of soil. Bare soil simply refers to soil with no vegetation or stone cover. No change made on this point.

Q: *“L387-389 What is the SE and number of slopes evaluated for the reported slope values of 0.17 and 0.19?”*

A: The computed slope values are based on an analysis of all hillslope pixels in the DEM (tens of millions of pixels). It is somewhat difficult to place a precise error estimate on this value. The mean value can be estimated very precisely because we have tens of millions of pixels to average and the error of the mean decreases as \sqrt{N} . There is, however, also a structural error associated with the fact that what is hillslope versus what is valley bottom cannot be defined with absolute precision. No change made on this point.

Q: *“L404-40 Why does so much of the terrain exhibit positive curvature values? This seems curious unless this corresponds to relief values and if so this is worthy of discussion. Also the deviation between grassland and shrubland curvature is correctly described in the figure caption as from ~ 10 to 300 m^2 rather than the 30 m^2 value reported in the text.”*

A: It may be that the community has become used to looking at numerical models of landscapes driven to topographic steady state, and that this has biased our view of the world. In such cases the vast majority of the landscape (i.e. essentially all hillslopes) will have negative curvature. In nature, however, toeslopes commonly have positive curvature and colluvial infilling often results in wide unchannelized valley bottoms with positive curvature. The numerical model results presented by Pelletier (2011) that were motivated by WGEW and do not assume topographic steady state (they were subjected to a pulse of uplift/base-level fall followed by stasis) show that much of the landscape has positive curvature. In the revision we changed the 30 m² reported value to 10 m² on line 406.

Q: “L407-417 (Figure 9) The use of the slope-area plots seem a bit ad hoc with the exception of the identification of the colluvial region of the data sets. What part of the slope area plots were used to generate the exponents of 0.15 and 0.18 for grasslands and shrublands respectively? Is there a reason for presenting the concavity index values as they are not discussed at all in the document except to note that they somewhat different? I’m also a bit flummoxed by the use of the shrubland mean rather than plotting individual watersheds as there is no way to tell if the watersheds are similar enough to warrant combining. And finally it is common to log-bin the data to reduce noise – I am unsure how the authors generated the concavity index values. Presumably using just the fluvial portion of the plots (> 50 m² for shrublands and > 100 m² for the grasslands) and perhaps log-binned? Also it is difficult to distinguish between the slope area plots since they both use a solid black line of the same weight. It appears as though the shrubland plot has depositional areas or knickpoints or something else leading to a non-smooth slope area relationship.”

A: Yes, the data were log-binned. We will take this figure and the associated text out of the revision. We just thought that this analysis was useful to show as part of a complete topographic analysis of the study area, but we have removed it since the results are not essential. Figure and text removed.

Q: “L428-431 Why rely on a diffusivity value from scarp degradation studies (which presumably are faster than hillslope diffusion) rather than the smaller diffusivity values from the western US? This is an area where the authors could explore the parameter space using a range of values in eqns 11 and 12.”

A: Scarp degradation is modeled as hillslope diffusion by Hanks (2000) and all of the papers he cites. As such, we are unsure what the reviewer is referring to when he/she suggests a difference between diffusivity values derived from scarp degradation and those derived from hillslope diffusion. Scarp degradation is hillslope diffusion so there is no difference. The scarp studies referenced by Hanks (2000) that led to the 1 m²/kyr value are all from the western U.S. and range between approximately 0.5 and 2 m²/kyr, hence they have a geometric mean of approximately 1 m²/kyr. No change made on this point.

Q: “L523-538 Holmgren (2003) is from a region far to the south of the study area. I seem to recall that more local paleoclimate reconstructions suggest a much later transition from grasslands to shrublands. While the new perspective afforded by the more recent paleo vegetation data is understandably exciting, more context is necessary to demonstrate why this new(ish) data is applicable to a region to the north of the midden data.”

A: The study sites of Holmgren et al. (2003) are directly east of WGEW (both are centered on 31.75°N). There is no more local paleovegetation reconstruction that exists. No change made on this point.

Q: “L571-578 How does the reduction in vegetative influence on soil stability translate to channels – as an increase in sediment supplied to channels could potentially armor the beds? Especially given that the channels are described as transport-limited.”

A: A difference in bed armoring between the shrubland and grassland sites would not result from a difference in sediment supply but rather from a significant difference in the texture of the bed material. Bed sediment texture is similar in Lucky Hills and Kendall. No change made on this point.

Q: “Section 4.4 The global compilations suggesting little correlation between MAP and erosion rates are increasingly becoming less surprising as increasing numbers of studies and conceptual models are starting to converge on periglacial temperatures and associated processes, rather than MAP, influencing erosion rates in regions mid-latitude terrain (e.g. Herman and Braun, 2008; Marshall et al., 2015; Savi et al., 2015; Schaller et al., 2002; Tucker et al., 2011). Consider modifying this section to constrain the weak linkages between MAP and erosion rates to unglaciated areas outside the influence of periglacial processes.”

A: We intended this discussion to focus on non-glaciated areas outside the dominant influence of periglacial processes. Sentence will be modified in revision to: “Recent work on the role of vegetation, and its changes through time, can provide a basis for understanding the relatively low correlation between erosion rates and MAP in unglaciated areas outside the dominant influence of periglacial processes and the complex relationship between erosion rates and climate in such areas more generally.” That said, while periglacial processes involve temperature cycling near the freezing point of water, they also involve water, hence MAP or other measures of water availability are still relevant to the discussion of peri/paraglacial process rates. Sentence modified as stated.

Q: “L601-605 Perhaps I missed it but I did not see evidence in the manuscript for erosion rates during an arid to humid transition – so while this discussion item is worth considering in general when evaluating the role of the magnitude of transient erosion states, I’m unsure if the study results can be applied to this outstanding argument.”

A: Correct – our study provides no data for the landscape response to arid-to-humid transitions. We will simply remove the sentence “That is, erosion rates can be larger during a humid-to-arid transition than during an arid-to-humid transition, even if the mean climatic conditions (averaged over the transition) are equal” in the revision. Sentence removed.

Q: “L607-609 This statement on vegetation as an erosion agent due to bioturbation seems to directly contradict the hypothesis stated in lines 571-578. And while trees are considered both soil dilators and mechanisms for bedrock detachment and transport, I am unaware of any studies that suggest grasses or shrubs are effective mechanisms for increasing erosion.”

A: We do not see a contradiction here. Lines 571-578 refer to the concept that less vegetation cover leads to an increase in drainage density, all else being equal. Drainage density is set by a competition between diffusive (colluvial) and advective (fluvial) processes. Less vegetation cover leads to a decrease in colluvial transport rates and an increase in fluvial transport rates, as

stated on lines 607-609, hence it increases drainage density by increasing the relative importance of advective processes relative to diffusive processes, as stated on lines 571-578.

Previous studies have shown that more vegetation cover drives higher rates of colluvial (i.e. bioturbative) transport (e.g. Hughes et al., 2009) while less vegetation cover drives higher rates of fluvial erosion (e.g. this paper and many references therein, including Abrahams et al., 1995), all else being equal. We agree there may be no studies in which an increase in grass or shrub density specifically has been shown to increase colluvial transport rates. However, there are certainly a number of studies (e.g. Hughes et al., 2009) that demonstrate that an increase in biomass generally increases sediment transport by bioturbation.

References:

Hughes, M.W., P.C. Almond, and J.J. Roering (2009), Increased sediment transport via bioturbation at the last glacial-interglacial transition, *Geology*, 37(10), 919-922, doi:10.1130/G30159A.1

No change made on this point.

The incorrect figure number on line 391 and the incorrect date on Holmgren et al. (2003) in the reference list will be fixed in the revised paper. We will also include labels E and F in Fig. 3. We appreciate the reviewer pointing out these errors.

Reviewer 2:

Q: *“The only issue I would like to raise is in the last paragraph of this review....The data analysis section of the paper is great and clearly shows associations between vegetation and morphology. The main issue I have is that an equilibrium model does not seem to be the right tool to test the hypothesis that “late Holocene vegetation changes can modulate drainage density, hillslope-scale relief, and watershed-scale erosion rates.” According to this paper all these changes might have happened approximately in the last 3K years. This hypothesis requires a transient model which would examine if the observed decadal erosion rates, when used with a conservation law, can modify landscapes such that an initially identical topography erodes faster, reducing relief and developing increased drainages that can be recognized on an evolved topography, which would have similar patterns to observed landscapes. I wonder why the authors specifically used an equilibrium model instead of using some of the existing models the lead author uses in his research. Exploring the hypothesis posed in the paper using a transient 2D or 3D model, perhaps in addition to the analytical model presented in the paper, would make this paper a lot stronger and more complete. I strongly recommend the authors to consider revising their papers with this in mind.”*

A: To address this issue we have explicitly added a transient component to the analysis. The added text is as follows:

In the Methods section:

“To estimate the time required for low-order channels to grow headward in response to hillslope vegetation changes, we modeled the channels as a diffusive system (e.g., Begin, 1988). In diffusive systems, the time scale, t_r , required for a response over a length scale, L , can be estimated using

$$t_r \sim \frac{L^2}{D} \quad (13)$$

where D is a diffusivity. The value of D in the diffusive model for alluvial channels is given by the ratio of the unit volumetric sediment flux to the along-channel slope:

$$D = \frac{Q_s}{wS}. \quad (14)''$$

In the Results section:

“ In order to test the hypothesis that 2-4 kyr is sufficient time for drainage density to have fully responded to the recent grassland/woodland-to-shrubland transition, we used equations (13) and (14) to estimate the response time of low-order channels. We used $L = 50$ m to represent the approximate distance that the valley heads migrated upslope in response to late Holocene vegetation changes. We used data associated with the outlets of watersheds 105 and 106 as representative of the low-order channels in the shrubland-dominated portions of WGEW. Using equation (14), the D value for these channels is approximately 3×10^3 m²/kyr based on the ~ 0.1 m³/yr sediment flux of the 105 and 106 watersheds (Fig. 10), a width of approximately 1 m, and slope of approximately 0.03. Substituting these values for L and D into equation (13) yields an estimate for t_r equal to 0.83 kyr, or ≈ 1 kyr in keeping with the approximate nature of this calculation. This time scale is significantly shorter than the age of the transition (2-4 kyr), suggesting that sufficient time has occurred for drainage density to have fully responded to the changes in vegetation cover in the lower elevations of WGEW.”

Also:

“The fact that curvature values are very similar between shrubland and grassland below spatial scales ~ 10 m² is consistent with the hypothesis that hillslopes in the lower elevations of WGEW have not yet fully adjusted to the increase in drainage density associated with the grassland-to-shrubland transition. However, as the analysis of equations (13) and (14) reveal, the drainage density itself (which is the focus of this paper) has likely had sufficient time to fully adjust to the vegetation changes.”

The reviewer suggested that we use more sophisticated landscape evolution models to investigate whether 3 kyr is sufficient time for the system to fully respond to hillslope vegetation changes at this study site. This is a very reasonable suggestion. However, we think the simpler approach we have included adequately addresses the issue he/she raised. The first author appreciates that the reviewer acknowledges that he is capable of performing more sophisticated landscape evolution modeling. He choose not to include such models in this paper because 1) he thought the models presented were adequate to address the problem, 2) he wants to take the time to carefully apply such models to WGEW as part of a future study that includes additional data (e.g. soil texture on hillslopes and channels), and 3) he is currently revamping his transport-limited landscape evolution models with more sophisticated solution techniques for a number of new applications and release to the community.

Q: “I also list a few papers below which are relevant to the subject studied in this paper and present similar vegetation controls on landscape morphology, which the authors may want to use for comparison with their findings.”

A: We wish to thank the reviewer for these suggestions. We decided to add the Yetemen et al. (2010) reference as this one seemed most relevant. While the others are also relevant, they appear to be more focused on slope aspect controls on vegetation and slope form, which is of great interest to us but is less directly relevant to our study (which does not deal with slope aspect). Sentence added:

“We also propose that the difference in mean curvatures between shrublands and grasslands between contributing areas of ~ 10 and ~ 300 m² partly reflects recent expansion of the drainage network in the shrublands of WGEW. This hypothesis is consistent with the fact that the

deviation of curvature values between the two sites begins at a contributing area comparable to the support area (i.e. the contributing area required to form a valley head) in the grasslands and complementary studies (e.g., Yetemen et al., 2010) that have quantified the influence of vegetation cover on curvature-area relationships.”

Tracked changes:

The influence of Holocene vegetation changes on topography and erosion rates: A case study at Walnut Gulch Experimental Watershed, Arizona

Jon D. Pelletier^{1*}, Mary H. Nichols², Mark A. Nearing²

¹*Department of Geosciences, University of Arizona, Gould-Simpson Building, 1040 East Fourth Street, Tucson, Arizona 85721-0077, USA*

²*USDA Southwest Watershed Research Center, 2000 East Allen Rd., Tucson, Arizona 85719, USA*

*corresponding author; email: jdpellet@email.arizona.edu

Abstract

Quantifying how landscapes have responded and will respond to vegetation changes is an essential goal of geomorphology. The Walnut Gulch Experimental Watershed offers a unique opportunity to quantify the impact of vegetation changes on landscape evolution over geologic time scales. The Walnut Gulch Experimental Watershed (WGEW) is dominated by grasslands at high elevations and shrublands at low elevations. Paleovegetation data suggest that portions of WGEW higher than approximately 1430 m a.s.l. have been grasslands and/or woodlands throughout the late Quaternary, while elevations lower than 1430 m a.s.l. changed from a

grassland/woodland to a shrubland c. 2-4 ka. Elevations below 1430 m a.s.l. have decadal time-scale erosion rates approximately ten times higher, drainage densities approximately three times higher, and hillslope-scale relief approximately three times lower than elevations above 1430 m. We leverage the abundant geomorphic data collected at WGEW over the past several decades to calibrate a mathematical model that predicts the equilibrium drainage density in shrublands and grasslands/woodlands at WGEW. We use this model to test the hypothesis that the difference in drainage density between the shrublands and grassland/woodlands at WGEW is partly the result of a late Holocene vegetation change in the lower elevations of WGEW, using the upper elevations as a control. Model predictions for the increase in drainage density associated with the shift from grasslands/woodlands to shrublands are consistent with measured values. Using modern erosion rates and the magnitude of relief reduction associated with the transition from grasslands/woodlands to shrublands, we estimate the timing of the grassland-to-shrubland transition in the lower elevations of WGEW to be approximately 3 ka, i.e., broadly consistent with paleovegetation studies. Our results provide support for the hypothesis that common vegetation changes in semi-arid environments (e.g. from grassland to shrubland) can change erosion rates by more than an order of magnitude, with important consequences for landscape morphology.

Keywords: landscape evolution, drainage density, vegetation cover, Walnut Gulch Experimental Watershed

1. Introduction

1.1. Problem statement

Understanding how climate change controls landscape evolution is a central problem in geomorphology. Climate changes are multifaceted, with changes in temperature (mean and variability), precipitation (mean and variability) and vegetation cover (type and density) often occurring simultaneously. The multifaceted nature of climatic changes can make it difficult to identify which aspects of climate change are most important in driving landscape modification in specific cases. Yet, given the accelerated climatic changes expected to occur in the coming decades, understanding how landforms are likely to respond to specific climatic drivers, acting alone or in concert, is critically important to society (e.g., Pelletier et al., 2015).

In the southwestern U.S. the existence of an extensive, regionally correlative fan and valley floor depositional unit (the Q3a unit of Bull, 1991) suggests that the Pleistocene-to-Holocene transition was a major perturbation in what are now semi-arid shrubland-dominated landscapes but which were predominantly pinyon-juniper woodlands at the Last Glacial Maximum (LGM). Where rates of aggradation in modern shrubland drainage basins have been measured, the Pleistocene-to-Holocene transition was associated with more than an order-of-magnitude increase above either LGM or mid-to-late Holocene rates (e.g., Fig. 3.11 of Weldon, 1986). The retreat of grasslands and woodlands to higher elevations and their replacement by shrublands likely exposed elevations of the southwestern U.S. between approximately 800 m and at least 1400 m a.s.l. to significant increases in percent bare ground, thus modifying the rainfall-runoff partitioning of hillslopes and their resistance to fluvial/slope-wash erosion. In especially arid areas of the southwestern U.S. such as the central Mojave Desert, the range of elevations affected by late Quaternary conversions of grasslands/woodlands to shrublands extends to elevations as high as 1800 m a.s.l. (Pelletier, 2014).

Given the strong correlation between percent bare ground and drainage density in the southwestern U.S. (Melton, 1957), it has been hypothesized that modern shrublands sufficiently high in elevation to have been grasslands or woodlands at LGM underwent large increases in drainage density during the Pleistocene-to-Holocene transition. Such an expansion of the fluvial drainage network could have mobilized hillslope deposits stored as colluvium during the last glacial epoch, mobilizing large volumes of sediment into the fluvial system during the transition to the present interglacial (Bull, 1991; Pelletier, 2014). This hypothesis is consistent with measured ages of the onset of aggradation in valley floor and alluvial fan depositional zones in the central Mojave Desert, in which aggradation occurs earliest (c. 15 ka) in depositional zones fed by source regions with relatively low elevations in the 800-1800 m a.s.l. range and progressively later in areas fed by eroding regions at higher elevations (Pelletier, 2014). Alternative explanations for the punctuated nature of late Quaternary aggradation in the southwestern U.S. invoke changes in the frequency and/or magnitude of floods (Antinao and McDonald, 2013b) and argue that hillslope vegetation changes played a limited role (Antinao and McDonald, 2013a). To date, tests of the paleovegetation change hypothesis in the southwestern U.S. have been limited to studies of the timing of deposition on valley-floor channels and alluvial fans. Erosion of the source drainage basins themselves has been relatively understudied.

The Walnut Gulch Experimental Watershed (WGEW) provides an excellent opportunity to test the paleovegetation change hypothesis in a drainage basin that has been extensively monitored for decades. The western, low-elevation portion of WGEW is currently a shrubland while the eastern portion is predominantly a grassland (a small area of woodland occupies the highest elevations). Paleovegetation data, however, indicate that all of WGEW was a grassland

or woodland until just a few thousand years ago. Scientists working at WGEW have measured rates of sediment export from watersheds using sediment samplers (e.g., Nichols et al., 2008) and rates of sediment redistribution within watersheds using anthropogenic radionuclides (e.g., ^{137}Cs) and rare-earth element tracers (e.g., Nearing et al., 2005; Polyakov et al., 2009). These data, in addition to the results of analyses of airborne lidar data presented here, make it possible to calibrate every parameter of a mathematical model that predicts the equilibrium drainage density of the landscape under different dominant vegetation types. In this paper we use this mathematical model to test the hypothesis that grassland/woodland-to-shrubland vegetation changes in the lower elevations of WGEW drove large increases in drainage density and erosion rates and a decrease in hillslope-scale relief.

1.2. Study Site

1.2.1. Geology and Soils

The Walnut Gulch Experimental Watershed (WGEW) is part of the U.S. Department of Agriculture (USDA) Agricultural Research Service's (ARS) Southwest Watershed Research Center (SWRC). WGEW is located at the boundary between the Chihuahuan and Sonoran Deserts and elevations of between approximately 1300 and 1600 m a.s.l. The approximately 150 km² watershed has a planar geometry at large spatial scales, dipping to the WSW with a slope of approximately 1.5%, i.e., approximately 230 m of elevation change over a distance of 15 km.

The bedrock geology of WGEW includes sedimentary, plutonic, and volcanic rocks of Precambrian to late Cenozoic age (Fig. 1). Due to the complex nature of the rock types exposed in the southern portion of the watershed, we focus this study on the northern portion of the watershed, which is dominated by the Gleason Road Conglomerate (GRC).

The GRC was derived primarily from erosion of the Dragoon Mountains to the east of WGEW and is estimated to be Plio-Pleistocene in age by Osterkamp (2008), who noted: “The upper part of the Gleeson Road Conglomerate is probably equivalent both stratigraphically and in age to the Plio-Pleistocene upper basin fill of Brown et al. (1966). To the west and northwest, along the axis of the San Pedro River Valley, the upper part of the Gleeson Road Conglomerate grades into fine-grained fluvial and lacustrine beds of the Plio-Pleistocene St. David Formation (Gray 1965, Melton 1965).” The GRC dips gently (5-8°) to the north and northwest (Gilluly, 1956, plate 5), a fact which may contribute to the generally longer and more gently sloping south-facing hillslopes relative to north-facing hillslopes in the watershed. The tilted strata of the GRC were beveled to a gently sloping topographic surface and incised into during Quaternary time. Incision of the GRC was driven by uplift/tilting of the fan and/or by incision of the San Pedro River, triggering headward erosion of Walnut Gulch and its tributaries.

The northern portion of WGEW is composed of the Whetstone Pediment of Bryan (1926) and is divided into two parts: the “dissected Whetstone Pediment” (DWP) at elevations below approximately 1430 m a.s.l. and the “upper Whetstone Pediment” (UWP) at higher elevations (Fig. 1). The DWP is distinguished from the UWP by its lower relief and less well-developed soils. Differences between the DWP and UWP have been attributed primarily to headward extension of tributaries resulting from late Quaternary incision of the San Pedro River (Cooley, 1968) as well as renewed river and tributary incision following livestock grazing beginning c. 1880 (Renard et al., 1993; Osterkamp, 2008). However, the boundary between the DWP and UWP coincides with the transition from the modern shrubland to the grassland (Fig. 2). As such, we hypothesize that vegetation cover and its changes over geologic time scales have also contributed to differences between the DWP and UWP.

Deep sandy gravel loams of the Blacktail-Elgin-Stronghold-McAllister-Bernardino Group occur in areas of the UWP. In the lower, western part of the watershed, soils are in the Luckyhills-McNeal Group. Soils of the Luckyhills-McNeal Group tend to be sandy and gravelly loams that are immature compared with soils of the Blacktail-Elgin-Stronghold-McAllister-Bernardino Group. The A horizon of the Luckyhills-McNeal Group is typically absent, having been removed by late Quaternary erosion (Breckenfeld, 1994). The boundary between these two soil groups coincides with the boundary between the DWP and UWP and the transition from the modern shrubland to the grassland.

1.2.3. Climate and Vegetation

Mean annual temperature at Tombstone (located in the western portion of WGEW at an elevation of 1384 m a.s.l.) is 17.6°C and mean annual precipitation is approximately 300 mm. There is both a winter and summer rainy season, but approximately 70% of rainfall occurs in the summer months and the greater intensity of summer storms means that runoff results almost exclusively during the summer season of July through September. Mean annual precipitation is approximately 10% higher [and rainfall erosivity \(which is based on rainfall intensity at 30 min duration\) is approximately 20% higher](#) at the highest elevations of the watershed relative to the lowest elevations (Nearing et al., 2015).

Modern vegetation cover in the U.S.-Mexico Borderlands region closely follows elevation, with desert scrub occurring below approximately 1500 m, grasslands from 1400–1700 m, lower encinal (“encina” is Spanish for “oak”) from 1700–2600 m, upper encinal from 1900–2600 m, and forest from 2200–2600 m (Wagner, 1977). For each zone, the highest elevations are reached on dry, south- and west-facing slopes and the lowest elevations on north-facing slopes and valley bottoms.

Paleovegetation records from the Borderlands region suggest that the western portions of WGEW have transitioned from a grasslands/woodland to a shrubland over the past few thousand years, while the eastern half of the watershed has been a grassland/woodland for at least the past 40,000 yr and likely since the penultimate interglacial period. Low stalagmite ^{18}O values at the Last Glacial Maximum (LGM) in the Cave of the Bells paleoclimate record indicate that conditions were much wetter and cooler during LGM (Wagner et al., 2010), in agreement with paleovegetation studies (Betancourt et al., 1990; Anderson, 1993; Betancourt et al., 2001; Arundel, 2002; Holmgren et al., 2005; Holmgren, 2005). Packrat midden records indicate the presence of grasslands and/or pinyon–juniper woodlands at LGM in what is currently Chihuahuan desert scrub at elevations of 1200-1400 m a.s.l. (Betancourt et al., 2001). Holmgren (2005) documented the presence of the primary grass species at WGEW (*Bouteloua eriopoda*) (currently abundant only at elevations above approximately 1430 m a.s.l.) at an elevation of 1287 m c. 4750 ^{14}C yr B.P. Elements of the modern shrubland, such as *Larrea tridentate*, appeared as late as 2190 ^{14}C yr B.P. at 1287 m a.s.l. based on macrofossils, but may have been present as early as 4095 ^{14}C yr B.P. based on pollen. As such, the latest Quaternary transition from grasslands/woodlands to shrublands in WGEW occurred gradually and was completed only within the past few thousand years.

1.2.4. Intensive monitoring sites

WGEW is home to two intensive monitoring sites, one in the shrubland of the DWP and the other in the grassland of the UWP. These sites provide the data necessary, in conjunction with the topographic analyses presented here, to calibrate a mathematical model that predicts the equilibrium drainage density as a function of vegetation cover and to test the hypothesis that differences in landscape morphology and erosion rates between the northwestern and

northeastern portions of WGEW are partly the result of a transition from grassland/woodland to shrubland within the past few thousand years in the northwestern portion of the drainage basin that did not occur in the northeastern portion.

Watersheds 63.102, 63.103, 63.104, 63.105, and 63.106 are located in shrublands at an elevation of approximately 1370 m a.s.l. in what is referred to as “Lucky Hills,” which has been the site of a variety of intensive scientific studies since the 1960s. Cover during the rainy season at Lucky Hills is approximately 25% bare soil, 25% canopy, and 50% erosion pavement (rocks). Dominant vegetation includes: Creosote (*Larrea tridentata*, shrub) and Whitethorn (*Acacia constricta*, shrub), with lesser populations of Desert Zinnia (*Zinnia acerosa*, shrub), Tarbush (*Flourensia cernua*, shrub), and sparse Black Grama (*Bouteloua eriopoda*, grass). The matrix material of surface layer is composed of 60% sand, 25% silt, and 15% clay. Sediment from the watershed is monitored with a supercritical flume with an automatic traversing slot sampler (Simanton et al., 1993). [The shrubland sites have a range of slope aspects and exhibit sediment yields approximately thirty times higher than the grassland sites across all slope aspects.](#)

Watershed 63.112 is located in the Kendall grassland site at WGEW, approximately 10 km east of Lucky Hills and at an elevation of approximately 1525 m. [Kendall drains to the west and has roughly equal areas of N- and S-facing hillslopes.](#) The site is predominantly covered by grass and forbs with some shrubs and succulents with a combined canopy cover of approximately 35%. Ground cover during the rainy season has been measured at 28% rock, 42% litter, and 14% plant basal cover (Nearing et al., 2007). Compared to the 25% bare soil at Lucky Hills, the bare soil exposed at Kendall is negligible (i.e. a few percent or less). Historically, the dominant desert grassland bunchgrasses at the site have been black grama (*Bouteloua eriopoda*), side-oats grama (*B. curtipendula*), three-awn (*Aristida sp.*), and cane beardgrass (*Bothriochloa*

barbinodis) (King et al., 2008), and more recently, Lehmann lovegrass (*Eragrostis lehmanniana*) (Moran et al., 2009).

Nearing et al. (2005) used spatially distributed ^{137}Cs measurements to quantify fluvial/slope-wash erosion and deposition rates within and from watersheds 63.103 (Lucky Hills) and 63.112 (Kendall). Nearing et al. (2005) found that in Lucky Hills the fraction of the drainage basin experiencing erosion was much higher (85%) than in Kendall (53%), where erosion and deposition rates were lower and approximately balanced such that the rate of net sediment exported from the Kendall watershed was more than a factor of ten lower than from the Lucky Hills watershed. These observations are consistent with sediment yields measured from 1995-2005 that are more than ten times higher in drainage basins of similar size at Lucky Hills (231 t $\text{km}^{-2} \text{yr}^{-1}$ in watershed 102 (area of 1.46 Ha)) than at Kendall (7 t $\text{km}^{-2} \text{yr}^{-1}$ in watershed 112 (area of 1.86 Ha)) (Nearing et al., 2007). Assuming a soil bulk density of 1500 kg m^{-3} , these sediment yields correspond to mean erosion rates of approximately $1.5 \times 10^{-4} \text{ m yr}^{-1}$ in the shrublands of Lucky Hills and $5 \times 10^{-6} \text{ m yr}^{-1}$ in the grasslands of Kendall (Table 1). Hillslopes are slightly steeper in Kendall, so if anything we would expect erosion rates to be higher at Kendall than at Lucky Hills if slope gradient were the dominant factor in controlling erosion rates. Nearing et al. (2005) interpreted the differences in erosion rates between Lucky Hills and Kendall to be primarily a function of vegetation cover, i.e. “hydrologic response differences as a function of vegetation differences are probably largely responsible for the differences in hillslope erosion rates between the two watersheds. If flows are more concentrated and vegetative cover is less, as on the Lucky Hills site, flow shear stresses and stream power will tend to be greater, resulting in a greater hydrologic potential for erosion. Also important is probably the higher litter cover and plant basal area cover on the grassland site that would have a direct protective effect

against erosion.” This interpretation is consistent with the conclusions of Abrahams et al. (1995) and Parsons et al. (1996), who emphasized the role of vegetation cover in controlling fluvial/slope-wash erosion rates in their plot-scale studies at WGEW. In this paper we explore the implications of these erosion rate differences for landscape morphology and topographic evolution of WGEW over geologic time scales.

2. Methods

2.1. Topographic analysis

In this section we describe the methods used to quantify the similarities and differences in landscape morphology between the modern grassland and shrubland sites, with an eye toward providing the data necessary to calibrate the mathematical model described in section 2.2. In all of the topographic analyses described in this section we used a 1 m pixel⁻¹ Digital Elevation Model (DEM) for WGEW derived from airborne laser swath mapping (Heilman et al., 2008). Prior to the analyses, the DEM was smoothed with an Optimal Weiner Filter (OWF), following the approach of Pelletier (2013), to remove small-scale variability related to errors related to distinguishing ground from vegetation points and other imperfections of the lidar-derived DEM. The smoothing did not significantly alter slope gradients but did significantly reduce anomalous curvature values related to DEM imperfections.

The drainage density in the shrubland areas appears to be much higher than in the grassland areas (Fig. 3). To quantify this difference we used the method developed by Pelletier (2013) to identify the network of valley bottoms (i.e. where water flow is localized and fluvial processes are responsible for the majority of erosion) in the vicinity of Lucky Hills and Kendall. In this method, the DEM is filtered using the OWF, the contour curvature is computed at every

pixel, and valley heads are identified as the areas closest to the divides where the contour curvature exceeds a user-defined threshold value. In Pelletier (2013) and in this paper a threshold contour curvature of 0.1 m^{-1} was used for valley head identification (i.e., the method of Pelletier (2013) was used without modification or parameter tuning). Once valley heads are identified, a multiple-flow-direction routing algorithm is used to route a unit of runoff from each valley head to identify the valley bottoms downstream. In this study we used the distance along flow lines from topographic divides to the first valley bottom as a measure of the extent of the drainage network. We computed the mean value of this hillslope length for all pixels entering valley heads. This mean value can be compared to the prediction of a mathematical model that computes the mean distance along flow lines from topographic divides to valley heads as a function of colluvial and fluvial transport coefficients. Hillslope length measured in this way is inversely related to drainage density (Horton, 1945). Its mean value contains information equivalent to drainage density, but it has the advantage that it is a mappable quantity (Tucker et al., 2001). The drainage network analysis was performed on representative $1.6 \times 1.6 \text{ km}$ areas in the vicinity of Lucky Hills and Kendall to quantify the difference in drainage density between shrubland and grassland areas within WGEW.

Relief was mapped as the difference between the highest elevation upstream from each pixel along flow lines. The results of the drainage network identification procedure described above were used to limit this analysis to the hillslope pixels only. We then computed the mean hillslope relief within 10 m elevation bins from 1320 m to 1550 m a.s.l. The resulting graph quantifies how the mean hillslope relief varies with elevation across the shrubland-to-grassland transition.

We also computed the mean value of the along-channel slope gradient and curvature (i.e. the Laplacian) as functions of contributing area. Differences in mean curvature as a function of contributing area provide a quantitative signature of how late Holocene vegetation changes have modified the landscape morphology in the vicinity of the hillslope-to-valley-bottom transition.

2.2. Mathematical modeling

In this section we describe the mathematical model used to quantify erosion over geologic time scales and its dependence on landscape morphology at WGEW. The mathematical model is used to predict the equilibrium drainage density, quantified as the mean distance along flow lines from divides to valley bottoms, in both shrublands and grasslands, in order to test the hypothesis that the difference in drainage density observed between the shrublands and grasslands at WGEW can be attributed, in part, to late Holocene vegetation changes in the shrubland portion of the watershed.

Erosion at WGEW over geologic time scales can be approximated as the sum of erosion due to colluvial and fluvial/slope-wash processes. Sediment transport by colluvial processes leads to a diffusion equation for topography if slope are uniformly soil-mantled and topographic gradients are modest (Culling, 1960):

$$E_c = -D\nabla^2 z \tag{1}$$

where E_c is the erosion rate by colluvial processes (defined to be positive if the landscape is eroding), D is diffusivity in $\text{m}^2 \text{yr}^{-1}$, and z is elevation in m. Equation (1) assumes that colluvial sediment flux is equal to the product of a coefficient (i.e., the diffusivity D) and the local slope gradient. This assumption is reasonable for WGEW, where hillslopes are uniformly soil mantled and the mean hillslope gradient is 7%.

We assume that fluvial/slope-wash processes in WGEW can be approximated as transport limited. We use the term fluvial/slope-wash to refer to all sediment transport by flowing water, wherever it occurs along the continuum from hillslopes (i.e., as sheet and rill flow) to channels (i.e., fluvial erosion). A transport-limited condition applies to landscapes in which most of the sediment is transported as bed-material load and sediment is readily deposited if the shear stress by flowing water declines with increasing distance along flow pathways. In the alternative detachment-limited end-member model of fluvial/slope-wash erosion, the shear stress required to detach sediment is much larger than the shear stress required to transport it, hence sediment redeposition is rare or nonexistent once detachment/entrainment occurs. Pelletier (2012) addressed the geomorphic conditions under which transport-limited versus detachment-limited conditions are likely to occur, taking into account data for the relative proportion of sediment transported as bed-material load versus wash load, among other factors, using WGEW as a case study. He concluded that among these two end-member models, WGEW is most accurately considered to be transport limited.

The assumption of transport-limited conditions implies that fluvial/slope-wash erosion, E_f , is equal to the divergence of the fluvial/slope-wash volumetric unit sediment flux, \mathbf{q}_s (the volumetric sediment flux per unit width of water flow):

$$E_f = \frac{1}{\varepsilon_0} \nabla \cdot \mathbf{q}_s . \quad (2)$$

where ε_0 is the grain packing density (assumed here to be 0.55, e.g., equivalent to a bulk density of 1500 kg m^{-3} for a grain density of 2700 kg m^{-3}). The divergence of the fluvial/slope-wash sediment flux can, within the valley network, be approximated by the derivative of the volumetric unit sediment flux in the along-valley direction, q_s , with respect to the distance downstream along flow lines, x . Adopting this approximation and summing the colluvial and

fluvial/slope-wash components, the total erosion rate at any point on the landscape is thus given by

$$E = \frac{1}{\varepsilon_0} \frac{\partial q_s}{\partial x} - D\nabla^2 z . \quad (3)$$

In this paper we use equation (3) to predict the equilibrium drainage density, quantified as the mean distance from divides to valley bottoms, associated with grasslands and shrublands at WGEW. Specifically, we substitute an empirical power-law relationship for the mean distance, x , along flow lines from divides to valley bottoms versus contributing area, A , into equation (3) and solve for the value of x where the fluvial erosion rate exceeds the colluvial deposition rate by an amount equal to E .

Next, we further parameterize the three terms in equation (3) in terms of measured proxies for erosion rate (e.g. decadal-scale sediment fluxes) and topographic parameters. The erosion rate, E , is constrained using the ratio of the volumetric sediment flux, Q_s , reported by Nearing et al. (2007) and contributing area, A :

$$E = \frac{1}{\varepsilon_0} \frac{Q_s}{A} . \quad (4)$$

The fluvial erosion term can be written in terms of Q_s using

$$\frac{\partial q_s}{\partial x} = \frac{\partial}{\partial x} \left(\frac{Q_s}{w} \right) . \quad (5)$$

where w is the valley-bottom width. We approximate the mean rate of colluvial deposition at valley heads by

$$D\nabla^2 z \Big|_{\text{heads}} \approx \frac{3DS_h}{w} , \quad (6)$$

where S_h is the mean slope gradient of toe slopes as they intersect the valley bottom at valley heads (Fig. 5). Equation (6) derives from the mass balance of a square segment of the valley

bottom in the vicinity of the valley head (Fig. 5). According to the assumption that soil-mantled hillslopes evolve diffusively, valley bottoms in the vicinity of the valley head receive a unit sediment flux equal to DS_h by colluvial transport processes from each of three adjacent hillslope segments. In the cross-sectional profile, the difference in sediment flux across the profile is equal to DS_h and that difference occurs over a distance of w . As such, the colluvial deposition rate, computed in a manner that is independent of DEM or pixel resolution, includes the valley width in the denominator (Pelletier, 2010a). Similarly, the divergence in the along-valley direction is approximately DS_h/w . The factor of 2 does not appear in the along-valley derivative because colluvial sediment flux enters the valley bottom segment only from the upslope direction. Colluvial sediment flux leaving the valley head is assumed to be negligible because the slope of the valley bottom is typically much smaller than that of the hillslope entering it from upslope.

Next, we introduce three power-law relationships that relate volumetric sediment flux to contributing area, channel width to contributing area, and contributing area to distance along flow lines from topographic divides. Sediment flux is a power-law function of contributing area at WGEW (Section 3.1):

$$\frac{1}{\varepsilon_0} Q_s = k_{Qs} A^p. \quad (7)$$

The coefficient k_{Qs} in equation (7) is a sediment transport efficiency parameter that is a function of runoff, sediment texture, vegetation cover, and potentially other factors. It takes on different values in the shrubland and the grassland (i.e., k_{Qss} and k_{Qsg}), reflecting the fact that sediment yield is a function of vegetation cover. Its value in shrublands, along with that of p , is obtained by a least-squares regression of equation (7) to Q_s values for the five shrubland drainage basins of Lucky Hills reported by Nearing et al. (2007). Its value for the grasslands is constrained by assuming that the same value of p applies to both shrublands and grasslands, and using the single

Q_s data point available for grasslands (i.e., for watershed 112) to constrain k_{Qs} using equation (7).

Miller (1995) measured 222 cross-sectional channel profiles in the field at WGEW and used those data to calibrate a power-law relationship between channel width and contributing area:

$$w = k_w A^l \quad (8)$$

where $k_w = 0.023 \text{ m}^{0.32}$ and $l = 0.34$. Contributing area has a power-law relationship with the mean distance, x , from divides along flow lines:

$$A = k_A x^c \quad (9)$$

In section 3 we report the values of k_{Qss} and k_{Qsg} , p , k_w , l , k_A , and c for WGEW. Substituting equations (4)-(9) into equation (3) yields

$$k_{Qs} k_A^{p-1} x^{c(p-1)} = \frac{k_{Qs}}{k_w} k_A^{p-l} c(p-l) x^{c(p-l)-1} - \frac{3DS_h}{k_w k_A^l x^{cl}} \quad (10)$$

Equation (10) is applied using vegetation-specific values for k_Q and S_h to solve for the mean distance from divides to valley heads in shrublands and grasslands, i.e., x_s and x_g , respectively.

That is, x_s is obtained by solving

$$k_{Qss} k_A^{p-1} x_s^{c(p-1)} = \frac{k_{Qss}}{k_w} k_A^{p-l} c(p-l) x_s^{c(p-l)-1} - \frac{3DS_{hs}}{k_w k_A^l x_s^{cl}} \quad (11)$$

and x_g is obtained by solving

$$k_{Qsg} k_A^{p-1} x_g^{c(p-1)} = \frac{k_{Qsg}}{k_w} k_A^{p-l} c(p-l) x_g^{c(p-l)-1} - \frac{3DS_{hg}}{k_w k_A^l x_g^{cl}} \quad (12)$$

Equation (3) is generally applicable within the fluvial network. Once the colluvial deposition rate is approximated using equation (6) (which makes use of S_h , the mean gradient of the hillslope toe slopes at valley heads), subsequent equations, including equations (10)-(12), apply only to valley heads. Equation (10)-(12) are a mathematical representation of the conceptual model, first

proposed by Tarboton et al. (1992), that erosion at valley heads is a competition between transport-limited fluvial erosion and colluvial deposition. That is, fluvial erosion rates must exceed colluvial deposition rates by an amount equal to the net erosion rate on the landscape in order to maintain an equilibrium drainage density.

To estimate the time required for low-order channels to grow headward in response to hillslope vegetation changes, we modeled the channels as a diffusive system (e.g., Begin, 1988). In diffusive systems, the time scale, t_r , required for a response over a length scale, L , can be estimated using

$$t_r \sim \frac{L^2}{D} \quad (13)$$

where D is a diffusivity. The value of D in the diffusive model for alluvial channels is given by the ratio of the unit volumetric sediment flux to the along-channel slope:

$$D = \frac{Q_s}{wS} \quad (14)$$

3. Results

3.1. Topographic analyses

Figure 3 illustrates the results of the drainage network identification for 1.6 km x 1.6 km examples of the landscape in the vicinity of the Lucky Hills and Kendall sites. The mean distance along flow lines from divides to valley bottoms is 18 m in the shrubland area and 50 m in grassland area, using the Pelletier (2013) algorithm. Figures 4A&4B illustrate that hillslopes in the shrubland area are more finely dissected with rills and gullies than in the grassland area. As part of this analysis we also measured the mean slope gradient of hillslopes immediately adjacent to valley heads, i.e. S_h in equation (10). We obtained $S_{hs} = 0.17$ m/m in shrublands and

$S_{hs} = 0.19$ m/m in grasslands. In lower-order valleys, along-channels slopes are approximately 0.03 m/m.

Mean hillslope relief increases substantially across the shrubland-to-grassland transition (Fig. 6). Between elevations of approximately 1320 and 1430 m a.s.l., mean relief is uniformly low (approx. 0.5-1 m). Above elevations of approximately 1450 m, hillslope relief increases abruptly and continues to increase with increasing elevation.

Contributing area follows a piece-wise power-law function of distance along flow lines from topographic divides (Fig. 7), with one set of values for k_A and c applicable on hillslopes and another set of values applicable within the valley network. Above contributing areas of approximately 50 m^2 (or, equivalently, distances from the divide equal to approximately 15 m), contributing area increases as the 2.5 power of distance from the divide for both grassland and shrubland areas, i.e. $k_A = 0.3 \text{ m}^{-0.5}$ and $c = 2.5$ in equation (9). Below contributing areas of 50 m^2 , $k_A = 2 \text{ m}^{-0.75}$ and $c = 1.25$. We used $k_A = 0.3 \text{ m}^{-0.5}$ and $c = 2.5$ when solving equations (11) and (12), since these values are most applicable to points within the valley network (i.e. at valley heads and points downstream). This is a self-consistent approach because the solutions to equations (11)&(12) are larger than 15 m (Section 3.2).

Plots of mean topographic curvature (i.e. the Laplacian of z) versus contributing area (Fig. 8) indicate that mean curvatures are nearly identical at small and large contributing areas but differ substantially within the range of contributing areas from ~ 30 to 300 m^2 .

~~Plots of mean along channel slope versus contributing area (Fig. 9) for the shrubland and grassland area follow typical patterns for fluvial topography, i.e. the data follow a power law relationship for relatively large contributing areas and deviate from power-law scaling at small contributing areas due to the predominance of colluvial processes at hillslope scales. The~~

exponents of the slope-area relationships differ somewhat between the shrubland and grassland areas, i.e. 0.15 for the shrubland and 0.18 for the grassland areas. The slope-area plots also deviate from power law scaling at different spatial scales. Data from shrublands maintain a power law down to contributing areas of approximately 50 m^2 , while data from the grassland areas deviate at larger spatial scales corresponding to contributing areas of approximately 100 m^2 . This finding is consistent with the higher drainage density measured in the shrubland-dominated portions of the landscape compared to the grassland portions.

3.2. Mathematical modeling

In this section we use the results of Section 3.1., together with analyses of the sediment yield reported by Nearing et al. (2007), to constrain the terms in equations (11)&(12) in order to solve for the mean distance from divides to valley heads in shrublands and grasslands. In order to constrain the absolute values of $k_{Q_{ss}}$ and p , we performed a least-squares minimization of equation (11) to the decadal-scale sediment yields reported by Nearing et al. (2007) for watersheds 102-106 (Lucky Hills) (Table 1). This regression yields $k_{Q_{ss}} = 2 \times 10^{-6} \pm \text{m}^{1.56} \text{ yr}^{-1}$ (with a range of values including one standard error from 2×10^{-7} to 2×10^{-5}), $p = 1.44 \pm 0.2$, and $R^2 = 0.93$ (Fig. 109). Assuming that the value of p derived from the shrubland watersheds also applies to the grassland watershed, the value of k_{Q_s} for the grassland is estimated to be $k_{Q_{sg}} = 6 \times 10^{-8} \text{ m}^{1.56} \text{ yr}^{-1}$. For D we adopt the value of $1 \times 10^{-3} \text{ m}^2 \text{ yr}^{-1}$ commonly inferred from scarp degradation studies in the southwestern U.S. (e.g., Hanks, 2000, Table 2 cites D values for the Basin and Range of the western U.S. of between 6.4×10^{-4} and $2 \times 10^{-3} \text{ m}^2 \text{ yr}^{-1}$ based on eight published studies). The full list of model parameters and their values is provided in Table 2.

We used equations (11)&(12) to predict the mean distance along flow lines from divides to valley bottoms in shrublands and grasslands. The predicted values are $x_s = 16 \text{ m}$ and $x_g = 66 \text{ m}$

(Table 3). In the topographic analysis shown in Figure 3, we measured 18 m in shrublands and 60 m in grasslands. As such, the model predicts mean distances from divides to valley bottoms within 10% of measured values.

Figure 410 plots the magnitude of the three terms for a range of possible mean distances from divide to valley bottom. The fluvial erosion rate is $\sim 10^{-2}$ m yr⁻¹ in shrublands and $\sim 10^{-3}$ m yr⁻¹ in grasslands, increasing with distance downstream, reflecting the nonlinear relationship between sediment flux and drainage area (Fig. 409). The colluvial deposition rate is $\sim 10^{-3}$ m yr⁻¹ for both shrublands and grasslands and decreases modestly with increasing distance from the divide as a result of the increase in channel width with increasing contributing area (equation (8)). Figure 410 demonstrates that the fluvial erosion rate must be quite large (approximately two orders of magnitude larger than the net erosion rate in this case) in order to counteract the effects of colluvial deposition and thus maintain a valley head.

We used the mean curvature versus contributing area data plotted in Figure 8 to reconstruct an average longitudinal profile from divides to valley bottoms in shrublands and grasslands in order to infer the approximate relief reduction associated with the late Holocene shift from grasslands/woodlands to shrublands in WGEW. Figure 421 illustrates the results of this integration. Integrating the curvature versus contributing area data in Figure 8 twice leads to two constants of integration, one of which is constrained by the requirement that the slope along flow pathways at divides is zero and the other by using a constant reference elevation at a contributing area of approximately 300 m². We chose $A \approx 300$ m² as the location to enforce the reference elevation because this is the contributing area below which the mean curvature begins to deviate significantly between the grassland and shrubland areas. The difference in elevation between the two profiles plotted in Figure 421 provides an estimate of the minimum erosion or

relief reduction associated with the shift from grassland to shrubland in the lower elevations of WGEW. We consider the results of Figure 4211 to be a minimum estimate because systematic differences in mean slope between the grassland and shrubland across a wide range of scales (Fig. 9) are not reflected (or not fully reflected) in curvature or any reconstruction of the longitudinal profile based on integrating the curvature. The results in Figure 4211 suggest that divides have lowered by a minimum of approximately 0.3 m as a result of Holocene vegetation changes. Given hillslope-scale erosion rates of approximately $\sim 10^{-4}$ m yr⁻¹ in shrublands and the much smaller erosion rate of $\sim 10^{-5}$ m yr⁻¹ in grasslands, it would take approximately 3 kyr following a transition from grasslands/woodlands to shrublands to erode the landscape by an amount 0.3 m greater in shrublands compared to grasslands/woodlands. This estimate is comparable to the age of the grassland/woodland-to-shrubland transition in the region at an elevation of 1287 inferred from paleovegetation studies in the region, i.e., 2-4 ¹⁴C kyr B.P. (Holmgren, 2005).

In order to test the hypothesis that 2-4 kyr is sufficient time for drainage density to have fully responded to the recent grassland/woodland-to-shrubland transition, we used equations (13) and (14) to estimate the response time of low-order channels. We used $L = 50$ m to represent the approximate distance that the valley heads migrated upslope in response to late Holocene vegetation changes. We used data associated with the outlets of watersheds 105 and 106 as representative of the low-order channels in the shrubland-dominated portions of WGEW. Using equation (14), the D value for these channels is approximately 3×10^3 m²/kyr based on the ~ 0.1 m³/yr sediment flux of the 105 and 106 watersheds (Fig. 10), a width of approximately 1 m, and slope of approximately 0.03. Substituting these values for L and D into equation (13) yields an estimate for t_r equal to 0.83 kyr, or ~ 1 kyr in keeping with the approximate nature of this

calculation. This time scale is significantly shorter than the age of the transition (2-4 kyr), suggesting that sufficient time has occurred for drainage density to have fully responded to the changes in vegetation cover in the lower elevations of WGEW.

4. Discussion

4.1. Uncertainty in parameter values and their impact on the model results

Equations (11)&(12) predict mean distances from divides to valley bottoms that are broadly similar to measured values. Several factors limit the accuracy of the comparison between measured and predicted values. First, we relied upon a regional value for the diffusivity D because we do not have a reliable means of calibrating this value locally. Second, decadal-scale erosion rates computed from sediment samplers may differ from long-term erosion rates. Nearing et al. (2007) found that for the six watersheds considered here, half of the sediment yield measured from 1995-2005 was derived from 6-10 events and that the largest events contributed between 9 and 11% of the total yield. Thus, while transport is highly episodic in WGEW, the most effective flood events have a sufficient number of recurrences to provide an estimate of the yield that does not depend sensitively on the time scale. This conclusion is consistent with the fact that erosion rates inferred from sediment sampling from 1995-2005 are similar to erosion rates measured over the post-bomb period using ^{137}Cs (Nearing et al., 2005). That said, the sediment yields reported by Nearing et al. (2007) may not include the most extreme drought conditions or other disturbances that could cause long-term sediment yields to be larger than those reported in Table 1.

In particular, Kirchner et al. (2001) demonstrated the potential pitfall of applying sediment yields or erosion rates measured at one time scale to another time scale by

demonstrating that rates over different time scales can differ by orders of magnitude. We agree that this is a significant potential concern. However, we believe that our decadal-scale sediment yields are an appropriate estimate for millennial-scale sediment yields based on three lines of argument. First, as noted above, published analyses have shown that fluvial sediment transport in WGEW, while highly episodic, is not dominated by just a few extreme events. The effective discharge (i.e. the discharge above which half of the total load is transported) occurs many times within a 30-year record. In a study of sediment transport in the 1995-2005 period, for example, Nearing et al. (2007) addressed this issue as follows: “For six of the seven watersheds, between 6 and 10 events produced 50% of the total sediment yields over the 11-year period.” That is, the effective discharge has a recurrence interval of between approximately 1 and 2 years. Second, sediment yields calculated from 1995-2005 by Nearing et al. (2007) closely match yields measured over approximately 50 years using ¹³⁷Cs (Nearing et al., 2005). This 50 year time scale includes significant droughts at WGEW, i.e. time periods when vegetation cover would have been anomalously low and erosion rates high. Third, recent work suggests that the order-of-magnitude increase in sediment yields measured by Kirchner et al. (2001) between interannual and millennial time scales may partly reflect the influence of high-severity (i.e. stand-replacing) forest fires in their study area in Idaho. Orem and Pelletier (2016) measured wildfire-affected and non-wildfire-affected erosion rates measured over interannual time scales together with erosion rates measured over millennial and million-year time scales in the forested Valles Caldera, New Mexico. Orem and Pelletier (2016) observed a similar increase in erosion rates to that of Kirchner et al. (2001). They were also able to demonstrate that the increase they measured was due to the episodic effects of high-severity wildfire. In contrast to forested areas, wildfires are of limited size and severity in shrublands. Similarly, grassland fires in Arizona typically

result in modest (if any) increases in runoff and erosion rates (e.g., Stone et al., 2003). As such, there is a basis for concluding that the large increase in erosion rates measured by Kirchner et al. (2001) between interannual and millennial time scales is unlikely to occur in WGEW.

4.2. Further discussion of the hypothesis of a vegetation-change-driven increase in drainage density in the shrublands of WGEW

We also propose that the difference in mean curvatures between shrublands and grasslands ~~between~~ within the range of contributing areas ~~offrom~~ ~10 and to $\sim 300 \text{ m}^2$ partly reflects recent expansion of the drainage network in the shrublands of WGEW. This hypothesis is consistent with the fact that the deviation of curvature values between the two sites begins at a contributing area comparable to the support area (i.e. the contributing area required to form a valley head) in the grasslands: and complementary studies (e.g., Yetemen et al., 2010) that have quantified the influence of vegetation cover on curvature-area relationships. As such, we propose that the shrubland areas previously had support areas comparable to the grassland areas and that drainage network expansion has influenced the drainage network and the morphology of the adjacent hillslopes down to spatial scales corresponding to contributing areas of $\sim 10 \text{ m}^2$. The fact that curvature values are very similar between shrubland and grassland below spatial scales $\sim 10 \text{ m}^2$ is consistent with the hypothesis that hillslopes in the lower elevations of WGEW have not yet fully adjusted to the increase in drainage density associated with the grassland-to-shrubland transition. However, as the analysis of equations (13) and (14) reveal, the drainage density itself (which is the primary focus of this paper) has likely had sufficient time to fully adjust to the vegetation changes.

Figure 5 demonstrates that mean hillslope relief increases substantially across the shrubland-to-grassland transition. We propose that some of this difference in hillslope relief is a

consequence of the difference in fluvial/slope-wash erosion rates, i.e. that higher erosion rates in the lower-elevation shrublands have caused relief reduction in the past few thousand years relative to the higher-elevation grasslands, and that this difference in fluvial/slope-wash erosion rates is the result of a geologically recent increase in drainage density in the shrublands of WGEW. However, it is likely that a portion of the difference in mean hillslope relief across the study site also reflects variable uplift rates, i.e. the fact that the uplift of any piedmont or foothill region tends to increase towards the mountain range. Flexural-isostatic response to erosion (which has been proposed to be an important component of late Cenozoic landscape evolution in southern Arizona (Menges and Pearthree, 1989; Pelletier, 2010b)) of the Dragoon Mountains can be expected to have caused eastward tilting of WGEW, i.e. higher uplift rates in the higher elevations of WGEW compared to the lower elevations. Tilting would not explain the abrupt increase in relief at elevations just above 1430 m a.s.l., however, since no faulting occurs in the vicinity of this contour. Therefore, it is likely that some of the difference in hillslope-scale relief across the shrubland-to-grassland transition at WGEW is a result of the difference in erosion rates between the shrublands and grasslands. While we can be certain of grassland-to-shrubland shift only during the present interglacial period, the Quaternary period has seen many interglacial periods broadly similar in climate to the current period, hence it is likely that the lower elevations of WGEW have seen grassland-to-shrubland conversions more than once over the past approximately 2 Myr. Each of these episodes could have contributed to relief reduction of the lower elevations of the study site relative to the higher elevations.

Previous studies at WGEW have emphasized the role of base-level lowering and vegetation changes within the past 130 years on the differences in erosion rate between Lucky Hills and Kendall (Nearing et al., 2007). However, recent paleovegetation studies have provided

a new perspective. Specifically, Holmgren's (2005) documentation of shrubland species in the region several thousand years before present at an elevation less than 100 m lower than Lucky Hills suggests that the lower elevations of WGEW likely shifted from a grassland/woodland to a shrubland prior to the 1880s. While base-level lowering has steepened hillslopes and channels close to the main-stem channel of Walnut Gulch, hillslope-scale relief ~~and slope gradients~~ are clearly larger at Kendall than at Lucky Hills (Figs Fig. 6&9), indicating that base-level lowering may be a dominant factor only for those areas within close proximity to the main channel. The magnitude of the differences in topography (i.e., drainage density and the magnitude of erosion than can be inferred from the change) is difficult to fit into a period as short as 130 years. Given erosion rates measured over the past sixty at Lucky Hills, approximately 2 cm of erosion can be expected to have occurred over the past 130 years. Figure 1211 suggests that erosion associated with a recent increase in drainage density is likely at least ten times this value, and thus more consistent with a vegetation change that occurred several thousand years before present.

4.3. Implications for our understanding of the controls on drainage density

The model of this paper contributes to our broader understanding of the controls on drainage density and it provides a mathematical model for predicting drainage density that may be useful in other study sites.

Previous studies have demonstrated that drainage density is controlled by relief (e.g. Montgomery and Dietrich, 1992; Tarboton, 1992; Tucker and Bras, 1999), climate (Melton, 1957; Abrahams and Ponczynski, 1984), parent material (e.g. Ray and Fischer, 1960; Day, 1980), and time (e.g. Ruhe, 1952; Dohrenwend et al., 1987). While many studies have demonstrated the importance of individual factors on drainage density, we lack a comprehensive model for drainage density that integrates all of these factors. Equation (10) represents one

possible candidate for such a model in soil-mantled, transport-limited landscapes. Time is not included in the model because it is based on an equilibrium mass-balance framework. Nevertheless, equation (10) predicts the drainage density to which a transient landscape will likely approach over time following a perturbation.

Relief enters the model via the erosion rate, E (quantified for the case of WGEW using multiple measurements of Q_s/A), and the mean toe slope gradient near valley heads, S_h . Climate and vegetation cover enters the model through the parameters D (which increases with increasing soil moisture and temperature cycling around 0°C (which together drive soil creep) and increasing vegetation cover (which drives bioturbation)) and k_{Qs} (which increases with rainfall and decreases with vegetation cover)). In addition, equation (10) explicitly includes channel width and its scaling with contributing area, factors that, to our knowledge, have not been included in previous mathematical models of drainage density.

Drainage density is most commonly found to be an inverse function of mean annual precipitation or effective precipitation. This finding is consistent with the conceptual model of this paper that vegetation cover is the predominant climate-related variable that influences drainage density, and that vegetation cover and drainage density are inversely related. Melton (1957), for example, documented an inverse correlation between drainage density and the precipitation-effectiveness (P-E) index at over eighty sites in the southwestern United States including arid (low elevation) and humid (high-elevation) climates. A similar negative correlation between drainage density and mean annual precipitation was found by Abrahams and Ponczynski (1984). Naively, one might expect more precipitation to result in greater channel incision and hence less contributing area, between divides and valley heads (and hence a higher drainage density), all else being equal. Melton (1957), however, proposed that greater aridity

results in a lower vegetation density and, hence, a reduction in the cohesive strength protecting soils on hillslopes, thus leading to a higher drainage density. One might also expect a lower vegetation density to increase the runoff-to-rainfall ratio and hence also increase drainage density, but runoff intensity varied by only a factor of two across Melton's sites while drainage density varied by nearly two orders of magnitude. Istanbuloglu and Bras (2007) provided theoretical support for Melton's interpretation, illustrating that a lower vegetation densities can lead to higher drainage densities through the cohesive or anchoring effect of plant roots.

4.4. Implications for our understanding of erosion-climate linkages

There has been an ongoing debate in the geomorphic literature regarding the importance of climate (not limited to but often defined as mean annual precipitation (MAP)) on erosion rates. Given the significant correlation between MAP and erosion rates in many studies within individual mountain ranges (e.g., Reiners et al., 2003; Bookhagen and Strecker, 2012), it is perhaps surprising how little correlation exists between MAP and erosion rates in global compilation/synthesis studies (e.g. von Blanckenburg, 2005; Portenga and Bierman, 2011). Even studies that emphasize the climatic control on erosion rates note that R^2 values between erosion rates and MAP are quite small (e.g., Yanites and Kesler, 2015).

Recent work on the role of vegetation, and its changes through time, can provide a basis for understanding the relatively low correlation between erosion rates and MAP in [global compilation studies](#) [unglaciated areas outside the dominant influence of periglacial processes](#) and the complex relationship between erosion rates and climate [in such areas](#) more generally. For example, Torres Acosta et al. (2015) recently documented a negative correlation between erosion rates and both vegetation cover and MAP in Kenya. They proposed that the primary effect of more humid conditions is to increase vegetation cover on hillslopes, thereby reducing erosion

rates on otherwise similar slope gradients. This concept is consistent with the classic Langbein and Schumm (1958) curve. Langbein and Schumm (1958) proposed that sediment yields are maximized in semi-arid climates (all else being equal) because such climates generate sufficient rainfall to detach and transport soil in overland/rill flow but insufficient vegetation cover to protect/anchor the soil. As MAP increases in this conceptual model, more precipitation is available to drive erosion, but this effect is more than offset by a decrease in the susceptibility of soil to erode due to the increased anchoring effect associated with greater plant cover/biomass. The results of this paper demonstrate further complexity in the erosion-climate relationship, i.e., that the change in climate (and hence of vegetation cover) can be as important or more important than its mean state. ~~That is, erosion rates can be larger during a humid to arid transition than during an arid to humid transition, even if the mean climatic conditions (averaged over the transition) are equal.~~ It is important to emphasize that the effect of vegetation on the rate of erosion by colluvial processes may be entirely different than its effect on fluvial/slope-wash processes. All else being equal, increased vegetation cover is likely to increase erosion rates by colluvial processes, since more plants can be expected to drive higher rates of bioturbation (e.g., Osterkamp et al., 2011). As such, it is crucial to consider colluvial and fluvial/slope-wash processes separately when considering the effects of vegetation on hillslope erosion.

5. Conclusions

In this study we leveraged all relevant data from a uniquely well-studied semi-arid watershed to test the hypothesis that late Holocene vegetation changes can modulate drainage density, hillslope-scale relief, and watershed-scale erosion rates. We documented that areas below 1430 m a.s.l. have decadal-scale erosion rates approximately ten times higher, drainage

densities approximately three times higher, and hillslope-scale relief approximately three times lower than elevations above 1430 m. We calibrated all the terms of a mathematical landscape evolution model and used the model to predict the equilibrium drainage density associated with shrublands and grasslands. Model predictions for the increase in drainage density associated with the shift from grasslands/woodlands to shrublands are broadly consistent with measured values. Using modern erosion rates and the magnitude of relief reduction associated with the transition from grasslands/woodlands to shrublands, we also estimated the timing of the grassland-to-shrubland transition in the lower elevations of WGEW to be approximately 3 ka, i.e., broadly consistent with constraints from paleovegetation studies. Our work provides a mathematical model for predicting equilibrium drainage density in transport-limited fluvial environments that may be applicable in other study sites.

Data Availability

The DEM used in this paper can be obtained upon request from the corresponding author. All other data used in the paper are in the published literature.

Acknowledgements

We wish to thank Tyson Swetnam for drafting Figure 2C. J.D.P. was partially supported by NSF award #1331408. [We thank two anonymous reviewers for thoughtful, constructive reviews that helped us improve the quality of the paper.](#)

References

Abrahams, A. D., Parsons, A. J., and Wainwright, J.: Effects of vegetation change on interrill erosion, Walnut Gulch, southern Arizona, *Geomorphology*, 13, 37–48, doi:10.1016/0169-555X(95)00027-3, 1995.

Abrahams, A. D., and Ponczynski, J. J.: Drainage density in relation to precipitation intensity in the U.S.A., *J. Hydrol.*, 75(1-4), 383–388, doi: 10.1016/0022-1694(84)90061-1, 1984.

Anderson, R. S.: A 35,000 Year Vegetation and Climate History from Potato Lake, Mogollon Rim, Arizona, *Quaternary Research*, 40, 351–359, doi: 10.1006/qres.1993.1088, 1993.

Antinao, J. - L., and McDonald, E.: A reduced relevance of vegetation change for alluvial aggradation in arid zones, *Geology*, doi:10.1130/G33623.1, 2013a.

Antinao, J. - L., and McDonald, E.: An enhanced role for the Tropical Pacific on the humid Pleistocene-Holocene transition in southwestern North America, *Quat. Sci. Rev.*, 78, 319–341, <http://dx.doi.org/10.1016/j.quascirev.2013.03.019>, 2013b.

Arundel, S. T.: Modeling climate limits of plants found in Sonoran Desert packrat middens, *Quaternary Research*, 58, 112–121, doi:10.1006/qres.2002.2349, 2002.

[Begin, Z. B. \(1988\), Application of a diffusion-erosion model to alluvial channels which degrade due to base-level lowering. *Earth Surf. Process. Landforms*, 13: 487–500. doi: 10.1002/esp.3290130603.](#)

Betancourt, J. L., Rylander, K. A., Penalba, C., and McVickar, J. L.: Late Quaternary vegetation history of Rough Canyon, south-central New Mexico, USA, *Palaeogeography Palaeoclimatology Palaeoecology*, 165, 71–95, doi:10.1016/S0031-0182(00)00154-1, 2001.

- Betancourt, J. L., Van Devender, T. R., and Martin, P. S.: Packrat Middens: The last 40,000 years of biotic change, The University of Arizona Press, Tucson, 467 p., 1990.
- Bookhagen, B., and Strecker, M. R.; Spatiotemporal trends in erosion rates across a pronounced rainfall gradient: Examples from the southern Central Andes: *Earth Planet. Sci. Lett.*, 327–328, 97–110, doi:10.1016/j.epsl.2012.02.005, 2012.
- Breckenfeld, D. J., Svetlik, W. A., and McGuire, C. E.: Soil Survey of Walnut Gulch Experimental Watershed, U.S. Dep. of Agric., Soil Conservation Service, Washington, D.C., 1995.
- Brown, S. G., Davidson, E. S., Kister, L. R., and Thomsen, B. W.: Water Resources of Fort Huachuca Military Reservation, Southeastern Arizona. U.S. Geological Survey Water-Supply Paper 1819-D, 57 pp., 1966.
- Bryan, K.: San Pedro Valley, Ariz., and the geographic cycle [abstract], *Geological Society of America Bulletin*, 54, 169–170, 1926.
- Bull, W. B.: *Geomorphic responses to climatic change*, Oxford, UK, Oxford University Press, 326 p., doi.wiley.com/10.1002/gea.3340080106, 1991.
- Cooley, M. E.: Some notes on the late Cenozoic drainage patterns in southeastern Arizona and southwestern New Mexico, in S. R. Titley, ed., *Southern Arizona Guidebook III*, Arizona Geological Society, Tucson, p. 75–78, 1968.
- Culling, W. E. H.: Analytical theory of erosion, *J. Geol.*, 68, 336–344, doi:10.1086/626663, 1960.
- Day, D. G.: Lithologic controls of drainage density: a study of six small rural catchments in New England, *NSW Catena*, 7, 339–351, doi:10.1016/S0341-8162(80)80024-5, 1980.

- Dohrenwend, J. C., A. D. Abrahams, and Turrin, B. D.: Drainage development on basaltic lava flows, Cima volcanic field, southeast California, and Lunar Crater volcanic field, south-central Nevada, *Geol. Soc. Am. Bull.*, 101(7), 982–986, doi:10.1130/0016-7606(1987)99<405:DDOBLF>2.0.CO;2, 1989.
- Gilluly, J.: *General Geology of Central Cochise County, Arizona*. U. S. Geological Survey Professional Paper 281, 189 p., 1956.
- Gray, R. S.: *Late Cenozoic Sediments in the San Pedro Valley near St. David, Arizona*. Unpublished Ph.D. dissertation, The University of Arizona, Tucson, 198 p., 1965.
- Hanks, T. C.: The age of scarplike landforms from diffusion-equation analysis, in *Quaternary Geochronology: Methods and Applications*, edited by J. S. Noller, J. M. Sowers and W. R. Lettis, pp. 313–338, American Geophysical Union, Washington D.C., doi:10.1029/RF004p0313, 2000.
- Holmgren, C. A.: *Late Quaternary Ecology, Climatology, and Biogeography of the northern Chihuahuan Desert from fossil packrat middens*, Ph.D. Dissertation, University of Arizona, Tucson, 156 p., Digital document available at <http://hdl.handle.net/10150/196087>, 2005.
- Holmgren, C. A., Penalba, M. C., Rylander, K. A., and Betancourt, J. L.: A 16,000 C-14 yr BP packrat midden series from the USA-Mexico Borderlands, *Quaternary Research* 60, 319–329, doi:10.1016/j.yqres.2003.08.001, 2005.
- Horton, R.E.: *Erosional Development of Streams and Their Drainage Basins; Hydrophysical Approach to Quantitative Morphology*, *Geological Society of America Bulletin*, 56 (3), 275-370, doi:10.1130/0016-7606(1945)56[275:EDOSAT]2.0.CO;2, 1945.

Istanbulluoglu, E., Bras, R. L.: Vegetation-modulated landscape evolution: Effects of vegetation on landscape processes, drainage density, and topography, *J. Geophys. Res.*, 110(F2), F02012, doi:10.1029/2004JF000249, 2007.

King, D. M., Skirvin, S. M., Collins, C. D. H., Moran, M. S., Biedenbender, S. H., Kidwell, M. R., Wertz, M. A., and Diaz-Gutierrez, A: Assessing vegetation change temporally and spatially in southeastern Arizona, *Water Resour. Res.*, 44, W05S15, doi:10.1029/2006WR005850, 2008.

[Kirchner, J. W., Finkel, R. C., Riebe, C. S., Granger, D. E., Clayton, J. L., King, J. G., and Megahan, W. F.: Mountain erosion over 10 yr, 10 k.y., and 10 m.y. time scales, *Geology*, 2, 9\(7\), 591–594, doi: 10.1130/0091-7613\(2001\)029<0591:MEOYKY>2.0, 2001.](#)

Langbein, W. B., and S. A. Schumm: Yield of sediment in relation to mean annual precipitation, *Eos Trans. AGU*, 39(6), 1076–1084, doi:10.1029/TR039i006p01076, 1958.

Melton, M. A.: An analysis of the relations among elements of climate, surface properties, and geomorphology, Dept. Geol. Columbia Univ. Tech. Rep. 11, Proj. NR 389-042, Off. of Nav. Res., New York, 1957.

Melton, M. A.: The geomorphic and paleoclimatic significance of alluvial deposits in southern Arizona, *J. Geol.*, 73(1), 1–38, doi:10.1086/627044, 1965.

Menges, C. M., and Pearthree, P. A.: Late Cenozoic tectonism in Arizona and its impact on regional landscape evolution, *Arizona Geological Society Digest*, 17, 649–680, doi: 1989.

Miller, S. N.: An analysis of channel morphology at Walnut Gulch: Linking field research with GIS applications, Unpublished M.S. Thesis, University of Arizona, Tucson Arizona, 167 p., 1995. Digital document available at <http://hdl.handle.net/10150/192090>.

Montgomery, D. R., and Dietrich, W. E.: Channel initiation and the problem of landscape scale, *Science*, 336, 232–234, doi:10.1126/science.255.5046.826, 1992.

Moran, M. S., Holifield Collins, C. D., Goodrich, D. C., Qi, J., Shannon, D. T., and Olsson, A.: Long-term remote sensing database, Walnut Gulch Experimental Watershed, Arizona, United States, *Water Resour. Res.*, 44, W05S10, doi:10.1029/2006WR005689, 2008.

Nearing, M. A., Kimoto, A., Nichols, M. H. and Ritchie, J. C.: Spatial patterns of soil erosion and deposition in two small, semiarid watersheds, *J. Geophys. Res.*, 110, F04020, doi:10.1029/2005JF000290, 2005.

Nichols, M. A., Stone, J. J., and Nearing, M. A.: Sediment database, Walnut Gulch Experimental Watershed, Arizona, United States, *Water Resour. Res.* 44:10.1029/2006WR005682, 2008.

Nearing, M. A., Unkrich, C. L., Goodrich, D. C., Nichols, M. H., and Keefer, T. O.: Temporal and elevation trends in rainfall erosivity on a 149 km² watershed in a semi-arid region of the American Southwest, *Inter. Soil Water Conserv. Res.* 3, 77-85, doi:10.1016/j.iswcr.2015.06.008, 2015.

[Orem, C. A., and Pelletier, J. D.: The predominance of post-wildfire erosion in the long-term denudation of the Valles Caldera, New Mexico. *J. Geophys. Res. Earth Surf.*, 120, doi:10.1002/2015JF003663, 2016.](#)

Osterkamp, W. R.: Geology, Soils, and Geomorphology of the Walnut Gulch Experimental Watershed, Tombstone, Arizona, *Journal of the Arizona-Nevada Academy of Science* 40(2):136-154, doi:[/10.2181/1533-6085-40.2.136](#), 2008.

- Osterkamp, W. R., Hupp, C. R., and Stoffel, M.: The interactions between vegetation and erosion: new directions for research at the interface of ecology and geomorphology, *Earth Surf. Process. Landf.*, 37, 23–36, doi:10.1002/esp.2173, 2012.
- Parsons, A., Abrahams, A. D., and Wainwright J.: Responses of interrill runoff and erosion rates to vegetation change in southern Arizona, *Geomorphology*, 14, 311–317, doi:10.1016/0169-555X(95)00044-6, 1996.
- Pelletier, J. D.: How do pediments form? A numerical modeling investigation with comparison to pediments in southern Arizona, USA, *Geological Society of America Bulletin*, 122, 1815–1829, doi:10.1130/B30128.1, 2010a.
- Pelletier, J. D.: Minimizing the grid-resolution dependence of flow routing algorithms for geomorphic applications, *Geomorphology*, 122, 91–98, doi:10.1016/j.geomorph.2010.06.001, 2010a.
- Pelletier, J. D.: Fluvial and slope-wash erosion of soil-mantled landscapes: detachment- or transport-limited? *Earth Surf. Process. Landf.*, 37, 37–51, doi: 10.1002/esp.2187, 2012.
- Pelletier, J. D.: The linkage between hillslope vegetation changes, elevation, and late-Quaternary fluvial-system aggradation in the Mojave Desert revisited, *ESurf*, 2, 455–468, doi: 10.5194/esurf-2-455-2014, 2014.
- Pelletier, J. D., Murray, A. B., Pierce, J. L., Bierman, P. R., Breshears, D. D., Crosby, B. T., Ellis, M., Foufoula-Georgiou, E., Heimsath, A. M., Houser, C., Lancaster, N., Marani, M., Merritts, D. J., Moore, L. J., Pederson, J. L., Poulos, M. J., Rittenour, T. M., Rowland, J. C., Ruggiero, P., Ward, D. J., Wickert, A. D. and Yager, E. M.: Forecasting the response of Earth's surface to future climatic and land use changes: A review of

- methods and research needs. *Earth's Future*, 3, 220–251. doi:10.1002/2014EF000290, 2015.
- Polyakov, V. O., Kimoto, A., Nearing, M. A., Nichols, M. H.: Tracing sediment movement on a semiarid watershed using rare earth elements, *Soil Sci. Soc. Am. J.*, 73, 1559–1565, doi:10.2136/sssaj2008.0378, 2009.
- Portenga, E. W., and Bierman, P. R.: Understanding Earth's eroding surface with ^{10}Be , *GSA Today*, 21(8), 4–10, doi:10.1130/G1111A.1, 2011.
- Ray, R. G., and Fischer, W. A.: Quantitative photography – a geologic research tool, *Photogr. Eng.*, 25, 143–150, 1960.
- Reiners, P. W., Ehlers, T. A., Mitchell, S. G., and Montgomery, D. R.: Coupled spatial variations in precipitation and long-term erosion rates across the Washington Cascades, *Nature*, 426, 645–647, doi:10.1038/nature02111, 2003.
- Renard, K. G., Lane, L. J., Simanton, J. R., Emmerich, W. E., Stone, J. J., Weltz, M. A., Goodrich, D. C., and Yakowitz, D. S.: Agricultural impacts in an arid environment: Walnut Gulch studies, *Hydrol. Sci. Technol.*, 9, 145–190, 1993.
- Ruhe, R. V.: Topographic discontinuities of the Des Moines lobe, *Am. J. Sci.*, 250, 46–56, 1952
- Simanton, J. R., Osterkamp, W. R., and Renard, K. G.: Sediment yield in a semiarid basin; sampling equipment impacts, *Proc. Yokohama, Japan Sym., Sediment problems: strategies for monitoring, prediction and control*, edited by R. F. Hadley and T. Mizuyama, July, IAHS Pub. No. 217, 3–9, 1993.
- Skirvin, S., Kidwell, M., Biedenbender, S., Henley, J. P. King, D., Collins, C. H., Moran, S., and Weltz, M.: Vegetation data, Walnut Gulch Experimental Watershed, Arizona, United States, *Water Resour. Res.*, 44, W05S08, doi:10.1029/2006WR005724, 2008.

[Stone, J. J. Wickre, J., Paige, G. B., Guertin, P., and Gottfried, G.: Post-wildfire runoff and erosion response on grassland and oak woodlands in southeastern Arizona, online document available at https://www.firescience.gov/projects/03-2-3-11/project/03-2-3-11_03_2_3_11_Deliverable_03.pdf, 2003.](https://www.firescience.gov/projects/03-2-3-11/project/03-2-3-11_03_2_3_11_Deliverable_03.pdf)

Tarboton, D. G., Bras, R. L., and Rodriguez-Iturbe, I.: A physical basis for drainage density, *Geomorphology*, 5(1-2), 59–76, doi:10.1016/0169-555X(92)90058-V, 1992.

Torres Acosta, V., Schildgen, T. F., Clarke, B. A., Scherler, D., Bookhagen, B., Wittmann, H., von Blanckenburg, F., Strecker, M. R.: Effect of vegetation cover on millennial-scale landscape denudation rates in East Africa, *Lithosphere*, 7, 408–420, doi:10.1130/L402.1, 2015.

Tucker, G. E., and Bras, R. L.: Hillslope processes, drainage density, and landscape morphology, *Water. Resour. Res.*, 34(10), 2751–2764, doi:10.1029/98WR01474, 1998.

Tucker, G. E., Catani, F., Rinaldo, A., and Bras, R. L.: Statistical analysis of drainage density from digital terrain data, *Geomorphology*, 36(3-4), 187–202, doi:10.1016/S0169-555X(00)00056-8, 2001.

von Blanckenburg, F.: The control mechanisms of erosion and weathering at basin scale from cosmogenic nuclides in river sediment, *Earth Planet. Sci. Lett.*, 237, 462–479, doi:10.1016/j.epsl.2005.06.030, 2005.

Wagner, W. L.: Floristic affinities of Animas Mountains, southwestern New Mexico, M.S. thesis, University of New Mexico, Albuquerque, 1977.

Wagner, J. D. M., J. E. Cole, J. W. Beck, P. J. Patchett, G. M. Henderson, and H. R. Barnett: Moisture variability in the southwestern United States linked to abrupt glacial climate change, *Nature Geoscience*, 3, 110–113, doi:10.1038/ngeo707, 2010.

Weldon, R. J.: The late Cenozoic geology of Cajon Pass: implications for tectonics and sedimentation along the San Andreas fault, Ph.D. Dissertation, California Institute of Technology. Digital Document Available at http://thesis.library.caltech.edu/3283/1/Weldon_rj_1986.pdf, 1986.

Yanites, B. J., and Kesler, S. E.: A climate signal in exhumation patterns revealed by porphyry copper deposits, *Nature Geos.*, 8, 462-465, doi:10.1038/ngeo2429, 2015.

[Yetemen O., Istanbuluoglu, E., and Vivoni, E. R.: The implications of geology, soils, and vegetation on landscape morphology: Inferences from semi-arid basins with complex vegetation patterns in Central New Mexico, USA. *Geomorphology*, 116, 246–263, 2010.](#)

Table 1. Sediment yield data and watershed characteristics. The erosion rate calculation assumes a soil bulk density of 1500 kg m^{-3}

Watershed ID	Predominant vegetation cover type	Contributing area (ha)	Sediment yield ($\text{t ha}^{-1} \text{ yr}^{-1}$)	Erosion rate (mm yr^{-1})
102	Shrub	1.46	2.31	0.154
103	Shrub	3.68	5.66	0.377
104	Shrub	4.53	1.36	0.091
105	Shrub	0.18	0.75	0.050
106	Shrub	0.34	0.80	0.053
112	Grass	1.86	0.07	0.005

Table 2. List of model parameters and values.

Symbol	Units	Description	Value
D	$\text{m}^2 \text{ yr}^{-1}$	topographic diffusivity	1×10^{-3}
S_{hs}	unitless	mean gradient of toe slopes (shrublands)	0.17
S_{hg}	unitless	mean gradient of toe slopes (grasslands)	0.19
l	unitless	exponent of width-area relationship	0.34
k_w	$\text{m}^{0.32}$	coefficient of width-area relationship	0.023
c	unitless	exponent of area-distance relationship	2.5
k_A	$\text{m}^{-0.5}$	coefficient of area-distance relationship	0.3
p	unitless	exponent of sediment-flux-area relationship	1.44
k_{Qss}	$\text{m}^{1.56} \text{ yr}^{-1}$	sediment transport coefficient (shrublands)	2×10^{-6}
k_{Qsg}	$\text{m}^{1.56} \text{ yr}^{-1}$	sediment transport coefficient (grasslands)	6×10^{-8}

Table 3. Measured (from DEM analysis) and predicted values (from equations (11)&(12)) for the mean distance from divides to valley bottoms in shrublands (x_s) and grasslands (x_g).

	Measured (m)	Predicted (m)
x_s	18	16
x_g	60	66




Figure 1. Maps of the bedrock geology and geomorphology of Walnut Gulch Experimental Watershed (WGEW). (A) Bedrock geology from Osterkamp (2008). Rectangle identifies the portion of WGEW that is the focus of this study. (B) Geomorphic map from Osterkamp (2008). The boundary between the Dissected Whetstone Pediment and Upper Whetstone Pediment marks a key transition in landscape morphology, soil type, and vegetation cover (Fig. 2).





Figure 2. Relationships among landscape morphology and vegetation cover in the study area. (A) Shaded relief image of the topography, illustrating the significant increase in hillslope-scale relief from the western to the eastern portion of the study area. (B) Grayscale map of topographic curvature (i.e., Laplacian), demonstrating generally lower absolute hillslope curvatures (i.e., more gray) in the western (shrubland) portion of the study area relative to the eastern (grassland) portion (more black). (C) Vegetation map, after Skirvin et al. (2008), identifying the areas that are primarily shrubland, grassland, and transitional between the two.






Figure 3. Drainage density is higher in shrubland areas than in grassland areas of the study site. Shaded relief images of representative 1.6 km x 1.6 km areas of (A) shrublands, including Lucky Hills watersheds 102-106 and (B) grasslands, including Kendall watershed 112. (C)&(D) Images of the drainage network identified using the Pelletier (2013) algorithm for the areas shown in (A)&(B), respectively. (E)&(F) Grayscale maps of the distance along flow lines from divides to the valley bottom. The mean of the mapped values in (E) at valley heads is 18 m while the mean of the mapped value in (F) at valley heads is 60 m.




Figure 4. Detailed shaded-relief images (locations shown in Fig. 3) illustrating the presence of parallel hillslope rills and gullies in the shrubland areas (shown in A). Grassland areas (shown in B) have fewer such features.




Figure 5. Schematic figure of a valley head. The profile shown along A-A' is the cross-section of the valley head where colluvial sediment flux from hillslopes of mean gradient S_h enter a segment of width w . The requirement that fluvial erosion rates must be greater than colluvial

deposition rates at the valley head provides a quantitative criterion for predicting the drainage density of landscapes.




Figure 6. Plot of mean hillslope relief as a function of elevation, illustrating the marked increase in relief above elevations of approximately 1430 m a.s.l. in the study area. Each data point represents the mean hillslope relief in 10-m-wide elevation bins.




Figure 7. Plots of contributing area versus mean distance from the divide for the shrubland (approximated as the portion of the study area below 1430 m a.s.l.) and grassland areas (above 1430 m). The dashed line plots the piece-wise power-law relationship (equation (9)) exhibited by the data.




Figure 8. Plots of mean topographic curvature as a function of contributing area (distance from divide also shown along x axis using data in Fig. 7). Topographic curvatures are similar in shrublands and grasslands at small and large contributing areas, with a minimum of 0.03 m^{-1} near divides. Within a range of contributing areas from ~ 10 to $\sim 300 \text{ m}^2$ the data show significant differences in mean curvature between shrublands and grasslands.




Figure 9. Plot of mean slope versus contributing area. Shrublands and grasslands both show a power-law relationship, with deviations from power-law behavior occurring at larger spatial scales in grasslands relative to shrublands.




Figure 10.




Figure 9. Plot of the decadal time-scale volumetric sediment fluxes in shrublands measured at the five watersheds of the Lucky Hills as a function of contributing area. The straight line is the result of a least-squares regression to the logarithms of both sides of equation (7), from which the values of $k_{Q_{ss}}$ and p were constrained.




Figure 410. Plots of the total erosion rate, E , the fluvial/slope-wash erosion rate, E_f , and the colluvial deposition rate, $-E_c$, as a function of distance along flow lines from divides, x . The values of x_s and x_g (where $E = E_f + E_c$) are also shown.




Figure 421. Plots of the mean longitudinal profile of hillslopes and valley bottoms in shrubland and grassland areas, constructed by integrating the mean curvature data in Figure 8.