

Response to review by Dr. Alan Trenhaile

We would like to thank Dr. Alan Trenhaile for his positive and constructive comments. Below are our responses to all of these comments including details of any changes we have made to the manuscript guided by the reviewer's suggestions. We have numbered the reviewer comments and they are coloured black, while our responses are coloured blue, and italicised where we are quoting directly from the revised manuscript.

5 1. Cosmogenic dating was initially greeted by rock coast workers, as a quasi-panacea, a long awaited solution to the question, that had arisen over more than a century, regarding the age of shore platforms, and especially whether they are primarily contemporary (Holocene) or inherited (from previous interglacials when sea level was similar to today's landforms. Related to this question and of more immediate practical concern was its apparent ability to provide reliable data on rates of cliff recession over lengthy periods during the Holocene, and to test predictive models and better predict the effects of climate change, and especially rising sea level. There have been a few useful ^{10}Be dating applications. They include the work of Choi et al. (2012), who suggested that the seaward portions of shore platforms in Korea are up to about 150 thousand years old, and that the modern platform is cutting into its interglacial predecessor. Conversely, Regard et al., (2012) opined that in northern France, the mean rate of cliff retreat since the mid-Holocene has been 15 $11\text{--}13 \text{ cm yr}^{-1}$, which would have been sufficient to create 'contemporary' intertidal shore platforms hundreds of metres in width. Despite these valuable contributions, I would contend that cosmogenic nuclide analysis, whether for dating landforms or quantifying their rates of erosion has failed, as yet, to revolutionize the study of rock coasts.

10 The two studies discussed by the reviewer above (plus another nice paper by Rogers et al. (2012)) have done an excellent job in demonstrating that there is significant potential for cosmogenic radionuclides (CRNs) to yield evidence for long-term processes and rates in rocky coast settings. As the reviewer points out, measurement of cosmogenic radionuclides has not been able to "revolutionise the study of rock coasts", nor would we expect it to. Rather, they provide an additional tool in the coastal geomorphologist's repertoire to help understand the rates and processes by which cliffted coasts and shore platforms have evolved in the past, and notably, over timescales longer than those at which observational data has been collected. We were motivated by reviewer's own conclusions when reviewing climate change impacts on rocky coasts that "... we must... acquire more precise and longer records of erosion rates and other expressions of process efficacy and their relationship to prevailing morphogenic conditions" (Trenhaile, 2014, p. 15). It remains to be seen whether cosmogenic radionuclides can improve the precision of erosion rate estimates, but it certainly has the potential to provide longer records (over centennial to millennial timescales) in erosive environments in which there is no alternative evidence of a previous state.

15 20 25 30 35 2. Given this, the present paper, which discusses and models the effect of several of the factors and assumptions that constrain the use of cosmogenic analysis on rock coasts, represents a welcome addition to the literature and one which will, no doubt, improve our ability to understand the longterm development of rock coasts. Nevertheless, this paper once again points out the inherent limitations of cosmogenic analysis on rocky coasts for which we have little longterm data on such factors as downwearing and backwearing rates (erosion in the vertical and horizontal planes, respectively) and historical and short-term (storm and calm condition) variations in sediment thickness, type, and extent.

30 35 40 45 We are pleased that the reviewer finds our work to be "a welcome addition to the literature" and we hope that it can "improve our ability to understand the long-term development of rock coasts" through the application of CRNs in such environments. We felt that it was a critical step to further explore assumptions that may be required in order for future studies to interpret concentrations of CRNs in terms of process rates on shore platforms so that sampling strategies might be better informed to test these assumptions. This is particularly important given the expense of making CRN measurements that may limit the number of samples that can be acquired in a single study. As an example our results suggest that there is the potential to estimate rates of block removal on stepped platforms through high density sampling and this would be an exciting avenue for future work. We have not made any modifications to the paper based on these opening statements as we feel the context and our aims are already set out clearly in the manuscript, if the reviewer had any specific modifications to our introductory discussion in mind then we would welcome further comment.

3. My main criticism of this paper is the almost complete neglect of downwearing by wearing processes, including by tidal wetting and drying and salt weathering.

We acknowledge this criticism and its importance, and are not suggesting that these processes are not important mechanisms of downwear on shore platforms. Previous studies that have tried to quantify cliff retreat rates have relied on the assumption of steady-state retreat whereby a constant shore platform morphology is translated landward through time. In this study we have applied a dynamic morphological evolution model, guided by the reviewer's own work (e.g. Trenhaile, 2000), that is capable of generating the sorts of behaviour we wished to explore (e.g. shore platforms that widen with time). We have thus gone further than any previous studies of cosmogenic radionuclides in coastal settings when considering the morphological evolution of the coast. All numerical models are abstractions of the real world at some level but our experiments allow us to explore a number of interesting behaviours such as both continuous and discontinuous distribution of erosion (gradual downwear vs block removal). Such a model, at first order, broadly recreates the patterns of downwear observed from short-term monitoring data, with downwear decreasing with decreasing elevation on the shore platform (see also response to reviewer comment #5).

4. The paper notes that models have indicated that shore platforms and cliffs trend towards a morphological steady state. There are certainly many examples of steady states in model predictions with constant sea level, although when one considers the effect of changing relative sea level, simulated longterm platform development is more complex (see for example: Trenhaile, A. S. 2001: Modeling the Quaternary evolution of shore platforms and erosional continental shelves . Earth Surface Processes and Landforms 26, pp. 1103-28, and Trenhaile, A. S. 2014. Modelling the effect of Pliocene-Quaternary changes in sea level on stable and tectonically active land masses. Earth Surface Processes and Landforms 39, 1221-35).

In this paper we do not explore the history of shore platform development throughout the quaternary but rather focus on the development of shore platforms during the Holocene (more specifically, the last 7-8 thousand years, the time over which eustatic sea level has been relatively stable). Where shore platforms are inherited features formed perhaps during a previous interglacial period, subsequent exposure would result in CRN concentrations substantially higher than those from contemporary platforms, as demonstrated by Choi et al. (2012). We have clarified this in the introduction by adding a more detailed description of the findings of Choi et al. (2012) and by clarifying that we are only concerned in this study with Holocene platforms stating:

"Some shore platforms may have formed during the Holocene and thus be entirely contemporaneous features, whilst others, particularly wide platforms in resistant lithologies, have been interpreted as inherited features, formed during previous sea level high stands and reoccupied during the Holocene (e.g. Blanco Chao et al., 2003). Evidence for the antiquity of shore platforms may be revealed by CRNs. For example, Choi et al. (2012) measured ^{10}Be concentrations in shore platforms cut into resistant lithologies on the Korean coast. High concentrations were consistent with modelled ages extending back as far as 142 ka, and importantly, these ages did not correct for weathering and erosion and thus should be considered as minimum ages. Thus Choi et al. (2012) conclude that these platforms are at least partially inherited features, originated in the Pleistocene."

"In this study we quantified the sensitivity of platform CRN concentrations to topographic shielding, various processes of platform erosion/downwear, the presence/absence of beach cover, and transience in shore profile evolution. In doing so we consider only the development of shore platforms during the Holocene, over the timescale during which eustatic sea level has been relatively stable (7 ka to present). We addressed this with a numerical model coupling cross-shore coastal evolution and ^{10}Be production to explore the potential for quantifying coastal retreat rates from ^{10}Be concentration measurements."

5. The paper quotes our work (Porter et al., 2010) in eastern Canada (on page 15 of the manuscript), but essentially ignores the relevant conclusions that relate to spatial variations across the intertidal zone. This paper showed, based on laboratory experiments lasting several years and about 2000 rock samples, together with about 200 transverse micro-erosion metre stations in the field, that, while weathering downwearing rates (isolated in the laboratory experiments) tend to be a maximum in the upper intertidal zone, there is no clear pattern in the field, possibly because of the effect of other erosional mechanisms with different elevational efficiencies.

5 We are grateful to the reviewer for highlighting their findings, and apologise that we did not do justice to this excellent piece of work that explores weathering and erosion processes on shore platforms both in the field and through controlled experiments. The lack of a “clear pattern in the field” meant that we were not able to use the data from their study to parameterise our model, however the model does reproduce the general trend they observed, that downwearing tends to be highest in the upper intertidal zone, declining offshore. Our intention in this study has been to explore the general case, parsimoniously, to better understand the sensitivity of CRN concentrations to factors such as the potential for non-uniform distribution of downwearing, and our analyses are by no means exhaustive. Future work that measure CRN concentrations at specific sites where there are good observational datasets will be a logical and important next step.

10 That said, there is a distinct scaling issue between their experiments which take place over at most a 6 year period, and the centennial to millennial timeframe over which we are required to model coastal evolution in order to explore the controls on the build-up of ^{10}Be . We do not feel it would be appropriate to take the results of a single site and extrapolate over several millennia for the purposes of this study, although site specific investigation using CRNs would be well advised to place their findings in the context of such observations. Nevertheless, Stephenson et al. (2012) found that rates of platform erosion averaged over a 32 year period were not significantly different from the two year record initially analysed at the same sites, concluding that low magnitude, high frequency events drive platform lowering at that site. It is not yet clear whether such short-term downwear rates can be extrapolated at other sites, in other lithologies, for different tidal regimes and different erosion processes. Ongoing work with micro erosion metres, combined with our growing ability to derive high resolution 3D surface models with LiDAR and photogrammetry will help to address this matter further and we would welcome such research.

15 20 Given all this, we have not carried out any additional experiments in light of the reviewer’s criticism. If the reviewer has any specific requirements he would like to see then we would welcome them. Rather, we have expanded our discussion of these issues in the paper to highlight that there is still plenty to be done as detailed below.

25 At the end of the introduction we have added:

“In this study we quantified the sensitivity of platform CRN concentrations to topographic shielding, various processes of platform erosion/downwear, the presence/absence of beach cover, and transience in shore profile evolution. Our intention is to explore the general case parsimoniously to better understand controls on ^{10}Be concentrations, rather than to make accurate predictions for any specific field site.”

30 In justifying the development of a dynamic morphological model we now state:

35 “Moreover, micro-erosion meter measurements of platform downwear suggest that downwear is not uniform across the shore profile, but tends to be faster in the upper intertidal zone and decline with depth (Porter et al., 2010), and to explore the influence of such a distribution on ^{10}Be concentrations, a dynamic morphological model was required.”

40 In the discussion we have expanded our brief mention of downwear measurements to say:

35 “Accounting for the distribution of downwear would be aided by data on the distribution of downwear rates from micro-erosion-meter measurements (e.g. Robinson, 1977; Porter et al., 2010) and there is evidence that rates measured over the short term (1-2 years) are consistent with rates measured over three decades (Stephenson et al., 2010, 2012). Nevertheless, it is still not clear whether it is appropriate to extrapolate these rates over centennial to millennial timescales required to accumulate measurable concentrations of ^{10}Be , nor is it yet clear whether this result is consistent across different shore platforms around the world. Future studies that measure the distribution of CRN concentrations (both across the shore platform and at depth) at sites with long records of observed downwearing rates would be an important next step in this line of enquiry.”

45 6. Whether there are weathering patterns or not, mean rates of downwearing recorded by numerous workers in different environments and on different types of rock, which generally range from almost 0 up to a few millimeters per year (and hence significant lowering per millennium), surely have an important effect on predicted erosion rates. The reliance on models of platform development (whether parallel or declining slope retreat) to represent rates of surface lowering (as opposed to field measurements and micro-erosion metre data) remains an inherent weakness in attempts to apply cosmogenic techniques to rocky coasts.

5 The reviewer has highlighted an interesting issue in the application of cosmogenic radionuclides to rocky coasts that will need to be addressed in future studies, namely the disconnect between short-term observations and long-term platform evolution, as we have already mentioned above. Our exploration of the sensitivity of CRN concentrations is not site specific and thus the use of spot measurements from a particular location, extrapolated back throughout most of the

10 Holocene does not seem appropriate. It strikes us that, despite the findings of Stephenson et al (2010; 2012) that rates observed over a few years are similar to those over a few decades for two field sites, the extrapolation of short-term measurements (3 decades at best) of shore platform erosion over millennial timescales is also inherently weak. Future work measuring ^{10}Be concentrations at sites with good long-term field records of surface lowering would be an excellent future extension of the application of CRNs to rock coasts, and we have called for this in the paper (see our response to reviewer comment #5). Moreover, our growing capacity to observe changes in platform morphology in a distributed way using terrestrial LiDAR and photogrammetry in order to identify individual erosion events will lead to better understanding of the efficacy and distribution of platform downwear processes, and we commend the reviewer for pioneering these sorts of investigations.

15 As the reviewer points out both here and in his final comment, we do not yet understand the spatial distribution of weathering on shore platforms.

7. The traditional distinction (classification) between sloping and subhorizontal (or to use Sunamura's terminology, type A and B platforms) belies the fact that there is a continuous spectrum of forms with gradients that reflect tidal range and rock hardness (resistance), as well as possibly sediment grain size and the effect of Holocene sea levels higher than today.

20 We thank the reviewer for highlighting that such a classification is an over simplification and we agree that these two forms are effectively end-members of a much wider array of shore platform morphologies. Our experiments were intended to explore such intermediate platform morphologies, we do not limit ourselves to type-A and type-B platforms. However the beginning of section 2 may have been misleading on this point. We now begin this section by saying:

25 "Cliffed, rocky coasts, are commonly fronted by shore platforms that have previously been classified into two types (Sunamura, 1992). Type-A platforms are characterised by a gently sloping erosional platform surface extending offshore beyond maximum low water. Type-B platforms are shallow gradient to sub-horizontal and terminate at their seaward edge at maximum low water through a scarp (Figure 2a). Numerical models of shore platform evolution have successfully recreated both of these end-member morphologies, but have revealed that there are a range of other possible morphologies in between, for example sloping platforms terminating in a scarp (e.g. Trenhaile, 2000; Walkden and Hall, 2005; Matsumoto et al., 2016)."

30 8. It is not clear to me why, on line 12 or page 4, abrasion is assumed to be dominant with a steady state that assumes platform downwearing is gradual and constant (especially since weathering is not considered). Sediment on shore platforms accumulates preferentially at the cliff foot and in structural depressions on the platform. In the latter case, abrasion rates with vary enormously across the intertidal zone in an essentially random manner. Abrasion towards the rear of the platform is spread over the intertidal surface as the cliff retreats, but constant abrasion assumes constant wave and sea level conditions as well as constant beach types and amounts.

35 We thank the reviewer for highlighting the overly simple nature of this explanation. Our assumption is that in the case of steady-state retreat that platform lowering is constant and gradual across the shore platform. Over centennial to millennial timescales we might reasonably expect that the sorts of processes the reviewer describes here will tend to occur uniformly over a platform. Indeed, the existence of generally flat or planar platforms in some settings suggest this must be the case.

40 We now state:

"A steady state approach assumes platform downwear is gradual and constant across the entire profile, and proportional to the rate of cliff retreat. We are therefore assuming that the combined mechanical and chemical processes that can cause shore platform lowering culminate to constant and gradual downwear in this case."

45 9. Page 4, line 20, many models do/did not emphasize erosion at the water surface but rather erosion due to wave generated bottom currents (which is only effective in clays and other 'soft' rocks).

We now state that: "We developed a simple numerical model for shore profile evolution (the ROck and BOttom COastal Profile [RoBoCoP] Model), broadly similar to those of Sunamura (1992), Anderson et al. (1999) and Trenhaile (2000). These models assume..."

5 10. An alternative to the Bruun Rule, which has been criticized by several workers and in any case is not designed for beaches with rigid (shore platform) foundations, is given by: Trenhaile, A. S. 2004. Modeling the accumulation and dynamics of beaches on shore platforms. *Marine Geology* 206, 55-72.

We thank the reviewer to drawing our attention to this alternative model for beach profile shape. We note that our model is not employing the criticised "Bruun Rule" that relates rates of shoreline retreat to relative sea level rise, but rather we use a "Bruun Profile" as a description of the shape of the beach profile. We chose a Bruun profile for its computational simplicity. Our goal was not to explicitly replicate beach behaviour but rather to explore the influence that shielding by sediment cover has on the accumulation of ^{10}Be in shore platforms. Thus we took a parsimonious approach to representing beach morphology. We note that Trenhaile (2004) takes a similarly simplistic approach, representing the shoreface as a planar slope of constant gradient, though that gradient varies depending on wave conditions and grain size.

15 11. The paper needs some proof-reading. There are other examples that could be given but note 'ditributed' on line 14 of page 10 and on line 15 it should be 'adjacent to the cliff'.

We apologise for the sloppy presentation despite our endeavours to catch all typological errors, and thank the reviewer for highlighting any such mistakes when they have occurred. We have corrected the blunders explicitly pointed out by all reviewers, and our redraft has undergone thorough proof reading by all authors.

20 12. On lines 12 to 13 on page 11 it is claimed that there is no downwearing on top of flat-topped steps. This is wrong - weathering certainly lowers the surface until (and in some cases before) it is eliminated by step retreat.

This is not a claim since we are referring specifically to the model simulations: "there is no vertical downwear in these simulations". Our simulations exploring block removal processes are exploratory and can be considered end-member scenarios. We sought to understand the influence that block removal processes might have on ^{10}Be concentrations at the surface. Were there some vertical downwear then we would expect the gradients in ^{10}Be concentrations across the flat-topped steps to be lower, i.e. concentrations would not increase as rapidly in an offshore direction. The result presented in this section demonstrates that block removal processes will cause deviations from the expected "hump" shaped distribution, but the general form of the hump still remains, and this is encouraging for potential future applications.

25 30 13. Lines 10 to 13 on page 13 emphasize errors due to differences in predicted downwearing rates according to the two evolutionary models. The reality is likely to be worse owing to weathering induced downwearing (with spatial patterns, if any, that we do not yet understand).

Yes, the point we are trying to make is that the more rapid the rate of downwearing, the more ^{10}Be -laden rock is removed from the platform surface and the lower the subsequent production rate will be; therefore, the lower the CRN concentrations will be. This will lead to an over prediction of cliff retreat rates if not factored in (i.e. if one were to blindly assume steady-state retreat of a planar shore platform). If weathering processes are leading to more rapid downwear then this problem will be amplified. We have stressed in the discussion that measuring CRN concentrations at sites with good records of downwear will be an important next step in developing our ability to estimate rates of cliff retreat, or platform downwear using CRNs. We have not added to the results section highlighted by the reviewer, but instead build on this issue in the discussion:

35 40 "In scenario (ii) cliff retreat rates tended toward a constant rate with the result that the distribution of ^{10}Be was approximately constant through time (Figure 14). The concentrations of ^{10}Be were not consistent with a steady state evolution scenario in which the platform gradient is fixed, which predicted roughly twice the amount of ^{10}Be for a particular position on the platform. The difference can be explained by the dissimilar platform morphology brought about by uneven distribution of platform downwear in the transient model simulations. Greater rates of downwear in the inter-tidal zone

lower the platform more rapidly in the near-shore, which removes ^{10}Be -laden rock and results in deeper water in the near-shore (and therefore reduced ^{10}Be production) relative to the steady state model runs that assume constant and uniform downwear. These differences may be exacerbated when taking into account the spatial distribution of other processes such as weathering, the spatial distribution of which is currently poorly understood (Porter et al., 2010)."

5 Response to review by Dr. Mark Dickson

We are grateful to Dr. Mark Dickson for his positive and constructive comments, and extend our thanks also to Hironori Matsumoto for his contributions to the review. We welcome the inclusion of aspiring PhD students in the peer review process. Below we detail our responses to all of these comments, highlighting any changes we have made to the manuscript guided by the reviewer's suggestions. We have numbered the reviewer comments and they are coloured black, while our responses are 10 coloured blue, and italicised where we are quoting directly from the revised manuscript.

1. This is a welcome addition to a rather slowly emerging literature on the application of cosmogenic dating to inform rocky shore evolution studies. The proposed model usefully extends the work described by Regard et al. (2012) and includes I think the majority of the factors required. In doing so, the paper highlights that the use of this method to determine 15 erosion rates on rocky shores, while conceptually simple, in practice is rather more complex. Nevertheless, the lack of quantitative measurements of rates of change has been a long-standing limitation in our field, so efforts in the direction of this paper are welcome.

We thank the reviewer for these positive comments and hope this work can be useful for guiding future applications of cosmogenic radionuclides in rock coast settings.

2. Below are some comments that the authors might wish to consider during any revision. Please note that these comments 20 integrate thoughts from Hironori Matsumoto who is currently making further developments to a rocky shore profile evolution model (Matsumoto, H., Dickson, M. E., and Kench, P. S.: An exploratory numerical model of rocky shore profile evolution, *Geomorphology*, 268, 98–109, doi:10.1016/j.geomorph.2016.05.017, 2016.)

Thank you for bringing this manuscript to our attention, we had already cited it in the manuscript. We thank Hironori Matsumoto for his contributions to the review, and more generally we welcome the consultation of PhD students in an 25 open review process.

3. Can you clarify how the model considers platform slope? Does it calculate the cliff toe position with a fixed platform geometry, similar to previous tide-less models (Sunamura, 1975)? Perhaps not, because later, in section 5.5, there is 30 reference to the gradient of the shore platform decreasing through time, as the platform widens. There is also reference to the emergence of a 'stepped' platform - what is the reason for this? Ultimately it is a little unclear how exactly the geometry is handled. The text states that the model is similar to the existing models of Trenhaile (2000), Walkden and hall (2005), and Matsumoto et al (2016), which all consider tide and vertical components. It would be useful to expand and clarify similarities and differences in that regard.

We apologise that the details of the morphological development were unclear and are pleased to have the opportunity to 35 clarify these points. There are two morphological models used in this study. The first is a steady-state model in which a constant shore profile characterised by a fixed platform gradient is translated landward through time, tracking the elevation of mean sea level. Therefore, in these steady-state scenarios, tides do not influence the morphological evolution of the platform. However tides still modify the production of ^{10}Be on the platform due to water shielding. In the second 40 modelling approach, a dynamic morphological model is used, and it is this model that is broadly similar to Trenhaile (2000) and Matsumoto et al. (2016). The dynamic model calculates the point of wave breaking and therefore the width of the surf zone, which are dependent on the instantaneous mean water level which is set by the tides. Horizontal erosion at the water level is proportional to the delivery of wave energy (equation 1), parameterised as the height of the breaking wave once it has crossed the surf zone (equation 4). The platform is also eroded at depth below the water line, with the amount of erosion declining exponentially with depth. Stepped platforms can develop because the total amount of

erosion integrated over a tidal cycle has two maxima as a function of elevation, one just below the upper tidal limit, and one just above the lower tidal limit (similar to Figure 1 in Trenhaile, 2000).

We have restructured the beginning of section 2 to be more explicit about the different approaches taken to morphological modelling:

5 *“Cliffed, rocky coasts, are commonly fronted by shore platforms that have previously been classified into two types (Sunamura, 1992). Type-A platforms are characterised by a gently sloping erosional platform surface extending offshore beyond maximum low water. Type-B platforms are shallow gradient to sub-horizontal and terminate at their seaward edge at maximum low water through a scarp (Figure 2a). Numerical models of shore platform evolution have successfully recreated both of these endmember morphologies, but have revealed that there are a range of other possible morphologies in between, for example sloping platforms terminating in a scarp (e.g. Trenhaile, 2000; Walkden and Hall, 2005; Matsumoto et al., 2016).”*

10 *“Numerical models of platform evolution demonstrate that shore platforms and adjacent sea cliffs tend towards a morphological steady state. Under such conditions the morphology may reflect the combination of RSL change, tides and wave energy availability. However, the assumption of steady state retreat may not always be applicable (Dickson et al., 2013). We expected the style of platform evolution to be important for the distribution of ^{10}Be across a shore platform. Moreover, micro-erosion meter measurements of platform downwear suggest that downwear is not uniform across the shore profile, but tends to be faster in the upper inter-tidal zone and decline with depth (Porter et al., 2010), and to explore the influence of such a distribution on ^{10}Be concentrations, a dynamic morphological model was required.”*

15 *“Therefore, two different morphological models are used in this study. The first assumes steady state evolution of the shore profile, such that coastal morphology does not change its form through time, and a constant cliff-platform geometry is translated landward through time. The second model is a dynamic shore platform evolution model similar to that of Trenhaile (2000) that can reproduce a range of platform geometries. These two approaches are described in more detail in the subsequent sections.”*

20 4. Did the authors consider prospects for using the model to test/discuss factors that affect ^{10}Be concentrations on other types of shore platform geometry beyond the sloping (type-A) platform investigated? The paper refers to type-B platforms, raising the question as to whether the model adds any new knowledge of likely differences in concentrations across different possible platform geometries.

25 Please see also our response to Alan Trenhaile’s similar comment (response #7 in the response to Alan Trenhaile’s review). We are not exclusively considering type-A and type-B platforms, these are end-member cases, and the dynamic morphological model used in this study produces alternative platform geometries such as stepped platforms. We now write:

30 *“Cliffed, rocky coasts, are commonly fronted by shore platforms that have previously been classified into two types (Sunamura, 1992). Type-A platforms are characterised by a gently sloping erosional platform surface extending offshore beyond maximum low water. Type-B platforms are shallow gradient to sub-horizontal and terminate at their seaward edge at maximum low water through a scarp (Figure 2a). Numerical models of shore platform evolution have successfully recreated both of these end-member morphologies, but have revealed that there are a range of other possible morphologies in between, for example sloping platforms terminating in a scarp (e.g. Trenhaile, 2000; Walkden and Hall, 2005; Matsumoto et al., 2016).”*

35 *“Therefore, two different morphological models are used in this study. The first assumes steady state evolution of the shore profile, such that coastal morphology does not change its form through time, and a constant cliff-platform geometry is translated landward through time. The second model is a dynamic shore platform evolution model similar to that of Trenhaile (2000) that can reproduce a range of platform geometries. These two approaches are described in more detail in the subsequent sections.”*

40 5. The model considers topographic shielding in the case of a constant cliff height. A further exploration of interest would be to consider a slowly increasing cliff height. What comes to mind is the example of progressive cliffing into hillslopes

rounded over long glacial periods. This could introduce further complications for ^{10}Be concentrations, because erosion into cliffs of increasing height might progressively increase beach thickness (5.4.1)... assuming no gradient in alongshore sediment flux. The number of potential model scenarios can quickly increase, but it would be interesting to have a somewhat expanded discussion of this factor.

5 On the first part of your comment, we agree that it is unlikely on many coastlines that cliff heights have been constant during platform development but will depend on the morphology of the landscape that is being eroded into. Generally, topographic shielding only has a minor influence on CRN concentrations, since the shielding factors increase rapidly and non-linearly toward unity moving away from the cliff (Figure 4). To illustrate this we have plotted below concentrations for a simple case of steady-state platform retreat at 10 cm yr^{-1} for a 1/100 gradient platform for cliff heights of 0, 25 and
10 50 m. The difference in peak concentrations between the 0 and 50m cliff are < 10%. A gradual changing of cliff height would therefore not make a significant difference to the predicted concentrations.

With the second part of your comment we acknowledge that we offered little discussion of the feedbacks between cliffs and beaches. It was intentional at this stage not to perform experiments exploring these feedbacks because as yet they are poorly represented in existing morphological models, particularly when considering alongshore transport.

15 6. There is some interesting discussion on the implications of platform downwear, but I agree with the other reviewer that this aspect of the modelling could be improved - he makes a number of points to consider in that regard which I have not expanded on here.

We have made a number of changes to the manuscript to address Alan Trenhaile's comments and point the reviewer to these for details (see responses to Alan Trenhaile's review: #3, #5, #6, #8 and #13). We have not changed our modelling
20 approach to include more specific processes related to downwear as we think this would distract from our intention to provide a heuristic exploration of the controls on shore platform ^{10}Be concentrations.

7. In concluding, is it possible to tabulate or summarise somehow the relative sensitivities of the different factors?

The reviewer has a good suggestion in theory; however, it is not practical to summarize the sensitivity of so many different factors in a simple and sensible manner. We investigated the possibility by conducting some additional experiments in
25 which model parameters were systematically varied and superimposed plots of ^{10}Be concentrations but found the relative sensitivities were still unclear. Our conclusion from this is that such a simple tabulation is not possible because the sensitivity of the many parameters cannot be generalized in a meaningful way. Instead, the modelling that we present will allow future workers to conduct such a sensitivity analysis for a specific platform.

8. Please double check that you have the exponent written correctly in equation 5 (i.e. '+' $hw(t)$)?

30 This should have been negative, thank you for noticing. We have removed the brackets so that the signs are now correct.

9. Please double check the wording of the last sentence on p7 (upper / lower?)

These were correct but we see how it might have been confusing. These changes are relative to a scenario with no tides at all. The upper platform experiences periodic submergence and emergence during a tidal cycle, but the net effect is a reduction in the overall ^{10}Be production, whilst on the lower platform the net effect is an increase in the overall ^{10}Be production. We have rewritten this section to clarify:

"Tides modify production in the platform by varying water depth hw and intermittently submerging and exposing the platform sub-aerially. Relative to a scenario with no tidal variation, Regard et al. (2012) demonstrated that tides have a net effect to reduce ^{10}Be production in the upper inter-tidal platform due to periodic platform submergence that reduces the net cosmic ray flux received, while ^{10}Be production in the lower platform increases due to periodic exposure."

40 10. two "and"s p4 line 28 - Clarify what is meant by significant (para 8, line 16)

Corrected

11. Mis-spelling of fund[e]mentally

Corrected

12. Grammar problem para 9 line 24

We were unable to find this error. We contacted the reviewer directly who was not able to clarify an error. If this mistake exists and has not been corrected by our redrafting and proof reading, we hope it will be caught during typesetting.

13. no full stop before Failure p15 l12

5 **Corrected**

14. Choi2012 p16

Corrected

References

Anderson, R.S., Densmore, A.L., and Ellis, M.A., 1999, The generation and degradation of marine terraces: *Basin Research*, v. 11, no. 1, p. 7–19, doi: 10.1046/j.1365-2117.1999.00085.x.

10 Chao, R.B., Casais, M.C., Cortizas, A.M., Alberti, A.P., and Trenhaile, A.S., 2003, Evolution and inheritance of a rock coast: Western Galicia, Northwestern Spain: *Earth Surface Processes and Landforms*, v. 28, no. 7, p. 757–775, doi: 10.1002/esp.496.

Choi, K.H., Seong, Y.B., Jung, P.M., and Lee, S.Y., 2012, Using Cosmogenic ^{10}Be Dating to Unravel the Antiquity of a Rocky 15 Shore Platform on the West Coast of Korea: *Journal of Coastal Research*, v. 282, no. 3, p. 641–657, doi: 10.2112/JCOASTRES-D-11-00087.1.

Dickson, M.E., Ogawa, H., Kench, P.S., and Hutchinson, A., 2013, Sea-cliff retreat and shore platform widening: Steady-state equilibrium? *Earth Surface Processes and Landforms*, v. 38, no. 9, p. 1046–1048, doi: 10.1002/esp.3422.

Lazarus, E.D., McNamara, D.E., Smith, M.D., Gopalakrishnan, S., and Murray, A.B., 2011, Emergent behavior in a coupled 20 economic and coastline model for beach nourishment: *Nonlinear Processes in Geophysics*, v. 18, no. 6, p. 989–999, doi: 10.5194/npg-18-989-2011.

Matsumoto, H., Dickson, M.E., and Kench, P.S., 2016, An exploratory numerical model of rocky shore profile evolution: *Geomorphology*, v. 268, p. 98–109, doi: 10.1016/j.geomorph.2016.05.017.

Porter, N.J., Trenhaile, A.S., Prestanski, K., and Kanyaya, J.I., 2010, Patterns of surface downwearing on shore platforms in 25 eastern Canada: *Earth Surface Processes and Landforms*, v. 35, no. 15, p. 1793–1810, doi: 10.1002/esp.2018.

Regard, V., Dewez, T., Bourlais, D.L., Anderson, R.S., Duperret, A., Costa, S., Leanni, L., Lasseur, E., Pedoja, K., and Maillet, G.M., 2012, Late Holocene seacliff retreat recorded by ^{10}Be profiles across a coastal platform: Theory and example from the English Channel: *Quaternary Geochronology*, v. 11, p. 87–97, doi: 10.1016/j.quageo.2012.02.027.

Robinson, L.A., 1977, Erosive processes on the shore platform of northeast Yorkshire, England: *Marine Geology*, v. 23, no. 4, 30 p. 339–361, doi: 10.1016/0025-3227(77)90038-X.

Rogers, H.E., Swanson, T.W., and Stone, J.O., 2012, Long-term shoreline retreat rates on Whidbey Island, Washington, USA: *Quaternary Research*, v. 78, no. 2, p. 315–322, doi: 10.1016/j.yqres.2012.06.001.

Stephenson, W.J., Kirk, R.M., Hemmingsen, S.A., and Hemmingsen, M.A., 2010, Decadal scale micro erosion rates on shore 35 platforms: *Geomorphology*, v. 114, no. 1–2, p. 22–29, doi: 10.1016/j.geomorph.2008.10.013.

Stephenson, W.J., Kirk, R.M., Kennedy, D.M., Finlayson, B.L., and Chen, Z., 2012, Long term shore platform surface lowering rates: Revisiting Gill and Lang after 32 years: *Marine Geology*, v. 299–302, p. 90–95, doi: 10.1016/j.margeo.2012.01.005.

Sunamura, T., 1975, A Laboratory Study of Wave-Cut Platform Formation: *The Journal of Geology*, v. 83, no. 3, p. 389–397, 40 doi: 10.1086/628101.

Sunamura, T., 1992, *Geomorphology of Rocky Coasts*: John Wiley and Sons Ltd.

Trenhaile, A.S., 2014a, Climate change and its impact on rock coasts: *Geological Society, London, Memoirs*, v. 40, no. 1, p. 7–17, doi: 10.1144/M40.2.

Trenhaile, A.S., 2004, Modeling the accumulation and dynamics of beaches on shore platforms: *Marine Geology*, v. 206, no. 1–4, p. 55–72, doi: 10.1016/j.margeo.2004.03.013.

Trenhaile, A.S., 2000, Modeling the development of wave-cut shore platforms: *Marine Geology*, v. 166, no. 1–4, p. 163–178, doi: 10.1016/S0025-3227(00)00013-X.

Trenhaile, A.S., 2001a, Modeling the effect of late Quaternary interglacial sea levels on wave-cut shore platforms: *Marine Geology*, v. 172, no. 3–4, p. 205–223, doi: 10.1016/S0025-3227(00)00136-5.

Trenhaile, A., 2014b, Modelling the effect of Pliocene-Quaternary changes in sea level on stable and tectonically active land masses: *Earth Surface Processes and Landforms*, v. 39, no. 9, p. 1221–1235, doi: 10.1002/esp.3574.

5 Trenhaile, A.S., 2001b, Modelling the quaternary evolution of shore platforms and erosional continental shelves: *Earth Surface Processes and Landforms*, v. 26, no. 10, p. 1103–1128, doi: 10.1002/esp.255.

Walkden, M.J.A., and Hall, J.W., 2005, A predictive Mesoscale model of the erosion and profile development of soft rock shores: *Coastal Engineering*, v. 52, no. 6, p. 535–563, doi: 10.1016/j.coastaleng.2005.02.005.

Controls on the distribution of cosmogenic ^{10}Be across shore platforms

Martin D. Hurst^{1,2}, Dylan H. Rood³, and Michael A. Ellis²

¹School of Geographical and Earth Sciences, University of Glasgow, Glasgow, G12 8QQ, UK.

²British Geological Survey, Nicker Hill, Keyworth, Nottingham, NG12 5GG, UK.

³Department of Earth Science and Engineering, Imperial College London, South Kensington Campus, London SW7 2AZ, UK

Correspondence to: Martin D. Hurst (Martin.Hurst@glasgow.ac.uk)

Abstract. Quantifying rates of erosion on cliffted coasts across a range of timescales is vital for understanding the drivers and processes of coastal change and for assessing risks posed by future cliff retreat. Historical records cover at best the last 150 years; Cosmogenic radionuclides, such as ^{10}Be could allow us to look further into past to assess coastal change at **millennial** **millennial** timescales. CRNs accumulate *in-situ* near the Earth surface and have been used extensively to quantify erosion

5 rates, burial dates and surface exposure ages in terrestrial landscapes over the last three decades. More recently, applications in rocky coast settings have quantified the timing of mass wasting events, determined long-term-averaged rates of cliff retreat and revealed the exposure history of shore platforms. In this contribution, we developed and explored a numerical model for the accumulation of ^{10}Be on eroding shore platforms. In a series of numerical experiments, we investigated the influence of topographic and water shielding, dynamic platform erosion processes, the presence and variation in beach cover, and heterogeneous distribution of erosion on the distribution of ^{10}Be across shore platforms. Results demonstrate that, taking into account relative sea level change and tides, the concentration of ^{10}Be is sensitive to rates of cliff retreat. Factors such as topographic shielding and beach cover, act to reduce ^{10}Be concentrations on the platform, and may result in overestimation of cliff retreat rates if not accounted for. The shape of the distribution of ^{10}Be across a shore platform can potentially reveal whether cliff retreat rates are declining or accelerating through time. Measurement of ^{10}Be in shore platforms has great potential to allow us
10 to quantify long-term rates of cliff retreat and platform erosion.

1 Introduction

There is societal need to assess the rates, and the change of rates, at which cliffted coastlines will erode in the face of changing sea-levels and wave-climates that may result in more energetic coasts (Bray and Hooke, 1997; Trenhaile, 2010; Ashton et al., 2011; Barkwith et al., 2014). The lack of long-term records of cliff and shore-platform erosion rates is a key problem to
20 address (Trenhaile, 2014). An emerging tool to assess past rates of cliff retreat comes from the accumulation of cosmogenic radionuclides (CRNs), such as ^{10}Be , across active marine platforms that are generated via cliff retreat (Regard et al., 2012; Choi et al., 2012). CRNs have the potential to yield rates of cliff recession and shore-platform erosion over **millennial** **millennial** times-scales. These records will provide long-term context for historical and present rates of cliff recession, and facilitate calibration of dynamic models of shore platform erosion and cliff recession (Trenhaile, 2000; Walkden and Hall, 2005; Ashton

et al., 2011; Matsumoto et al., 2016). The erosion of rocky coastlines is the result of a dynamic suite of processes that operate at the coast (Sunamura, 1992; Kennedy et al., 2014). The production of CRNs in the shore platform is in turn influenced by these processes and the resultant morphological evolution of the coast (Regard et al., 2012). This paper explores the factors that influence the spatial patterns of ^{10}Be across emergent platforms, with the objective of allowing us to better constraining 5 quantitative measures of cliff retreat rate.

Estimates of sea cliff retreat rates over decadal to centennial timescales have been made by observing the change in position of the cliff top, derived from historical maps, aerial photography, satellite imagery, and field surveys (e.g. Bray and Hooke, 1997; Costa et al., 2004; Dornbusch et al., 2006; Brooks and Spencer, 2010; Katz and Mushkin, 2013). Observations are limited by the length of historical records, which span at best the late 1800s to present. Sunamura (2015) demonstrated that longer-term 10 records are needed. The return period of coastal mass wasting events, the principal mechanisms of coastal cliff retreat, may in some cases be much longer than these historical records (Recorbet et al., 2010). Methods to estimate long-term rates of cliff retreat are required in order to time average rates across multiple failure events. It is important that modern observations of sea cliff retreat can be placed in the context of long-term coastal evolution, over time-spans unaffected by human intervention at the coast. Long-term records of cliff retreat in response to relative sea level (RSL) change are required in order to predict how 15 coastal erosion may proceed into the future in the face of anticipated RSL rise and increased storminess. Yet, by their very nature, eroding coastlines leave scant evidence of any former state, and their form reflects little about their long-term erosional trajectories (Matsumoto et al., 2016). Cosmogenic radionuclides (CRNs) have the potential to reveal the long-term history of coastal change and to quantify process rates along rocky coastlines (Recorbet et al., 2010; Choi et al., 2012; Regard et al., 2012; Rogers et al., 2012).

20 CRNs are isotopes produced by the interaction of cosmic rays with target elements in the upper few metres of the Earth surface. Measurements of the abundance of CRNs (particularly ^{10}Be) produced in rock and soil samples (*in situ*) provides a versatile geochronometer to quantify how long a sample has been exposed at/near the Earth surface, or interpret how rapidly it has been eroded (Balco et al., 2008; Dunai, 1995; Granger et al., 2013). Recently, these techniques have been used for application in rocky coast settings. Recorbet et al. (2010) used CRNs to demonstrate that a calcareous sandstone sea cliff in 25 south eastern France last failed around 3.5 ka (thousand years before present), suggesting a long return period for sea cliff retreat events at that site. Rogers et al. (2012) assessed long-term shoreline recession in Washington, USA by using CRNs to date large boulders, released and abandoned on the shore platform during recession of till bluffs.

The measurement of ^{10}Be concentrations from shore platform samples to estimate long-term rates of sea cliff retreat was 30 pioneered by Regard et al. (2012) working on the flint-bearing chalk coastline near Mesnil-Val, France. Regard et al. developed a numerical model to predict the concentration of ^{10}Be on shore platforms as a function of the rate of cliff retreat. They minimised residuals between model results and a transect of eight ^{10}Be concentration measurements from across the shore platform in order to estimate long-term average sea cliff retreat rates that were similar to estimates from historical observations; however, the uncertainties on the analyses were large and limited the resolution/confidence of the comparison.

35 ~~The Some shore platforms may have formed during the Holocene and thus be entirely contemporaneous features, whilst others, particularly wide platforms in resistant lithologies, have been interpreted as inherited features, formed during previous~~

sea level high stands and reoccupied during the Holocene (e.g. Chao et al., 2003). Evidence for the antiquity of shore platforms may be revealed by CRNs. For example, Choi et al. (2012) measured ^{10}Be concentrations in shore platforms cut into resistant lithologies on the Korean coast. High concentrations were consistent with modelled ages extending back as far as 142 ka, and importantly, these ages did not correct for weathering and erosion and thus should be considered as minimum ages. Thus 5 Choi et al. (2012) conclude that these platforms are at least partially inherited features, originated in the Pleistocene.

On a contemporary shore platform (formed during the Holocene), the theoretical distribution of ^{10}Be concentrations on an eroding shore platform generally increases and then decreases with distance from the modern cliff (Figure 1). In the nearshore, ^{10}Be concentrations increase offshore because the shore platform has been exposed for longer. However, the rate of ^{10}Be production decreases offshore because cover by sea water attenuates the cosmic ray flux, hence the 10 amount of cosmic radiation received by the platform decreases with increased water depth. Additionally, downwear of the coastal platform surface removes the highest concentration ^{10}Be rock from the surface and exhumes lower concentrations that were previously below the surface (Figure 1). The combined result of these factors is a "humped" distribution of ^{10}Be concentrations. The magnitude of the maximum concentration is predicted to be proportional to the long-term averaged rate of sea cliff retreat (Regard et al., 2012); higher rates of sea cliff retreat result in lower ^{10}Be concentrations in the platform (Figure 1). 15 Tides modify these predictions by altering the distribution of ^{10}Be production across the shore platform due to attenuation of cosmic rays in the water column. Regard et al. (2012) showed that increasing tidal range causes the cross-shore position of the peak concentration to migrate seaward. Regard et al. also demonstrated that whilst rising RSL increases water depth, counter-intuitively it can result in higher ^{10}Be concentrations on the shore platform due to associated reduction in platform downwear rates.

20 Several other factors may influence the accumulation of ^{10}Be in shore platforms that have not been accounted for in previous coastal studies. Shore platforms may erode through gradual (e.g. weathering and abrasion) or episodic (e.g. quarrying) processes (Dornbusch and Robinson, 2011; Naylor et al., 2016), and the relative influence of these processes on the distribution of ^{10}Be has yet to be explored. Similarly, the assumption that shore platforms evolve in morphological steady state (i.e. landward 25 translation of a constant shore profile morphology through time that tracks relative sea level change), which is sometimes also referred to as equilibrium retreat, may not always be appropriate (Dickson et al., 2013). The style of coastal evolution is expected to influence the distribution of ^{10}Be across a shore platform. Modelling studies suggest that shore platform gradients may decline through time and platforms widen (Trenhaile, 2000; Walkden and Hall, 2005). Talus cones and beaches are often found fringing coastal cliffs, covering shore platforms and therefore shielding them from cosmic rays. ^{10}Be concentrations in the shore platform would be reduced (Regard et al., 2012). Coastal cliffs may shield the platform from a portion of the 30 incoming cosmic ray flux, thus also reducing ^{10}Be production in the nearshore, and shielding is proportional to cliff height.

In this study we quantified the sensitivity of platform CRN concentrations to topographic shielding, various processes of platform erosion/downwear, the presence/absence of beach cover, and transience in shore profile evolution. Our intention is 35 to explore the general case parsimoniously to better understand controls on ^{10}Be concentrations, rather than to make accurate predictions for any specific field site. In doing so, we only considered the development of shore platforms during the Holocene,

over the timescale during which eustatic sea level has been relatively stable (7 ka to present). We addressed this with a numerical model coupling cross-shore coastal evolution and ^{10}Be production to explore the potential for quantifying coastal retreat rates from ^{10}Be concentration measurements.

2 Numerical model-Model for shore platform evolutionShore Platform Evolution and ^{10}Be Production

5 Cliffted, rocky coasts, are commonly fronted by shore platforms that ~~are have previously been~~ classified into two types (Sunamura, 1992). Type-A platforms are characterised by a gently sloping erosional platform surface extending offshore beyond maximum low water. Type-B platforms are shallow gradient to sub-horizontal and terminate at their seaward edge at maximum low water through a scarp (Figure 2a). Numerical models of shore platform evolution have ~~succesfully successfully~~ recreated both of these end-member morphologies, ~~but have revealed that there are a range of other possible morphologies in between, for example sloping platforms terminating in a scarp~~ (e.g. Trenhaile, 2000; Walkden and Hall, 2005; Matsumoto et al., 2016).

10 Numerical models of platform evolution demonstrate that shore platforms and adjacent sea cliffs tend towards a morphological steady state, ~~such that coastal morphology does not change its form through time, and a constant cliff-platform geometry is translated landward through time~~. Under such conditions the morphology may reflect the combination of RSL change, tides and wave energy availability. However, the assumption of steady state retreat may not always be applicable (Dickson et al., 2013).

15 We expected the style of platform evolution to be important for the distribution of ^{10}Be across a shore platform. ~~Therefore we performed experiments not only assuming steady state coastal retreat, but also using a dynamic model for platform evolution. Moreover, micro-erosion meter measurements of platform downwear suggest that downwear is not uniform across the shore profile, but tends to be faster in the upper inter-tidal zone and decline with depth (Porter et al., 2010), and to explore the influence of such a distribution on ^{10}Be concentrations, a dynamic morphological model was required.~~

20 Therefore, two different morphological models are used in this study. The first assumes steady state evolution of the shore profile, such that coastal morphology does not change its form through time, and a constant cliff-platform geometry is translated landward through time. The second model is a dynamic shore platform evolution model similar to that of Trenhaile (2000) that can reproduce a range of platform geometries. These two approaches are described in more detail in the subsequent sections.

2.1 Steady State Coastal Retreat

25 We initially assumed that coastal cliff and shore platform evolution can be considered a steady-state process: a constant coastal cross-section profile is translated landward through time, with the elevation of the cliff-platform junction tracking RSL (Figure 2b). As such, platform downwear was assumed proportional to the product of cliff retreat rate and platform gradient α (Regard et al., 2012).

30 A steady state approach assumes platform downwear is gradual and constant (~~which implies abrasion is the dominant process~~) across the entire profile, and proportional to the rate of cliff retreat. ~~However, several~~ We are therefore assuming that the combined mechanical and chemical processes that can cause shore platform lowering culminate to constant and gradual downwear in this case. Wave energy, tidal range and the types of processes operating on the shore platform do not factor in

to the morphological development explicitly but rather are assumed to combine to cause steady state retreat. Several coastal platforms have been observed to erode due to quarrying and block removal (e.g. Dornbusch and Robinson, 2011; Naylor et al., 2016). In order to explore the potential implication of these erosion processes for the accumulation of ^{10}Be in a shore platform, we also evolved a series of stepped platforms by steady state retreat.

5 2.2 Numerical Model for Dynamic Platform Evolution

In order to explore ^{10}Be concentrations across a transient (i.e. not steady state) shore platform, we developed a simple numerical model for shore profile evolution (the ROck and BOttom COastal Profile [RoBoCoP] Model), broadly similar to those of Sunamura (1992), Anderson et al. (1999), Trenhaile (2000) and Walkden and Hall (2005) and Trenhaile (2000). These models assume that horizontal erosion at the water level is proportional to the availability of wave energy (or by proxy, wave height). The shore profile was considered as a regularly spaced vertical stack of cells with horizontal position x [L] (all dimensions denoted in square brackets as [L]ength, [M]ass and [T]ime). The change in position of the coast at the water level x_w [L] through time t [T] was assumed to be linearly proportional to the height of breaking waves reaching the shore H_c [L]:

$$\frac{dx_w}{dt} = K \rho_w g H_c \quad (1)$$

In Equation 1, K [$\text{L}^2 \cdot \text{T} \cdot \text{M}^{-1}$] is a coefficient related to the resistance of bedrock to erosion, ρ_w [$\text{M} \cdot \text{L}^{-3}$] is the density of water and ~~and~~ g [$\text{L} \cdot \text{T}^{-2}$] is acceleration due to gravity. The water depth h_w [L] for the initiation of wave breaking h_b [L] was related to wave height H [L], such that breaking wave height H_b [L] is the wave height that exceeds a water depth-dependent threshold:

$$H_b = 0.78 h_b \quad (2)$$

Breaking wave height was determined by iterating deep water wave conditions from deep to shallow water and calculating h_w according to linear wave theory (e.g. Hurst et al., 2015), or following an empirical relationship relating breaking wave conditions to offshore wave conditions (e.g. Komar and Gaughan, 1972):

$$H_b = 0.39 g^{1/5} T^{2/5} H_0^{4/5} \quad (3)$$

In Equation 3, T [T] is wave period and H_0 [L] is deep water wave height. If Equation 3 is used to predict H_b , Equation 2 can be inverted to determine the water depth at which wave breaking begins. Following wave breaking, wave height is assumed to decay exponentially with distance across the platform, such that the wave height at the water line can be described as:

$$H_c = H_b e^{-k W_s} \quad (4)$$

In Equation 4, W_s [L] is the width of the surf zone, measured from $x(h_w = h_b)$ to $x(h_w = 0)$, k is a dimensionless constant that represents the rate of breaking wave energy dissipation, which reflects bed roughness in the surf zone (Trenhaile, 2000); we used $k = 0.02$ throughout. These properties are all time dependent, because the elevation of the water surface varies due to tides (superimposed onto any RSL change). The distribution of erosion across the platform was integrated across the tidal cycle 5 of period T_t [T]. Platform erosion below the water line was assumed to decline exponentially with water depth h_w . Combining Equations 1-4, the governing equation for evolution of the shore platform becomes:

$$\frac{dx}{dt} = K \rho_w g \int_{t=0}^{T_t} H_b(t) e^{-(k W_s(t) + h_w(t)) h_w(t) - k W_s(t)} \quad \{h_w \in \mathbb{R} : h_w \geq 0\} \quad (5)$$

2.3 Beach Cover

In order to explore the influence of beach cover on the accumulation of ^{10}Be on the coastal platform, we approximated the 10 profile morphology of beaches using a power-law function (Figure 3; Bruun, 1954):

$$z_b = z_{b0} - Ax_b^m \quad (6)$$

In Equation 6, z_b [L] is the elevation of the beach, z_{b0} [L] is the elevation of the beach at the top of the berm, A [$\text{L}^{1/3}$] is a scaling parameter that relates to the size of beach material, and m is a dimensionless exponent that represents the distribution of wave energy dissipation on the shoreface. We used a value of $A = 0.12$ suitable for gravel, and a shape ~~exponent~~^{exponent} 15 $m = 2/3$, consistent with a number of studies (Dean and Darlymple, 2002). The beach profile extends from the position of top of the berm described by the beach width B_w [L] and berm height B_h [L] (Figure 3).

3 Numerical model for ^{10}Be Production in the Shore Platform

The production of ^{10}Be ~~fundamentally~~^{fundamentally} depends on how long the surface has been exposed, and the rate at which material is removed through erosion (Gosse and Phillips, 2001). On a shore platform, exposure is modulated by topographic 20 shielding, beach cover and water cover (Regard et al., 2012). Erosion of the shore platform may take place through ~~abrasion~~^{gradual downwear processes} or plucking, and may not be spatially and temporally uniform (Dickson et al., 2013).

3.1 Production of ^{10}Be in Rock

The concentration of ^{10}Be N [atoms M^{-1}] changes through time according to:

$$\frac{dN}{dt} = P(h_r) - \lambda N \quad (7)$$

25 In Equation 7, $P(h_r)$ [atoms $\cdot \text{M}^{-1} \cdot \text{T}^{-1}$] is the depth-dependent production rate of ^{10}Be , h_r [L] is depth below the rock surface, and $\lambda = 4.99 \times 10^{-7}$ is the ^{10}Be radioactive decay constant (Chmeleff et al., 2010; Korschinek et al., 2010). The production of

^{10}Be *in-situ* at and near the Earth surface declines exponentially with depth (self-shielding) as the cosmic ray flux attenuates (e.g. Gosse and Phillips, 2001; Balco et al., 2008; Mudd et al., 2016):

$$P(h_r) = \sum_i P_s(i) e^{-(\frac{h_r}{h^*(i)})} \quad (8)$$

In Equation 8, P_s [atoms·M $^{-1}$ ·T $^{-1}$] refers to the production rate at the rock surface for the production pathway i . ^{10}Be is predominantly produced by neutron spallation, with minor contribution from fast and slow muon interactions. Muogenic production penetrates much deeper into the Earth surface and therefore may source a significant part of observed ^{10}Be concentrations. Similar to West et al. (2014), we used a single exponential curve to integrate fast and slow muogenic production pathways (Braucher et al., 2011, 2013). The attenuation of spallation and muogenic reactions declines according to an attenuation length scale h^* [L] that is dependent on the density of bedrock ρ_r [M·L $^{-3}$] and a pathway dependent attenuation factor Λ [M·L $^{-2}$] (for spallation $\Lambda = 1600 \text{ kg m}^{-2}$; and for muogenic production $\Lambda = 42000 \text{ kg m}^{-2}$):

$$h^*(i) = \frac{\rho_r}{\Lambda(i)} \quad (9)$$

3.2 Topographic Shielding

Rock surfaces may be shielded from a portion of the incoming cosmic ray flux by local topography. At the coast, sea cliffs can block a significant portion of the sky and therefore partially shields the platform from cosmic rays. Typically shielding factors can be quantified by field surveys (Dunne et al., 1999) or from topographic data (Codilean, 2006; Mudd et al., 2016). However, when the sea cliff is retreating, the amount of topographic shielding at a fixed location will change through time with increasing distance to the position of the cliff.

In order to model the influence of topographic shielding by the cliff on the accumulation of ^{10}Be , we idealised that the cliff is vertical and of height C_H [L], and assumed the cliff line to be straight in planformplan-form. For a given distance from the cliff x_c we defined the viewshed generated by the cliff as the angle in the sky θ from horizontal made by the top of the cliff in the azimuth direction ϕ for each observation j :

$$\theta(j) = \tan^{-1} \left(\frac{C_H \cos \phi(j)}{x_c} \right) \quad (10)$$

Following Dunne et al. (1999), a shielding factor S_{cliff} is the ratio of cosmic ray flux given the viewshed F to the maximum cosmic ray flux for an unobstructed flat surface F_{max} :

$$S_{cliff} = \frac{F}{F_{max}} = \frac{\Delta\phi}{2\pi} \sum_{j=1}^n \sin^{m+1} \theta(j) \quad (11)$$

where n is the total number of viewshed observations, that span the azimuth range $\phi = 0 - 360^\circ$. A straight, vertical cliff line yields $S_{cliff} = 0.5$ at $x_c = 0$ since 50% of the sky is obstructed.

Topographic ~~shielding~~ can also be modelled ~~explicitly~~ from a digital elevation model (DEM) following Codilean (2006). However these shielding values only apply to the current platform and cliff morphology. We compared Equation 11 to shielding factors calculated from a DEM for shore platforms on the coast of East Sussex, UK (1 m resolution airborne LiDAR; data courtesy of Channel Coast Observatory; www.channelcoast.org; accessed 25th May 2014), using 5 software published by Mudd et al. (2016) [https://github.com/LSDtopotools/LSDTopoTools_CRNBasinwide].

3.3 Water Shielding

A shore platform is periodically exposed or submerged by the sea due to tides. The cosmic ray flux attenuates exponentially with water depth h_w [L] so that the production rate at the surface of the platform decreases according to:

$$P_s(x, i) = P_0(i) e^{\frac{-h_w(x)}{h_w^*(i)}} \quad (12)$$

10 In Equation 12, P_0 [atoms·M⁻¹·T⁻¹] is the reference production rate at the water surface, which is a function of latitude and altitude (e.g. Lal, 1991; Dunai, 2000; Stone, 2000; Desilets et al., 2006; Lifton et al., 2005). Equation 9 is used to calculate z_w^* by substituting water density ρ_w [M·L⁻³] for ρ_r . Thus, the production by each pathway can be adjusted for attenuation in the water column, however this adjustment depends on the tidal cycle at the site of interest.

3.3.1 Tides

15 Tides modify production in the platform by varying water depth h_w and intermittently ~~submerging and~~ exposing the platform sub-aerially. ~~Relative to a scenario with no tidal variation,~~ Regard et al. (2012) demonstrated that tides have a net effect to reduce ¹⁰Be production in the upper inter-tidal platform due to periodic platform submergence that reduces the net cosmic ray flux received, while ¹⁰Be production in the lower platform increases ~~relatively~~ due to periodic exposure. Here we extended this analysis to explore the influence of tidal regime on ¹⁰Be production.

20 Predictions of the tide are made as the sum of its harmonic constituents (e.g. Pugh and Woodworth, 2014):

$$z_w(t) = \sum_n H_n \cos(\sigma_n t - g_n) \quad (13)$$

In Equation 13, z_w [L] is the elevation of the mean water surface, the subscript n refers to the tidal constituent, H_n is the amplitude [L] of that constituent, σ_n is the angular speed [°·T⁻¹], and g_n is the phase lag [T]. Predictions of tidal elevation can be converted to water depth across the platform:

$$25 \quad h_w(x, t) = \begin{cases} z_w(t) - z_r(x, t) & \text{if } h_w \geq 0 \\ 0 & \text{if } h_w(t) < 0 \end{cases} \quad (14)$$

In Equation 14, z_r [L] is the elevation of the platform surface; h_w must be positive (i.e. if the water level falls below the platform elevation the water depth at this location is zero). Combining Equation 12 and 14, the production rate at the platform

surface averaged over a tidal cycle (T) can then be calculated as:

$$P_s(x, i) = \frac{1}{T} \sum_{t=0}^T P_0(i) e^{-\frac{h_w(x, t)}{h_w^*(i)}} \quad (15)$$

We used Equations 13-15 to compare the distribution of platform surface production rates across the shore platform for hypothetical diurnal, mixed and semi-diurnal tidal regimes, generated using the tidal constituents listed in Table 1. These 5 hypothetical tidal regimes were designed to cover a similar range in water levels across roughly a single lunar duration. We also compared the predicted P_s for a simple semi-diurnal tide to the effective production averaged over a full year of tidal records from the a tide gauge at Newhaven, East Sussex [available from www.channelcoast.org; accessed 2nd October 2014].

3.3.2 Relative Sea Level Change

For model experiments driven by steady state retreat, RSL rise results in higher concentrations of ^{10}Be across the shore 10 platform. This may seem counter-intuitive at first, however less vertical downwear of the platform surface is required to maintain the steady state profile during rising RSL and less material with high ^{10}Be concentrations is removed by downwear (Regard et al., 2012). In this study, we used constant rates of RSL change, since Holocene sea-levels have been relatively stable over the last 7 ka; however, for application to specific sites, RSL histories derived from local sedimentary records or regional crustal flexure models (e.g. Bradley et al., 2011) are recommended.

15 4 Experimental Setup

We conducted a number of experiments to explore the influence of specific processes on the distribution of ^{10}Be across a shore platform. Here we describe the model setup used to investigate (i) the influence of platform erosion process; (ii) the influence of beach cover; (iii) the influence of tides; and (iv) the influence of transient profile evolution on the accumulation of ^{10}Be in the shore platform. Global parameters consistent across all experiments are listed in Table 2.

20 4.1 Block Removal Processes

To investigate the influence of block removal processes, we evolved a steady state profile with a fixed average slope $\alpha = 1/100$ consisting of offset horizontal surfaces akin to horizontal bedding planes. The separation between surfaces was varied from 0 25 m (uniform sloped platform, downwear inferred to be by abrasion) to 0.8 m. Steps in the steady state profile were allowed to migrate landward through time in concert with a prescribed constant cliff retreat rate of $\epsilon = 0.1 \text{ m yr}^{-1}$. RSL was held constant and a diurnal tide with $H_n = 1.0 \text{ m}$ was used.

4.2 Beach Cover

Beaches cover the shore platforms in the ~~nearshore~~near-shore, which partially shields the platform from cosmic rays and, as a result, reduces ^{10}Be production in the platform surface. The presence of beach cover may not be consistent, but will depend

on the supply of material from the adjacent cliffs, the supply and removal of sediment due to both alongshore and cross-shore sediment transport, and the rate at which beach material is physically and chemically weathered. We conducted a series of experiments designed to investigate the influence of beach cover on the accumulation of ^{10}Be on a shore platform. Firstly, we considered a simple case where beach morphology is held constant through time, and explore the influence ~~of that~~ that constant 5 beach width ($B_w = 0, 10, 20, 50, 100$ m) on the concentration of ^{10}Be in the shore platform. Secondly, we considered the condition that beach width may vary through time, with beach width varying as a sinusoidal function over decadal timescales (wavelength = 100 years), with an average $B_W = 50$ m and amplitude of 30 m. Thirdly, we considered the condition that beaches may be ~~being~~ lost through time, as beach material is transported away more rapidly than it is supplied by alongshore transport and cliff erosion. These scenarios all evolved the shore profile by steady state retreat with a cliff retreat rate of $\epsilon = 0.1$ 10 m yr^{-1} . RSL was held constant and a diurnal tide with $H_n = 1.0$ m was used.

4.3 Steady state and Transient Shore Platform Evolution

Previous CRN studies have assumed that shore platform evolution proceeds in steady state such that a constant shore profile morphology is translated landward through time, and tracks the trajectory of RSL change (Regard et al., 2012). Here, we explored the influence of RSL rise on shore platform morphology and the resulting distribution of ^{10}Be across shore platforms 15 with RoBoCoP (see Section 2.2). We compared the results of dynamic shore platform evolution experiments to those of steady state shore profile retreat to investigate the influence that the assumption steady retreat may have on the accumulation of ^{10}Be and our ability to interpret rates of cliff retreat. The initial and boundary conditions for the model were held constant (see Table 3). The model is initialised with a sloped platform profile and forward modelled for 10 Ka. In our first set of experiments, we held RSL constant whilst during the second set of experiments RSL rose at a constant rate (0.5 mm yr^{-1}). RSL rise is 20 known from previous experiments to influence the rate of cliff retreat and how it changes through time, the morphology of the platform, and the amount and distribution of platform downwear (Trenhaile, 2000; Walkden and Hall, 2005). We compared the resultant ^{10}Be concentrations to predictions that assume steady state retreat in order to determine whether this assumption allows ~~estimation~~ estimation of average transient retreat rates.

5 Results

25 5.1 Topographic Shielding

The topographic shielding scaling factor S_T is predicted to have a value of 0.5 immediately adjacent to the cliff, increasing ~~nonlinearly~~ non-linearly with distance from the cliff. (Figure 4a). Cliff height principally controls the rate at which S_T increase, such that taller cliffs result in a greater degree of shielding for a greater distance offshore. S_T approaches unity asymptotically 30 by $x_c \approx 4 \cdot C_H$; there is little influence of the cliff on ^{10}Be production beyond. Our approach (Equation 11) assumes a straight cliff line with constant cliff height. We compared this approach to the distribution of topographic shielding factors modelled for a real stretch of coastline in East Sussex, UK, following Codilean (2006) (Figure 4b). Again, these ~~distributed~~ distributed S_T

values are smallest immediately adjacent to the cliff (0.5) and increase nonlinearly-non-linearly with distance across the shore platform. The values calculated from the DEM are in good agreement with Equation 11 as demonstrated by three example transects, each with different cliff heights (Figure 4c). Differences between observed and model S_T values are due to variation in the gradient of the cliff face and the planform-plan-form geometry of the cliff line. Since it is unlikely in most settings to 5 have information on how C_H and the planform-plan-form geometry may have changed through time, we used Equation 11 to represent topographic shielding, assuming constant C_H .

5.2 Water Shielding by Tides

Variation in water level at hourly timescales due to tides modifies the cosmic ray flux delivered to the platform. The tidal regime at a site may influence distribution of ^{10}Be production across a shore platform, and so we explored the effects of 10 hypothetical diurnal, mixed and semi-diurnal tidal regimes (Figure 5a-f) on ^{10}Be concentrations. The tidal constituents used to model these are shown in Table 1. Figure 5g shows the distribution of relative spallation production P_s/P_0 across a planar platform defined by $\alpha = 1/100$ under these six different hypothetical tidal regimes. Tides act to reduce the production in the nearshore-near-shore/upper intertidal-inter-tidal zone due to periodic submergence of the upper platform, whilst production in the lower intertidal-inter-tidal zone is increased due to periodic exposure of the platform. Simple diurnal/semidiurnal 15 semi-diurnal tides show the strongest modification of ^{10}Be production according to our experiments, but this is strongly dependent on H_n for the additional constituents. Different tidal regimes may result in differences in the production of ^{10}Be across the coastal platform and therefore it is important to consider the tidal regime at a particular site to model ^{10}Be production in the platform.

Figure 6 shows the distribution of the effective production rate at the platform surface for purely diurnal tides with an 20 amplitude of 2.4 m, compared to the tide gauge water level record at Newhaven, East Sussex, UK. Incorporation of a tidal record with the full range of harmonic components dampens production very slightly in the nearshore-near-shore region and slightly increases production further offshore relative to a simple the semi-diurnal tide model. These differences do not significantly influence ^{10}Be production. A simple diurnal/semidiurnal-semi-diurnal tide model with a single representative tidal amplitude is appropriate at some sites.

25 5.3 Block Removal Processes

Model results of the migration of bedrock steps due to block removal processes across a shore platform that evolves through steady state retreat is shown in Figure 7a for a range of step sizes. The landward migration of steps results in the sudden exposure of platform that was previously the depth of the step size below the platform surface. Since ^{10}Be production rates decline exponentially with depth into the rock (Equation 8), block removal unearths rock with significantly lower concentration, 30 resulting in sudden drops in ^{10}Be concentration with distance across the platform (Figure 7b). However, the flat-topped steps are subject to no vertical downwear in these simulations and so they continue to accumulate ^{10}Be such that, between bedrock steps, concentrations increase with distance from the cliff. The result is a saw-toothed distribution of concentrations, superimposed

on the "humped" distribution expected when there is no block removal. The magnitude and wavelength of the variability in concentrations is controlled by the size and frequency of the bedrock steps.

5.4 Beach Cover Control on ^{10}Be Concentrations

Beaches were represented by a Bruun Profile (Equation 6) that extends seaward from a prescribed beach width and berm height 5 (Figure 3). Beach cover may not be constant through time. We explored the influence of beach width, variable beach width, and declining beach width on shore platform ^{10}Be concentrations.

5.4.1 Influence of Beach Cover

Beach cover is expected to partially shield the ~~nearshore~~near-shore platform from cosmic rays and therefore reduce concentrations of ^{10}Be in the platform surface. Wider beaches should result in lower ^{10}Be concentrations. We ran model experiments with 10 different values of B_W , which was held constant for the duration of each experiment (Figure 8a). The predicted distributions of ^{10}Be concentrations increase and then decrease in the offshore direction, and the magnitude of the "hump" is reduced when beaches are wider (Figure 8b). The position of the peak with respect to the cliff moves offshore as production in the ~~nearshore~~near-shore is reduced by beach cover. The black line corresponds to the scenario with no beach cover. The concentrations of ^{10}Be at the platform surface decrease in a linear fashion with wider beaches (Figure 9), to the extent that a 50 m wide beach 15 results in a 18% reduction in the magnitude of the ^{10}Be concentrations. If not accounted for, this would lead to over prediction of cliff retreat rates when inverting measured ^{10}Be concentrations, since, in the absence of beaches, lower concentrations of ^{10}Be suggest faster cliff retreat rates (Regard et al., 2012).

5.4.2 Influence of Variable Beach Cover

Beach cover may not be constant through time, however there is rarely any evidence available revealing how beaches may 20 have changed during the Holocene on eroding coastlines, since eroded material tends to be removed. We explored how variable beach cover modifies ^{10}Be concentrations by modelling B_W through time as a sinusoidal function with wavelength 100 years and amplitude 30 m about an average of $B_W = 50$ m (Figure 10a). Thus, the beach progrades and regresses every 100 years, maintaining an equilibrium beach profile (Figure 10b). The resultant distribution of ^{10}Be concentrations are compared to simulations with no beach cover and a constant $B_W = 50$ m (Figure 10c). We found that there is very little difference in ^{10}Be 25 concentrations between a constant and variable beach width scenario.

We also explored scenarios in which beach cover was reduced through time. Figure 11 shows the morphological evolution and concentrations of ^{10}Be in the shore platform surface predicted for the condition where beaches have thinned from a constant width of 50, 100 or 200 m over the last 1000 years. The thinning of beaches, and thus the presence of more beach cover in the past reduces the expected concentrations of ^{10}Be in the shore platform. This will result in over-prediction of cliff retreat rates 30 when not accounted for, since platform ^{10}Be concentrations will be lower, as otherwise associated with more rapid rates of cliff retreat.

5.5 Transient Shore Platform Evolution

5.5.1 Constant sea level

We modelled the evolution of a shore profile given constant RSL and the consequential concentrations of ^{10}Be across the modelled shore profile. The gradient of the shore platform decreases through time, as the platform widens (Figure 12a). A 5 stepped platform emerges with the step at the lower tidal limit, reflected by a downward step in ^{10}Be concentrations on the shore platform (Figure 12b). Wave energy attenuation increases as the platform widens such that cliff retreat rates decline as the model simulation proceeds (Figure 12c). Reduction in the mean gradient of the platform is shown in Figure 12d. The distribution of ^{10}Be concentrations is humped, similar in form to predictions of a steady state retreat model (Regard et al., 2012). The magnitude of the hump increases through time, as cliff retreat rates slow.

10 In order to compare results directly, we then ran simulations where the morphology evolved in steady state (cliff retreat rate was held constant, using the median values reported in Figure 12b, platform gradient was taken as the instantaneous average platform gradient in each profile in Figure 12a (measured between platform elevations $z = 0.5$ and $z = -1$ m). These simulations ran for long enough that 1000 m of cliff retreat had occurred. Comparison of the distribution of ^{10}Be concentrations between the transient (solid lines) and steady state (dashed lines) model runs is shown in Figure 13a, shaded by their retreat rate as listed 15 in the legend in Figure 12b. The magnitude of the peak in concentrations were in this case similar between transient and steady state simulations (Figure 13b), however, the position of the peak was further offshore in the steady state simulations (Figure 13c).

5.5.2 Rising Relative Sea Level

We modelled the transient evolution of a shore profile and ^{10}Be concentrations with sea level rise of 0.5 mm yr^{-1} (Figure 14). 20 The shore platform evolves rapidly to form a low gradient shore platform ramp (Figure 14a). The elevation of the platform/cliff junction increases through time and tracks the high tide level, superimposed on the trajectory of RSL rise. Cliff retreat rates are initially rapid and decline towards a constant rate of 0.32 cm yr^{-1} (Figure 14c). Having attained a constant rate of cliff retreat, concentrations of ^{10}Be in the platform as a function of distance from the cliff are constant, consistent with the concept of steady-state morphological retreat. Due to low platform gradients, and the absence of significant platform downwear below the 25 intertidal inter-tidal zone, concentrations of ^{10}Be continue to increase offshore and so no "hump" in concentrations is observed.

We compared these predictions to those of a steady state retreat model with cliff retreat rate (32 cm yr^{-1}) and platform gradient ($1/500$) observed in the above transient experiment. The results are plotted as the dashed line in Figure 14b. The steady state retreat model similarly predicts concentrations that increase across the entire shore platform due to minimal platform downwear in the presence of sea level rise. However, the steady state model predicts significantly higher concentrations of ^{10}Be across 30 the shore platform, by about a factor of two, due to differences in the distribution of downwear between the two models. In the transient model runs, the amount of platform downwear declines exponentially with water depth, leading to rapid erosion in the mid to upper intertidal inter-tidal zone, where ^{10}Be production rates are most rapid. This is evidenced by the concave-up shore profile nearest to the cliff. Therefore not only does more downwear occur on the upper part of the transient profile, but as a

result, water depths increase more quickly offshore, and increased water shielding reduces ^{10}Be production and concentrations on the shore platform as a whole. Thus, if we were to assume steady state retreat and a planar shore platform in a setting where there is more platform downwear in the ~~intertidal~~ inter-tidal zone, we are likely to overestimate retreat rates since ^{10}Be concentrations will be lower than expected for a given retreat rate.

5 6 Discussion

We have identified that concentrations of ^{10}Be on shore platforms are sensitive to (i) shielding by cliffs; (ii) the type of process (abrasion gradual downwear vs quarrying); (iii) the tidal regime; (iv) the nature of beach cover and how it has changed through time; and (v) the style (steady state *vs.* transient) by which the platform evolves, particularly in the face of RSL change.

Nevertheless, these results demonstrate that there is great potential for CRN measurements to provide first-order estimates 10 of long-term rates of sea cliff retreat (Regard et al., 2012; Rogers et al., 2012; Choi et al., 2012). Quantifying these factors, and how they may have changed over the millennial timescales required by CRN studies may not always be possible. Some of these factors can be accounted for explicitly in site specific studies, such as cliff shielding, given the height of the modern cliff, tides, informed by nearby tide gauges, and RSL change, from proxy records or glacio-isostatic adjustment models. We are still required to assume that these factors have not changed through time; it is always necessary, to some extent, to extrapolate the 15 modern coastal configuration back into the past. We have little or no information about the paleaeotopography offshore from a modern cliff. Observations of beach cover can be made from historical data (e.g. Dornbusch et al., 2008), but are limited to the length of the historical record (c. 150 years maximum) and may be influenced by human intervention at the coast. Many of these factors act to reduce predictions of ^{10}Be in the shore platform, but maintain the overall "humped" shape of the distribution. It appears not possible to distinguish from the ^{10}Be concentrations what the history of beach cover is if different from present. It 20 may be possible to identify when a platform is evolving transiently (see below). If factors that reduce ^{10}Be concentrations were not considered then we would expect to ~~overpredict~~ over-predict rates of cliff retreat. Consequently, cliff retreat rates derived from CRN studies might be considered maximum estimates. Below we discuss some of these issues further in the context of quantifying long-term rates of cliff retreat.

6.1 Block removal processes

25 Our simplified experiments demonstrate that where block removal processes are an important process for platform evolution, the expected distribution of ^{10}Be becomes more variable, with higher frequency variation superimposed on the expected 'humped' distribution (Figure 7). Consequently, CRNs can still reveal rates of cliff retreat, but a careful sampling strategy will be required to account for block removal processes. Observations of platform erosion process should be made when sampling for CRNs, and the size of steps or blocks will be important to record. Where available, information on the rate at which steps 30 migrate and blocks are removed will help to inform sampling and interpretation (e.g. Dornbusch and Robinson, 2011; Naylor et al., 2016). Measuring ^{10}Be concentrations on shore platforms where block removal processes are dominant may allow the rate of step migration to be determined in the presence of large bedrock steps (Figure 7). This will require high density sam-

pling, focused on an individual bedrock step surface. If sampling on a platform with a stepped profile, data on the size of steps should be recorded and position of samples relative to steps will also be important.

6.2 Transient shore platform development

Previous studies had assumed steady state profile retreat was an adequate description of the morphological evolution of the shore platform in order to predict cliff retreat rates from ^{10}Be concentrations in the shore platform over millennial timescales (Regard et al., 2012). Dynamic shore profile evolution models (e.g. Sunamura, 1992; Anderson et al., 1999; Trenhaile, 2000; Walkden and Hall, 2005; Matsumoto et al., 2016) predict that coasts tend towards steady state whereby rapid cliff retreat widens shore platforms and the resultant increased wave energy dissipation reduces cliff retreat rates and increases erosion of the shore platform. We coupled predictions of ^{10}Be production to a dynamic shore profile evolution model (RoBoCoP; see section 2.2) in scenarios with (i) constant RSL, such that cliff retreat rates gradually reduced through time due to widening of the shore platform; and (ii) rising RSL, such that cliff retreat rates tended to a constant rate in time.

In scenario (i) the concentrations of ^{10}Be on the shore platform increased through time as cliff retreat rates declined (Figure 12). Comparison to predictions of steady state profile retreat (Figure 13) revealed that the position of the peak in ^{10}Be concentrations was further offshore when assuming steady state retreat. The results of Regard et al. (2012) using a steady state retreat model demonstrated that the position of the peak is sensitive to the tidal range and the platform gradient. Tidal range was held constant for all of our simulations, however platform gradient declined during transient platform evolution simulations. Comparison of field measurements of ^{10}Be concentrations and model runs that assume steady state retreat may therefore reveal when platform gradients are declining through time based on a mismatch in the position of the observed and modelled peak concentrations. In this case, the magnitude of the peaks suggests similar cliff retreat rates whether assuming steady state or transient shore profile evolution.

In scenario (ii) cliff retreat rates tended toward a constant rate with the result that the distribution of ^{10}Be was approximately constant through time (Figure 14). The concentrations of ^{10}Be were not consistent with a steady state evolution scenario in which the platform gradient is fixed, which predicted roughly twice the amount of ^{10}Be for a ~~particular~~ particular position on the platform. The difference can be explained by the dissimilar platform morphology brought about by uneven distribution of platform downwear in the transient model simulations. Greater rates of downwear in the ~~intertidal~~ inter-tidal zone lower the platform more rapidly in the ~~nearshore~~ near-shore, which removes ^{10}Be -laden rock and results in deeper water in the ~~nearshore~~ near-shore (and therefore reduced ^{10}Be production) than in the steady state model runs that assume constant α . These differences may be exacerbated when taking into account the spatial distribution of other processes such as weathering, the spatial distribution of which is currently poorly understood (Porter et al., 2010). In attempting to reconstruct cliff retreat rates the elevation profile of the platform should be accounted for such that the distribution of downwear across the platform is taken into account. ~~This could be aided by data on the distribution of downwear rates from micro-erosion-meter measurements (e.g. Robinson, 1977; Porter et al., 2010; Stephenson et al., 2010, 2012)~~ Failure to do so would result in ~~overprediction~~ over-prediction of erosion rates since faster retreat rates would be required to match the lower concentrations observed on the platform. Accounting for the distribution of downwear would be aided by data on the

distribution of downwear rates from micro-erosion-meter measurements (e.g. Robinson, 1977; Porter et al., 2010) and there is evidence that rates measured over the short term (1–2 years) are consistent with rates measured over three decades (Stephenson et al., 2010, 2011). Nevertheless, it is still not clear whether it is appropriate to extrapolate these rates over centennial to millennial timescales required to accumulate measurable concentrations of ^{10}Be , nor is it yet clear whether this result is consistent across different shore platforms around the world. Future studies that measure the distribution of CRN concentrations (both across the shore platform and at depth) at sites with long records of observed downwearing rates would be an important next step in this line of enquiry.

6.3 Beach cover

Beach cover in the ~~nearshore~~–~~near-shore~~ partially shields the underlying platform from ^{10}Be accumulation and results in a reduction in the magnitude of the hump. Yet we suggest that the significant reductions are only observed for relatively high beach widths (≥ 50 m), that are only likely to persist on slowly eroding coastlines, since they may absorb wave energy and partially protect the ~~clifffline~~~~cliff-line~~. Theoretical considerations suggest a dynamic relationship between beach material and cliff retreat; beaches can act both to provide abrasive tools to enhance cliff erosion or to protect the cliff from wave energy (e.g. Limber and Murray, 2011). ~~Similarly, cliffs are an important supply of material that makes up beaches; accelerated cliff retreat might allow beaches to build and shield the platform from cosmic rays.~~ We did not model this ~~dynamism~~~~dynamism~~, but instead favoured exploration of simple relationships between beach dynamics and the accumulation of CRNs in the platform in order to try and understand first order controls. We found that variation in beach cover through time was not important for CRN concentrations on a shore platform when compared to a scenario with a constant and representative average beach width. ~~However, our~~~~Our~~ approach treated cliff retreat rate as constant ~~and gradual~~, thus not capturing feedbacks between beach cover and cliff retreat, ~~nor exploring the influence that stochastic mass wasting processes might have on the variability of beach cover.~~

6.4 Inheritance of ^{10}Be

Predicted concentrations of ^{10}Be in shore platforms are relatively low compared to typical applications in geomorphic studies. Cosmogenic ^{10}Be produced at the surface is dominated by spallation reactions, but muogenic production penetrates deeper into the Earth surface (Heisinger et al., 2002b, a; Braucher et al., 2013). Muons only account for a small fraction of the production once the platform is exposed, but platforms may already contain an appreciable concentration of ^{10}Be prior to exposure formed by deep-penetrating muons over much longer timescales. The amount of "inherited" ^{10}Be will decline with the depth below the top of the cliff (i.e. the cliff height) but is also influenced by the rate of surface lowering at the cliff top (Lal, 1991). It is important to collect ~~nearshore~~–~~near-shore~~ samples to quantify how much inherited ^{10}Be is in the rock prior to platform exposure.

Platforms may also be geomorphically inherited landforms, having formed during a previous inter-glacial sea level high stand (e.g. the Eemian; 130–115 ka) and reoccupied by the sea during the Holocene (e.g. Trenhaile, 2001; Chao et al., 2003; Choi et al., 2012). If shore platforms are contemporary features, ^{10}Be concentrations will be low, and their distribution controlled primarily by cliff retreat rate, and other factors explored in this paper. If shore platforms are inherited features then ^{10}Be con-

centrations will be substantially higher reflecting subaerial exposure during the last glacial period Choi2012 (Choi et al., 2012). The location of a sudden increase in ^{10}Be concentrations on the shore platform may reveal the location of the paleo-cliffline formed the previous time the shore platform was occupied, allowing a long-term average cliff retreat rate to be determined from the difference to the modern cliff position (Regard et al., 2012).

5 6.5 Interpreting rates of cliff retreat from ^{10}Be concentrations

In order to determine rates of cliff retreat from ^{10}Be concentrations on shore platforms, results of a coupled morphological and CRN production model are compared to measured concentrations in order to statistically determine the most likely combinations of parameters and retreat rates that yield close fit between modelled and observed concentrations (Regard et al., 2012). In addition to high rates of cliff retreat, topographic shielding (Figure 4) due to adjacent sea cliffs and the presence of beaches 10 (Figure 8) that may have previously been more extensive (Figure 11) all act to reduce the amount of ^{10}Be in the shore platform, compared to a scenario with no cliffs and no beaches. Therefore, not accounting for the presence of beach material and topographic shielding is likely to lead to overestimation of rates of sea cliff retreat. Regard et al. (2012) didn't did not account for beach cover and how it may have changed through time, but observed beaches are relatively narrow and thin, and the cliffs at 15 their field site were small enough to have negligible effect on their estimated cliff retreat rates, given uncertainties in their ^{10}Be concentration measurements.

Observations of beach widths and berm heights may be made from both modern and historical data (e.g. Dornbusch et al., 2008), but there is little information about how beaches may have changed over millennial timescales. However, exposed shore platforms are most likely to be associated with locations with little or declining beach cover, since if beaches were accumulating the platform would not be exposed and cliff retreat rates might be expected to drop. The model predicts minimal differences in 20 the distribution of ^{10}Be concentrations in the shore platform surface for simulations with constant versus variable beach cover (Figure 10). A representative average beach width guided by historical observations may suffice when interpreting cliff retreat rates from ^{10}Be concentrations.

7 Conclusions

We find that the accumulation of ^{10}Be in the shore platform is primarily sensitive to the rate of cliff retreat. Concentrations 25 of ^{10}Be in the shore platform are also influenced by a number of other factors, including topographic shielding by sea cliffs, shielding due to beach and talus cover, water shielding due to tides and relative sea level change, the type of processes eroding the platform, and the style of platform evolution (steady state vs transient). These factors generally tend to reduce the production of ^{10}Be in the shore platform, particularly in the nearshore near-shore, nearest to cliffs and where the platform is most likely 30 to be covered by a beach. Nevertheless, comparison of measured ^{10}Be concentrations to model simulations that include these factors should allow determination of long-term average cliff retreats. If these factors are not adequately considered then there will be a tendency to overpredict over-predict cliff retreat rates and so cliff retreat rates derived from cosmogenic ^{10}Be might be considered as maximum estimates. The shape of the distribution of ^{10}Be across a shore platform can reveal whether cliff

retreat rates are declining or accelerating through time. We conclude that measurement of ^{10}Be in shore platforms has great potential to allow us to quantify long-term rates of cliff retreat and platform erosion.

Author contributions. MDH, DHR and MAE concieved the study; MDH wrote the software and performed the analyses; MDH wrote the manuscript with contributions from all authors.

5 *Acknowledgements.* The topographic and tide data used in this paper is available from the Channel Coast Observatory at www.channelcoast.org. All code used in this analysis is open source and can be downloaded from https://github.com/mdhurst1/RoBoCoP_CRN. We are grateful to Robert Anderson, Uwe Dornbusch, Zuzanna Swirad and Nick Rosser for discussion that shaped the development of this study.

References

Anderson, R. S., Densmore, A. L., and Ellis, M. A.: The generation and degradation of marine terraces, *Basin Research*, 11, 7–19, doi:10.1046/j.1365-2117.1999.00085.x, 1999.

Ashton, A. D., Walkden, M. J. A., and Dickson, M. E.: Equilibrium responses of cliffted coasts to changes in the rate of sea level rise, *Marine Geology*, 284, 217–229, doi:10.1016/j.margeo.2011.01.007, 2011.

Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J.: A complete and easily accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements, *Quaternary Geochronology*, 3, 174–195, doi:10.1016/j.quageo.2007.12.001, 2008.

Barkwith, A., Hurst, M. D., Thomas, C. W., Ellis, M. A., Limber, P. L., and Murray, A. B.: Coastal vulnerability of a pinned, soft-cliff coastline, II: Assessing the influence of sea walls on future morphology, *Earth Surface Dynamics*, 2, 233–242, doi:10.5194/esurf-2-233-2014, 2014.

Bradley, S. L., Milne, G. A., Shennan, I., and Edwards, R.: An improved glacial isostatic adjustment model for the British Isles, *Journal of Quaternary Science*, 26, 541–552, doi:10.1002/jqs.1481, 2011.

Braucher, R., Merchel, S., Borgomano, J., and Bourlès, D. L.: Production of cosmogenic radionuclides at great depth: A multi element approach, *Earth and Planetary Science Letters*, 309, 1–9, doi:10.1016/j.epsl.2011.06.036, 2011.

Braucher, R., Bourlès, D., Merchel, S., Vidal Romani, J., Fernandez-Mosquera, D., Marti, K., Léanni, L., Chauvet, F., Arnold, M., Aumaître, G., and Keddadouche, K.: Determination of muon attenuation lengths in depth profiles from in situ produced cosmogenic nuclides, *Nuclear Instruments and Methods in Physics Research, Section B: Beam Interactions with Materials and Atoms*, 294, 484–490, doi:10.1016/j.nimb.2012.05.023, 2013.

Bray, M. J. and Hooke, J. M.: Prediction of soft-cliff retreat with acceleration sea-level rise, *Journal of Coastal Research*, 13, 453–467, 1997.

Brooks, S. M. and Spencer, T.: Temporal and spatial variations in recession rates and sediment release from soft rock cliffs, Suffolk coast, UK, *Geomorphology*, 124, 26–41, doi:10.1016/j.geomorph.2010.08.005, 2010.

Bruun, P.: Coast Erosion and the Development of Beach Profiles, U.S. Army Beach Erosion Board Technical Memorandum 44, 44, 79, 1954.

Chao, R. B., Casais, M. C., Cortizas, A. M., Alberti, A. P., and Trenhaile, A. S.: Evolution and inheritance of a rock coast: Western Galicia, Northwestern Spain, *Earth Surface Processes and Landforms*, 28, 757–775, doi:10.1002/esp.496, 2003.

Chmeleff, J., von Blanckenburg, F., Kossert, K., and Jakob, D.: Determination of the ^{10}Be half-life by multicollector ICP-MS and liquid scintillation counting, *Nuclear Instruments and Methods in Physics Research, Section B: Beam Interactions with Materials and Atoms*, 268, 192–199, doi:10.1016/j.nimb.2009.09.012, 2010.

Choi, K. H., Seong, Y. B., Jung, P. M., and Lee, S. Y.: Using Cosmogenic ^{10}Be Dating to Unravel the Antiquity of a Rocky Shore Platform on the West Coast of Korea, *Journal of Coastal Research*, 282, 641–657, doi:10.2112/JCOASTRES-D-11-00087.1, 2012.

Codilean, A. T.: Calculation of the cosmogenic nuclide production topographic shielding scaling factor for large areas using DEMs, *Earth Surface Processes and Landforms*, 31, 785–794, doi:10.1002/esp.1336, 2006.

Costa, S., Delahaye, D., Freire-Diaz, S., Di Nocera, L., Davidson, R., and Plessis, E.: Quantification of the Normandy and Picardy chalk cliff retreat by photogrammetric analysis, *Geological Society, London, Engineering Geology Special Publications*, 20, 139–148, doi:10.1144/GSL.ENG.2004.020.01.11, 2004.

Dean, R. G. and Darlymple, R. A.: *Coastal Processes with Engineering Applications*, Cambridge University Press, 2002.

Desilets, D., Zreda, M., and Prabu, T.: Extended scaling factors for in situ cosmogenic nuclides: New measurements at low latitude, *Earth and Planetary Science Letters*, 246, 265–276, doi:10.1016/j.epsl.2006.03.051, 2006.

Dickson, M. E., Ogawa, H., Kench, P. S., and Hutchinson, A.: Sea-cliff retreat and shore platform widening: Steady-state equilibrium?, *Earth Surface Processes and Landforms*, 38, 1046–1048, doi:10.1002/esp.3422, 2013.

Dornbusch, U. and Robinson, D. A.: Block removal and step backwearing as erosion processes on rock shore platforms: A preliminary case study of the chalk shore platforms of south-east England, *Earth Surface Processes and Landforms*, 36, 661–671, doi:10.1002/esp.2086, 5 2011.

Dornbusch, U., Robinson, D. a., Moses, C., Williams, R., and Costa, S.: Retreat of Chalk cliffs in the eastern English Channel during the last century, *Journal of Maps*, 2, 71–78, doi:10.4113/jom.2006.46, <http://www.tandfonline.com/doi/abs/10.4113/jom.2006.46>, 2006.

Dornbusch, U., Robinson, D. A., Moses, C. A., and Williams, R. B. G.: Temporal and spatial variations of chalk cliff retreat in East Sussex, 1873 to 2001, *Marine Geology*, 249, 271–282, doi:10.1016/j.margeo.2007.12.005, 2008.

10 Dunai, T. J.: *Cosmogenic nuclides*, Cambridge University Press, Cambridge, UK ; New York, 1995.

Dunai, T. J.: Scaling factors for production rates of in situ production cosmogenic nuclides: a critical reevaluation., *Earth and Planetary Science Letters*, 176, 157–169, 2000.

Dunne, J., Elmore, D., and Muzikar, P.: Scaling factors for the rates of production of cosmogenic nuclides for geometric shielding and attenuation at depth on sloped surfaces, *Geomorphology*, 27, 3–11, doi:10.1016/S0169-555X(98)00086-5, 1999.

15 Gosse, J. C. and Phillips, F. M.: Terrestrial in situ cosmogenic nuclides:theory and application, *Quaternary Science Reviews*, 20, 1475–1560, doi:10.1016/S0277-3791(00)00171-2, 2001.

Granger, D. E., Lifton, N. A., and Willenbring, J. K.: A cosmic trip: 25 years of cosmogenic nuclides in geology, *Bulletin of the Geological Society of America*, 125, 1379–1402, doi:10.1130/B30774.1, 2013.

Heisinger, B., Lal, D., Jull, A., Kubik, P., Ivy-Ochs, S., Knie, K., and Nolte, E.: Production of selected cosmogenic radionuclides by muons: 20 2. Capture of negative muons, *Earth and Planetary Science Letters*, 200, 357–369, 2002a.

Heisinger, B., Lal, D., Jull, J., Kubik, P., Ivy-Ochs, S., Neumaier, S., Knie, K., Lazarev, V., and Nolte, E.: Productioin of selected cosmogenic radionuclides by muons 1. Fast muons, *Earth and Planetary Science Letters*, 200, 345–355, 2002b.

Hurst, M. D., Barkwith, A., Ellis, M. A., Thomas, C. W., and Murray, A. B.: Exploring the sensitivities of crenulate bay shorelines to wave climates using a new vector-based one-line model, *Journal of Geophysical Research F: Earth Surface*, 120, 2586–2608, 25 doi:10.1002/2015JF003704, 2015.

Katz, O. and Mushkin, A.: Characteristics of sea-cliff erosion induced by a strong winter storm in the eastern Mediterranean, *Quaternary Research (United States)*, 80, 20–32, doi:10.1016/j.yqres.2013.04.004, 2013.

Kennedy, D. M., Stephenson, W. J., and Naylor, L. a.: Chapter 1 Introduction to the rock coasts of the world, *Geological Society, London, Memoirs*, 40, 1–5, doi:10.1144/M40.1, <http://mem.lyellcollection.org/cgi/doi/10.1144/M40.1>, 2014.

30 Komar, P. D. and Gaughan, M. K.: Airy Wave Theory and Breaker Height Prediction, *Proceedings 13th Conference of Coastal Engineering*, 1972, 405–418, doi:10.1061/9780872620490.023, 1972.

Korschinek, G., Bergmaier, A., Faestermann, T., Gerstmann, U. C., Knie, K., Rugel, G., Wallner, A., Dillmann, I., Dollinger, G., von Gostomski, C. L., Kossett, K., Maiti, M., Poutivtsev, M., and Remmert, A.: A new value for the half-life of ^{10}Be by Heavy-Ion Elastic Recoil Detection and liquid scintillation counting, *Nuclear Instruments and Methods in Physics Research, Section B: Beam Interactions with Materials and Atoms*, 268, 187–191, doi:10.1016/j.nimb.2009.09.020, 2010.

35 Lal, D.: Cosmic-ray labeling of erosion surfaces - insitu nuclide production-rates and erosion models, *Earth and Planetary Science Letters*, 104, 424–439, doi:doi:10.1016/0012-821X(91)90220-C |, 1991.

Lifton, N. A., Bieber, J. W., Clem, J. M., Duldig, M. L., Evenson, P., Humble, J. E., and Pyle, R.: Addressing solar modulation and long-term uncertainties in scaling secondary cosmic rays for in situ cosmogenic nuclide applications, *Earth and Planetary Science Letters*, 239, 140–161, doi:10.1016/j.epsl.2005.07.001, 2005.

5 Limber, P. W. and Murray, A. B.: Beach and sea-cliff dynamics as a driver of long-term rocky coastline evolution and stability, *Geology*, 39, 1147–1150, doi:10.1130/G32315.1, 2011.

Matsumoto, H., Dickson, M. E., and Kench, P. S.: An exploratory numerical model of rocky shore profile evolution, *Geomorphology*, 268, 98–109, doi:10.1016/j.geomorph.2016.05.017, 2016.

10 Mudd, S. M., Harel, M.-A., Hurst, M. D., Grieve, S. W. D., and Marrero, S. M.: The CAIRN method: Automated, reproducible calculation of catchment-averaged denudation rates from cosmogenic radionuclide concentrations, *Earth Surface Dynamics*, pp. 655–674, doi:10.5194/esurf-4-655-2016, 2016.

Naylor, L. A., Stephenson, W. J., Smith, H. C. M., Way, O., Mendelsohn, J., and Cowley, A.: Geomorphological control on boulder transport and coastal erosion before, during and after an extreme extra-tropical cyclone, *Earth Surface Processes and Landforms*, 41, 685–700, doi:10.1002/esp.3900, 2016.

15 Porter, N. J., Trenhaile, A. S., Prestanski, K., and Kanyaya, J. I.: Patterns of surface downwearing on shore platforms in eastern Canada, *Earth Surface Processes and Landforms*, 35, 1793–1810, doi:10.1002/esp.2018, 2010.

Pugh, D. and Woodworth, P.: *Sea-level science: understanding tides, surges, tsunamis and mean sea-level changes*, Cambridge University Press, 2014.

Recorbet, F., Rochette, P., Braucher, R., Bourlès, D., Benedetti, L., Hantz, D., and Finkel, R. C.: Evidence for active retreat of a coastal cliff between 3.5 and 12 ka in Cassis (South East France), *Geomorphology*, 115, 1–10, doi:10.1016/j.geomorph.2009.04.023, 2010.

20 Regard, V., Dewez, T., Bourlès, D. L., Anderson, R. S., Duperret, A., Costa, S., Leanni, L., Lasseur, E., Pedoja, K., and Maillet, G. M.: Late Holocene seacliff retreat recorded by ^{10}Be profiles across a coastal platform: Theory and example from the English Channel, *Quaternary Geochronology*, 11, 87–97, doi:10.1016/j.quageo.2012.02.027, 2012.

Robinson, L. A.: Erosive processes on the shore platform of northeast Yorkshire, England, *Marine Geology*, 23, 339–361, doi:10.1016/0025-3227(77)90038-X, 1977.

25 Rogers, H. E., Swanson, T. W., and Stone, J. O.: Long-term shoreline retreat rates on Whidbey Island, Washington, USA, *Quaternary Research* (United States), 78, 315–322, doi:10.1016/j.yqres.2012.06.001, 2012.

Stephenson, W. J., Kirk, R. M., Hemmingsen, S. A., and Hemmingsen, M. A.: Decadal scale micro erosion rates on shore platforms, *Geomorphology*, 114, 22–29, doi:10.1016/j.geomorph.2008.10.013, 2010.

30 Stephenson, W. J., Kirk, R. M., Kennedy, D. M., Finlayson, B. L., and Chen, Z.: Long term shore platform surface lowering rates: Revisiting Gill and Lang after 32 years, *Marine Geology*, 299–302, 90–95, doi:10.1016/j.margeo.2012.01.005, 2012.

Stone, J. O.: Air pressure and cosmogenic isotope production, *Journal of Geophysical Research*, 105, 23 753, doi:10.1029/2000JB900181, 2000.

Sunamura, T.: *Geomorphology of Rocky Coasts*, J. Wiley, 1992.

35 Sunamura, T.: Review Rocky coast processes: with special reference to the recession of soft rock cliffs, *Proceedings of the Japanese Academy, Series B*, 91, 481–500, doi:10.2183/pjab.91.481, 2015.

Trenhaile, A. S.: Modeling the development of wave-cut shore platforms, *Marine Geology*, 166, 163–178, doi:10.1016/S0025-3227(00)00013-X, 2000.

Trenhaile, A. S.: Modeling the effect of late Quaternary interglacial sea levels on wave-cut shore platforms, *Marine Geology*, 172, 205–223, doi:10.1016/S0025-3227(00)00136-5, 2001.

Trenhaile, A. S.: The effect of Holocene changes in relative sea level on the morphology of rocky coasts, *Geomorphology*, 114, 30–41, doi:10.1016/j.geomorph.2009.02.003, 2010.

5 Trenhaile, A. S.: Climate change and its impact on rock coasts, *Geological Society, London, Memoirs*, 40, 7–17, doi:10.1144/M40.2, <http://mem.lyellcollection.org/content/40/1/7.full>, 2014.

Walkden, M. J. A. and Hall, J. W.: A predictive Mesoscale model of the erosion and profile development of soft rock shores, *Coastal Engineering*, 52, 535–563, doi:10.1016/j.coastaleng.2005.02.005, 2005.

West, A. J., Hetzel, R., Li, G., Jin, Z., Zhang, F., Hilton, R. G., and Densmore, A. L.: Dilution of ^{10}Be in detrital quartz by earthquake-induced
10 landslides: Implications for determining denudation rates and potential to provide insights into landslide sediment dynamics, *Earth and Planetary Science Letters*, 396, 143–153, doi:10.1016/j.epsl.2014.03.058, 2014.




Figure 1. Schematic diagram of a rocky coast showing the expected distribution of ^{10}Be across a shore platform. Cliff retreat exposes pristine platform with low ^{10}Be concentrations. Exposure time increases with distance from the cliff (increasing ^{10}Be concentrations), but platform downwear removes ^{10}Be -rich rock, and increased water shielding reduces ^{10}Be production offshore. The result is a hump-shaped distribution where the magnitude of the hump is inversely proportional to rate of cliff retreat.




Figure 2. (a) End-member types of shore platform as defined by Sunamura (1992). (b) Illustration of steady state shore profile retreat subject to relative sea level rise.




Figure 3. Example of beach model used in modelling ^{10}Be accumulation in a shore platform. Beach is defined by a beach width B_w and berm height B_h and a Bruun profile (Bruun, 1954) seaward of the defined beach width.




Figure 4. Topographic Shielding due to sea cliff. (a) Analytical model for topographic shielding on a shore platform as a function of distance from cliff measured according to Equation 11 for cliff heights ranging from 5 to 100 m. (b) Example map of the distribution of topographic shielding across a shore platform at Beachy Head, East Sussex, UK. (c) Comparison of Equation 11 to values measured using distributed shielding routines (Codilean, 2006; Mudd et al., 2016).




Figure 5. Influence of tidal regime on the distribution of ^{10}Be Production across a shore platform. (a)-(f) Tidal water levels over a 28 day period for the hypothetical tidal regimes explored here (see Table 1). (g) Production rates relative to the case where there is no water shielding ($h_w = 0$) as a function of distance offshore on a shore platform with gradient $\alpha = 1/100$ for the different tidal regimes plotted in (b)-(g).




Figure 6. Distribution of ^{10}Be Production across a shore platform with gradient $\alpha = 1/100$ for a simple diurnal tide model with amplitude 2.4 m and for real tide data from Newhaven, East Sussex, UK. Note there is very little difference in production between the two, suggesting a diurnal model with representative amplitude is sufficient for the purpose of modelling ^{10}Be production.




Figure 7. (a) Evolution of stepped shore platform profiles over 10 ka for step sizes (bedding thickness) up to 1 m. (b) Corresponding distribution of ^{10}Be as a function of step size. Large steps due to block removal processes result in a saw-toothed distribution superimposed on the expected humped distribution. Variability in the platform surface concentration increases with step size.




Figure 8. (a) Evolution of shore platform profiles over 10 ka with constant beach cover in the [nearshore](#)-[near-shore](#) described by a Bruun profile (Equation 6). (b) Corresponding distribution of ¹⁰Be as a function of beach width. Beach cover reduces the concentrations of ¹⁰Be in the platform surface, and wider beaches result in lower ¹⁰Be concentrations and cause the position of the peak in concentrations to be further out from the cliff. Note that quite extreme beach cover $B_W \geq 50 \text{ m}$ is required to significantly ($\geq 15\%$) reduce platform ¹⁰Be concentrations.




Figure 9. Reduction in the magnitude of peak concentration of ^{10}Be with increasing beach width. Black symbols shows the peak concentrations taken from Figure 8. The relationship is approximately linear as shown by the dashed line fit by least squares regression. Note that quite extreme beach cover $B_W \geq 50$ m is required to significantly ($\geq 15\%$) reduce platform ^{10}Be concentrations.




Figure 10. (a) Variation in beach width through time modelled as a sinusoidal function with a beach width $B_w = 50 \pm 30 \text{ m}$ and wavelength 100 years. (b) Evolution of shore platform profiles over 10 ka with variable beach cover in the [nearshore](#)-[near-shore](#) described by a Bruun profile (Equation 6). (c) Corresponding distribution of ^{10}Be for the case with no beach cover, fixed $B_w = 50 \text{ m}$, and variable beach cover. Note that there is little difference in ^{10}Be concentrations between a constant beach cover and variable beach cover.




Figure 11. (a) Evolution of shore platform profiles over 10 ka with variable beach cover in the [nearshore](#) [near-shore](#) described by a Bruun profile (Equation 6). Beach width reduces to $B_w = 0 \text{ m}$ during the last 1000 yrs of the model runs. (b) Corresponding distribution of ^{10}Be for the case with no beach cover and thinning beaches during the last 100 yrs. The presence of beaches that dissapeared over the last 1000 yrs would result in lower concentrations of ^{10}Be in the platform surface.




Figure 12. Transient simulations of shore profile evolution and ^{10}Be concentrations for the case with no sea level rise. (a) Evolution of shore platform profiles over 10 ka driven by RoBoCoP (colour-coded in 1 ka interval). (b) Corresponding ^{10}Be concentration predictions, normalised to the position of the cliff, (colour-coding also corresponds to the median retreat rates listed). Retreat rates are initially rapid and decline through time (c) and show a humped distribution, similar to steady state model predictions. The magnitude of the hump increases through time as the length of exposure increases and cliff retreat rates and platform downwear rates decline, resulting in a decrease in platform gradient through time (d).




Figure 13. Comparison of transient simulations from Figure 12 to predictions assuming steady state profile retreat. Steady state models are run with constant retreat rate taken as the median values in the transient simulations (Figure 12). Color coding corresponds to the median retreat rates reported in Figure 12b, and the instantaneous mean platform gradient for each profile in 12a. (a) Distribution of ¹⁰Be concentration predictions. The magnitude of the hump is similar as shown in (b), but the location of the peak in concentration is further offshore when steady state retreat is assumed, as also shown in (c).




Figure 14. Transient simulations of shore profile evolution and ^{10}Be concentrations for the case with 0.5 mm yr^{-1} relative sea level rise. (a) Evolution of shore platform profiles over 10 ka driven by RoBoCoP (1 ka interval). Retreat rates are initially rapid but steady to 32 cm yr^{-1} after 1 K yrs. Platform gradient declines through time. (b) Corresponding ^{10}Be concentration predictions, normalised to the position of the cliff. Note that ^{10}Be concentrations are consistent with distance from the cliff throughout the model runs, consistent with steady-state morphological retreat. The dashed line shows the predictions of a steady-state morphological model with retreat rate of 32 cm yr^{-1} , platform gradient of 1/500 and relative sea level rise rate of 0.5 mm yr^{-1} . Steady state model predicts higher concentrations than transient simulations by about a factor of 2.

Table 1. Tidal constituents used to generate synthetic tides to explore variation in ^{10}Be production under different tidal regimes.

Tidal Constituent	Tide A		Tide B		Tide C		Tide D		Tide E		Tide F	
	H_n	σ_n										
$K1$	2.5	15.041	-	-	0.7	15.041	2.0	15.041	-	-	0.4	15.041
$K2$	-	-	-	-	-	-	-	-	-	-	0.1	30.082
$M1$	-	-	-	-	-	-	-	-	-	-	0.1	14.492
$M2$	-	-	2.5	28.984	1.8	28.984	-	-	2.0	28.984	1.5	28.984
$N2$	-	-	-	-	-	-	-	-	0.5	28.440	0.5	28.440
$O1$	-	-	-	-	-	-	0.5	13.943	-	-	0.2	13.043

Table 2. Global parameters used in modelling ^{10}Be accumulation in a shore platform

Description	Symbol	Value
Acceleration due to Gravity	g	9.81 m s ⁻²
Density of sea water	ρ_w	1025 kg m ⁻³
Density of rock	ρ_r	1800 kg m ⁻³
Surface production rate (spallogenic)	$P_0(s)$	4.0 atoms g ⁻¹ yr ⁻¹
Surface production rate (muogenic)	$P_0(\mu)$	0.028 atoms g ⁻¹ yr ⁻¹
^{10}Be decay constant	λ	4.99×10^{-7}
Attenuation rate (spallogenic)	Λ_s	1600 kg m ⁻²
Attenuation rate (muogenic)	Λ_μ	42000 kg m ⁻²

Table 3. Parameters used in modelling transient shore platform evolution

Description	Symbol	Value
Initial platform gradient	α	0.1 m m ⁻¹
Offshore Wave Height	H_0	1.0 m
Wave Period	T	6 s
Resistance Coefficient	K	10 ⁻⁴ m s kg ⁻¹
Wave energy dissipation coefficient	k	0.2 (Dimensionless)
Tidal amplitude	H_n	1.0 m
Angular speed	σ_n	28.984 ° hour ⁻¹