

## ***Interactive comment on “SHORT COMMUNICATION: Massive Erosion in Monsoonal Central India Linked to Late Holocene Landcover Degradation” by Liviu Giosan et al.***

**Y. KULKARNI**

kulyogesh19@gmail.com

Received and published: 21 August 2017

The short communication is based on the landcover, soil and its erosion from two different zones of lithologies present in Godavari Drainage Basin (GDB) which is also previously discussed by Kulkarni et al. (2015) based on the mineral magnetic studies of Godavari sediments and Bay of Bengal sediments. Their study show the general increasing trend in ferrimagnetic concentration in middle Bay of Bengal sediments suggesting to the dominance of Deccan source over quartzo-feldspathic source for Late Holocene. They suggested the combine effect of distinct lithological units, geomorphic setup and spatial distribution of monsoonal rainfall plays an important role in sediment

Printer-friendly version

Discussion paper



generation in GDB. Previously Sangode et al. (2001) also suggested the dominance of Peninsular Source over Himalayan source in Bay of Bengal sediments. Kulkarni et al (2014) based on mineral magnetic study inferred the dominant Deccan basaltic source over the floodplains of Godavari River for entire stretch as a result of intense weathering of Deccan plateau.

Recently Cui et al. (2017) based on organic carbon and mineral magnetic analysis of well dated (AMS 14C) Godavari delta core 'CY' suggested the increasing Deccan basaltic source during Late Holocene particularly after nearly 6 cal ka BP. Their study also show the significant increase in ferrimagnetic minerals from nearly 3.2 to 3.1 cal. ka BP attested to severe decline in vegetation in Deccan plateau. Majority of CMZ is represented by Pranhita River basin rather than upper Godavari and the mixing may have a complex role in Godavari onwards is not considered. Besides this, the Bikshamaiah and Subramaniyan (1980) although given detail account on chemical and sediment mass transfer in GDB and established some controlling factors for same, they did not classify the geology of GDB as two major geological units. The part of Godavari River in the Godavari graben is flowing above the Gondwana and Purana sedimentary units which although accounts for about 11% of basin (Biksham and Subramaniyan, 1988) but acts as major lithounit in main channel as well as in Pranhita River (A major tributary of Godavari).

## References:

1. Cui, M, Wang, Zhanghua ,Nageshwar Rao, Kakani , Sangode, S J, Saito, Yoshiki, Ting, Chen, Kulkarni Y.R., G naga Kumar, K Ch V, Demudu (2017) A mid-to-late Holocene record of vegetation decline and erosion triggered by monsoon weakening and human adaptations in the south –east Indian Peninsula; The Holocene, 0959683617715694.
2. Kulkarni, Y.R., Sangode, S.J., Bloemandal, J., Meshram, D.C.,Suresh, N. (2015) Mineral Magnetic Characterization of the Godavari River and Western Bay of Bengal

Printer-friendly version

Discussion paper



Sediments: Implications to Source to Sink Relations; Journal of Geological Society of India, v. 85, pp. 71-78

3. Kulkarni, Y.R., Sangode, S.J., Meshram, D.C., Patil, S.K. and Dutt, Yatindra (2014) Mineral magnetic characterization of the Godavari River sediments; Journal of Geological Society of India, v. 81, pp. 376-384

4. Sangode, S J, Suresh, N and Bagati, T N (2001) Godavari source in the Bengal fan sediments: results from magnetic susceptibility dispersal pattern. Current Science, 660-664.

5. Biksham, G. and Subramanian, V. (1988) Elemental composition of Godavari sediments (central and southern Indian subcontinent). Chemical Geology, 70 275-286 275

---

Interactive comment on Earth Surf. Dynam. Discuss., <https://doi.org/10.5194/esurf-2017-35>, 2017.

Printer-friendly version

Discussion paper

