

28 Abstract

29 Soil erosion plays a crucial role in transferring sediment and carbon from land to sea, yet
30 little is known about the rhythm and rates of soil erosion prior to the most recent few
31 centuries. Here we reconstruct a Holocene erosional history from central India, as
32 integrated by the Godavari River in a sediment core from the Bay of Bengal. We quantify
33 terrigenous fluxes, fingerprint sources for the lithogenic fraction and assess the age of the
34 exported terrigenous carbon. Taken together, our data show that the monsoon decline in
35 the late Holocene significantly increased soil erosion and the age of exported organic
36 carbon. This acceleration of natural erosion was later exacerbated by the Neolithic
37 adoption and Iron Age extensification of agriculture on the Deccan Plateau. Despite a
38 constantly elevated sea level since the middle Holocene, this erosion acceleration led to a
39 rapid growth of the continental margin. We conclude that in monsoon conditions aridity
40 boosts rather than suppresses sediment and carbon export acting as a monsoon erosional
41 pump modulated by landcover conditions.

42

43 1. Soil Erosion in the Holocene

44

45 On decadal to millennial timescales, climate is the principal natural control on soil
46 erosion via changes in temperature and precipitation as well as their impact on vegetation
47 type and cover (Allen and Breshears, 1998; Reichstein et al., 2013). Global sediment
48 budgets for the Holocene indicate that humans surpassed these natural controls and
49 became the main driver of soil erosion by at least 2000 years ago (Montgomery, 2007;
50 Wilkinson and McElroy, 2007; Dotterweich, 2013). Transfer of sediment, carbon and
51 solutes from land to ocean is of crucial importance for understanding continental margin
52 architecture as well as carbon and other elemental cycles. For example, soils contain
53 about two times more carbon than the atmosphere and, as a result, small changes in the
54 residence time of organic carbon in soils can significantly affect the atmospheric
55 inventory of carbon dioxide (Lal, 2004). Besides heterotrophic microbial respiration,
56 erosion is the principal process that releases carbon from soils. Eroded carbon can
57 subsequently be degraded/reburied along the aquatic continuum to the ocean (Stallard,
58 1998; Aufdenkampe et al., 2011; van Oost et al., 2012).

59 In the absence of historical documentation of human impacts, the complexity of soil
60 erosion hampers the reconstruction of carbon transfer processes prior to the last few
61 centuries (e.g., Hoffmann et al. 2013; Dotterweich, 2013; Vanwalleghem et al., 2017).
62 Consequently, global carbon budgets implicitly assume steady state conditions for lateral
63 transport and carbon degradation along the aquatic continuum in pre-industrial times
64 (Battin et al., 2009; Regnier et al., 2013; Chappell et al., 2015). In contrast, abundant
65 archaeological and geological evidence (e.g., van Andel et al., 1990; Bork and Lang,
66 2003; Bayon et al., 2012; Dotterweich, 2013) as well as modeling (Kaplan et al., 2010;
67 Wang et al., 2017) suggests widespread impacts of early human landuse on continental
68 landscapes, soil erosion and associated carbon transfer processes.

69 Here we present a soil erosion history from the Indian peninsula recorded in a sediment
70 core retrieved near the mouth of Godavari River (Fig. 1) in the Bay of Bengal (NGHP-
71 01-16A at 16°35.6'N, 82°41.0'E; 1,268 m water depth; Collett et al., 2015). The age
72 model for the core based on 11 radiocarbon dates on mixed planktonic foraminifera was
73 previously published by Ponton et al. (2012). The Godavari basin was not affected by
74 tectonics at the Holocene time scale, or glacial/snow meltwater and strong orographic
75 precipitation, which augment and complicate the water and sediment discharge of the
76 larger Himalayan rivers like the Ganges or Brahmaputra. Instead, it integrates rainfall
77 from the core monsoon zone (CMZ), the region of central India that is representative for
78 both the mean monsoon regime and its fluctuations over the peninsula (see Ponton et al.,
79 2012 and references therein). Consequently, over 90% of Godavari's water discharge into
80 the Bay of Bengal derives from summer monsoon precipitation (Rao et al., 2005), making
81 its sedimentary deposits a prime target for continental climate reconstructions and a

82 repository for sedimentary proxies of erosion prior and after the Neolithic adoption of
83 agriculture in central India.

84

85 2. The Godavari Sediment System

86 Originating at an elevation of 920 m in the Sahyadri coastal range (*aka* Western Ghats)
87 near the Arabian Sea coast, the Godavari crosses the entire Indian peninsula toward the
88 Bay of Bengal (Fig. 1a). Currently the water discharge of the river is $\sim 85 \text{ km}^3/\text{yr}$ with a
89 sediment load of $\sim 175 \text{ MT}/\text{yr}$ (Syvitski and Saito, 2007). Because the coastal range limits
90 penetration of the Arabian Sea moisture delivered by the monsoon, precipitation in the
91 Godavari basin primarily originates from the Bay of Bengal (Gunnell et al. 2007). As a
92 result, the climate is most humid at the coast (i.e., Eastern Ghats range) and becomes
93 increasingly arid toward the interior on the Deccan Plateau (Fig. 1b). The natural
94 vegetation reflects this gradual decrease in moisture: the headwaters on the Deccan are
95 dominated by C_4 -plant thornbush savannah adapted to dry conditions, whereas C_3 -flora
96 (deciduous forests) are dominant in the Eastern Ghats (Asouti and Fuller, 2008; Fig. 1c).

97 Sediments transported by the Godavari are sourced from two major geological units
98 (Bikshamaiah and Subramanian, 1980). The upper river basin developed on the Deccan
99 Traps, a large igneous province consisting of relatively young flood basalts (Cretaceous
100 to early Neogene) that largely span the Deccan Plateau. The lower river basin developed
101 over old Proterozoic to Archaean crystalline igneous/metamorphic rocks of the Indian
102 Craton (Fig. 1d). The relatively young Deccan basalts retain a highly radiogenic mantle-
103 derived Nd isotope composition (ϵNd of $+1\pm 5$) while the old continental crust of the
104 Indian Craton has a relatively unradiogenic isotopic composition (ϵNd of -35 ± 8), yielding
105 a sharp contrast between geological end-members. Thus, the sediment provenance for the
106 Godavari sediments can be deduced from the Nd isotopic signatures of the detrital
107 inorganic fraction in our core because the Nd signal remains unmodified through bedrock
108 weathering processes (McLennan and Hemming, 1992; DePaolo, 1988).

109 Black soils cover the Deccan Plateau whereas red soils are generally typical for the
110 Eastern Ghats (Bhattacharyya et al., 2003; Fig. 1d). Although both types of soils have
111 been affected by landuse since prehistorical times, the black soils of the arid to semi-arid
112 Deccan appear to be the most degraded at present (Singh et al., 1992). Intense erosion
113 within the basin is reflected by the inordinately large sediment load of the Godavari
114 (Bikshamaiah and Subramanian, 1980) similar to other monsoonal rivers (Summerfield
115 and Hulton, 1994). In contrast to the dynamic Himalayan rivers of the Indo-Gangetic
116 alluvial plain, Godavari and its tributaries are incised in rock or alluvium and have
117 relatively stable sandy channels. As for other rivers affected by storms (Edwards and
118 Owens, 1991; Hilton et al, 2008), extreme rainfall events are disproportionately important

119 for erosion in the Godavari watershed and in subsequent transport of sediments to the
120 ocean (Kale, 2003). Given their incised morphology, shifts in channel position in
121 response to floods are however rare above the Godavari delta (Kale, 2002). Floodplains
122 are limited in extent (2% of the basin; Bikshamaiah and Subramanian, 1980), and loss of
123 sediments to overbank deposition is minor (Kale, 2002). Therefore storage is minimal in
124 these intermediate alluvial reservoirs that normally would increase the residence time of
125 sediments, including particulate organic carbon.

126 Once reaching the Bay of Bengal, sediment delivered by the Godavari has constructed a
127 large Holocene delta (Rao et al. 2005; Cui et al., 2017). Offshore from the Godavari
128 mouth, a persistent sediment plume extends over 300 km during the monsoon season
129 when over 90% of the fluvial sediment is discharged (Sridhar et al., 2008). Because the
130 shelf in front of the delta is unusually narrow (i.e., under 10 km at our core location)
131 copious sediment deposition takes place directly on the continental slope, resulting in
132 sediment accumulation rates as high as 250 cm/kyr; Ponton et al., 2012). Owing to the
133 narrow shelf, changes in sea level would also have minimal effects on sediment
134 deposition at our site, especially after the early Holocene when the global sea level
135 reached within a few meters of modern values (Lambeck et al., 2014). For these reasons
136 our core located close to the river mouth (~35 km) is unlikely to contain any significant
137 contributions from other sediment sources, in agreement with previous studies (e.g.,
138 Bejugam and Nayak, 2017).

139 The relatively simple sedimentary regime of the Godavari system in combination with the
140 monsoon-dominated climatology and simple geology of the Godavari basin allows for
141 relatively straightforward interpretation of sediment sources and transfer processes. The
142 monsoon washload is rapidly and directly delivered to the continental margin without
143 significant trapping in intermediate depocenters along the river. As the suspended load
144 makes up over 95% of the total sediment transported by the Godavari (Syvitski and Saito,
145 2007), the washload-derived terrestrial proxies are representative for the production of
146 fine-grained sediment in the basin. Potential contributions from resuspension of shelf
147 sediments cannot be excluded but are likely minor due to the narrowness of the shelf;
148 furthermore, given the large sedimentation rates on the shelf itself (Forsberg et al., 2007),
149 the resuspended sediment is expected to be quasi-contemporaneous with sediments
150 arriving on the slope directly from the river plume.

151 3. Hydroclimate in the Core Monsoon Zone

152 We have previously reconstructed the Holocene paleoclimate using the same sediment
153 core discussed herein (Ponton et al., 2012; Zorzi et al., 2015). Terrestrial reconstructions
154 were based on the carbon isotopic compositions of higher plant leaf-wax biomarkers (i.e.,
155 long-chain *n*-alkanoic acids C₂₆₋₃₂) and pollen, whereas contemporaneous sea surface
156 paleoceanographic conditions in front of the Godavari delta came from the oxygen

157 isotopic composition of planktonic foraminifer *Globigerinoides ruber*. Sedimentary leaf
158 waxes provide an integrated $\delta^{13}\text{C}$ record of the flora in the CMZ that document an
159 increase in aridity-adapted vegetation (C_4 plants) after the monsoon maximum in the
160 early Holocene (Ponton et al., 2012; Fig. 2). The overall trend of the $\delta^{13}\text{C}$ leaf wax record
161 agrees with the view that the seasonality of Northern Hemisphere insolation (Ponton et
162 al., 2012) led to progressively weaker monsoons over the Holocene. However, two clear
163 aridification steps are evident: between ~ 5000 and 4500 years ago, and $\sim 1,700$ years ago
164 (Fig. 2). Pollen from the same core (Zorzi et al., 2015) reinforces these conclusions:
165 coastal forest and mangrove pollen (Fig. 2) that are typical for the more humid coastal
166 regions of the Eastern Ghats and Godavari delta declined over the Holocene.

167 Dryness-adapted thornbush pollen from the Deccan Plateau increased substantially after
168 the second aridification step ~ 1700 years ago, overlapping well with the maximum
169 contribution of C_4 plant-derived leaf waxes (see Zorzi et al., 2015). For the same time
170 interval, the ice volume-corrected oxygen isotopic composition of planktonic foraminifer
171 *Globigerinoides ruber* documented a series of low values interpreted as high salinity
172 events at the Godavari mouths (see Ponton et al., 2012). Together these continental and
173 oceanic records suggest that the CMZ aridification intensified in the latest Holocene via a
174 series of short drier episodes (Ponton et al., 2012). This interpretation is reinforced by
175 speleothem-derived records from central and northern India for the past thousand years
176 (Sinha et al., 2011), and the overall evolution of the CMZ hydroclimate as seen from our
177 core is supported by local reconstructions from the Lonar crater lake in central India
178 (Prasad et al., 2014; Sarkar et al., 2015), Godavari delta (Cui et al., 2017) and other
179 records from the larger Indian monsoon domain (Gupta et al., 2003; Fleitmann et al.,
180 2003; Prasad and Enzel, 2006; Berkelhammer et al., 2012; Dixit et al., 2014).

181 4. Erosion in the Godavari Basin

182 The Holocene sediment flux at our core location (Fig. 2) is representative for the
183 Godavari continental slope (Mazumdar et al., 2009; Ramprasad et al. 2011; Joshi et al.,
184 2014) and is driven by changes in the siliciclastic sedimentation rate as dilution by
185 biogenic carbonates is less than 5% (Johnson et al., 2014). Despite a lower sea level at
186 the time, the flux was relatively small in the early Holocene ($\sim 25 \text{ g/cm}^2/\text{kyr}$) but began to
187 increase after 6000 years ago ($\sim 40 \text{ g/cm}^2/\text{kyr}$), as soon as the monsoon started to decline
188 but well before the adoption of Neolithic agriculture and settlement of the savannah zone
189 of the central peninsula (~ 4500 years ago; Fuller, 2011). Between 4000 and 3500 years
190 ago permanent agricultural settlements spread throughout the Deccan Plateau. The
191 associated small-scale metallurgy (copper-working) requiring firewood together with the
192 agricultural intensification probably also affected erosion via widespread deployment of
193 two cropping seasons (Kajale, 1988; Fuller and Madella, 2001). As the climate remained
194 arid, sediment fluxes stayed high despite a phase of agricultural abandonment and

195 depopulation between ~3,200 and 2,900 years ago (Dhavalikar, 1984; Roberts et al.,
196 2016).

197 A further step increase in the sediment flux (~90 g/cm²/kyr on average) occurred after
198 ~3000 years ago, this time with no apparent concurrent change in climate. The Nd
199 isotopic signal points to an increase in the Deccan sedimentary output at the time, after a
200 muted variability earlier in the Holocene when the Indian Craton consistently provided
201 ~50-60% of the sediments (Fig. 2; see Supplementary Materials). Ferrimagnetic minerals
202 interpreted as originating from the Deccan (Sangode et al., 2001; Kulkarni et al., 2014)
203 also increase in late Holocene sediments in the Godavari delta (Cui et al., 2017) and Bay
204 of Bengal (Kulkarni et al., 2015) supporting our interpretation. Augmented Deccan input
205 was suggested for the Godavari delta even earlier after ~6000 years ago (Cui et al., 2017),
206 in step with the initial aridification.

207 New improvements in agricultural technology became widespread in the Deccan Plateau,
208 including use of iron agricultural tools (Mohanty and Selvakumar, 2001) that required
209 firewood-fuelled smelting (Fuller, 2008). A new phase of agricultural settlement began in
210 the middle Godavari basin (eastern Maharashtra) between ~3000 to ~2800 years ago
211 (Brubaker 2000). However, the largest boost in sediment flux occurred after ~2000 years
212 ago, when the monsoon reached its driest phase and when further increases in population
213 occurred resulting in the founding of towns and the first cities of the region at the
214 beginning of the Historic period (Allchin 1995; Parabrahma Sastry 2003). This doubling
215 in sediment flux relative to the early Holocene values involved a basin-wide increase in
216 erosion. The contribution from the Deccan Plateau, although at its maximum according to
217 the Nd isotope mixing model, only accounts for a 15% shift in sediment source (Fig. 2).

218 Overall, watersheds with high precipitation have higher discharge and discharge
219 magnitude is considered a primary regulator for sediment and carbon erosional fluxes to
220 the ocean (e.g., Summerfield and Hulton, 1994; Ludwig et al., 1996; Galy et al., 2015).
221 However, our Godavari record shows that erosional output is maximized by aridity
222 because significant rain and seasonal floods still occur during the summer monsoon
223 season (Mujumdar et al., 1970; Kale, 2003). Aridification and/or agricultural expansion
224 lead to changes in vegetation type (i.e., forest decrease in favour of savannah) and cover
225 (i.e., shrinking of naturally vegetated lands in favour of agricultural and/or degrading arid
226 lands) that exacerbate soil erosion (i.e., Langbein and Schum, 1958; Dunne, 1979;
227 Walling and Webb, 1983; Istanbuluoglu and Bras, 2005; Vanacker et al., 2007; Collins
228 and Bras, 2008).

229 5. Carbon Export from the Godavari Basin

230 The terrigenous organic carbon exported by rivers consists of a mixture of dissolved and
231 particulate components derived from contemporary vegetation and of carbon stored in

232 bedrock, soils and fluvial sediments that may be significantly pre-aged (Smittenberg et
233 al., 2006; Galy and Eglinton, 2011; Feng et al., 2013). On the Godavari slope, the
234 terrigenous fraction dominates the total organic carbon (TOC) in marine sediments
235 (Johnson et al., 2014). In agreement with this, TOC radiocarbon ages in our core have
236 been previously found to be remarkably similar to co-located ages of the strictly
237 terrigenous higher plant leaf wax fraction (Ponton, 2012). This age similarity also
238 excludes interferences from within-river biological productivity (e.g., Eglinton and
239 Hamilton, 1967; Eglinton and Eglinton, 2008). To assess the variability of the terrigenous
240 carbon age exported by Godavari River based on this understanding we used high
241 resolution TOC radiocarbon measurements to calculate radiocarbon age offsets relative to
242 the atmosphere (Soulet et al., 2016; see Supplementary Materials). Over the Holocene,
243 these biospheric organic carbon radiocarbon age offsets in our core mirror the history of
244 erosion in the basin (Fig. 2).

245 As a first order observation, TOC ages (Fig. 2 and Supplementary Materials) are
246 significantly older (~200 to 2000 ¹⁴C years) than their depositional age in our Godavari
247 core. Before 5000 years ago the bulk organic carbon radiocarbon age offset were ~600
248 ¹⁴C years old on average. In contrast, the highly erosional regime under both climatic and
249 early human pressure in the late Holocene led to the export of significantly older carbon
250 from the terrestrial biosphere, i.e., ~1300 ¹⁴C on average. This increase in radiocarbon
251 age offset occurred largely during the two aridification steps identified by Ponton et al.
252 (2012): more abruptly between ~5000 and 4500 years ago and more gradually after
253 ~1700 years ago (Fig. 2).

254 In the absence of significant storage in alluvial sediments in the Godavari catchment,
255 several processes can explain the doubling in age offset over the Holocene: an overall
256 slowing of soil carbon turnover in the drying climate of central India, a decrease in TOC
257 contribution from contemporaneous vegetation relative to older (pre-aged) soil carbon
258 input and/or deeper exhumation of soils contributing increasingly older carbon. Given the
259 drastic changes in vegetation cover and increase in erosion in the Godavari basin, a
260 decrease in soil turnover is unlikely during the Holocene aridification process (Carvalhais
261 et al. 2014). In turn, the good agreement between the pollen and leaf-wax $\delta^{13}\text{C}$ records in
262 our core (Ponton et al., 2012; Zorzi et al., 2015) with independent monsoon
263 reconstructions suggests sustained delivery of recently fixed biospheric organic carbon to
264 the delta. Thus, the doubling in age offset over the Holocene is best explained by
265 increasing contributions from an older soil component, which could only come through
266 deeper erosion. Because the age of soil organic carbon in soil profiles increases with
267 depth (Trumbore, 2009), older mixtures imply a deeper soil erosion, whether uniform, or
268 through gullies, which are common especially on the Deccan Plateau (Kothyari, 1996).

269 6. The Monsoon Erosional Pump

270 Overall, these multiple lines of evidence indicate that soil erosion in the CMZ, as
271 integrated by the Godavari River, increased throughout the basin immediately as climate
272 began to dry at the end of mid Holocene, and was further enhanced by Deccan
273 agricultural activities in the late Holocene. The likely mechanism for this erosion
274 acceleration is the extreme seasonal distribution of the rainfall that characterizes the
275 monsoon (Wang and Ding, 2008), which promoted erosion on the more sparsely
276 vegetated landscapes (Molnar, 2001; DiBiase and Whipple, 2011; Plink-Bjorklund,
277 2015). Our findings thus point to a veritable “monsoon erosional pump” that accelerates
278 during minimum landcover conditions when the protective role of vegetation is reduced,
279 whether naturally or by humans. The volume of total eroded sediments since the mid
280 Holocene must have been considerable as the continental margin growth accelerated with
281 the shelf edge aggrading ~80 meters in the last ~2000 years alone (Forsberg et al., 2007).

282 This “landcover-mode” of the monsoon erosional pump must have been active before the
283 Holocene as well, affecting the transfer of terrigenous sediment, solutes and carbon from
284 land to the ocean. The beat of monsoon precipitation on orbital timescales is not well
285 constrained but considered to be modulated by at a combination of precession and
286 obliquity frequencies based on monsoon wind reconstructions (e.g., Clemens and Prell,
287 2003). Such complex variability did not inevitably follow the sea level cyclicity (e.g.,
288 Goodbred and Kuehl, 2000), which is usually assumed to control most of the sediment
289 transfer from land to the deep ocean (see Blum and Hattier-Womack, 2009 and references
290 therein for an analysis underlining the increased recognition for a climate role). Thus
291 untangling the effects of the monsoon is difficult especially during the Quaternary (e.g.,
292 Phillips et al., 2014), but may be easier to discern earlier when the sea level change
293 magnitude was reduced. Landcover effects are less likely to occur in the upper basins of
294 Himalayan monsoonal rivers where there are other sources of water such as snow or
295 glaciers and where elevation (i.e., temperature) and orographic precipitation promote
296 ecological stability (Galy et al., 2008a). The erosional pump in these high, steep regions
297 is still active due to monsoonal seasonality but in a “topographic-mode” dominated
298 primarily by landslides (Montgomery and Brandon, 2002; but see Olen et al., 2016 for an
299 alternative viewpoint). However, the landcover-mode for the erosional pump should still
300 be active in their lower basins where aridity controls vegetation type and cover (e.g.,
301 Galy et al. 2008b).

302 Recent coupled erosion-carbon cycling modeling suggests that long-term anthropogenic
303 acceleration of erosion has had a significant impact on the global carbon cycle by
304 intensifying the burial of terrigenous carbon (Wang et al., 2017). Prior to damming, the
305 monsoon domain supplied ~70% of the sediment load coming from large rivers (Syvitski
306 and Saito, 2007), although it only covers ~15% of the Earth’s surface (Hsu et al., 2011).
307 Therefore, we suspect that the cumulative effect of the monsoon erosional pump on the
308 carbon budget was substantial in augmenting the burial of terrigenous carbon during the

309 Holocene and needs to be estimated for inclusion in assessments of the net soil–
310 atmosphere carbon exchange.

311

312

313 Acknowledgments

314 We thank colleagues from the NGHP-01 expedition for intellectual interactions leading
315 to this work. We also thank Daniel Montlucon assistance in producing lab data. This
316 study was supported by grants from Woods Hole Oceanographic Institution, the National
317 Science Foundation (OCE-0841736 and OCE-0623766) and Swiss National Science
318 Foundation (“CAPS LOCK” 200021-140850 and “CAPS-LOCK2” 200021-163162).

319 References

- 320 Allchin, F. R.: Early cities and states beyond the Ganges Valley, in: *The Archaeology of*
321 *Early Historic South Asia: The Emergence of Cities and States*, Allchin, F. R. (Ed.),
322 Cambridge University Press, Cambridge, UK, 123-151, 1995.
- 323 Allen, C. D. and Breshears, D. D.: Drought-induced shift of a forest–woodland ecotone:
324 rapid landscape response to climate variation, *Proc. Natl. Acad. Sci. USA*, 95, 14839-
325 14842, 1998.
- 326 Asouti, E. and Fuller, D. Q.: *Trees and Woodlands in South India: Archaeological*
327 *Perspectives*, Left Coast Press, Walnut Creek, California, 2008.
- 328 Aufdenkampe, A. K., Mayorga, E., Raymond, P. A., Melack, J. M., Doney, S. C., and
329 Alin, S. R.: Riverine coupling of biogeochemical cycles between land, oceans, and
330 atmosphere, *Front. Ecol. Environ.*, 9, 53-60, doi:10.1890/100014, 2011.
- 331 Battin, T. J., Luysaert, S., Kaplan, L. A., Aufdenkampe, A. K., Richter, A. and Tranvik,
332 L. J.: The boundless carbon cycle, *Nat. Geosci.*, 2, 598-600, 2009.
- 333 Bayon, G., Dennielou, B., Etoubleau, J., Ponzevera, E., Toucanne, S. and Bermell, S.:
334 Intensifying weathering and land use in Iron Age Central Africa, *Science*, 335, 1219-
335 1222, 2012.
- 336 Bejugam, P., Nayak G. N.: Source and depositional processes of the surface sediments
337 and their implications on productivity in recent past off Mahanadi to Pennar River
338 mouths, western Bay of Bengal, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 483, 58–69,
339 2017.
- 340 Berkelhammer, M., Sinha, A., Stott, L., Cheng, H., Pausata, F. S. R., and Yoshimura, K.:
341 An abrupt shift in the Indian monsoon 4000 years ago, in: *Climates, Landscapes, and*
342 *Civilizations*, Geophysical Monograph, 198, Giosan, L., Fuller, D. Q., Nicoll, K., Flad,
343 R. K., and Clift, P. D. (Eds.), American Geophysical Union, Washington D.C., 75–87,
344 doi:10.1029/2012GM001207, 2012.
- 345 Bhattacharyya, T., Pal, D. K., Mandal, C., Chandran, P., Ray, S. K., Sarkar, D.,
346 Velmourougane, K., Srivastava, A., Sidhu, G. S., Singh, R. S., Sahoo, A. K., Dutta, D.,
347 Nair, K. M., Srivastava, R., Tiwary, P., Nagar, A. P., and Nimkhedkar, S. S.: Soils of
348 India: historical perspective, classification and recent advances, *Curr. Sci. India*, 104,
349 1308-1323, 2003.
- 350 Bikshamaiah, G. and Subramanian, V.: Sediment transport of the Godavari River basin
351 and its controlling factors, *J. Hydrol.*, 101, 275-290, 1988.

- 352 Blum, M. D. and Hattier-Womack, J.: Climate change, sea-level change, and fluvial
353 sediment supply to deepwater depositional systems, in: External Controls on Modern
354 Clastic Turbidite Systems, SEPM Special Publication, 92, Kneller, B., Martinsen, O. J.,
355 and McCaffrey, B. (Eds), Tulsa, Oklahoma, 15-39, 2009.
- 356 Bork, H.R. and Lang, A.: Quantification of past soil erosion and land use/land cover
357 changes in Germany, in: Long Term Hillslope and Fluvial System Modelling, Lang, A.,
358 Hennrich, K. P., and Dikau, R. (Eds.), Springer, Berlin, Germany, 231-239, 2003.
- 359 Brubaker, R.: Aspects of mortuary variability in the South Indian Iron Age, Bull. Deccan
360 College Res. Inst., 60, 253-302, 2000.
- 361 Carvalhais, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., Mu, M.,
362 Saatchi, S., Santoro, M., Thurner, M., Weber, U., Ahrens, B., Beer, C., Cescatti, A.,
363 Randerson, J. T., and Reichstein, M.: Global covariation of carbon turnover times with
364 climate in terrestrial ecosystems, *Nature*, 514, 213-217, 2014.
- 365 Chappell, A., Baldock, J. and Sanderman, J.: The global significance of omitting soil
366 erosion from soil organic carbon cycling schemes, *Nat. Clim. Change*, 6, 187-191,
367 2016.
- 368 Clemens, S. C. and Prell, W. L.: A 350,000 year summer-monsoon multi-proxy stack
369 from the Owen Ridge, Northern Arabian Sea, *Mar. Geol.*, 201, 35–51, 2003.
- 370 Collett, T., Riedel, M., Cochran, J., Boswell, R., Presley, J., Kumar, P., Sathe, A., Sethi,
371 A., Lall, M., and the NGHP Expedition Scientists: Indian National Gas Hydrate
372 Program Expedition 01 report, U.S. Geological Survey Scientific Investigations Report
373 2012–5054, 1442 pp., 2015.
- 374 Collins, D. B. G. and Bras, R. L.: Climatic control of sediment yield in dry lands
375 following climate and land cover change, *Water Resour. Res.*, 44, W10405,
376 doi:10.1029/2007WR006474, 2008.
- 377 Contreras-Rosales, L. A., Jennerjahn, T., Tharammal, T., Meyer, V., Lückge, A., Paul, A.
378 and Schefuß, E.: Evolution of the Indian Summer Monsoon and terrestrial vegetation in
379 the Bengal region during the past 18 ka, *Quat. Sci. Rev.*, 102, 133-148, 2014.
- 380 Cui, M., Wang, Z., Rao, N. K., Sangode, S. J., Saito, Y., Chen, T., Kulkarni, Y. R.,
381 Kumar, K. C. V., and Demudu, G.: A mid-to-late Holocene record of vegetation decline
382 and erosion triggered by monsoon weakening and human adaptations in the south-east
383 Indian Peninsula, *The Holocene*, doi.org/10.1177/0959683617715694, 2017.

- 384 DePaolo, D. J.: Neodymium Isotope Geochemistry, An Introduction, Springer-Verlag,
385 Berlin, 187 pp., 1988.
- 386 Dhavalikar, M. K.: Towards an ecological model for chalcolithic cultures of central and
387 western India, *J. Anthropol. Archaeol.*, 3, 133-158, 1984.
- 388 DiBiase, R. A. and Whipple, K. X.: The influence of erosion thresholds and runoff
389 variability on the relationships among topography, climate, and erosion rate, *J.*
390 *Geophys. Res.-Earth*, 116, F04036, doi:10.1029/2011JF002095, 2011.
- 391 Dixit, Y., Hodell, D. A., and Petrie, C. A.: Abrupt weakening of the summer monsoon in
392 northwest India similar to 4100 yr ago, *Geology*, 42, 339-342, 2014.
- 393 Dotterweich, M.: The history of human-induced soil erosion: Geomorphic legacies, early
394 descriptions and research, and the development of soil conservation—A global
395 synopsis, *Geomorphology*, 201, 1–34. doi:10.1016/j.geomorph.2013.07.021, 2013.
- 396 Dunne, T.: Sediment yield and land use in tropical catchments, *J. Hydrol.*, 42, 281–300,
397 doi:10.1016/0022-1694(79)90052-0, 1979.
- 398 Edwards, W. M. and Owens, L. B.: Large storm effects on total soil erosion, *J. Soil Water*
399 *Conserv.*, 46, 75–78, 1991.
- 400 Eglinton, G. and Hamilton, R. J.: Leaf epicuticular waxes, *Science*, 156, 1322-1335,
401 1967.
- 402 Eglinton, T. I. and Eglinton, G.: Molecular proxies for paleoclimatology, *Earth Planet.*
403 *Sci. Lett.*, 275, 1-16, 2008.
- 404 Feng, X. J., Vonk, J. E., van Dongen, B. E., Gustafsson, O., Semiletov, I. P., Dudarev, O.
405 V., Wang, Z. H., Montlucon, D. B., Wacker, L., and Eglinton, T. I.: Differential
406 mobilization of terrestrial carbon pools in Eurasian Arctic river basins, *Proc. Natl.*
407 *Acad. Sci. USA*, 110, 14168-14173, 2013.
- 408 Fleitmann, D., Burns, S. J., Mudelsee, M., Neff, U., Kramers, J., Mangini, A., and Matter,
409 A.: Holocene forcing of the Indian monsoon recorded in a stalagmite from Southern
410 Oman, *Science*, 300, 1737-1739, 2003.
- 411 Forsberg, C. F., Solheim, A., Kvalstad, T. J., Vaidya, R., and Mohanty, S.: Slope
412 instability and mass transport deposits on the Godavari river delta, east Indian margin
413 from a regional geological perspective, *Submarine Mass Movements and Their*
414 *Consequences* 27, 19-27, 2007.

- 415 Fuller, D. Q.: Asia, South: Neolithic cultures, in: *Encyclopedia of Archaeology*, Pearsall,
416 D. (Ed.), Elsevier Academic Press, San Diego, California, 756-768, 2008.
- 417 Fuller, D. Q.: Finding plant domestication in the Indian subcontinent. *Current*
418 *Anthropology*, 52, Suppl. 4, 2011.
- 419 Fuller, D. Q. and Madella, M.: Issues in Harappan Archaeobotany: Retrospect and
420 Prospect., in: *Indian Archaeology in Retrospect, Volume II. Protohistory*, Settar, S., and
421 Korisettar, R. (Eds.), Indian Council for Historical Research, New Delhi, 47-80, 2001.
- 422 Galy, V. and Eglinton, T. I.: Protracted storage of biospheric carbon in the Ganges-
423 Brahmaputra basin, *Nat. Geosci.*, 4, 843-847, 2011.
- 424 Galy, V., François, L., France-Lanord, C., Faure, P., Kudrass, H., Palhol, F. and Singh,
425 S.K.: C4 plants decline in the Himalayan basin since the Last Glacial Maximum, *Quat.*
426 *Sci. Rev.*, 27, 1396-1409, 2008a.
- 427 Galy, V., France-Lanord, C., and Lartiges, B.: Loading and fate of particulate organic
428 carbon from the Himalaya to the Ganga–Brahmaputra delta, *Geochim. Cosmochim.*
429 *Acta*, 72, 1767-1787, 2008b.
- 430 Galy, V., Peucker-Ehrenbrink, B. and Eglinton, T.: Global carbon export from the
431 terrestrial biosphere controlled by erosion. *Nature*, 521, 204-207, 2015.
- 432 Goodbred, S. L. and Kuehl, S. A.: Enormous Ganges-Brahmaputra sediment discharge
433 during strengthened early Holocene monsoon, *Geology*, 28, 1083-1086, 2000.
- 434 Gunnell, Y., Anupama, K., and Sultan, B.: Response of the South Indian runoff
435 harvesting civilization to northeast monsoon rainfall variability during the last 2000
436 years: instrumental records and indirect evidence, *Holocene*, 17, 207-215, 2007.
- 437 Gupta, A. K., Anderson, D. M., and Overpeck, J. T.: Abrupt changes in the Asian
438 southwest monsoon during the Holocene and their links to the North Atlantic Ocean,
439 *Nature*, 421, 354-357, 2003.
- 440 Hilton, R. G., Galy, A., Hovius, N., Chen, M. C., Horng, M. J., and Chen, H.: Tropical-
441 cyclone-driven erosion of the terrestrial biosphere from mountains, *Nat. Geosci.*, 1,
442 759-762, 2008.
- 443 Hoffmann, T., Mudd, S. M., van Oost, K., Verstraeten, G., Erkens, G., Lang, A.,
444 Middelkoop, H., Boyle, J., Kaplan, J. O., Willenbring, J., and Aalto, R.: Humans and
445 the missing C-sink: Erosion and burial of soil carbon through time, *Earth Surf. Dynam.*,
446 1, 45-52, 2013.

- 447 Hsu, P. C., Li, T., and Wang, B.: Trends in global monsoon area and precipitation over
448 the past 30 years. *Geophys. Res. Lett.*, 38, L08701, doi: 10.1029/2011GL046893, 2011.
- 449 Istanbuluoglu, E. and Bras, R. L., Vegetation-modulated landscape evolution: Effects of
450 vegetation on landscape processes, drainage density, and topography: *J. Geophys. Res.-*
451 *Earth*, 110, F02012, doi:10.1029/2004jf000249, 2005.
- 452 Johnson, J. E., Phillips, S. C., Torres, M. E., Piñero, E., Rose, K. K., and Giosan, L.:
453 Influence of total organic carbon deposition on the inventory of gas hydrate in the
454 Indian continental margins, *Mar. Pet. Geol.*, 58, 06-424, 2014.
- 455 Joshi, R. K., Mazumdar, A., Peketi, A., Ramamurty, P. B., Naik, B. G., Kocherla, M.,
456 Carvalho, M. A., Mahalakshmi, P., Dewangan, P., and Ramana, M. V.: Gas hydrate
457 destabilization and methane release events in the Krishna–Godavari Basin, Bay of
458 Bengal. *Mar. Pet. Geol.*, 58, 476-489, 2014.
- 459 Kajale, M. D.: Plant Economy, in: Excavations at Inamgaon, Vol. 1, Part 2, Dhavalikar,
460 M. K., Sankalia, H. D., and Ansari, Z. D. (Eds.), Deccan College Postgraduate and
461 Research Institute, Pune, 727–821, 1988.
- 462 Kale, V. S.: Fluvial geomorphology of Indian rivers: an overview, *Prog. Phys. Geog.*, 26,
463 400–433, 2002.
- 464 Kale, V. S.: *Geomorphic Effects of Monsoon Floods on Indian Rivers, Natural Hazards,*
465 28, Springer, New York, 2003.
- 466 Kaplan, J. O., Krumhardt, K. M., Ellis, E. C., Ruddiman, W. F., Lemmen, C., and Klein
467 Goldewijk, K.: Holocene carbon emissions as a result of anthropogenic land cover
468 change, *The Holocene*, 21, 775 -791, doi:10.1177/0959683610386983, 2010.
- 469 Kothyari, U. C.: Erosion and Sedimentation Problems in India, in: *Erosion and Sediment*
470 *Yield: Global and Regional Perspectives (Proceedings of the Exeter Symposium, July*
471 *1996)*, IAHS Publ. 236, Walling, D. E. and Webb, B. W. (Eds.), IAHS Press,
472 Wallingford, Oxfordshire, UK, 531-540, 1996.
- 473 Kulkarni, Y. R., Sangode, S. J., Meshram, D. C., Patil, S. K., and Dutt, Y.: Mineral
474 magnetic characterization of the Godavari River sediments; *J. Geol. Soc. India*, 81,
475 376-384, 2014.
- 476 Kulkarni, Y. R., Sangode, S. J., Bloemandal, J., Meshram, D. C., and Suresh, N.: Mineral
477 magnetic characterization of the Godavari River and western Bay of Bengal sediments:
478 Implications to source to sink relations; *J. of Geol. Soc. India*, 85, 71-78, 2015.

- 479 Lal, R.: Soil carbon sequestration impacts on global climate change and food security,
480 Science, 304, 1623-1627, 2004.
- 481 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambradge, M.: Sea level and global
482 ice volumes from the Last Glacial Maximum to the Holocene, Proc. Natl. Acad. Sci.
483 USA, 111, 15296-15303, 2014.
- 484 Langbein, W. B. and Schumm, S. A.: Yield of sediment in relation to mean annual
485 precipitation, EOS Trans. Amer. Geophys. Union, 39, 1076-1084, 1958.
- 486 Ludwig, W., Probst, J. L., and Kempe, S.: Predicting the oceanic input of organic carbon
487 by continental erosion, Global Biogeochem. Cycles, 10, 23-41, 1996.
- 488 Mazumdar, A., Dewangan, P., Joao, H. M., Peketi, A., Khosla, V. R., Kocherla, M.,
489 Badesab, F. K., Joshi, R. K., Roxanne, P., Ramamurty, P. B., Karisiddaiah, S. M., Patil,
490 D. J., Dayal, A. M., Ramprasad, T., Hawkesworth, C. J., and Avanzinelli, R.: Evidence
491 of paleo-cold seep activity from the Bay of Bengal, offshore India, Geochem. Geophys.
492 Geosys., 10, Q06005, doi:10.1029/2008GC002337, 2009.
- 493 McLennan, S.M. and Hemming, S.: Samarium/neodymium elemental and isotopic
494 systematics in sedimentary rocks, Geochim. Cosmochim. Acta 56, 887–898,1992.
- 495 Mohanty, R. K. and Selvakumar, V.: The archaeology of the Megaliths in India: 1947-
496 1997, in: Indian Archaeology in Retrospect, 1. Prehistory, Settar, S., and Korisetar, R.
497 (Eds.), Manohar, New Delhi, 313-352, 2001.
- 498 Molnar, P.: Climate change, flooding in arid environments, and erosion rates, Geology,
499 29, 1071-1074, 2001.
- 500 Montgomery, D. R.: Soil erosion and agricultural sustainability, Proc. Natl. Acad. Sci.
501 USA, 104, 13268-13272, 2007.
- 502 Montgomery, D. R. and Brandon, M. T.: Topographic controls on erosion rates in
503 tectonically active mountain ranges, Earth Planet. Sci. Lett. 201, 481-489, 2002.
- 504 Mujumdar, G. G., Rajaguru, S. N., and Papu, R. S.: The recent Godavari Flood
505 (September 1969) and its relevance to prehistoric archeology, Bull. Deccan College
506 Res. Inst. 24, 1–17, 1970.
- 507 NBSS & LUP (National Bureau of Soil Survey & Land Use Planning of India): Soils of
508 India, (Suborder Associations), Map scale 1: 6,300,000, NBSS & LUP, Nagpur, 1983.

- 509 Olen, S. M., Bookhagen, B., and Strecker, M. R.: Role of climate and vegetation density
510 in modulating denudation rates in the Himalaya, *Earth Planet. Sci. Lett.*, 445, 57-67,
511 2016.
- 512 Parabrahma Sastry, P. V.: The Early Historic Transition, in: Pre- and Protohistoric
513 Andhra Pradesh up to 500 BC, Murty, M. L. K. (Ed.), Orient Longman, New Delhi,
514 139-147, 2003.
- 515 Phillips, S. C., Johnson, J. E., Giosan, L., and Rose, K.: Monsoon-influenced variation in
516 productivity and lithogenic sediment flux since 110 ka in the offshore Mahanadi Basin,
517 northern Bay of Bengal, *Mar. Pet. Geol.*, 58, 502-525, 2014.
- 518 Plink-Björklund, P.: Morphodynamics of rivers strongly affected by monsoon
519 precipitation: Review of depositional style and forcing factors. *Sed. Geol.*, 323, 110-
520 147, 2015.
- 521 Ponton, C.: Aridification of the Indian subcontinent during the Holocene: implications for
522 landscape evolution, sedimentation, carbon cycle, and human civilizations, Doctoral
523 dissertation, Massachusetts Institute of Technology, Cambridge, MA, USA,
524 doi:10.1575/1912/5291, 2012.
- 525 Ponton, C., Giosan, L., Eglinton, T.E., Fuller D. Q., Johnson, J. E., Kumar, P., and
526 Collett, T. S.: Holocene Aridification of India, *Geophys. Res. Lett.*, 39, L03704,
527 doi:10.1029/2011GL050722, 2012.
- 528 Prasad, S., Anoop, A., Riedel, N., Sarkar, S., Menzel, P., Basavaiah, N., Krishnan, R.,
529 Fuller, D., Plessen, B., Gaye, B., and Röhl, U.: Prolonged monsoon droughts and links
530 to Indo-Pacific warm pool: A Holocene record from Lonar Lake, central India. *Earth
531 Planet. Sci. Lett.*, 391, 171-182, 2014.
- 532 Prasad, S. and Enzel, Y.: Holocene paleoclimates of India, *Quat. Res.*, 66, 442-453, 2006.
- 533 Ramprasad, T., Dewangan, P., Ramana, M. V., Mazumdar, A., Karisiddaiah, S. M.,
534 Ramya, E. R., and Sriram, G.: Evidence of slumping/sliding in Krishna–Godavari
535 offshore basin due to gas/fluid movements, *Mar. Pet. Geol.*, 28, 1806-1816, 2011.
- 536 Rao, K. N., Sadakata, N., Malini, B. H., and Takayasu, K.: Sedimentation processes and
537 asymmetric development of the Godavari delta, India, in: *River Deltas, Concepts,
538 Models and Examples*, SEPM Special Publication, 83, Giosan, L. and Bhattacharya, J.
539 P. (Eds.), Tulsa, Oklahoma, 433-449, 2005.
- 540 Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A.,
541 Laruelle, G. G., Lauerwald, R., Luysaert, S., Andersson, A. J., Arndt, S., Arnosti, C.,

- 542 Borges, A. V., Dale, A. W., Gallego-Sala, A., Godd ris, Y., Goossens, N., Hartmann,
543 J., Heinze, C., Ilyina, T., Joos, F., LaRowe, D. E., Leifeld, J., Meysman, F. J. R.,
544 Munhoven, G., Raymond, P. A., Spahni, R., Suntharalingam, P., and Thullner, M.:
545 Anthropogenic perturbation of the carbon fluxes from land to ocean, *Nat. Geosci.*, 6,
546 597-607, 2013.
- 547 Reichstein M., Bahn, M., Ciais, P., Frank, D., Mahecha, M. D., Seneviratne, S. I.,
548 Zscheischler, J., Beer, C., Buchmann, N., Frank, D. C., Papale, D., Rammig, A., Smith,
549 P., Thonicke, K., van der Velde, M., Vicca, S., Walz, A., and Wattenbach, M.: Climate
550 extremes and the carbon cycle, *Nature*, 500, 287–295, 2013.
- 551 Roberts, P., Boivin, N., Petraglia, M., Masser, P., Meece, S., Weisskopf, A., and Fuller,
552 D. Q.: Local diversity in settlement, demography and subsistence across the southern
553 Indian Neolithic-Iron Age transition: site growth and abandonment at Sanganakallu-
554 Kupgal, *Archaeol. Anthropol. Sci.*, 8, 575-599, 2016.
- 555 Sangode, S. J., Suresh, N., and Bagati, T. N.: Godavari source in the Bengal fan
556 sediments: results from magnetic susceptibility dispersal pattern, *Curr. Sci, India*, 660-
557 664, 2001.
- 558 Sarkar, S., Prasad, S., Wilkes, H., Riedel, N., Stebich, M., Basavaiah, N., and Sachse, D.:
559 Monsoon source shifts during the drying mid-Holocene: Biomarker isotope based
560 evidence from the core ‘monsoon zone’(CMZ) of India, *Quat. Sci. Rev.*, 123,144-157,
561 2015.
- 562 Singh, G., Babu, R., Narain, P., Bhushan, L. S., and Abrol, A. P.: Soil erosion rates in
563 India. *J. Soil Water Cons.*, 47, 97-99, 1992.
- 564 Sinha, A., Stott, L., Berkelhammer, M., Cheng, H., Edwards, R. L., Buckley, B.,
565 Aldenderfer, M., and Mudelsee, M.: A global context for megadroughts in monsoon
566 Asia during the past millennium, *Quat. Sci. Rev.*, 30, 47-62, 2011.
- 567 Soulet, G., Skinner, L.C., Beaupr , S.R., and Galy, V.: A note on reporting of reservoir
568 ¹⁴C disequilibria and age offsets, *Radiocarbon*, 58, 205-211, 2016.
- 569 Smittenberg, R. H., Eglinton, T. I., Schouten, and Damste, J. S. S.: Ongoing buildup of
570 refractory organic carbon in boreal soils during the Holocene, *Science*, 314, 1283-1286,
571 2006.
- 572 Sridhar, P. N., Ali, M. M., Vethamony, P., Babu, M. T., Ramana, I. V., and Jayakumar,
573 S.: Seasonal Occurrence of Unique Sediment Plume in the Bay of Bengal, *EOS Trans.*
574 *Amer. Geophys. Union*, 89, 22-23, 2008

- 575 Stallard, R. F.: Terrestrial sedimentation and the carbon cycle: Coupling weathering and
576 erosion to carbon burial. *Global Biogeochem. Cycles*, 12, 231-257, 1998.
- 577 Summerfield, M. A. and Hulton, N. J.: Natural controls of fluvial denudation rates in
578 major world drainage basins, *J. Geophys. Res.*, 99, 13871–13883,
579 doi:10.1029/94JB00715, 1994.
- 580 Syvitski, J. P. M. and Saito, Y.: Morphodynamics of deltas under the influence of
581 humans, *Glob. Planet. Chang.*, 57, 261–282, 2007.
- 582 Trumbore, S.: Radiocarbon and Soil Carbon Dynamics, *Ann. Rev. Earth Planet. Sci.*, 37,
583 47-66, 2009.
- 584 Vanacker, V., von Blanckenburg, F., Govers, G., Molina, A., Poesen, J., Deckers, J., and
585 Kubik, P.: Restoring dense vegetation can slow mountain erosion to near natural
586 benchmark levels, *Geology*, 35, 303–306, doi:10.1130/G23109A.1, 2007.
- 587 van Andel, T. H., Zangger, E., and Demitrack, A.: Land use and soil erosion in
588 prehistoric and historical Greece, *J. Field Archaeol.*, 17, 379-396, doi: 10.2307/530002,
589 1990.
- 590 van Oost, K., Verstraeten, G., Doetterl, S., Notebaert, B., Wiaux, F., Broothaerts, N., and
591 Six, J.: Legacy of human-induced C erosion and burial on soil-atmosphere C exchange,
592 *Proc. Natl. Acad. Sci. USA*, 109, 19492-19497, 2012.
- 593 Vanwalleghem, T., Gómez, J. A., Infante Amate, J., González de Molina, M.,
594 Vanderlinden, K., Guzmán, G., Laguna, A., and Giráldez, J. V.: Impact of historical
595 land use and soil management change on soil erosion and agricultural sustainability
596 during the Anthropocene, *Anthropocene* 17, 13–29, 2017.
- 597 Walling D. E. and Webb, B.W.: Patterns of sediment yields, in: *Background to*
598 *Paleohydrology*, Gregory, K. J. (Ed.), John Wiley & Sons, London, 69-100, 1983.
- 599 Wang, B. and Ding, Q.: Global monsoon: Dominant mode of annual variation in the
600 tropics, *Dyn. Atmos. Oceans*, 44, 165-183, 2008.
- 601 Wang, Z., Hoffmann, T., Six, J., Kaplan, J. O., Govers, G., Doetterl, S., and van Oost, K.:
602 Human-induced erosion has offset one-third of carbon emissions from land cover
603 change, *Nat. Clim. Change*, 7, 345–349, 2017.
- 604 Wilkinson, B. H. and McElroy, B. J.: The impacts of humans on continental erosion and
605 sedimentation, *Geol. Soc. Am. Bull.*, 119, 140–156, 2007.

606 Zorzi, C., Goñi, M. F. S., Anupama, K., Prasad, S., Hanquiez, V., Johnson, J., and
607 Giosan, L.: Indian monsoon variations during three contrasting climatic periods: The
608 Holocene, Heinrich Stadial 2 and the last interglacial–glacial transition, *Quat. Sci. Rev.*,
609 125, 50-60, 2015.




Figure 1. Godavari River drainage basin in its (a) physiographical, (b) hydroclimatic (Asouti and Fuller, 2008), (c) ecological (Asouti and Fuller, 2008), (d) geological (Bikshamaiah and Subramanian, 1980) and (e) soil cover context (NBSS&LUP, 1983). Core NGHP-01-16A location is indicated in (a) by the red dot. Average bedrock ϵ_{Nd} values are shown in (d).

610


611

612

613

614

615 Figure 2. Paleoenvironmental reconstructions from core NGHP-01-16A for the CMZ as
 616 integrated by the Godavari River: (a) Sediment fluxes as mass accumulation rates and
 617 sediment sources from Nd isotope fingerprinting (Deccan Trap sediment contribution is
 618 estimated from a two-end member model; see text and Supplementary Materials); (b)
 619 TOC radiocarbon age offset relative to depositional age; (c) Hydroclimate and ecology
 620 from pollen (Zorzi et al., 2015) and leaf wax carbon isotopes (Ponton et al., 2012).
 621



622
 623
 624
 625
 626