

1 Response to RC1

2 (responses in blue)

3
4 Mutz et al. look to use paleoclimate GCMs to identify drivers of past geomorphic change. This is a topic
5 for which I hold great interest, and I feel that the authors have crafted a very useful set of model results
6 that they leave underutilized. As such, I feel inclined to accept the paper on the basis of the useful results,
7 but to request major revisions such that they do their own work justice.

8
9 My major concerns, which will become clear in the line-by-line comments (please feel free to respond to
10 similar comments en masse) are as follows: 1. The paper is motivated by denudation rates and landscape
11 evolution, but really includes this as a speculative wrapper that is not substantiated. I suggest that instead
12 you propose testable hypotheses surrounding your findings.

13 2. Related to #1, much of the text is a litany of "temperature was X here ...". I find such statements of
14 results useful only insofar as they expand upon a figure (associated with a supplementary data set) that
15 presents the results. While these sections are written clearly, I would suggest that the authors focus on a set
16 of geomorphic questions (if this be their motivation) and how the model-data set informs those questions.

17 3. Many of the discussions of model results are of ice-covered regions, yet no consideration of direct
18 glacial erosion is given. Furthermore, no reference to the changes in the statistics of discharge or
19 catchment area in ice-covered regions is given. This seems a disservice to this relatively high-resolution
20 paleoclimate AGCM: the geologic setting **must** be considered, otherwise it seems that the authors'
21 pushing on the modeling end has not been matched by a simple geological history sanity check. I would
22 suggest that either significantly glaciated regions and the catchments that they feed be masked out, or that
23 glacial erosion and its associated processes be included in the discussion.

24 4. (Discussed only here): You have not compared your models against any data. I understand that this may
25 be simply a modeling exercise that you do compare to other models. However, I think that such a
26 comparison could assuage skepticism about your results and lend support to your case, especially if you
27 include it as part of a local case study (see the third point below). The core of these three points is that,
28 with a bit more care, I think your results could say something really useful to the geomorphic community.
29 Currently, the paper seems to be more a statement of, "this is important to geomorphology", followed by a
30 long list of the model results. I challenge the authors to demonstrate (rather than simply stating) the
31 importance of their work to geomorphology in a way that includes how it may impact the way scientists
32 view Quaternary landscape evolution. Ideas include:

- 33 • Changes in means (done)
34 • Changes in statistical distributions of temperature and precipitation – think extreme events, frost-cracking
35 window, etc.
36 • A focus on a few iconic regions while **explicitly** ignoring significantly ice-covered domains (I think
37 this would be easiest, though obviously would be thrilled if you decided to tackle glacial processes)
38 • Using this focus to build a template for how to use paleoclimate GCM outputs to advance the field of
39 geomorphology. Currently, I think that the work is acceptable following changes for internal consistency
40 and geological accuracy (see #1 and #3), but I think that you could be selling yourselves short if you don't
41 dig just a tiny bit deeper to investigate your forcings and their impact on geomorphology. I hope that you
42 find these comments helpful in continuing to craft an insightful piece of work out of what seems to be a
43 strong modeling approach.

44
45 We thank Prof. Andrew Wickert for his highly valuable review of our manuscript. Many important points
46 were raised in the review and we hope that our appreciation for the input is sufficiently reflected in the
47 revisions we made in response to it. We also encourage him to see our response to the second reviewer,
48 where we provide additional geologic relevance of this study by now including a comparison of available
49 terrestrial proxy data to our model results. We also explain throughout our response to the 2nd reviewer
50 why an application of the predicted climate change to predict denudation rate changes is a large
51 undertaking that can not be meaningfully conducted in this paper, but warrants more detailed applications
52 of the models to individual areas (a topic of ongoing work/application for us). We refrain from using the
53 model predicted runoff in the global GCM (even though it's conducted at relatively high
54 resolution compared to previous work) to calculate changes in fluvial incision. This would be
55 better done by mapping the predicted precipitation changes onto higher resolution (<90 m)
56 DEMs and solving the kinematic wave equation for each fluvial erosion in each catchment, for

57 the changes in precipitation. However, as we repeatedly mention above, this is not possible to
58 include in this manuscript without first characterising how the precipitation has changed in each
59 region (the current manuscript goals). Work in progress we are conducting is trying to apply the
60 kinematic wave equation and palaeoprecipitation to selected areas, but it's proving difficult to
61 implement meaningfully without temporally continuous (e.g. LGM to present) simulations of
62 precipitation change. We hope this brings to the reviewers attention the complications associated
63 with doing full erosion history calculations based on these results. We have expanded the last
64 paragraph in the instruction to convey the above perspective better, and more clearly articulate
65 (and justify) the scope and limitations of the manuscript.

66
67 We appreciate the importance of addressing specific sets of geomorphic questions and hypotheses (1 & 2)
68 and we are currently taking an in-depth look at quantifying the potential for erosion by a variety of
69 processes. These include different methods of quantifying frost cracking intensity and extreme
70 precipitation events and how these changed over time. However, in order to include those in this
71 manuscript, we fear that would have to seriously compromise the thoroughness with which we investigate
72 these questions at the moment. Instead, we hope that we can convey the usefulness of our consistently set
73 up palaeoclimate simulations as a framework for addressing any of these particular questions in detail, and
74 have modified sections of this manuscript accordingly. This includes, but is not restricted to, extensive
75 compilations of proxy-based precipitation reconstructions for our two larger study sites (South Asia and
76 western South America) and comparison of this data to our model output. With this, we hope that we were
77 also able to address the concerns raised in point 4. In order to address the important point raised about
78 glaciated areas (3), we added an ice cover layer on all of our difference plots, included global maps of ice
79 extent (as used for our simulations) in the supplementary material, and discussed where the large
80 differences in temperature and precipitation we highlight in the manuscript are accompanied by changes in
81 ice cover. Thus we hope to prevent that interpretations of the implications of our results are made without
82 consideration of changes in ice cover (and consequently shifts in the process domain).

83
84 Line-by-line:

85 23. US Pacific Northwest Pacific → drop second "Pacific"

86 This has been corrected. Thank you for catching that.

87

88 29. future observational studies interested in quantifying → future observational studies that quantify
89 (studies can't be interested in things, strictly speaking)

90 That is right of course. It has been corrected as suggested.

91

92 53. orogen scale → orogen-scale

93 It has been corrected as suggested.

94

95 ~57. A couple of recent studies from the climate science community shed light on the impacts of the
96 Andes (first ref below) and continents in general (second ref below). In case these are interesting to you,
97 I'm pasting the bibliographic information here:

98 Maroon, E. A., D. M. W. Frierson, and D. S. Battisti (2015), The tropical precipitation response to Andes
99 topography and ocean heat fluxes in an aquaplanet model, *J. Clim.*, 28(1), 381–398, doi:10.1175/JCLI-D-
100 14-00188.1.

101 Maroon, E. A., D. M. W. Frierson, S. M. Kang, and J. Scheff (2016), The precip itation response to an
102 idealized subtropical continent, *J. Clim.*, 29(12), 4543–4564, doi:10.1175/JCLI-D-15-0616.1.

103 Many thanks for the references. These are indeed of much interest to us and we included them in the
104 revised manuscript.

105

106 73-75. "Furthermore, recent controversy exists concerning the spatial and temporal scales over which
107 geologic and geochemical observations can record climate-driven changes in weathering and erosion [e.g.
108 Whipple, 2009; von Blanckenburg et al., 2015; Braun, 2016].": I see that you do not return to this point
109 later, so could you describe the controversy for those who are not familiar with it?

110 Thank you for pointing this out. We described the controversy briefly for those unfamiliar with it after the
111 sentence quoted above.

112

113 81. I see that later you discuss a little about what an AOGCM may do, but I will be looking for justification

114 about how an AGCM may suffice. Is this in part because you prescribe the b.c.'s and you are running it for
115 17 years only? If so, could you discuss potential systematic variations between this and an AOGCM?
116 We prescribe sea surface temperature reconstructions (SSTs) boundary conditions, which allows us to
117 bypass the computationally expensive coupled simulations. Because these are fixed climatologies (though
118 with seasonality preserved), the simulation of fewer years suffices. As a consequence, however, we do not
119 expect to see decadal scale variability as we would in case of coupled models or prescribed SSTs that vary
120 from year to year (such as present day simulations using AMIP SST's). We discuss this in the revised
121 manuscript.

122
123 89. "PLIO to the Last Glacial Maximum": as you include no time-slices between these, I suggest making
124 these part of the list and dropping the "to the".
125 This has been corrected as suggested.

126
127 147. "This section describes the clustering method used in this study." You could drop this sentence – the
128 section title should be enough for even an inattentive reader!
129 We followed this suggestion and dropped that sentence.

130
131 176-178. I was wondering how you picked the number of clusters: I am glad to see that you performed a
132 thorough search.

133 Thank you. We added some text explaining this. We systematically increased the number of clusters from
134 3 to 10 and assessed the distinctiveness or similarities of resulting climate clusters. Once the increase in the
135 number of cluster no longer resulted in the addition of another cluster that was distinctly different from the
136 others, we used this as a cut off point and used the cluster number of the previous iteration as the optimal
137 cluster number.

138
139 Section 3: Much of this is information that I find better communicated through figures than with text. It is
140 clearly written, however, and I am reluctant to suggest a rewrite for brevity in a length-unconstrained
141 journal so long as the text can be co-located with the figure.

142 190-192. I see you have another "This section describes..." sentence. If this is your preferred way to write,
143 you may keep it; here, the second sentence is not such a good topic-sentence replacement.
144 Thank you. We kept the sentence in this instance as it also immediately draws attention to the relevant
145 figures, which may also serve in addressing the previous point you raised.

146
147 197-198. i.e. over the ice sheets. (This applies to other regions as well, and should be important to point
148 out if you are going to then discuss fluvial processes in orogens)
149 Many thanks for pointing this out. We are more mindful of this in the revised discussion.

150
151 203-paragraph: Also because of local ice loss, presumably. So I think that the two prior paragraphs could
152 have a new summary that "The greatest changes in temperature is observed where the greatest change in
153 local ice extent occurs."
154 Thank you. We followed your suggestion.

155
156 214-215. Have you considered discussions of the African Humid Period?
157 We had not considered discussion of precipitation changes in North Africa, since it lies outside the regions
158 we focus on. However, we appreciate that Holocene precipitation changes in the region are important and
159 may be of interest to many readers. We therefore included a short discussion of Holocene precipitation
160 changes in Northern Africa in the revised manuscript.

161
162 373-374. If you are looking at the influence of temperature and precipitation on erosion, and you are not
163 including subglacial erosion, then your preceding text must indicate where your changes really are
164 indicative of ice extent – both as a separate process domain and as a driver of fluvial processes and
165 potential changes in the statistics of river discharge.
166 Thank you. We are more mindful of this in our discussion.

167
168 Section 4.1. Your first paragraph (weathering) differs from the content (comparing your model results with
169 those published). These should be in different subsections, and the weathering paragraph may need to be
170 expanded. Your "weathering and erosion" paragraph also neglects direct effects of glaciers, ice caps, and
171 ice sheets, which were globally significant.

172 Thank you for this suggestion. We re-structured this section as suggested and took ice extent into
173 consideration in the erosion section.

174

175 Section 4.2. Once again, your discussion is often of formerly (or currently) ice-covered regions without
176 explicitly acknowledging that this is a different process domain. In addition, as with the previous section,
177 the body paragraphs are mostly about model comparison and regional changes with sparse link to the
178 landscape-evolution factors indicated in the topic paragraph.

179 As above, we restructured this section, took ice extent into account and chose a more fitting section title.

180

181 416-423. Please discuss the direct influence of glaciers on the erosion orogens in the context of changing
182 precipitation (and therefore mass balance). Is it significant or not?

183 Although it is challenging to sufficiently quantify changes in glacier-related erosion due to differences in
184 precipitation, we now include this point in our discussion.

185

186 433-434. "Coastal North America"? Doesn't look like it: seems to be most of NA south of the ice margin.

187 That is correct. We revised our descriptions accordingly.

188

189 Section 4.3. The authors describe the results here, but I find the connection to erosion rates to be
190 insufficiently described compared to how they are highlighted in the topic sentence, as well as in the
191 abstract. I would like you to go one step beyond "ought to be considered" and actually posit how you
192 expect the erosion rates – and therefore, the balance between erosion and exhumation and perhaps the
193 equilibrium shapes of the mountains and their rivers – to vary. Otherwise, you are suggesting future work
194 rather than actually describing the possible geomorphic significance – and I think underutilizing your
195 results in a paper that is clearly targeted towards geomorphologists.

196 As described in our response above, we believe that trying to address specific problems such as these or
197 quantifying how differences would be expressed as erosion rates would be beyond the scope of this
198 manuscript and come at the cost of not being able to address these as thoroughly as we are currently
199 attempting in other ongoing work.

200

201 498. "which may favour frost driven weathering during glacial climate states" – the St. Elias range was
202 covered by glaciers! Yes, there can be some frost-cracking around the ice, but don't you think this is
203 important too? http://instaar.colorado.edu/groups/QGISL/ak_paleoglacier_atlas/gallery/index.html

204 Thank you for drawing our attention to this. We consider this in our discussion now and revised the
205 manuscript accordingly.

206

207 508. "enhanced sediment production driven by frost processes" – same as above. Glaciers were there.
208 Consider them.

209 As for the comment above, we also considered glaciers here in the revised manuscript.

210

211 Conclusions: Comparison to other models: is this match surprising or no? Did you (mostly) use the same
212 inputs and simply increase the grid resolution? If so, could you comment on how the improved grid and
213 possible variations in inputs and use of them ocean as a boundary condition may have affected (or not)
214 your results as compared to those of earlier studies? This would be more useful to include in the discussion
215 than a simple list of "Our temperature in place Y was T_0 , and X et al. wrote that they found it was T_1 ,
216 which is close to T_0 . Think big-picture, in both process and numerics!

217 Due to model-specific parameterisation, deviation is possible. In the revised manuscript, we comment on
218 this as well as on the model resolution and implications of using ocean as boundary conditions instead of
219 an ocean model.

220

221 533. Did your 8-10 degC changes occur significantly over areas that would be affected by hillslope or
222 fluvial processes? (i.e. unglaciated areas?)

223 Some unglaciated areas experience large differences in temperature, but the maxima of 8-10 degC
224 geographically coincide with ice cover changes. We acknowledge and discuss this in the revised
225 manuscript.

226

Response to RC2

(responses in blue)

The authors simulate the pre-industrial, mid-Holocene, LGM, and Pliocene climates using the ECHAM5 general circulation model. The motivation is to understand how past climatic states may change regional climatologies, particularly over mountainous areas that have been the focus of much erosion and geomorphic work. The authors find that past climatic states (particularly the LGM and Pliocene) do produce changes in absolute temperature and precipitation and in the annual ranges of these two climatic variables.

Overall, I did not find the manuscript to be a particularly useful addition to the literature. Though the motivation is potentially novel, the analysis is not complete and needs substantially more work. Most of the manuscript focuses on simply describing climatic changes, while neglecting novel analyses. Consequently, at this point, I recommend reject, though perhaps with substantial work (including more modeling, comparison with existing models, and/or comparison with data), it may become publishable in the future in *ESurf*.

We thank the reviewer for their constructive feedback on our manuscript. We found the reviewer's comment very useful in the improvement of this manuscript and hope that the modifications we made do the reviewer's thorough review justice. Most importantly, we followed the reviewer's suggestion to deepen the comparison of our model simulation results with data, specifically proxy-based reconstructions, as we believe this suggestion is of most relevance to the earth surface science community targeted by *ESurf*. While it was not possible to create a complete overview of all proxy-based studies everywhere and for all time periods, we included an extensive compilation of site specific reconstructions of precipitation for our largest areas (South America and Tibet). In prioritizing the compilation of precipitation reconstructions, we also hope to address the reviewer's concerns about the GCM's performance regarding this variable, and are happy to report that the model shows satisfactory to good performance in predicting the direction of changes in MH and LGM.

First, the authors need to decide what the point of the paper is. Most of the paper reads as a description of climatological changes for 3 periods in the geologic past (MH, LGM, and Plio) (indeed, most of the text is written this way). However, this work has already been done, most prominently by the PMIP and PlioMIP set of model intercomparisons. What does this manuscript offer that these model intercomparisons have not already analyzed? A case could be made that these model intercomparisons are typically of a global nature (though there has been some work on changes in the Asian Monsoon systems using both PlioMIP and PMIP results (Jiang et al., 2013; Jiang and Lang, 2010; Zhang et al., 2013)), so that the analysis of the orogens in this study is useful. However, given that GCMs have difficulty simulating precipitation and in particular simulating precipitation over complex topography, the usefulness of simply describing changes over the Himalaya, Andes, Cascades, and St. Elias ranges is somewhat muted. For example, why should readers believe that ECHAM5 produces reasonable results over the St. Elias range? Why not use (or at least compare with) the existing model intercomparisons to look at changes in these locations? Most of the PlioMIP simulations are of a lower resolution than the model simulations presented here, though not all (Haywood et al., 2013). Many of the newer PlioMIP2 simulations are being run at a higher resolution and permit at least some comparison with the data here (Chandan and Peltier, 2017). Having myself tried to access PlioMIP data, I understand that it can be difficult to get access to the PlioMIP output, but if the point of the paper is to quantify actual climatological change, then comparison with other models is a must (or at least a thorough treatment of possible boundary condition uncertainties and additional ECHAM5 model runs to establish the sensitivity of precipitation/temperature in these areas); otherwise, we have no reason to believe that ECHAM5 presents anything resembling a proper picture of climatic change in the past.

Summary Response: We appreciate this assessment and hope to have enhanced the usefulness of the manuscript with modifications we've made throughout the results and discussion sections. While GCM simulations for the time slices we chose already exist and individual studies were also conducted at a relatively high resolution, we believe (contrary to the reviewer's suggestion) that the usefulness of our results lies in both the high resolution combined with the consistency in model choice, resolution, output frequency (and methods of descriptive statistics). We emphasise this point in the revised manuscript. We acknowledge in the text (methods section 2.1 and introduction) the shortcomings of GCMs in predicting orographic precipitation in the discussion and in the revised text now compare the simulated precipitation with readily available proxy-based reconstructions for specific locations in Tibet and South America where we were able to compile data (this was a large undertaking). The revised manuscript now demonstrates the shortcomings of GCMs and provides an additional data compilation that *ESurf* readers interested in

285 palaeoclimate effects on denudation may find useful. The changes we have made to address this reviewers
286 suggestions are contained in the revised introduction, and extensively throughout the Discussion section
287 (4.0), as well as the addition of two new (concluding figures).
288

289 **In More Detail:** While we appreciated the reviewer's comments, we note that what he/she is asking for in
290 the above comments is a manuscript that either focuses on an inter-model comparisons, or a comparison to
291 proxy data. We see merits in both, but both are not possible in a single manuscript. An inter-model
292 comparison is not really well suited to the aims of this journal and our target community to bring attention to
293 the magnitude of palaeoclimate changes that have occurred in different active orogens. Our intended
294 audience is not the palaeoclimate modelling community (and journals associated with it), but rather the
295 surface processes community. We specifically chose this journal to provide this community with
296 palaeoclimate predictions that may be of interest to them. In our own experience in trying to interpret
297 palaeodenudation rates from data we produce, our first goal is always trying to find predictions or
298 observations of climate change for a region. Thus, we set out to write this paper to provide what we find in
299 our research as the first, most useful, step in understanding the surface process history of a region.
300 Furthermore, one major aspect of this study that we want to retain is the statistical analysis of the
301 climatology of each time slice as this is what is most useful for the ESurf community in terms of knowing if
302 a region they are working is has experienced a signal change in climate through time.
303

304 Thus, we are left with the conundrum that what the review requests and we want would normally be three
305 full manuscripts that include: 1) a statistical analysis of climatological changes over active orogens (our
306 previous focus of the text), or 2) a comparison of a suite of similarly set up and common code (ECHAM5)
307 simulations of palaeoclimate time slices to other GCM models (one of the suggestions by this reviewer), or
308 3) a model-data comparisons for model evaluation at the time slices investigated (also suggested by this
309 reviewer).
310

311 ***In an attempt to hopefully reach a compromise with this reviewer – the revised manuscript is now***
312 ***structured around points 1 and 3 above (statistical analysis of climatological changes, and model-data***
313 ***comparison where possible).*** These changes have been manifested in a now expanded discussion section,
314 and through the addition of model-data comparison figures at the end of the text (see section 4.1.2 , 4.2.2,
315 and 4.4). These changes hopefully reach a happy middle-ground between our aims and the reviewer's
316 suggestions. The changes have significantly expanded the manuscript, and we hope that the reviewer also
317 recognizes that implementation of all the suggested changes is not possible within a single manuscript of
318 typical length for this journal. The revised manuscript now has 18 figures (including 1 in the supplemental
319 section) and 21 pages of manuscript text (not including references / captions).
320

321 Some parts of the paper address actual causes for climatic change (for example, discussion of the Pacific
322 North American Teleconnection (no citation given). Is this related to the PDO? Lines 493-495). Again, if this
323 is the primary point of the paper, then substantially more work needs to be done to address why
324 precipitation, for example, increases in the Himalaya in the Pliocene. If this was the point of the paper, it
325 would obviate the need to compare with other model simulations (see paragraph above), but would then
326 require substantially more work to identify how various atmospheric phenomenon change through time.
327 A generalized description of changes in the past is not particularly useful (most of this information can be
328 communicated fairly effectively with figures), so addressing the causes of these changes (or comparison
329 with proxies – see below) is worthwhile.
330

331 Parts of the discussion showcase comparisons with terrestrial proxy data, though this is limited to citing
332 previous work and stating that there is general agreement with previous, proxy-model work. If a proxy-model
333 comparison is the point of the paper, then more work needs to be done actually compiling the proxies and
334 doing a proper statistical test to see if there is agreement between modelled precip/temperature changes in
335 each of these orogens and existing data. This would be a useful contribution to the literature, but, again,
336 would require substantially more work.
337

338 We address the suggestions in the previous 2 paragraphs in the following ways: **First**, as was mentioned
339 earlier, we maintain that for this journal and the community that reads it there is value in demonstrating
340 climate change events in commonly studied active orogens where denudation studies (sensitive to climate
341 change) are conducted. This requires that we maintain our current statistical characterisations of the climate
342 change. With this approach, it then becomes intractable to focus on the climate dynamics associated with
343 change in each region because typically a thorough investigation of the climate dynamics associate with

344 change in each region could be a paper for each region in itself, and the broader picture of change in a range
345 would be missed. We try to meet the reviewer half way in this suggestion by including additional comments
346 on possible causes for climatic change in different regions.
347

348 The PNA (citation now included) is related to the PDO, and Alaskan precipitation and temperature is to
349 some degree controlled by it. In order to properly address these questions, however, further analyses looking
350 at dominant atmospheric variability modes, trajectories and other aspects of atmosphere would have to be
351 carried out, which are beyond the scope of this study. We have modified the manuscript in text at the end of
352 section 2.1 to clarify (and justify) why we do not conduct an inter model comparison, as well as some of the
353 caveats associated with our approach.
354

355 However, (**second**), we like the reviewer's suggestions of adding additional comparisons to proxy data, and
356 we heavily invested time in compiling this information in this revised version for locations where spatially
357 distributed data are available for the time slices available.

358 Unfortunately, changes from proxy studies are often reported in terms of relative changes compared to
359 modern (e.g., wetter/drier, warmer/colder) and are sometimes contradictory for the same location, making
360 the application of otherwise suitable statistical tests (e.g. t-test and even non-parametric tests) difficult.
361 There are, in some cases (see final figures in paper), even contradictions between the proxy data themselves
362 in neighbouring locations, a reality often underappreciated by the surface processes community.
363 Furthermore, terrestrial proxy data are not available for all the time slices and locations we investigate. This
364 is particularly true for the Pliocene time slice. **With these limitations of available proxies, we do the best
365 we can to compare available observations to proxy data. These comparisons are now provided in
366 revisions to the discussion section, and the addition of 2 figures at the end of the manuscript.**
367

368 We note that for the Alaska and Cascadia (NW USA) study areas, which are heavily and repeatedly
369 glaciated, that limited - to no proxy data are available (due to poor preservation of proxies over glacial-
370 interglacial cycles). Although some marine proxy records and records of past glaciations have been
371 published for these regions, they are not useful for our purposes because they do not record if terrestrial
372 locations in the regions were wetter/drier or warmer/colder during our time slices. Marine proxy records are
373 of limited use for comparison to the terrestrial changes investigated in this study. Thus, despite the lack of
374 terrestrial proxy comparisons available for two of the regions (Alaska, Cascadia), we maintain that
375 presentation of the model predicted changes for these data poor regions is extremely useful because the
376 model predictions augment existing terrestrial data gaps, and provide a starting point for future studies to
377 formulate testable hypotheses of climate change and potential denudation impacts.
378

379
380 Much of the motivation for the manuscript appears to be to understand how climatic changes may change
381 denudation/geomorphic analyses, but this is done in only a superficial way. If this is the point of the paper,
382 then, again, much more work needs to be done, rather than simply stating that erosion depends on climate
383 (lines 416-423 are a good example of statements that serve to motivate a paper, but don't provide any
384 actual analysis). For example, can the authors take some of the climate model output and, given a
385 potential 2000 mm/yr change in precipitation in the Himalaya since the Pliocene, actually re-interpret some
386 of the existing exhumation/denudation data? If not, why not? What additional data is necessary? And if such a
387 reanalysis isn't possible, then how does knowledge of such a change in precipitation facilitate future work?
388

389 This is a great suggestion and we are indeed doing this as ongoing work. But what the reviewer is asking
390 for here simply cannot be included in this paper without making it an extremely long manuscript. For
391 example, in the surface processes community, comparisons of river profiles to modern precipitation, or
392 temperature changes to frost cracking histories for **one** catchment / study area (e.g. <100x100 km²) is
393 typically a full manuscript in itself (e.g. Marhsall et al., 2015 Science Advances; Schaller et al. 2004
394 Journal of Geology). So, we respectfully disagree with the reviewer that this should be done within this
395 manuscript. It's simply not possible to do in a way that would be convincing to the surface processes
396 community.
397

398 I agree with the generalized statements made throughout the paper (i.e., that denudation and landscape
399 evolution depend on climate), but these are somewhat self-evident and, as currently written, the manuscript
400 does not make a fundamental contribution to improving our knowledge on this subject except to state that
401 climate changed in the past. Assumptions of stationarity are indeed a problem in interpreting modern datasets
402 that have a component of geologic history, but a really nice contribution of this paper would be to show how
403 these assumptions can be mitigated when one knows the history of precipitation or temperature.

404

405 We agree with the reviewer that it is “self-evident” that palaeoclimate can impact the denudation history of
406 an orogen, but how many palaeodenudation rate studies actually make a comparison to spatial distributed
407 predictions of palaeoclimate? Very few in our experience, and we cite the robust ones in the manuscript.
408 What most palaeodenudation rate studies do is compare some set of observation to the nearest set of proxy
409 observations, but this approach does not provide a sense of spatial variability in terrestrial climate change
410 that models predict (furthermore, proxy observations are often far removed from the denudation rate
411 observations). In our manuscript, we specifically avoid picking on previous publications that follow this
412 ‘conventional’ approach to show how this commonly used approach ignores spatial variability in climate
413 change. We don’t think this is a productive way to advance the science. Instead, we maintain that providing
414 spatially continuous model predictions at different time slices in a self-consistent set of simulations is
415 extremely useful for documenting the magnitude of climate change (which is not self-evident), and for
416 formulating testable hypotheses for future work and to identify where the best locations are in the world to
417 investigate climate change impacts on a regions denudation history. Thus, the way in which our research
418 group usually conducts palaeodenudation rate studies is to first run model simulations to formulate testable
419 hypotheses, and then to investigate/test these hypotheses with field studies and geochemical observations of
420 denudation histories. This approach to investigating palaeoclimate-palaeodenudation interactions is
421 definitely not self-evident in the literature in our opinion.

422 We very much agree that in order for our simulations to ultimately be useful in actually quantifying
423 denudation much more work needs to be done. Translating any of the changes observed here (or by other
424 studies) to erosion rates remains a big problem in the Earth surface science community. It is one that we are
425 actively working on and includes the application and comparison of different models for quantifying frost
426 cracking [Anderson et al., 1998, Hales and Roering, 2007, and Andersen et al. 2015] and possible
427 improvements on them, different measures for precipitation extremes and testing how well they are captured
428 in modern day simulations (by comparison with indices derived from observation based datasets), etc..
429 However, the format and length of a typical article does not allow thorough investigation of any of these as
430 part of this manuscript and we believe that an arbitrary selection of these efforts would ultimately be very
431 misleading. Instead, we hope to offer these consistently set up simulations as a useful framework for the
432 earth surface science community to build on. We added further descriptions and discussion to convey this.

433
434 Because ESurf is not a climate modeling journal, more discussion needs to be given as to the limitations of
435 ECHAM5 in a way that Earth surface process folks can understand. For example, what are the
436 uncertainties associated with simulating orographic precipitation? Though T159 is high-resolution, it still
437 requires substantially smoothing topography, which presumably introduces some uncertainty in to the results.
438 What uncertainties are associated with the PRISM reconstruction? (on a side note, which PRISM
439 reconstruction is used? PRISM3D? PRISM2? I mention this because the topographies between different
440 PRISM reconstructions are substantially different.)
441

442 We thank the reviewer for raising this important point. We emphasised the problem of simulating
443 orographic precipitation, and included the recommendation for downscaling where it may be required. We
444 used the PRISM3D reconstruction (as in Haywood et al. 2010), and commented on some of its uncertainties.
445

446
447 I found the use of the cluster analysis to be not intuitively helpful. If the authors want to keep using it, then the
448 authors need to at least walk the reader through an example of how to understand Figures 6, 9, 12, and 15.
449 Is C1 always the same climate zone in each figure and in each time-slice? If so, why are different colors
450 used? Why don’t the authors use something more intuitive, like Köppen’s climate classification scheme (Peel
451 et al., 2007) to classify climates? As best I understand it, the clustering analysis is used to show the spatial
452 extent of a given climate in a given time-slice and in a given location, but it’s not clear how one should
453 interpret these results.
454

455 We have modified the text in the results and discussion sections to help clarify this. In short, the clustering
456 analysis essentially fulfills a similar synoptic purpose, but optimises classification and is more fine-tuned to
457 this study’s purpose in its selection of variables. This is now reflected in text changes we made to the end of
458 the methods section 2.2. The climate clusters do indeed show the spatial extent of a given climate (described
459 by the mean vectors represented graphically in the raster plots and by numbers in the table included in the
460 supplementary material). The idea is to provide an overview of regional climate without the need to study
461 maps of individual variables, on which these patterns and the climatic homogeneity may not be seen as
462 easily. Each plot represents an optimal classification and thus cluster 1, for example, is not always described
463 by the same mean vector (though usually is usually very similar). The different colours are used to avoid the

464 interpretation that cluster 1, for example, is always characterised by the exact same mean vector. We have
465 included a more elaborate explanation in our revised manuscript to avoid confusion. While readers may be
466 more familiar with the Köppen climate classification scheme, we are more interested in providing an
467 overview not forcefully tied to the categories of this classification scheme. Clustering by various methods
468 (such as this one or PCA) as a synoptic tool are not uncommon (e.g. Paeth 2004, Mannig et al. 2013), but we
469 acknowledge that many readers may not be use to these tools and therefore elaborate explanations.
470

471
472 Minor Comments: Are the topographies for any of these ranges modified at all (it's unclear from the PRISM
473 reference (Line 131), whether this has been done).

474 **Yes, the topographies are different and we now specify the reconstruction we use (PRISM3D) in the**
475 **methods section.**

476
477 Lines 57-59: "Cold-temperature island" is not a climatic term in widespread use. What
478 precisely do you mean? Also, Boos and Kuang, 2010 specifically refute the idea that Plateau surface
479 elevation matters for the South Asian Monsoon and rather focus on the Himalaya instead.

480 **Yes, this matters only regionally, but not for the South Asian Monsoon. We have corrected this sentence.**
481

482 Line 61: Zhisheng et al. (2001) don't actually present any new geological data. Instead, it is all from cited
483 literature. The focus of their study was GCM results. Dettman et al. (2003) is not the only study to look at
484 this. Please see the following studies (which are just a sampling): (Caves et al., 2017; Kent-Corson et al.,
485 2006; Lechler et al., 2013; Lechler and Niemi, 2011; Licht et al., 2016; Methner et al., 2016; Mulch et al.,
486 2015,
487 2008; Pingel et al., 2016).

488 **Thank you for pointing out this inaccuracy and pointing us to additional studies we ought to list here. We**
489 **modified the text accordingly.**

490
491 Lines 77 and 78: "documenting the magnitude" appears twice.
492 **Thank you for pointing this out. We corrected this.**
493

494 Lines 96-97: Though, importantly, several recent studies have run ECHAM5 at a higher resolution ((Feng et
495 al., 2016; Feng and Poulsen, 2016)).
496 **That is true, of course. We mention these in our revised manuscript.**
497

498 Lines 102-103: This statement is somewhat odd, since the authors are specifically investigating
499 climatological changes over mountain ranges, where resolution typically tends to matter.
500 **Thank you for noticing this apparent inconsistency. We corrected the text accordingly.**
501

502 Lines 114-133: For all simulations, stating the pCO₂ used in the experiment would be most helpful,
503 particularly since it won't take up much room. Also, how is the land- surface treated? For example, the
504 authors state that they are using vegetation reconstructions, but it's unclear if this is then being fed into a
505 "built-in" land-surface model or if they are explicitly using JSBACH.
506 **This is a good point. We included these values in the text. We used the built-in land surface scheme (LSS) and**
507 **clarified this in the revised text.**
508

509 Lines 126: "for the" used twice.
510 **This has been corrected.**
511

512 Lines 203-204: Changes in Greenland and Antarctica are almost certainly unreliable. Because PRISM uses
513 a reconstructed ice-sheet extent, changes in temperature in Greenland and Antarctica are almost certainly
514 reflecting the imposed boundary condition, which itself has quite a bit of uncertainty. It's hard to get around
515 this, except to note that the change in temperature is entirely dependent upon the ice-sheet boundary
516 condition (see discussion in and of de Boer et al. (2015)).
517 **Thank you for pointing this out. We comment on the Pliocene uncertainty in ice sheet reconstructions in the**
518 **method section of the revised manuscript.**
519

520 Lines 416-423: Are runoff changes in these models coupled to precipitation changes? In all cases, does P-E
521 (precipitation minus evaporation) scale with changes in precipitation. I'm not particularly familiar with
522 JSBACH (presuming this is the land-surface model used), but if it has a CO₂ fertilization parameterization,
523 then runoff may be decoupled from precip. Some of these erosion processes may depend more on runoff
524 than precip.
525 **We use the built in LSS and the runoff is coupled to precipitation. Also, ECHAM5's runoff is not**
526 **particularly useful in river discharge modelling (see Weiland et al. 2011), which would be of interest in**
527 **context of erosion. Given this, we refrain from using the model predicted runoff in the global GCM (even**

528 though it's conducted at relatively high resolution compared to previous work) to calculate changes in
529 fluvial incision. This would be better done by mapping the predicted precipitation changes onto higher
530 resolution (<90 m) DEMs and solving the kinematic wave equation for each fluvial erosion in each
531 catchment, for the changes in precipitation. However, as we repeatedly mention above, this is not possible to
532 include in this manuscript without first characterising how the precipitation has changed in each region (the
533 current manuscript goals). Work in progress we are conducting is trying to apply the kinematic wave
534 equation and palaeoprecipitation to selected areas, but it's proving difficult to implement meaningfully
535 without temporally continuous (e.g. LGM to present) simulations of precipitation change. We hope this
536 brings to the readers attention the complications associated with doing full erosion history calculations based
537 on these results. We have expanded the last paragraph in the instruction to convey the above perspective
538 better, and more clearly articulate (and justify) the scope and limitations of the manuscript.

539
540
541 Figure 1: Would be nice to also plot the topography of the St. Elias range and the
542 Cascades.

543 This is a good idea. We included ECHAM5 topographies for Alaska and the Pacific Northwest in Fig. 1 in
544 the revised manuscript.

545
546 Figure 7b-Precip-PLIO: Why does precipitation appear to follow a wave-like pattern over tropical South
547 America? Is this due to the spectral nature of ECHAM5?

548
549 This may indeed be due to the spectral nature of ECHAM5.

550
551
552 References used in review:

553
554 Thank you for being thoughtful enough to provide these. We have added many of them to the revised text.

555
556 An, Z., Kutzbach, J.E., Prell, W.L., Porter, S.C., 2001. Evolution of Asian monsoons and phased uplift of
557 the Himalaya-Tibetan plateau since Late Miocene times. *Nature* 411, 62–66. doi:10.1038/35075035

558
559 Caves, J.K., Bayshashov, B.U., Zhamangara, A., Ritch, A.J., Ibarra, D.E., Sjostrom, D.J., Mix, H.T.,
560 Winnick, M.J., Chamberlain, C.P., 2017. Late Miocene uplift of the Tian Shan and Altai and reorganization of
561 Central Asia climate. *GSA Today* 27, 19–26.

562
563 Chandan, D., Peltier, W.R., 2017. Regional and global climate for the mid-Pliocene using CCSM4 and
564 PlioMIP2 boundary conditions. *Clim. Past* 13, 919–942. doi:10.5194/cp-2017-21

565
566 de Boer, B., Dolan, A.M., Bernales, J., Gasson, E., Goelzer, H., Golledge, N.R., Sutter, J., Huybrechts, P.,
567 Lohmann, G., Rogozhina, I., Abe-Ouchi, A., Saito, F., van de Wal, R.S.W., 2015. Simulating the Antarctic
568 ice sheet in the Late-Pliocene warm period: PLISMIP-ANT, an ice-sheet model intercomparison project.
569 *Cryosph.* 9, 881–903. doi:10.5194/tc-9-881-2015

570
571 Dettman, D.L., Fang, X., Garzzone, C.N., Li, J., 2003. Uplift-driven climate change at
572 12 Ma: a long $\delta^{18}O$ record from the NE margin of the Tibetan plateau. *Earth Planet. Sci. Lett.* 214, 267–
573 277. doi:10.1016/S0012-821X(03)00383-2

574
575 Feng, R., Poulsen, C.J., 2016. Refinement of Eocene lapse rates, fossil-leaf altimetry, and North American
576 Cordilleran surface elevation estimates. *Earth Planet. Sci. Lett.* doi:10.1016/j.epsl.2015.12.022

577
578 Feng, R., Poulsen, C.J., Werner, M., 2016. Tropical circulation intensification and tectonic extension
579 recorded by Neogene terrestrial $\delta^{18}O$ records of the western United States. *Geology* 44.
580 doi:10.1130/G38212.1

581
582 Haywood, A.M., Hill, D.J., Dolan, A.M., Otto-Bliesner, B.L., Bragg, F., Chan, W.L., Chandler, M.A.,
583 Contoux, C., Dowsett, H.J., Jost, A., Kamae, Y., Lohmann, G., Lunt, D.J., Abe-Ouchi, A., Pickering, S.J.,
584 Ramstein, G., Rosenbloom, N.A., Salzmann, U., Sohl, L., Stepanek, C., Ueda, H., Yan, Q., Zhang, Z.,
585 2013. Large-scale features of Pliocene climate: Results from the Pliocene Model Intercomparison Project.
586 *Clim. Past* 9, 191–209. doi:10.5194/cp-9-191-2013

587
588 Jiang, D., Lang, X., 2010. Last glacial maximum East Asian monsoon: Results of PMIP
589 simulations. *J. Clim.* 23, 5030–5038. doi:10.1175/2010JCLI3526.1

590
591 Jiang, D., Lang, X., Tian, Z., Ju, L., 2013. Mid-Holocene East Asian summer monsoon strengthening:
592 Insights from Paleoclimate Modeling Intercomparison Project (PMIP) simulations. *Palaeogeogr.*
593 *Palaeoclimatol. Palaeoecol.* 369, 422–429. doi:10.1016/j.palaeo.2012.11.007

- 594
595 Kent-Corson, M.L., Sherman, L.S., Mulch, A., Chamberlain, C.P., 2006. Cenozoic topographic and
596 climatic response to changing tectonic boundary conditions in Western North America. *Earth Planet. Sci.*
597 *Lett.* 252, 453–466. doi:10.1016/j.epsl.2006.09.049
- 598
599 Lechler, A.R., Niemi, N.A., 2011. Sedimentologic and isotopic constraints on the Paleogene
600 paleogeography and paleotopography of the southern Sierra Nevada, California. *Geology* 39, 379–382.
601 doi:10.1130/G31535.1
- 602 Lechler, A.R., Niemi, N. a., Hren, M.T., Lohmann, K.C., 2013. Paleoelevation estimates for the northern and
603 central proto-Basin and Range from carbonate clumped isotope thermometry. *Tectonics* 32.
604 doi:10.1002/tect.20016
- 605
606 Licht, A., Quade, J., Kowler, A., Santos, M. de los S., Hudson, A., Schauer, A., Huntington, K., Copeland,
607 P., Lawton, T., 2016. Impact of the North American Monsoon on isotope paleoaltimeters: Implications for the
608 paleoaltimetry of the American Southwest. *Am. J. Sci.* 317.
- 609
610 Methner, K., Fiebig, J., Wacker, U., Umhoefer, P., Chamberlain, C.P., Mulch, A., 2016. Eo-Oligocene proto-
611 Cascades topography revealed by clumped ($\Delta 47$) and oxygen isotope ($\delta 18O$) geochemistry (Chumstick
612 Basin, WA, USA). *Tectonics* 35, 546–564. doi:10.1002/2015TC003984
- 613
614 Mulch, A., Chamberlain, C.P., Cosca, M.A., Teyssier, C., Methner, K., Hren, M.T., Graham, S.A., 2015.
615 Rapid change in high-elevation precipitation patterns of western North America during the Middle Eocene
616 Climatic Optimum (MECO). *Am. J. Sci.* 315,
617 317–336. doi:10.2475/04.2015.02
- 618
619 Mulch, A., Sarna-Wojcicki, A.M., Perkins, M.E., Chamberlain, C.P., 2008. A Miocene to Pleistocene climate
620 and elevation record of the Sierra Nevada (California). *Proc. Natl. Acad. Sci. U. S. A.* 105, 6819–6824.
621 doi:10.1073/pnas.0708811105
- 622
623 Peel, M.C., Finlayson, B.L., McMahon, T.A., 2007. Updated world map of the Köppen-Geiger climate
624 classification. *Hydrol. Earth Syst. Sci.* 11, 1633–1644.
- 625
626 Pingel, H., Mulch, A., Alonso, R.N., Cottle, J., Hynek, S.A., Poletti, J., Rohrmann, A., Schmitt, A.K., Stockli,
627 D.F., Strecker, M.R., 2016. Surface uplift and convective rainfall along the southern Central Andes
628 (Angastaco Basin, Argentina). *Earth Planet. Sci. Lett.* 440, 33–42. doi:10.1016/j.epsl.2016.02.009
- 629
630 Zhang, R., Yan, Q., Zhang, Z.S., Jiang, D., Otto-Bliesner, B.L., Haywood, a. M., Hill, D.J., Dolan, a. M.,
631 Stepanek, C., Lohmann, G., Contoux, C., Bragg, F., Chan, W.-L.,
- 632 Chandler, M.a., Jost, a., Kamae, Y., Abe-Ouchi, a., Ramstein, G., Rosenbloom, N. a., Sohl, L., Ueda, H.,
633 2013. East Asian monsoon climate simulated in the PlioMIP. *Clim. Past Discuss.* 9,1135-1164.
634 doi:10.5194/cpd-9-1135-2013

635

636

637

638

639

640

641

642 List of Major Changes

643 Overall, we found the reviewer's comments refreshingly useful. We greatly appreciate the time the editors have taken to
644 find thoughtful reviewers, and the reviewers time in providing constructive comments. We were able to implement most
645 of the suggested changes. However, some of the requested changes by reviewer 2 would require multiple manuscripts,
646 and in our response to this reviewers comments we explain a (hopefully) happy middle ground between their requests
647 and the reality of what we think can be realistically implemented in a manuscript for the ESurf community of readers.
648 **The most extensive changes to the manuscript are in the introduction and discussion sections, as well as in the**
649 **modification of previously presented figures (Fig. 1, 2, 3, 7, 10, 13)** - since modification of these sections and figures
650 most directly addresses the reviewers concerns and they did not identify errors in our results required new simulations.

651
652 The general concern that resonated in both reviewers comments was the need to focus the manuscript more and add
653 'meat' to our arguments in one direction or another. We have taken this to heart, and **the revised manuscript presents**
654 **an in-depth (and very time consuming) compilation of terrestrial proxy data observations for two of the regions**
655 **(South Asia, South America)** which were not extensively glaciated and have good records available. These proxy data
656 were compared to the model results and provide a new and focused dimension to the manuscript. **As a results of these**
657 **efforts, we included two additional figures (Fig. 16, 17).** As detailed in our response to reviewer 2 - we do not present
658 a detailed inter-model comparison, although we have expanded our comparison to previous published work throughout
659 the text (mostly in the discussion section). We also provide additional justification for our approach, and highlight more
660 explicitly what is novel about our presentation of a series of experiments conducted with the same model at the same
661 resolution.

662
663 The marked-up manuscript below highlights changes made to the manuscript in more detail.

664
665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

690 **Where is Late Cenozoic climate change most likely to impact**
691 **denudation?**

692

693 Sebastian G. Mutz¹, Todd A. Ehlers¹, Martin Werner², Gerrit Lohmann², Christian Stepanek²,

694 Jingmin Li^{1,3}

695 ¹Department of Geosciences, University Tübingen, D-72074 Tübingen, Germany

²Department of Paleoclimate Dynamics, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, D-27570 Bremerhaven, Germany

³[now at](#) Institute for Geography and Geology, University of Würzburg, Würzburg, D-97074 Germany

696

697 *Correspondence to:* Sebastian G. Mutz (sebastian.mutz@uni-tuebingen.de)

698

699 **Abstract**

700 The denudation history of active orogens is often interpreted in the context of modern climate and vegetation gradients.

701 Here we address the validity of this approach and ask the question: what are the spatial and temporal variations in

702 palaeo-climate for a latitudinally diverse range of active orogens? We do this using high-resolution (T159, ca. 80 x 80

703 km at the equator) palaeo-climate simulations from the ECHAM5 global Atmospheric General Circulation Model and a

704 statistical cluster analysis of climate over different orogens (Andes, Himalaya, SE Alaska, Pacific NW USA). Time

705 periods and boundary conditions considered include the Pliocene (PLIO, ~3 Ma), the Last Glacial Maximum (LGM,

706 ~21 ka), Mid Holocene (MH, ~6 ka) and Pre-Industrial (PI, reference year 1850). The regional simulated climates of

707 each orogen are described by means of cluster analyses based on the variability of precipitation, 2m air temperature, the

708 intra-annual amplitude of these values, and monsoonal wind speeds where appropriate. Results indicate the largest

709 differences to the PI climate ~~are observed~~ existed for the LGM and PLIO climates in the form of widespread cooling and

710 reduced precipitation in the LGM and warming and enhanced precipitation during the PLIO. The LGM climate shows

711 the largest deviation in annual precipitation from the PI climate, and shows enhanced precipitation in the temperate

712 Andes, and coastal regions for both SE Alaska and the US Pacific Northwest [Pacific](#). Furthermore, LGM precipitation

713 is reduced in the western Himalayas and enhanced in the eastern Himalayas, resulting in a shift of the wettest regional

714 climates eastward along the orogen. The cluster-analysis results also suggest more climatic variability across latitudes

715 east of the Andes in the PLIO climate than in other time-slice experiments conducted here. Taken together, these results

716 highlight significant changes in Late Cenozoic regional climatology over the last ~3 Ma. [Comparison of simulated](#)

717 [climate with proxy-based reconstructions for the MH and LGM reveal satisfactory to good performance of the model in](#)

718 [reproducing precipitation changes, although in some cases discrepancies between neighbouring proxy observations](#)
719 [highlight contradictions between proxy observations themselves](#). Finally, we document regions where the largest
720 magnitudes of Late Cenozoic changes in precipitation and temperature occur and offer the highest potential for future
721 observational studies ~~interested in quantifying~~ that quantify the impact of climate change on denudation and weathering
722 rates.

723

724 **Keywords:** Cenozoic climate, ECHAM5, Last Glacial Maximum, Mid-Holocene, Pliocene, cluster analysis, Himalaya,
725 Tibet, Andes, Alaska, Cascadia

726

727 1. Introduction

728 Interpretation of orogen denudation histories in the context of climate and tectonic interactions is often hampered
729 by a paucity of terrestrial palaeo-climate proxy data needed to reconstruct spatial variations in palaeo-climate. [While it](#)
730 [is self-evident that palaeoclimate changes could influence palaeodenudation rates, it is not always self-evident what the](#)
731 [magnitude of climate change over different geologic time scales is, or what geographic locations offer the greatest](#)
732 [potential to investigate palaeoclimate impacts on denudation](#). Palaeoclimate reconstructions are particularly beneficial
733 when denudation rates are determined using geo- and thermo-chronology techniques that integrate over timescales of
734 10^3 - 10^6 years (e.g. cosmogenic radionuclides or low-temperature thermochronology) [e.g., Kirchner et al., 2001;
735 Schaller et al., 2002; Bookhagen et al., 2005; Moon et al., 2011; Thiede and Ehlers, 2013; Lease and Ehlers, 2013].
736 However, few studies using denudation rate determination methods that integrate over longer timescales have access to
737 information about past climate conditions that could influence these palaeo-denudation rates. Palaeo-climate modelling
738 offers an alternative approach to sparsely available proxy data for understanding the spatial and temporal variations in
739 precipitation and temperature in response to changes in orography [e.g. Takahashi and Battisti, 2007a, b; Insel et al.,
740 2010; Feng et al., 2013] and global climate change events [e.g. Salzmann, 2011; Jeffery et al., 2013]. In this study, we
741 characterize the climate at different times in the [Late](#) Cenozoic, and the magnitude of climate change for a range of
742 active orogens. Our emphasis is on identifying changes in climate parameters relevant to weathering and catchment
743 denudation to illustrate the potential importance of various global climate change events on surface processes.

744 Previous studies of orogen-scale climate change provide insight into how different tectonic or global climate
745 change events influence regional climate change. For example, sensitivity experiments demonstrated significant
746 changes in regional and global climate in response to landmass distribution and topography of the Andes, including
747 changes in moisture transport, [the north-south asymmetry of the Intertropical Convergence Zone and the north-south](#)
748 [asymmetry of the Inter-Tropical Convergence Zone](#) [e.g. Takahashi and Battisti, 2007a, ; Insel et al., 2010] [and](#)

749 [\(tropical\) precipitation](#) [Maroon et al., 2015, ; ~~Maroon et al.~~ 2016]. [Another example is the regional](#) and global climate
750 changes induced by the Tibetan Plateau surface uplift due to its role as a ~~cold-temperature island and~~ physical obstacle
751 to circulation [Raymo and Ruddiman, 1992; Kutzbach et al., 1993; Thomas, 1997; Bohner, 2006; Molnar et al., 2010;
752 Boos and Kuang, 2010]. The role of tectonic uplift in long term regional and global climate change remains a focus of
753 research and continues to be assessed with geologic datasets [e.g. ~~Zhisheng, 2001~~; Dettman et al., 2003; [Caves et al.,](#)
754 [2017](#); [Kent-Corson et al., 2006](#); [Lechler et al., 2013](#); [Lechler and Niemi, 2011](#); [Licht et al., 2016](#);
755 [Methner et al., 2016](#); [Mulch et al., 2015, 2008](#); [Pingel et al., 2016](#)] and climate modelling [e.g. Kutzbach et
756 al., 1989; Kutzbach et al., 1993; ~~Zhisheng, 2001~~; Bohner, 2006; Takahashi and Battisti, 2007a; Ehlers and Poulsen,
757 2009; Insel et al., 2010; Boos and Kuang, 2010]. Conversely, climate influences tectonic processes through erosion
758 [e.g. Molnar and England, 1990; Whipple et al., 1999; Montgomery et al., 2001; Willett et al., 2006; Whipple, 2009].
759 Quaternary climate change between glacial and interglacial conditions [e.g. Braconnot et al., 2007; Harrison et al.,
760 2013] resulted in not only the growth and decay of glaciers and glacial erosion [e.g. Yanites and Ehlers, 2012; Herman
761 et al., 2013; Valla et al., 2011] but also global changes in precipitation and temperature [e.g. Otto-Bliesner et al., 2006;
762 Li et al., 2017] that could influence catchment denudation in non-glaciated environments [e.g. Schaller and Ehlers,
763 2006; Glotzbach et al., 2013; Marshall et al., 2015]. These dynamics highlight the importance of investigating how
764 much climate has changed over orogens that are [the](#) focus of studies of climate-tectonic interactions and their impact on
765 erosion.

766 Despite recognition by previous studies that climate change events relevant to orogen denudation are prevalent
767 throughout the Late Cenozoic, few studies have critically evaluated how different climate change events may, or may
768 not, have affected the orogen climatology, weathering and erosion. Furthermore, recent controversy exists concerning
769 the spatial and temporal scales over which geologic and geochemical observations can record climate-driven changes in
770 weathering and erosion [e.g. Whipple, 2009; von Blanckenburg et al., 2015; Braun, 2016]. [For example, the previous](#)
771 [studies highlight that although palaeoclimate impacts on denudation rates are evident in some regions and measurable](#)
772 [with some approaches, they are not always present \(or detectable\) and the spatial and temporal scale of climate change](#)
773 [influences our ability to record climate sensitive denudation histories.](#) This study contributes to our understanding of
774 the interactions between climate, weathering, and erosion by bridging the gap between the palaeoclimatology and
775 surface processes communities by documenting the magnitude and distribution of climate change over tectonically
776 active orogens. ~~Our focus is on documenting the magnitude of paleoclimateclimate and climate change in different~~
777 ~~locations with the intent of informing past and ongoing paleodenudation studies of these regions. The application of~~
778 ~~these results to predicted changes in denudation rates is beyond the scope of this study and the focus of future work.~~

779 ~~We~~ [In this study, we](#) employ the ECHAM5 global Atmospheric General Circulation Model and document
780 climate [and climate](#) change for time slices ranging between the Pliocene (PLIO, ~3 Ma) to pre-industrial (PI) times for

781 the St. Elias Range of South East Alaska, the US Pacific Northwest (Olympic and Cascade Range), western South
782 America (Andes) and South Asia (incl. parts of Central- and East Asia). Our approach is two-fold and includes:

783 | 1. An empirical characterizsation of palaeo-climates in these regions based on the covariance and spatial
784 clustering of monthly precipitation and temperature, the monthly change in precipitation and temperature magnitude,
785 and wind speeds where appropriate.

786 | 2. Identification of changes in annual mean precipitation and temperature in selected regions ~~over in the~~
787 ~~following time, specifically from the~~ for four time periods: (PLIO, ~~to the~~ Last Glacial Maximum (LGM), the Mid-
788 Holocene (MH) and PI): [and subsequent validation of the simulated precipitation changes for MH and LGM.](#)

789 Our focus is on documenting climate and climate change in different locations with the intent of informing past and
790 ongoing [palaeodenudation studies of these regions. The results presented here also provide a means for future work to](#)
791 [formulate testable hypotheses and investigations into whether or not regions of large palaeoclimate change produced a](#)
792 [measurable signal in denudation rates. In this study, we intentionally refrain from applying predicted palaeoclimate](#)
793 [changes to predict denudation rate changes. Such a prediction is beyond the scope of this study because a convincing](#)
794 [\(and meaningful\) calculation of climate-driven transients in fluvial erosion \(e.g. via the kinematic wave equation\),](#)
795 [variations in frost cracking intensity, or changes in hillslope sediment production and transport at the large regional](#)
796 [scales considered here is not tractable within a single manuscript, and instead is the focus of our ongoing work. Instead,](#)
797 [our emphasis lies on addressing the first question we are confronted with in our own research into denudation rate](#)
798 [studies around world, namely - where is Late Cenozoic climate change most likely to impact denudation?](#)

799

800 | 2. Methods: Climate modelling and cluster analyses for climate characteriszation

801

802 2.1 ECHAM5 simulations

803 The global Atmospheric General Circulation Model ECHAM5 [Roeckner et al., 2003] has been developed at the
804 Max Planck Institute for Meteorology and is based on the spectral weather forecast model of the ECMWF [Simmons et
805 al., 1989]. In the context of palaeoclimate applications, the model has been used mostly at lower resolution (T31,
806 [approximately](#)ca. 3.75°x3.75°; [T63, ca. 1.9°x1.9° in case of Feng et al. \[2016\] and T106 in the case of Li et al. \[2016\]](#)
807 [and Feng and Poulsen \[2016\]](#)). The performed studies are not limited to the last millenium [e.g. Jungclaus et al., 2010]
808 but also include research in the field of both warmer and colder climates, at orbital [e.g. Gong et al., 2013; Lohmann et
809 al., 2013; Pfeiffer and Lohmann, 2016; Zhang et al., 2013a; Zhang et al., 2014; Wei and Lohmann, 2012] and tectonic
810 time scales [e.g. Knorr et al., 2011; Stepanek and Lohmann, 2012], and under anthropogenic influence [Gierz et al.,
811 2015].

812 Here, the ECHAM5 simulations were conducted at a T159 spatial resolution (horizontal grid size ca. 80 km x 80

813 km at the equator) with 31 vertical levels (between the surface and 10hPa). This high model resolution is admittedly not
814 required for [all of the climatological questions investigated in this study, and it should be noted that the skill of GCM's](#)
815 [in predicting orographic precipitation remains limited at this scale \[e.g. Meehl et al. 2007\]](#). However, simulations were
816 conducted at this resolution so that future work can apply the results [in combination with different dynamical and](#)
817 [statistical downscaling methods](#) to quantify changes at large catchment to orogen scales. [The output frequency is](#)
818 [relatively high \(1 day\) to enhance the usefulness of our simulations as input for landscape evolution and other models](#)
819 [that may benefit from daily input.](#)-The simulations were conducted for five different time periods: present-day (PD), PI,
820 MH, LGM and PLIO.

821 A PD simulation (not shown here) was used to establish confidence in the model performance before conducting
822 [palaeo-simulations](#) and has been compared with the following observation-based datasets: European Centre for
823 Medium-Range Weather Forecasts (ECMWF) re-analyses [ERA40, Uppala et al., 2005], National Centers for
824 Environmental Prediction and National Center for Atmospheric Research (NCEP/NCAR) re-analyses [Kalnay et al.,
825 1996; Kistler et al., 2001], NCEP Regional Reanalysis (NARR) [Mesinger et al., 2006], the Climate Research Unit
826 (CRU) TS3.21 dataset [Harris et al., 2013], High Asia Refined Analysis (HAR30) [Maussion et al., 2014] and the
827 University of Delaware dataset (UDEL v3.01) [Legates et al., 1990]. (See Mutz et al. [2016] for a detailed comparison
828 with a lower resolution model).

829 The PI climate simulation is an ECHAM5 experiment with PI (reference year 1850) boundary conditions. Sea
830 Surface Temperatures (SST) and Sea Ice Concentration (SIC) are derived from transient coupled ocean-atmosphere
831 simulations [Lorenz and Lohmann, 2004; Dietrich et al., 2013]. Following Dietrich et al. [2013], greenhouse gas
832 (GHG) concentrations ([CO₂: 280 ppm](#)) are taken from ice core based reconstructions of CO₂ [Etheridge et al., 1996],
833 CH₄ [Etheridge et al., 1998] and N₂O [Sowers et al., 2003]. Sea surface boundary conditions for MH originate from a
834 transient, low-resolution, coupled atmosphere-ocean simulation of the mid (6 ka) Holocene [Wei and Lohmann, 2012;
835 Lohmann et al, 2013], where the GHG concentrations ([CO₂: 280 ppm](#)) are taken from ice core reconstructions of
836 GHG's by Etheridge et al. [1996], Etheridge et al. [1998] and Sowers et al. [2003]. GHG's concentrations for the LGM
837 ([CO₂: 185 ppm](#)) have been prescribed following Otto-Bliesner et al. [2006]. Orbital parameters for MH and LGM are
838 set according to Dietrich et al. [2013] and Otto-Bliesner et al. [2006], respectively. LGM land-sea distribution and ice
839 sheet extent and thickness are set based on the PMIP III ([Palaeoclimate Modelling Intercomparison Project, phase 3](#))
840 guidelines (elaborated on by Abe-Ouchi et al [2015]). Following Schäfer-Neth and Paul [2003], SST and SIC for the
841 LGM are based on GLAMAP [Sarnthein et al. 2003] and CLIMAP [CLIMAP project members, 1981] reconstructions
842 for the [for the](#) Atlantic and Pacific/Indian Ocean, respectively. Global MH and LGM vegetation are based on maps of
843 plant functional types by the BIOME 6000 / Palaeovegetation Mapping Project [Prentice et al., 2000; Harrison et al.,
844 2001; Bigelow et al., 2003; Pickett et al., 2004] and model predictions by Arnold et al. [2009]. Boundary conditions for

845 | the PLIO simulation, including GHG concentrations ([CO2: 405](#)), orbital parameters and surface conditions (SST, SIC,
846 | sea land mask, topography and ice cover) are taken from the PRISM (Pliocene Research, Interpretation and Synoptic
847 | Mapping) project [Haywood et al., 2010; Sohl et al., 2009; Dowsett et al., 2010], [specifically PRISM3D](#). The PLIO
848 | vegetation boundary condition was created by converting the PRISM vegetation reconstruction to the JSBACH plant
849 | functional types as described by Stepanek and Lohmann [2012], [but the built-in land surface scheme was used](#).

850 | [SST reconstructions can be used as an interface between oceans and atmosphere \[e.g. Li et al. 2016\] instead of](#)
851 | [conducting the computationally more expensive fully coupled Atmosphere-Ocean GCM experiments. While the use of](#)
852 | [SST climatologies comes at the cost of capturing decadal-scale variability, and the results are ultimately biased towards](#)
853 | [the SST reconstructions the model is forced with, the simulated climate more quickly reaches an equilibrium state and](#)
854 | [the means of atmospheric variables used in this study do no change significantly after the relatively short spin-up](#)
855 | [period](#). The palaeoclimate simulations (PI, MH, LGM, PLIO) using ECHAM5 are [therefore](#) carried out for 17 model
856 | years, of which the first two years are used for model spin up. The monthly long-term averages (multi-year means for
857 | individual months) for precipitation, temperature, as well as precipitation and temperature amplitude, i.e. the mean
858 | difference between the hottest and coldest months, have been calculated from the following 15 model years for the
859 | analysis presented below.

860 | For further comparison between the simulations, the investigated regions were subdivided (Fig. 1). Western
861 | South America was subdivided into four regions: parts of tropical South America (80°-60° W, 23.5°-5° S), temperate
862 | South America (80°-60° W, 50°-23.5° S), tropical Andes (80°-60° W, 23.5°-5° S; high-pass filtered), i.e. most of the
863 | Peruvian Andes, Bolivian Andes and northernmost Chilean Andes, and temperate Andes (80°-60° W, 50°-23.5° S,
864 | high-pass filtered). South Asia was subdivided into three regions: tropical South Asia (40°-120°E, 0°-23.5°N),
865 | temperate South Asia (40°-120°E, 23.5°-60°N), and high altitude South Asia (40°-120°E, 0°-60°N; high-pass filtered).

866 | [Our approach of using a single GCM \(ECHAM5\) for our analysis is motivated by, and differs from, previous](#)
867 | [studies where inter-model variability exists from the use of different GCMs due to different parameterisations in each](#)
868 | [model. The variability in previous inter-model GCM comparisons exists despite the use of the same forcings \[e.g. see](#)
869 | [results highlighted in IPCC AR5\]. Similarities identified between these palaeoclimate simulations conducted with](#)
870 | [different GCMs using similar boundary conditions can establish confidence in the models when in agreement with](#)
871 | [proxy reconstructions. However, differences identified in inter-model GCM comparisons highlight biases by all or](#)
872 | [specific GCMs, or reveal sensitivities to one changed parameter, such as model resolution. Given these limitations of](#)
873 | [GCM modelling, we present in this study a comparison of a suite of ECHAM5 simulations to proxy-based](#)
874 | [reconstructions \(where possible\) and, to a lesser degree, comment on general agreement or disagreement of our](#)
875 | [ECHAM5 results with other modelling studies. A detailed inter-model comparison of our results with other GCMs is](#)
876 | [beyond the scope of this study, and better suited for a different study in a journal with a different focus and audience.](#)

877 [Rather, by using the same GCM and identical resolution for the time slice experiments, we reduce the number of](#)
878 [parameters \(or model parameterisations\) varying between simulations and thereby remove potential sources of error or](#)
879 [uncertainty that would otherwise have to be considered when comparing output from different models with different](#)
880 [parameterisations of processes, model resolution, and in some cases model forcings \(boundary conditions\).](#)
881 [Nevertheless, the reader is advised to use these model results with the GCM's shortcoming and uncertainties in](#)
882 [boundary condition reconstructions in mind. For example, precipitation results may require dynamical or statistical](#)
883 [downscaling to increase accuracy where higher resolution precipitation fields are required. Furthermore, readers are](#)
884 [advised to familiarise themselves with the palaeogeography reconstruction initiatives and associated uncertainties. For](#)
885 [example, while Pliocene ice sheet volume can be estimated, big uncertainties pertaining to their locations remain](#)
886 [\[Haywood et al. 2010\].](#)

887

888 **2.2 Cluster analysis to document temporal and spatial changes in climatology**

889 ~~This section describes the clustering method used in this study.~~ The aim of the clustering approach is to group
890 climate model surface grid boxes together based on similarities in climate. Cluster analyses are statistical tools that
891 allow elements (i) to be grouped by similarities in the elements' attributes. In this study, those elements are spatial units,
892 the elements' attributes are values from different climatic variables, and the measure of similarity is given by a
893 statistical distance. The four basic variables used as climatic attributes of these spatial elements are: near-surface (2m)
894 air temperature, seasonal 2m air temperature amplitude, precipitation rate, and seasonal precipitation rate amplitude.
895 Since monsoonal winds are a dominant feature of the climate in the South Asia region, near surface (10m) speeds of u-
896 wind and v-wind (zonal and meridional wind components, respectively) during the monsoon season (July) and outside
897 the monsoon season (January) are included as additional variables in our analysis of that region. Similarly, u-wind and
898 v-wind speeds during (January) and outside (July) the monsoon season in South America are added to the list of
899 considered variables to take into account the South American Monsoon System (SASM) in the cluster analysis for this
900 region. The long-term monthly means of those variables are used in a hierarchical clustering method, followed by a
901 non-hierarchical k-means correction with randomized re-groupment [Mutz et al., 2016; Wilks, 2011; Paeth, 2004;
902 Bahrenberg et al., 1992].

903 The hierarchical part of the clustering procedure starts with as many clusters as there are elements (ni), then
904 iteratively combines the most similar clusters to form a new cluster using centroids for the linkage procedure for
905 clusters containing multiple elements. The procedure is continued until the desired number of clusters (k) is reached.
906 One disadvantage of a pure hierarchical approach is that elements cannot be re-categorized once they are assigned to a
907 cluster, even though the addition of new elements to existing clusters changes the clusters' defining attributes and could
908 warrant a re-categorization of elements. We address this problem by implementation of a (non-hierarchical) k-means

909 clustering correction [e.g. Paeth, 2004]. Elements are re-categorized based on the multivariate centroids determined by
910 the hierarchical cluster analysis in order to minimize the sum of deviations from the cluster centroids. The
911 Mahalanobis distance [e.g. Wilks, 2011] is used as a measure of similarity or distance between the cluster centroids,
912 since it is a statistical distance and thus not sensitive to different variable units. The Mahalanobis distance also accounts
913 for possible multi-collinearity between variables.

914 The end results of the cluster analyses are subdivisions of the climate in the investigated regions into k
915 subdomains or clusters based on multiple climate variables. The region-specific k has to be prescribed before the
916 analyses. A large k may result in redundant additional clusters describing very similar climates, thereby defeating the
917 purpose of the analysis to identify and describe the dominant, distinctly different climates in the region and their
918 geographical coverage. Since it is not possible to know a priori the ideal number of clusters, k was varied between 3 and
919 10 for each region and the results presented below identify the optimal number of visibly distinctly different clusters
920 from the analysis. Optimal k was determined by assessing the distinctiveness and similarities between the climate
921 clusters in the systematic process of increasing k from 3 to 10. Once an increase in k no longer resulted in the addition
922 of another cluster that was climatologically distinctly different from the others, and instead resulted in at least two
923 similar clusters, k of the previous iteration was chosen as the optimal k for the region.

924 The cluster analysis ultimately results in a description of the geographical extent of a climate (cluster)
925 characterised by a certain combination of mean values for each of the variables associated with the climate. For
926 example, climate cluster 1 may be the most tropical climate in a region and thus be characterised by a high precipitation
927 values, high temperature values and low seasonal temperature amplitude. Each of the results (consisting of the
928 geographical extent of climates and mean vectors describing the climate) can be viewed as an optimal classification for
929 the specific region and time. It serves primarily as a means for providing an overview of the climate in each of the
930 regions at different times, reduces dimensionality of the raw simulation output, and identify regions of climatic
931 homogeneity that is difficult to notice by viewing simple maps of each climate variable. Its synoptic purpose is similar
932 to that of the widely known Köppen-Geiger classification scheme [Peel et al., 2007], but we allow for optimal
933 classification rather than prescribe classes, and our selection of variables is more restricted and made in accordance with
934 the focus of this study.

935

936 **3. Results**

937 Results from our analysis are first presented for general changes in global temperature and precipitation for the
938 different time slices (Fig. 2, 3), which is then followed by an analysis of changes in the climatology of selected orogens.
939 A more detailed description of temperature and precipitation changes in our selected orogens is presented in subsequent
940 subsections (Fig. 4 and following). All differences in climatology are expressed relative to the PI control run. Changes

941 relative to the PI rather than PD conditions are presented to avoid interpreting an anthropogenic bias in the results and
942 focusing instead on pre-anthropogenic variations in climate. For brevity, near-surface (2m) air temperature and total
943 precipitation rate are referred to as temperature and precipitation.

944

945 **3.1 Global differences in mean annual temperature**

946 This section describes the differences between simulated MH, LGM, and PLIO annual mean temperature anom-
947 alies with respect to PI shown in Fig. 2b, and PI temperature absolute values shown in Fig. 2a. Most temperature differ-
948 ences between the PI and MH climate are within -1°C to 1°C . Exceptions to this are the Hudson Bay, Weddell Sea and
949 Ross Sea regions which experience warming of $1-3^{\circ}\text{C}$, $1-5^{\circ}\text{C}$ and $1-9^{\circ}\text{C}$ respectively. Continental warming is mostly
950 restricted to low-altitude South America, Finland, western Russia, the Arabian Peninsula ($1-3^{\circ}\text{C}$) and subtropical north
951 Africa ($1-5^{\circ}\text{C}$). Simulation results show that LGM and PLIO annual mean temperature deviate from the PI means the
952 most. The global PLIO warming and LGM cooling trends are mostly uniform in direction, but the magnitude varies re-
953 gionally. The strongest LGM cooling is concentrated in [regions where the greatest change in ice extent occurs \(as indic-](#)
954 [ated on Fig. 2\), i.e.](#) Canada, Greenland, the North Atlantic, Northern Europe and Antarctica. Central Alaska shows no
955 temperature changes, whereas coastal South Alaska experiences cooling of $\leq 9^{\circ}\text{C}$. Cooling in the US Pacific northwest
956 is uniform and between 11 and 13°C . Most of high-altitude South America experiences mild cooling of $1-3^{\circ}\text{C}$, $3-5^{\circ}\text{C}$ in
957 the central Andes and $\leq 9^{\circ}\text{C}$ in the south. Along the Himalayan orogen, LGM temperature values are $5-7^{\circ}\text{C}$ below PI
958 values. Much of central Asia and the Tibetan plateau cools by $3-5^{\circ}\text{C}$, and most of India, low-altitude China and south-
959 east Asia by $1-3^{\circ}\text{C}$.

960 In the PLIO climate, parts of Antarctica, Greenland and the Greenland Sea experience the greatest temperature
961 increase ($\leq 19^{\circ}\text{C}$). Most of southern Alaska warms by $1-5^{\circ}\text{C}$ and $\leq 9^{\circ}\text{C}$ near McCarthy, Alaska. The US Pacific northw-
962 est warms by $1-5^{\circ}\text{C}$. The strongest warming in South America is concentrated at the Pacific west coast and the Andes
963 ($1-9^{\circ}\text{C}$), specifically between Lima and Chiclayo, and along the Chilean-Argentinian Andes south of Bolivia ($\leq 9^{\circ}\text{C}$).
964 Parts of low-altitude South America to the immediate east of the Andes experience cooling of $1-5^{\circ}\text{C}$. The Himalayan
965 orogen warms by $3-9^{\circ}\text{C}$, whereas Myanmar, Bangladesh, Nepal, northern India and northeast Pakistan cool by $1-9^{\circ}\text{C}$.

966

967 **3.2 Global differences in mean annual precipitation**

968 Notable differences occur between simulated MH, LGM, PLIO annual mean precipitation anomalies with re-
969 spect to PI shown in Fig. 3b, and the PI precipitation absolute values shown in Fig. 3a. Of these, MH precipitation devi-
970 ates the least from PI values. The differences between MH and PI precipitation on land appear to be largest in northern
971 tropical Africa (increase ≤ 1200 mm/a) and along the Himalayan orogen (increase ≤ 2000 mm/a) and in central Indian
972 states (decrease) ≤ 500 mm. The biggest differences in western South America are precipitation increases in central Chile

973 between Santiago and Puerto Montt. The LGM climate shows the largest deviation in annual precipitation from the PI
974 climate, and precipitation on land mostly decreases. Exceptions are increases in precipitation rates in North American
975 coastal regions, especially in coastal South Alaska (≤ 2300 mm/a) and the US Pacific Northwest (≤ 1700 mm/a). Further
976 exceptions are precipitation increases in low-altitude regions immediately east of the Peruvian Andes (≤ 1800 mm/a),
977 central Bolivia (≤ 1000 mm/a), most of Chile (≤ 1000 mm/a) and northeast India (≤ 1900 mm/a). Regions of notable pre-
978 cipitation decrease are northern Brazil (≤ 1700 mm/a), southernmost Chile and Argentina (≤ 1900 mm/a), coastal south
979 Peru (≤ 700 mm/a), central India (≤ 2300 mm/a) and Nepal (≤ 1600 mm/a).

980 Most of the precipitation on land in the PLIO climate is higher than those in the PI climate. Precipitation is en-
981 hanced by ca. 100-200 mm/a in most of the Atacama desert, by ≤ 1700 mm/a south of the Himalayan orogen and by
982 ≤ 1400 mm/a in tropical South America. Precipitation significantly decreases in central Peru (≤ 2600 mm), southernmost
983 Chile (≤ 2600 mm) and from eastern Nepal to northernmost northeast India (≤ 2500 mm).

984

985 | 3.3 Palaeoclimate characterization from the cluster analysis and changes in regional climatology

986 In addition to the above described global changes, the PLIO to PI regional climatology changes substantially in
987 the four investigated regions of: South Asia (section 3.3.1), the Andes (section 3.3.2), South Alaska (section 3.3.3) and
988 the Cascade Range (section 3.3.4). Each climate cluster defines separate distinct climate that is characterized by the
989 mean values of the different climate variables used in the analysis. The clusters are calculated by taking the arithmetic
990 means of all the values (climatic means) calculated for the grid boxes within each region. The regional climates are
991 referred to by their cluster number C_1, C_2, \dots, C_k , where k is the number of clusters specified for the region. The clusters
992 for specific palaeo-climates are mentioned in the text as $C_{i(t)}$, where i corresponds to the cluster number ($i=1, \dots, k$) and
993 t to the simulation time period ($t=PI, MH, LGM, PLIO$). The descriptions first highlight the similarities and then the
994 differences in regional climate. The cluster means of seasonal near-surface temperature amplitude and seasonal
995 precipitation amplitude are referred to as temperature and precipitation amplitude. The median, 25th percentile, 75th
996 percentile, minimum and maximum values for annual mean precipitation are referred to as $P_{md}, P_{25}, P_{75}, P_{min}$ and P_{max}
997 respectively. Likewise, the same statistics for temperature are referred to as $T_{md}, T_{25}, T_{75}, T_{min}$ and T_{max} . These are
998 presented as boxplots of climate variables in different time periods. When the character of a climate cluster is described
999 as “high”, “moderate” and “low”, the climatic attribute’s values are described relative to the value range of the specific
1000 region in time, thus high PLIO precipitation rates may be higher than high LGM precipitation rates. The character is
1001 presented a raster plots, to allow compact visual representation of it. The actual mean values for each variable in every
1002 time-slice and region-specific cluster are included in tables in the supplementary material.

1003

1004 | 3.3.1 Climate change and palaeoclimate characterization in South Asia, Central- and East Asia

1005 This section describes the regional climatology of the four investigated Cenozoic time slices and how
1006 precipitation and temperature changes from PLIO to PI times in tropical, temperate and high altitude regions. LGM and
1007 PLIO simulations show the largest simulated temperature and precipitation deviations (Fig. 4b) from PI temperature and
1008 precipitation (Fig. 4a) in the South Asia region. LGM temperatures are 1-7°C below PI temperatures and the direction
1009 of deviation is uniform across the study region. PLIO temperature is mostly above PI temperatures by 1-7°C. The
1010 cooling of 3-5°C in the region immediately south of the Himalayan orogen represents one of the few exceptions.
1011 Deviations of MH precipitation from PI precipitation in the region are greatest along the eastern Himalayan orogeny,
1012 which experiences an increase in precipitation (≤ 2000 mm/a). The same region experiences a notable decrease in
1013 precipitation in the LGM simulation, which is consistent in direction with the prevailing precipitation trend on land
1014 during the LGM. PLIO precipitation on land is typically higher than PI precipitation.

1015 Annual means of precipitation and temperature spatially averaged for the regional subdivisions and the different
1016 time slice simulations have been compared. The value range P_{25} to P_{75} of precipitation is higher for tropical South Asia
1017 than for temperate and high altitude South Asia (Fig. 5 a-c). The LGM values for P_{25} , P_{md} and P_{75} are lower than for the
1018 other time slice simulations, most visibly for tropical South Asia (ca. 100 mm/a). The temperature range (both $T_{75}-T_{25}$
1019 and $T_{max}-T_{min}$) is smallest in the hot (ca. 21°C) tropical South Asia, wider in the high altitude (ca. -8°C) South Asia, and
1020 widest in the temperate (ca. 2°C) South Asia region (Fig. 5 d-f). T_{md} , T_{25} and T_{75} values for the LGM are ca. 1°C, 1-2°C
1021 and 2°C below PI and MH temperatures in tropical, temperate and high altitude South Asia respectively, whereas the
1022 same temperature statistics for the PLIO simulation are ca. 1°C above PI and MH values in all regional subdivisions
1023 (Fig. 5 d-f). With respect to PI and MH values, precipitation and temperature are generally lower in the LGM and
1024 higher in the PLIO in tropical, temperate and high altitude South Asia.

1025 In all time periods, the wettest climate cluster C_1 covers an area along the southeastern Himalayan orogen (Fig. 6
1026 a-d) and is defined by the highest precipitation amplitude (dark blue, Fig. 6 e-h). $C_{5(PI)}$, $C_{3(MH)}$, $C_{4(LGM)}$ and $C_{5(PLIO)}$ are
1027 characterized by (dark blue, Fig. 6e-h) the highest temperatures, u-wind and v-wind speeds during the summer monsoon
1028 in their respective time periods, whereas $C_{4(PI)}$, $C_{5(MH)}$, and $C_{6(LGM)}$ are defined by low temperatures and highest
1029 temperature amplitude, u-wind and v-wind speeds outside the monsoon season (in January) in their respective time
1030 periods (Fig. 6 e-h). The latter 3 climate classes cover much of the more continental, northern landmass in their
1031 respective time periods and represents a cooler climate affected more by seasonal temperature fluctuations (Fig. 6 a-d).
1032 The two wettest climate clusters C_1 and C_2 are more restricted to the eastern end of the Himalayan orogen in the LGM
1033 than during other times, indicating that the LGM precipitation distribution over the South Asia landmass is more
1034 concentrated in this region than in other time slice experiments.

1035
1036 | **3.3.2 Climate change and palaeoclimate characterization in the Andes, Western South America**

1037 This section describes the cluster analysis based regional climatology of the four investigated Late Cenozoic
1038 time slices and illustrates how precipitation and temperature changes from PLIO to PI in tropical and temperate low-
1039 and high altitude (i.e. Andes) regions in western South America (Fig. 7-9).

1040 LGM and PLIO simulations show the largest simulated deviations (Fig. 7b) from PI temperature and
1041 precipitation (Fig. 7a) in western South America. The direction of LGM temperature deviations from PI temperatures is
1042 negative and uniform across the region. LGM temperatures are typically 1-3°C below PI temperatures across the region,
1043 and 1-7°C below PI values in the Peruvian Andes, which also experience the strongest and most widespread increase in
1044 precipitation during the LGM (≤ 1800 mm/a). Other regions, such as much of the northern Andes and tropical South
1045 America, experience a decrease of precipitation in the same experiment. PLIO temperature is mostly elevated above PI
1046 temperatures by 1-5°C. The Peruvian Andes experience a decrease in precipitation (≤ 2600 mm), while the northern
1047 Andes are wetter in the PLIO simulation compared to the PI control simulation.

1048 PI, MH, LGM and PLIO precipitation and temperature means for regional subdivisions have been compared.
1049 The P_{25} to P_{75} range is smallest for the relatively dry temperate Andes and largest for tropical South America and the
1050 tropical Andes (Fig. 8 a-d). P_{max} is lowest in the PLIO in all four regional subdivisions even though P_{md} , P_{25} and P_{75} in
1051 the PLIO simulation are similar to the same statistics calculated for PI and MH time slices. P_{md} , P_{25} and P_{75} for the LGM
1052 are ca. 50 mm/a lower in tropical South America and ca. 50 mm/a higher in the temperate Andes. Average PLIO
1053 temperatures are slightly warmer and LGM temperatures are slightly colder than PI and MH temperatures in tropical
1054 and temperate South America (Fig. 8 e and f). These differences are more pronounced in the Andes, however. T_{md} , T_{25}
1055 and T_{75} are ca. 5°C higher in the PLIO climate than in PI and MH climates in both temperate and tropical Andes,
1056 whereas the same temperatures for the LGM are ca. 2-4°C below PI and MH values (Fig. 8 g and h).

1057 For the LGM, the model computes drier-than-PI conditions in tropical South America and tropical Andes,
1058 enhanced precipitation in the temperate Andes, and a decrease in temperature that is most pronounced in the Andes. For
1059 the PLIO, the model predicts precipitation similar to PI, but with lower precipitation maxima. PLIO temperatures
1060 generally increase from PI temperatures, and this increase is most pronounced in the Andes.

1061 The climate variability in the region is described by six different clusters (Fig. 9 a-d), which have similar
1062 attributes in all time periods. The wettest climate C_1 is also defined by moderate to high precipitation amplitudes, low
1063 temperatures and moderate to high u-wind speeds in summer and winter in all time periods (dark blue, Fig. 9 e-h). $C_{2(PI)}$,
1064 $C_{2(MH)}$, $C_{3(LGM)}$ and $C_{2(PLIO)}$ are characterized by high temperatures and low seasonal temperature amplitude (dark blue,
1065 Fig. 9 e-h), geographically cover the north of the investigated region, and represent a more tropical climate. $C_{5(PI)}$,
1066 $C_{5(MH)}$, $C_{6(LGM)}$ and $C_{6(PLIO)}$ are defined by low precipitation and precipitation amplitude, high temperature amplitude and
1067 high u-wind speeds in winter (Fig. 9 e-h), cover the low-altitude south of the investigated region (Fig. 9 a-d) and
1068 represent dry, extra-tropical climates with more pronounced seasonality. In the PLIO simulation, the lower-altitude east

1069 of the region has four distinct climates, whereas the analysis for the other time slice experiments only yield three
1070 distinct climates for the same region.

1071

1072 | 3.3.3 Climate change and palaeoclimate characterization in the St. Elias Range, Southeast Alaska

1073 This section describes the changes in climate and the results from the cluster analysis for South Alaska (Fig. 10-12). As
1074 is the case for the other study areas, LGM and PLIO simulations show the largest simulated deviations (Fig. 10b) from
1075 PI temperature and precipitation (Fig. 10a). The sign of LGM temperature deviations from PI temperatures is negative
1076 and uniform across the region. LGM temperatures are typically 1-9°C below PI temperatures, with the east of the study
1077 area experiencing largest cooling. PLIO temperatures are typically 1-5°C above PI temperatures and the warming is
1078 uniform for the region. In comparison to the PI simulation, LGM precipitation is lower on land, but higher ($\leq 2300\text{mm}$)
1079 in much of the coastal regions of South Alaska. Annual PLIO precipitation is mostly higher ($\leq 800\text{mm}$) than for PI.

1080 P_{md} , P_{25} , P_{75} , P_{min} and P_{max} for South Alaskan mean annual precipitation do not differ much between PI, MH and
1081 PLIO climates, while P_{md} , P_{25} , P_{75} and P_{min} decrease by ca. 20-40 mm/a and P_{max} increases during the LGM (Fig. 11a).
1082 The Alaskan PLIO climate is distinguished from the PI and MH climates by its higher (ca. 2°C) regional temperature
1083 means, T_{25} , T_{75} and T_{md} (Fig. 11b). Mean annual temperatures, T_{25} , T_{75} , T_{min} and T_{max} are lower in the LGM than in any
1084 other considered time period (Fig. 11b), and about 3-5°C lower than during the PI and MH.

1085 Distinct climates are present in the PLIO to PI simulations for Southeast Alaska. Climate cluster C_1 is always
1086 geographically restricted to coastal southeast Alaska (Fig. 12 a-d) and characterized by the highest precipitation,
1087 precipitation amplitude, temperature, and by relatively low temperature amplitude (dark blue, Fig. 12 e-h). Climate C_2 is
1088 characterized by moderate to low precipitation, precipitation amplitude, temperature, and by low temperature amplitude.
1089 C_2 is either restricted to coastal southeast Alaska (in MH and LGM climates) or coastal southern Alaska (in PI and
1090 PLIO climates). Climate C_3 is described by low precipitation, precipitation amplitude, temperature, and moderate
1091 temperature amplitude in all simulations. It covers coastal western Alaska and separates climate C_1 and C_2 from the
1092 northern C_4 climate. Climate C_4 is distinguished by the highest mean temperature amplitude, by low temperature and
1093 precipitation amplitude, and by lowest precipitation.

1094 The geographical ranges of PI climates C_1 - C_4 and PLIO climates C_1 - C_4 are similar. $C_{1(\text{PI/PLIO})}$ and $C_{2(\text{PI/PLIO})}$ spread
1095 over a larger area than $C_{1(\text{MH/LGM})}$ and $C_{2(\text{MH/LGM})}$. $C_{2(\text{PI/PLIO})}$ are not restricted to coastal southeast Alaska, but also cover the
1096 coastal southwest of Alaska. The main difference in characterization between PI and PLIO climates C_1 - C_4 lies in the
1097 greater difference (towards lower values) in precipitation, precipitation amplitude and temperature from $C_{1(\text{PLIO})}$ to
1098 $C_{2(\text{PLIO})}$ compared to the relatively moderate decrease in those means from $C_{1(\text{PI})}$ to $C_{2(\text{PI})}$.

1099

1100 | 3.3.4 Climate change and palaeoclimate characterization in the Cascade Range, US Pacific Northwest

1101 This section describes the character of regional climatology in the US Pacific Northwest and its change over time
1102 (Fig. 13-15). The region experiences cooling of typically 9-11°C on land during the LGM, and warming of 1-5°C
1103 during the PLIO (Fig. 13b) when compared to PI temperatures (Fig. 13a). LGM precipitation increases over water,
1104 decreases on land by ≤ 800 mm/a in the North and in the vicinity of Seattle and increases on land by ≤ 1400 mm/a on
1105 Vancouver Island, around Portland and the Olympic Mountains, whereas PLIO precipitation does not deviate much
1106 from PI values over water and varies in the direction of deviation on land. MH temperature and precipitation deviation
1107 from PI values are negligible.

1108 P_{md} , P_{25} , P_{75} , P_{min} and P_{max} for the Cascade Range do not notably differ between the four time periods (Fig. 14a).
1109 The LGM range of precipitation values is slightly larger than that of the PI and MH with slightly increased P_{md} , while
1110 the respective range is smaller for simulation PLIO. The T_{md} , T_{25} , T_{75} and T_{max} values for the PLIO climate are ca. 2°C
1111 higher than those values for PI and MH (Fig. 14b). All temperature statistics for the LGM are notably (ca. 13°C) below
1112 their analogues in the other time periods (Fig. 14b).

1113 PI, LGM and PLIO clusters are similar in both their geographical patterns (Fig. 15 a, c, d) and their
1114 characterization by mean values (Fig. 15 e, g, h). C_1 is the wettest cluster and shows the highest amplitude in
1115 precipitation. The common characteristics of the C_2 cluster are moderate to high precipitation and precipitation
1116 amplitude. C_4 is characterized by the lowest precipitation and precipitation amplitudes, and the highest temperature
1117 amplitudes. Regions assigned to clusters C_1 and C_2 are in proximity to the coast, whereas C_4 is geographically restricted
1118 to more continental settings.

1119 In the PI and LGM climates, the wettest cluster C_1 is also characterized by high temperatures (Fig 10 e, g).
1120 However, virtually no grid boxes were assigned to $C_{1(LGM)}$. $C_{1(MH)}$ differs from other climate state's C_1 clusters in that it is
1121 also described by moderate to high near surface temperature and temperature amplitude (Fig 10 f), and in that it is
1122 geographically less restricted and, covering much of Vancouver Island and the continental coastline north of it (Fig 10
1123 b). Near surface temperatures are highest for C_2 in PI, LGM and PLIO climates (Fig 10 e, g, h) and low for $C_{2(MH)}$ (Fig
1124 10 f). $C_{2(MH)}$ is also geographically more restricted than C_2 clusters in PI, LGM and PLIO climates (Fig 10 a-d). $C_{2(PI)}$,
1125 $C_{2(MH)}$ and $C_{2(LGM)}$ have a low temperature amplitude (Fig 10 e-g), whereas $C_{2(PLIO)}$ is characterized by a moderate
1126 temperature amplitude (Fig 10 h).

1127

1128 4. Discussion

1129 In the following, we synthesize our results and compare to previous studies that investigate the effects of
1130 temperature and precipitation change on erosion. [Since our results do not warrant merited discussion of subglacial](#)
1131 [processes without additional work that is beyond the scope of this study, we instead advise caution in interpreting the](#)
1132 [presented precipitation and temperature results in an erosional context where the regions are covered with ice. For](#)

1133 [convenience, ice cover is indicated on figures 2,3,4,7,10 and 13, and a summary of ice cover used as boundary](#)
1134 [conditions for the different time slice experiments is included in the supplemental material.](#) Where possible, we relate
1135 the magnitude of climate change predicted in each geographical study area with terrestrial proxy data.

1136

1137 **4.1 Synthesis of temperature changes and implications for weathering and erosion**

1138

1139 **4.1.1 Temperature changes and implications for weathering and erosion**

1140 Changes in temperature can affect physical weathering due to temperature-induced changes in periglacial
1141 processes and promote frost cracking and frost creep [e.g., Matsuoka, 2001; Schaller et al., 2002; Matsuoka and
1142 Murton, 2008; Delunel et al., 2010; [Andersen et al., 2015](#); Marshall et al., 2015], and also biotic weathering and erosion
1143 [e.g. Moulton et al., 1998; Banfield et al., 1999; Dietrich and Perron, 2006]. Quantifying and understanding past
1144 changes in temperature is thus vital for our understanding of denudation histories. In the following, we highlight regions
1145 in the world where future observational studies might be able to document significant warming or cooling that would
1146 influence temperature related changes in physical and chemical weathering over the last ~3 Ma.

1147 Simulated MH temperatures show little deviation (typically < 1°C) from PI temperatures in the investigated
1148 regions (Fig. 2b), suggesting little difference in MH temperature-related weathering. [The LGM experiences widespread](#)
1149 [cooling, which is accentuated at the poles. LGM cooling is accentuated at the poles, in general agreement with studies](#)
1150 [such as Otto-Bliesner et al. \[2006\] and Braconnot et al. \[2007\], and , increases in](#) the equator-to-pole pressure gradient
1151 and consequently strengthens global atmospheric circulation. Despite this global trend, cooling in coastal South Alaska
1152 is higher ($\leq 9^\circ\text{C}$) than in central Alaska ($0\pm 1^\circ\text{C}$). [The larger temperature difference in South Alaska geographically](#)
1153 [coincides with ice cover \(Fig. 10b\), and should thus be interpreted in context of a different erosional regime.](#) Cooling in
1154 most of the lower-latitude regions in South America and central to southeast Asia is relatively mild. [The greatest](#)
1155 [temperature differences in South America are observed for western Patagonia, which was mostly covered by glaciers.](#)
1156 The Tibetan plateau experiences more cooling ($3\text{-}5^\circ\text{C}$) than adjacent low-altitude regions ($1\text{-}3^\circ\text{C}$) during the LGM.

1157 The PLIO simulation [is generally warmer, and temperature differences are shows little to no warming in the](#)
1158 [tropics and accentuated warming at the poles, as do findings of Salzmann et al. \[2011\] and Robinson \[2009\] and](#)
1159 [Ballantyne \[2010\] respectively. This would reduce the equator-to-pole sea and land surface temperature gradient, as](#)
1160 [also reported by Dowsett et al. \[2010\], and also weaken global atmospheric circulation. Agreement with proxy-based](#)
1161 [reconstructions, as is the case of the relatively little warming in lower latitudes, is not surprising given that sea surface](#)
1162 [temperature reconstructions are prescribed in this uncoupled atmosphere simulation. It should be noted that coupled](#)
1163 [ocean-atmosphere simulations do predict more low-latitude warming \[e.g. Stepanek and Lohmann 2012; Zhang et al.](#)
1164 [2013b\].](#) Warming in simulation PLIO is [present](#) greatest in parts of Canada, [and](#) Greenland and Antarctica (up to

1165 19°C), [which geographically coincides with the presence of ice in the PI reference simulation and thus may be](#)
1166 [attributed to differences in ice cover. It should therefore also be regarded as areas in which process domain shifted from](#)
1167 [glacial to non-glacial, and consistent with values based on multi-proxy studies \[Ballantyne et al., 2010\]. Due to a](#)
1168 [scarcity of paleo-botanical proxies in Antarctica, reconstruction-based temperature and ice-sheet extent estimates for a](#)
1169 [PLIO climate have high uncertainties \[Salzmann et al., 2011\], making model validation difficult. Furthermore,](#)
1170 [controversy about relatively little warming in the south polar regions compared to the north polar regions remains \[e.g.](#)
1171 [Hillenbrand and Fütterer, 2002; Wilson et al., 2002\]. Mid-latitude PLIO warming is mostly in the 1-3°C range with](#)
1172 [notable exceptions of cooling in the northern tropics of Africa and on the Indian subcontinent, especially south of the](#)
1173 [Himalayan orogen. The warming in simulation PLIO in South Alaska and the US Pacific northwest is mostly uniform](#)
1174 [and in the range of 1-5°C. As before, changes in ice cover reveal that the greatest warming may be associated with the](#)
1175 [absence of glaciers relative to the PI simulation, whereas](#) warming in South America is concentrated at the Pacific
1176 west coast and the Andes between Lima and Chiclayo, and along the Chilean-Argentinian Andes south of Bolivia (\leq
1177 9°C).

1178 Overall, annual mean temperatures in the MH simulation show little deviation from PI values. The more
1179 significant temperature deviations of the colder LGM and of the warmer PLIO simulations are accentuated at the poles
1180 leading to higher and lower equator-to-pole temperature gradients respectively. The largest temperature-related changes
1181 (relative to PI conditions) in weathering and subsequent erosion, [in many cases through a shift in the process domain](#)
1182 [from glacial to non-glacial or vice versa](#), are therefore to be expected in the LGM and PLIO climates.

1183

1184 **4.1.2 Temperature comparison to other studies**

1185 [LGM cooling is accentuated at the poles, thus increases the equator-to-pole pressure gradient and consequently](#)
1186 [strengthens global atmospheric circulation, and is in general agreement with studies such as Otto-Bliesner et al. \[2006\]](#)
1187 [and Braconnot et al. \[2007\]. The PLIO simulation shows little to no warming in the tropics and accentuated warming at](#)
1188 [the poles, as do findings of Salzmann et al. \[2011\] and Robinson \[2009\] and Ballantyne \[2010\] respectively. This would](#)
1189 [reduce the equator-to-pole sea and land surface temperature gradient, as also reported by Dowsett et al. \[2010\], and also](#)
1190 [weaken global atmospheric circulation. Agreement with proxy-based reconstructions, as is the case of the relatively](#)
1191 [little warming in lower latitudes, is not surprising given that sea surface temperature reconstructions \(derived from](#)
1192 [previous coarse resolution coupled ocean-atmosphere models\) are prescribed in this uncoupled atmosphere simulation.](#)
1193 [It should be noted that coupled ocean-atmosphere simulations do predict more low-latitude warming \[e.g. Stepanek and](#)
1194 [Lohmann 2012; Zhang et al. 2013b\]. The PLIO warming in parts of Canada and Greenland \(up to 19°C\) and consistent](#)
1195 [with values based on multi-proxy studies \[Ballantyne et al., 2010\]. Due to a scarcity of palaeobotanical proxies in](#)
1196 [Antarctica, reconstruction-based temperature and ice-sheet extent estimates for a PLIO climate have high uncertainties](#)

1197 | [\[Salzmann et al., 2011\]](#), making model validation difficult. Furthermore, controversy about relatively little warming in
1198 | [the south polar regions compared to the north polar regions remains \[e.g. Hillenbrand and Fütterer, 2002; Wilson et al.,](#)
1199 | [2002\]. Mid-latitude PLIO warming is mostly in the 1-3°C range with notable exceptions of cooling in the northern](#)
1200 | [tropics of Africa and on the Indian subcontinent, especially south of the Himalayan orogen.](#)

1201

1202 | **4.2 Synthesis of precipitation changes ~~and implications for orogen denudation~~**

1203

1204 | **[4.2.1 Precipitation and implications for weathering and erosion](#)**

1205 | Changes in precipitation affects erosion through river incision, sediment transport, and erosion due to extreme
1206 | precipitation events and storms [e.g. Whipple and Tucker, 1999; Hobbey et al., 2010]. Furthermore, vegetation type and
1207 | cover also co-evolve with variations in precipitation and with changes in geomorphology [e.g. Marston 2010; Roering
1208 | et al., 2010]. These vegetation changes in turn modify hillslope erosion by increasing root mass and canopy cover, and
1209 | decreasing water-induced erosion via surface runoff [e.g. Gyssels et al., 2005]. Therefore, understanding and
1210 | quantifying changes in precipitation in different palaeo-climates is necessary for a more complete reconstruction of
1211 | orogen denudation histories. A synthesis of predicted precipitation changes is provided below, and highlights regions
1212 | where changes in river discharge and hillslope processes might be impacted by climate change over the last ~3 Ma.

1213 | [Most of North Africa is notably wetter during the MH, which is characteristic of the African Humid Period](#)
1214 | [\[Sarnthein 1978\]. This pluvial regional expression of the Holocene Climatic Optimum is attributed to sudden changes in](#)
1215 | [the strength of the African monsoon caused by orbital-induced changes in summer insolation \[e.g. deMenocal et al.](#)
1216 | [2000\]. Southern Africa is characterised by a wetter climate to the east and drier climate to the west of the approximate](#)
1217 | [location of the Congo Air Boundary \(CAB\), the migration of which has previously been cited as a cause for](#)
1218 | [precipitation changes in East Africa \[e.g. Juninger et al. 2014\].](#) In contrast, simulated MH precipitation rates show little
1219 | deviation from the PI in most of the investigated regions, suggesting little difference in MH precipitation-related
1220 | erosion. The Himalayan orogen is an exception and shows a precipitation increase of [≤ up to 2000 mm/a](#). The climate's
1221 | enhanced erosion potential, that could result from such a climatic change, should be taken into consideration when
1222 | palaeo-erosion rates estimated from the geological record in this area are interpreted [e.g. Bookhagen et al., 2005].
1223 | Specifically, higher precipitation rates (along with differences in other rainfall-event parameters) could increase the
1224 | probability of mass movement events on hillslopes, especially where hillslopes are close to the angle of failure [e.g.
1225 | Montgomery, 2001], and modify fluxes to increase shear stresses exerted on river beds and increase stream capacity to
1226 | enhance erosion on river beds (e.g. by abrasion).

1227 | Most precipitation on land is decreased during the LGM due to large-scale cooling and decreased evaporation
1228 | over the tropics, resulting in an overall decrease in inland moisture transport [e.g., Braconnot et al. 2007]. [Coastal](#)-North

1229 America, south of the continental ice sheets, is an exception and experiences increases in precipitation. For example,
1230 the investigated US Pacific Northwest and the southeastern coast of Alaska are exceptions in that there is experience
1231 experience strongly enhanced precipitation of ≤ 1700 mm/a and ≤ 2300 mm/a, respectively. These changes

1232 geographically coincide with differences in ice extent. An increase in precipitation in these regions may have had direct
1233 consequences on the glaciers' mass balance and equilibrium line altitudes, where the glaciers' effectiveness in erosion is
1234 highest [e.g. Egholm et al., 2009; Yanites and Ehlers, 2012]. The differences in the direction of precipitation changes,
1235 and accompanying changes in ice cover. Reduced precipitation in other parts of southern Alaska result in a stronger
1236 south-to-north drying gradient than in the PI simulation. This could would likely result in more regionally differentiated
1237 variations in precipitation-specific erosional processes in the St. Elias Range rather than causing systematic offsets for
1238 the LGM. Although precipitation is significantly reduced along much of the Himalayan orogen (≤ 1600 mm/a), which is
1239 consistent with findings by, e.g., Braconnot et al. [2007], northeast India experiences strongly enhanced precipitation
1240 (≤ 1900 mm/a). This could have large implications for studies of uplift and erosion at orogen syntaxes, where highly
1241 localized and extreme denudation has been documented [e.g. Koons et al., 2013; Bendick and Ehlers, 2014].

1242 Overall, the PLIO climate is wetter than the PI climate, in particular in the (northern) mid-latitudes, and possibly
1243 related to a northward shift of the northern Hadley cell boundary that is ultimately the result of a reduced equator-to-
1244 pole temperature gradient [e.g. Haywood et al. 2000, 2013; Dowsett et al. 2010]. A reduction of this gradient by ca. 5°C
1245 is indeed present in the PLIO simulation of this study (Fig. 2b). Most of the PLIO precipitation over land increases
1246 during the PLIO. This finding agrees well with simulations performed at a lower spatial model resolution [cf. Stepanek
1247 and Lohmann, 2012]. PLIO precipitation significantly increases, esp. at the Himalayan orogen by ≤ 1400 mm/a, and
1248 decreases from eastern Nepal to Namcha Barwa (≤ 2500 mm/a). Most of the Atacama Desert experiences an increase in
1249 precipitation by 100-200 mm/a, which may have to be considered in erosion and uplift history reconstructions for the
1250 Andes. A significant increase (~ 2000 mm/a) in precipitation from simulation PLIO to modern conditions is simulated
1251 for the eastern margin of the Andean Plateau in Peru and for northern Bolivia. This is consistent with recent findings of
1252 a pulse of canyon incision in these locations in the last ~ 3 Ma [Lease and Ehlers, 2013].

1253 Overall, the simulated MH precipitation varies least from PI precipitation. The LGM is generally drier than the
1254 PI simulation, even though pockets of a wetter-than-PI climate do exist, such as much of coastal North America. Extra-
1255 tropical increased precipitation of the PLIO simulation and decreased precipitation of the LGM climate may be the
1256 result of decreased and increased equator-to-pole temperature gradients, respectively.

1257

1258 **4.2.2 Precipitation comparison to other studies**

1259 The large scale LGM precipitation decrease on land, related to cooling and decreased evaporation over the
1260 tropics, and greatly reduced precipitation along much of the Himalayan orogeny, is consistent with previous studies by,

1261 [\(for example\) Braconnot et al. \[2007\]. The large scale PLIO precipitation increase due to a reduced equator-to-pole](#)
1262 [temperature gradient, has previously been pointed out by e.g. Haywood et al. \[2000, 2013\] and Dowsett et al. \[2010\]. A](#)
1263 [reduction of this gradient by ca. 5°C is indeed present in the PLIO simulation of this study \(Fig. 2b\). This precipitation](#)
1264 [increase over land agrees well with simulations performed at a lower spatial model resolution \[cf. Stepanek and](#)
1265 [Lohmann, 2012\]. Section 4.4 includes a more in-depth discussion of how simulated MH and LGM precipitation](#)
1266 [differences compare with proxy-based reconstructions in South Asia and South America.](#)

1267

1268 **4.3 Trends in Late Cenozoic changes in regional climatology**

1269 This section describes the major changes in regional climatology and highlights their possible implications on
1270 erosion rates.

1271

1272 *Himalaya-Tibet, South Asia*

1273 In South Asia, cluster-analysis based categorization and description of climates (Fig. 6) remains similar
1274 throughout time. However, the two wettest climates (C₁ and C₂) are geographically more restricted to the eastern
1275 Himalayan orogen in the LGM simulation. Even though precipitation over the South Asia region is generally lower, this
1276 shift indicates that rainfall on land is more concentrated in this region and that the westward drying gradient along the
1277 orogen is more accentuated than during other time periods investigated here. While there is limited confidence in the
1278 global Atmospheric General Circulation Model's abilities to accurately represent meso-scale precipitation patterns [e.g.
1279 Cohen 1990], the simulation warrants careful consideration of possible, geographically non-uniform offsets in
1280 precipitation in investigations of denudation and uplift histories.

1281 MH precipitation and temperature in tropical, temperate and high-altitude South Asia is similar to PI
1282 precipitation and temperature, whereas LGM precipitation and temperatures are generally lower (by ca. 100 mm/a and
1283 1-2°C respectively), possibly reducing precipitation-driven erosion and enhancing frost-driven erosion in areas pushed
1284 into a near-zero temperature range during the LGM.

1285

1286 *Andes, South America*

1287 Clusters in South America (Fig. 9), which are somewhat reminiscent of the Köppen and Geiger classification
1288 [Kraus, 2001], remain mostly the same over the last 3 Ma. In the PLIO simulation, the lower-altitude east of the region
1289 is characterized by four distinct climates, which suggests enhanced latitudinal variability in the PLIO climate compared
1290 to PI with respect temperature and precipitation.

1291 The largest temperature deviations from PI values are derived for the PLIO simulation in the (tropical and
1292 temperate) Andes, where temperatures exceed PI values by 5°C. On the other hand, LGM temperatures in the Andes are

1293 ca. 2-4°C below PI values in the same region (Fig 7 g and h). In the LGM simulation, tropical South America
1294 experiences ca. 50 mm/a less precipitation, the temperate Andes receive ca. 50 mm/a more precipitation than in PI and
1295 MH simulations. These latitude-specific differences in precipitation changes ought to be considered in attempts to
1296 reconstruct precipitation-specific palaeo-erosion rates in the Andes on top of longitudinal climate gradients highlighted
1297 by, e.g., Montgomery et al. [2001].

1298

1299 *St. Elias Range, South Alaska*

1300 South Alaska is subdivided into two wetter and warmer clusters in the south, and two drier, colder clusters in the
1301 north. The latter are characterized by increased seasonal temperature variability due to being located at higher latitudes
1302 (Fig. 12). The different equator-to-pole temperature gradients for LGM and PLIO may affect the intensity of the Pacific
1303 North American Teleconnection (PNA) [Barnston and Livzey, 1987], which has significant influence on temperatures
1304 and precipitation, especially in southeast Alaska, and may in turn result in changes in regional precipitation and
1305 temperature patterns and thus on glacier mass balance. Changes in the Pacific Decadal Oscillation, which is related to
1306 the PNA pattern, has previously been connected to differences in Late Holocene precipitation [Barron and Anderson,
1307 2011]. While this climate cluster pattern appears to be a robust feature for the considered climate states, and hence over
1308 the recent geologic history, the LGM sets itself apart from PI and MH climates by generally lower precipitation (20-40
1309 mm) and lower temperatures (3-5°C, Fig. 10, 11), which may favour frost driven weathering during glacial climate
1310 states [e.g. Andersen et al., 2015; Marshall et al. 2015] in unglaciated areas, whereas glacial processes would have
1311 dominated most of this region as it was covered by ice. Simulation PLIO is distinguished by temperatures that exceed
1312 PI and MH conditions by ca. 2°C, and by larger temperature and precipitation value ranges, possibly modifying
1313 temperature- and precipitation-dependent erosional processes in the region of South Alaska.

1314

1315 *Cascade Range, US Pacific Northwest*

1316 In all time slices, the geographic climate patterns, based on the cluster analysis (Fig. 15), represents an increase
1317 in the degree of continentality from the wetter coastal climates to the further inland located climates with greater
1318 seasonal temperature amplitude and lower precipitation and precipitation amplitude (Fig 15 e-h). The most notable
1319 difference between the time slices is the strong cooling during the LGM, when temperatures are ca. 13°C (Fig. 13, 14)
1320 below those of other time periods. Given that the entire investigated region was covered by ice (Fig 13), we can assume
1321 a shift to glacially dominated processes. possibly leading to enhanced sediment production driven by frost processes, as
1322 proposed for parts of the Pacific Northwest by Marshall et al. [2015].

1323

1324 **4.4 Comparison of simulated and observed precipitation differences**

1325 The predicted precipitation differences reported in this study were compared with observed (proxy record)
1326 palaeoprecipitation change. Proxy based precipitation reconstructions for the MH and LGM are presented for South
1327 Asia and South America for the purpose of assessing ECHAM5 model performance, and for identifying inconsistencies
1328 between neighbouring proxy data. Due to the repeated glaciations, detailed terrestrial proxy records for the time slices
1329 investigated here are not available, to the best of our knowledge, for the Alaskan and Pacific NW USA studies.
1330 Although marine records and records of glacier extent are available in these regions, the results from them do not
1331 explicitly provide estimates of wetter/drier, or colder/warmer conditions that can be spatially compared to the
1332 simulation estimates. For these two areas with no available records, the ECHAM5 predicted results therefore provide
1333 predictions from which future studies can formulate testable hypotheses to evaluate.

1334 The palaeoclimate changes in terrestrial proxy records compiled here are reported as “wetter than today”, “drier
1335 than today” or “the same as today” for each of the study locations, and plotted on top of the simulation-based difference
1336 maps as upward facing blue triangles, downward facing red triangles and grey circles respectively (Fig. 16, 17). The
1337 numbers listed next to those indicators are the ID numbers assigned to the studies compiled for this comparison and are
1338 associated with a citation provided in the figure captions.

1339 In South Asia, 14/26 results from local studies agree with the model predicted precipitation changes for the MH.
1340 The model seems able to reproduce the predominantly wetter conditions on much of the Tibetan plateau, but predicts
1341 slightly drier conditions north of Chengdu, which is not reflected in local reconstructions. The modest mismatch
1342 between ECHAM5 predicted and proxy-based MH climate change in south Asia was also documented by Li et al.,
1343 [2017], whose simulations were conducted at a coarser (T106) resolution. Despite these model-proxy differences, we
1344 note that there are significant discrepancies between the proxy data themselves in neighbouring locations in the MH,
1345 highlighting caution in relying solely upon these data for regional palaeoclimate reconstructions. These differences
1346 could result from either poor age-constraints in the reported values, or systematic errors in the transfer functions used to
1347 convert proxy measurements to palaeoclimate conditions. The widespread drier conditions on the Tibetan Plateau and
1348 immediately north of Laos are confirmed by 7/7 of the palaeoprecipitation reconstructions. 23/39 of the reconstructed
1349 precipitation changes agree with model predictions for South America during the MH. The model predicted wetter
1350 conditions in the central Atacama desert, as well as the drier conditions northwest of Santiago are confirmed by most of
1351 the reconstructions. The wetter conditions in southernmost Peru and the border to Bolivia and Chile cannot be
1352 confirmed by local studies. 11/17 of the precipitation reconstructions for the LGM are in agreement with model
1353 predictions. These include wetter conditions in most of Chile. The most notable disagreement can be seen in northeast
1354 Chile at the border to Argentina and Bolivia, where model predicted wetter conditions are not confirmed by reported
1355 reconstructions from local sites.

1356 Model performance is, in general, higher for the LGM than for the MH and overall satisfactory given that it

1357 [cannot be expected to resolve sub-grid scale differences in reported palaeoprecipitation reconstructions. However, as](#)
1358 [mentioned above, it should be noted that some locations \(MH of south Asia, and MH of norther Chile\) discrepancies](#)
1359 [exist between neighbouring proxy samples and highlight the need for caution in how these data are interpreted. Other](#)
1360 [potential sources of error resulting in disagreement of simulated and proxy-based precipitation estimates are the model's](#)
1361 [shortcomings in simulating orographic precipitation at higher resolutions, and uncertainties in palaeoclimate](#)
1362 [reconstructions at the local sites. In summary, although some differences are evident in both the model-proxy data](#)
1363 [comparison and between neighbouring proxy data themselves, the above comparison highlights an overall good](#)
1364 [agreement between the model and data for the south Asia and South American study areas. Thus, although future](#)
1365 [advances in GCM model parameterisations and new or improved palaeoclimate proxy techniques are likely, the](#)
1366 [palaeoclimate changes documented here are found to be in general robust and provide a useful framework for future](#)
1367 [studies investigating how these predicted changes in palaeoclimate impact denudation.](#)

1368

1369 | **4.45 Conclusions**

1370 | We present a [statistical](#) cluster-analysis-based description of the geographic coverage of possible distinct
1371 regional expressions of climates from four different time slices (Fig. 6, 9, 12, 15). These are determined with respect to
1372 a selection of variables that characterize the climate of the region and may be relevant to weathering and erosional
1373 processes. While the geographic [distribution of](#) climate [patterns](#) remains similar throughout time (as indicated by results
1374 of four different climate states representative for the climate of the last 3 Ma), results for the PLIO simulation suggests
1375 more climatic variability east of the Andes (with respect to near-surface temperature, seasonal temperature amplitude,
1376 precipitation, seasonal precipitation amplitude and seasonal u-wind and v-wind speeds). Furthermore, the wetter
1377 climates in the South Asia region retreat eastward along the Himalayan orogen for the LGM simulation, this is due to
1378 decreased precipitation along the western part of the orogen and enhanced precipitation on the eastern end, possibly
1379 signifying more localised high erosion rates.

1380 Most global trends of the high-resolution LGM and PLIO simulations conducted here are in general agreement
1381 with previous studies [Otto-Bliesner et al., 2006; Braconnot et al., 2007; Wei and Lohmann, 2012; Lohmann et al.,
1382 2013; Zhang et al., 2013b, 2014; Stepanek and Lohmann, 2012]. The MH does not deviate notably from the PI, the
1383 LGM is relatively dry and cool, while the PLIO is comparably wet and warm. While the simulated regional changes in
1384 temperature and precipitation usually agree with the sign ([or direction](#)) of the simulated global changes, there are
1385 region-specific differences in the magnitude and direction. For example, the LGM precipitation of the Tropical Andes
1386 does not deviate significantly from PI precipitation, whereas LGM precipitation in the Temperate Andes is enhanced.

1387 | [Comparisons to local, proxy-based reconstructions of MH and LGM precipitation in South Asia and South](#)
1388 [America reveal satisfactory performance of the model in simulating the reported differences. The model performs better](#)

1389 [for the LGM than the MH. We note however that compilations of proxy data such as we present here, also identify](#)
1390 [inconsistencies between neighbouring proxy data themselves, warranting caution in the extent to which both proxy data](#)
1391 [and palaeoclimate models are interpreted for MH climate change in south Asia, and western South America.](#)

1392 The changes in regional climatology presented here are manifested, in part, by small to large magnitude changes
1393 in fluvial and hillslope relevant parameters such as precipitation and temperature. For the regions investigated here we
1394 find that precipitation differences between the PI, MH, LGM, and PLIO are in many areas around +/- 200-600 mm/yr,
1395 and locally can reach maximums of +/- 1000-2000 mm/yr (Figs. 4, 7, 10, 13). [In areas where significant precipitation](#)
1396 [increases are accompanied by changes in ice extent, such as parts of southern Alaska during the LGM, we would expect](#)
1397 [a shift in the erosional regime to glacier dominated processes.](#) Temperature differences between these same time periods
1398 are around 1-4 °C in many places, but reach maximum values of 8-10 °C. [Many of these maxima in the temperature](#)
1399 [differences geographically coincide with changes in ice sheet extent and must therefore be interpreted as part of a](#)
1400 [different erosional process domains. However, we also observe large temperature differences \(~5°C\) in unglaciated](#)
1401 [areas that would be affected by hillslope, frost cracking, and fluvial processes.](#) The magnitude of these differences are
1402 not trivial, and will likely impact fluvial and hillslope erosion and sediment transport, as well as biotic and abiotic
1403 weathering. The regions of large magnitude changes in precipitation and temperature documented here (Figs. 4, 7, 10,
1404 13) offer the highest potential for future observational studies interested in quantifying the impact of climate change on
1405 denudation and weathering rates.

1406

1407

1408 **Acknowledgements**

1409 [The model simulations presented in this study are freely available to interested persons by contacting S. Mutz or T.](#)
1410 [Ehlers. We note however that the data files are very large \(~4 TB, and too large to archive in journal supplementary](#)
1411 [material\) and require familiarity in reading/plotting NetCDF formatted files. Support from](#) European Research Council
1412 (ERC) Consolidator Grant number 615703 provided support for S. Mutz. Additional support is acknowledged from the
1413 German science foundation (DFG) priority research program 1803 (EarthShape: Earth Surface Shaping by Biota; grants
1414 EH329/14-1 and EH329/17-1). We thank B. Adams and J. Starke for constructive discussions. The DKRZ is thanked
1415 for computer time used for some of the simulations presented here. C. Stepanek, M. Werner, and G. Lohmann
1416 acknowledge funding by the Helmholtz Climate Initiative Reklim and the Alfred Wegener Institute's research
1417 programme Marine, Coastal and Polar Systems.

1418

1419

1420 **Figure Captions**

1421

1422 **Figure 1** Topography for regions (a) tropical South Asia, (b) temperate South Asia, (c) high altitude South Asia, (d)
1423 temperate South America, (e) tropical South America, (f) temperate Andes, (g) tropical Andes-, [SE Alaska and Cas-](#)
1424 [cadia](#).

1425 **Figure 2** Global PI annual mean near-surface temperatures (a), and deviations of MH, LGM and PLIO annual mean
1426 near-surface temperatures from PI values (b). Units are °C and insignificant ($p < 99\%$) differences (as determined by
1427 a t-test) are greyed out.

1428 **Figure 3** Global PI annual mean precipitation (a), and deviations of MH, LGM and PLIO annual mean near-surface
1429 temperatures from PI values (b). Units are mm/yr.

1430 **Figure 4** PI annual mean near-surface temperatures (a), and deviations of MH, LGM and PLIO annual mean near-sur-
1431 face temperatures from PI values (b) for the South Asia region. Insignificant ($p < 99\%$) differences (as determined
1432 by a t-test) are greyed out.

1433 **Figure 5** PI, MH, LGM and PLIO annual mean precipitation in (a) tropical South Asia, (b) temperate South Asia, and
1434 (c) high-altitude South Asia; PI, MH, LGM and PLIO annual mean temperatures in (d) tropical South Asia, (e) tem-
1435 perate South Asia, and (f) high-altitude South Asia. For each time slice, the minimum, lower 25th percentile, median,
1436 upper 75th percentile and maximum are plotted.

1437 **Figure 6** Geographical coverage and characterization of climate classes C_1 - C_6 based on cluster-analysis of 8 variables
1438 (near surface temperature, seasonal near surface temperature amplitude, total precipitation, seasonal precipitation
1439 amplitude, u-wind in January and July, v-wind in January and July) in the South Asia region. The geographical cov-
1440 erage of the climates C_1 - C_6 is shown on the left for PI (a), MH (b), LGM (c) and PLIO (d); the complementary,
1441 time-slice specific characterization of C_1 - C_6 for PI (e), MH (f), LGM (g) and PLIO (h) is shown on the right.

1442 **Figure 7** PI annual mean near-surface temperatures (a), and deviations of MH, LGM and PLIO annual mean near-sur-
1443 face temperatures from PI values (b) for western South America. Insignificant ($p < 99\%$) differences (as determined
1444 by a t-test) are greyed out.

1445 **Figure 8** PI, MH, LGM and PLIO annual mean precipitation in (a) tropical South America, (b) temperate South Amer-
1446 ica, (c) tropical Andes, and (d) temperate Andes; PI, MH, LGM and PLIO annual mean temperatures in (e) tropical

1447 South America, (f) temperate South America, (g) tropical Andes, and (h) temperate Andes. For each time slice, the
1448 minimum, lower 25th percentile, median, upper 75th percentile and maximum are plotted.

1449 **Figure 9** Geographical coverage and characterization of climate classes C₁- C₆ based on cluster-analysis of 8 variables
1450 (near surface temperature, seasonal near surface temperature amplitude, precipitation, seasonal precipitation amp-
1451 litude, u-wind in January and July, v-wind in January and July) in western South America. The geographical cover-
1452 age of the climates C₁- C₆ is shown on the left for PI (a), MH (b), LGM (c) and PLIO (d); the complementary, time-
1453 slice specific characterization of C₁- C₆ for PI (e), MH (f), LGM (g) and PLIO (h) is shown on the right.

1454 **Figure 10** PI annual mean near-surface temperatures (a), and deviations of MH, LGM and PLIO annual mean near-sur-
1455 face temperatures from PI values (b) for the South Alaska region. Insignificant (p < 99%) differences (as determined
1456 by a t-test) are greyed out.

1457 **Figure 11** PI, MH, LGM and PLIO annual mean precipitation (a), and mean annual temperatures (b) in South Alaska.
1458 For each time slice, the minimum, lower 25th percentile, median, upper 75th percentile and maximum are plotted.

1459 **Figure 12** Geographical coverage of climate classes C₁- C₄ based on cluster-analysis of 4 variables (near surface tem-
1460 perature, seasonal near surface temperature amplitude, total precipitation, seasonal total precipitation amplitude) in
1461 southern Alaska. The geographical coverage of the climates C₁- C₄ is shown on the left for PI (a), MH (b), LGM (c)
1462 and PLIO (d); the complementary, time-slice specific characterization of C₁- C₆ for PI (e), MH (f), LGM (g) and
1463 PLIO (h) is shown on the right.

1464 **Figure 13** PI annual mean near-surface temperatures (a), and deviations of MH, LGM and PLIO annual mean near-sur-
1465 face temperatures from PI values (b) for the US Pacific Northwest. Insignificant (p < 99%) differences (as determ-
1466 ined by a t-test) are greyed out.

1467 **Figure 14** PI, MH, LGM and PLIO annual mean precipitation (a), and annual mean temperatures (b) in the Cascades,
1468 US Pacific Northwest. For each time slice, the minimum, lower 25th percentile, median, upper 75th percentile and
1469 maximum are plotted.

1470 **Figure 15** Geographical coverage and characterization of climate classes C₁- C₄ based on cluster-analysis of 4 variables
1471 (near surface temperature, seasonal near surface temperature amplitude, total precipitation, seasonal total precipita-
1472 tion amplitude) in the Cascades, US Pacific Northwest. The geographical coverage of the climates C₁- C₄ is shown
1473 on the left for PI (a), MH (b), LGM (c) and PLIO (d); the complementary, time-slice specific characterization of C₁-

1474 C₆ for PI (e), MH (f), LGM (g) and PLIO (h) is shown on the right.

1475 **Figure 16** [Simulated annual mean precipitation deviations of MH \(left\) and LGM \(right\) from PI values in South Asia,](#)
1476 [and temporally corresponding proxy-based reconstructions, indicating wetter \(upward facing blue triangles\), drier](#)
1477 [\(downward facing red triangles\) or similar \(grey circles\) conditions in comparison with modern climate. MH proxy-](#)
1478 [based precipitation differences are taken from Mügler et al. \(2010\) \(66\), Wischnewski et al. \(2011\) \(67\), Mischke et](#)
1479 [al. \(2008\), Wischnewski et al. \(2011\), Herzsuh et al. \(2009\) \(68\), Yanhong et al. \(2006\) \(69\), Morrill et al. \(2006\)](#)
1480 [\(70\), Wang et al. \(2002\) \(71\), Wuennemann et al. \(2006\) \(72\), Zhang et al. \(2011\), Morinaga et al. \(1993\),](#)
1481 [Kashiwaya et al. \(1995\) \(73\), Shen et al. \(2005\) \(74\), Liu et al. \(2014\) \(75\), Herzsuh et al. \(2006\) \(76\), Zhang and](#)
1482 [Mischke \(2009\) \(77\), Nishimura et al. \(2014\) \(78\), Yu and Lai \(2014\) \(79\), Gasse et al. \(1991\) \(80\), Van Campo et](#)
1483 [al. \(1996\) \(81\), Demske et al. \(2009\) \(82\), Kramer et al. \(2010\) \(83\), Herzsuh et al. \(2006\) \(84\), Hodell et al.](#)
1484 [\(1999\)\(85\), Hodell et al. \(1999\) \(86\), Shen et al. \(2006\) \(87\), Tang et al. \(2000\) \(88\), Tang et al. \(2000\) \(89\), Zhou](#)
1485 [et al. \(2002\) \(90\), Liu et al. \(1998\) \(91\), Asashi \(2010\)\(92\), Kotila et al. \(2009\) \(93\), Kotila et al. \(2000\) \(94\), Wang](#)
1486 [et al. \(2002\) \(95\), Hu et al. \(2014\) \(96\), Hodell et al. \(1999\) \(97\), Hodell et al. \(1999\) \(98\).](#)

1487 **Figure 17** [Simulated annual mean precipitation deviations of MH \(left\) and LGM \(right\) from PI values in South Amer-](#)
1488 [ica, and temporally corresponding proxy-based reconstructions, indicating wetter \(upward facing blue triangles\),](#)
1489 [drier \(downward facing red triangles\) or similar \(grey circles\) conditions in comparison with modern climate. MH](#)
1490 [proxy-based precipitation differences are taken from Bird et al. \(2011\) \(1\), Hansen et al \(1994\) \(2\), Hansen et al](#)
1491 [\(1994\) \(3\), Hansen et al \(1994\) \(4\), Hansen et al \(1994\) \(5\), Hansen et al \(1994\) \(6\), Hillyer et al. \(2009\) \(7\),](#)
1492 [D’Agostino et al. \(2002\) \(8\), Baker et al. \(2001\) \(9\), Schwalb et al \(1999\) \(10\), Schwalb et al \(1999\) \(11\), Schwalb](#)
1493 [et al \(1999\) \(12\), Schwalb et al \(1999\) \(13\), Moreno et al \(2009\) \(14\), Pueyo et al \(2011\) \(15\), Mujica et al \(2015\)](#)
1494 [\(16\), Fritz et al. \(2004\) \(17\), Gayo et al. \(2012\) \(18\), Latorre et al. \(2006\) \(19\), Latorre et al. \(2003\) \(20\), Quade et al](#)
1495 [\(2008\) \(21\), Bobst et al. \(2001\) \(22\), Grosjean et al. \(2001\) \(23\), Betancourt et al. \(2000\) \(24\), Latorre et al. \(2002\)](#)
1496 [\(25\), Rech et al. \(2003\) \(26\), Diaz et al. \(2012\) \(27\), Maldonado et al \(2005\) \(28\), Diaz et al. \(2012\) \(29\), Lamy et](#)
1497 [al. \(2000\) \(30\), Kaiser et al. \(2008\) \(31\), Maldonado et al. \(2010\) \(32\), Villagrán et al. \(1990\) \(33\), Méndez et al.](#)
1498 [\(2015\) \(34\), Maldonado et al. \(2006\) \(35\), Lamy et al. \(1999\) \(36\), Jenny et al. \(2002\) \(37\), Jenny et al. \(2002b\)](#)
1499 [\(38\), Villa-Martínez et al. \(2003\) \(39\), Bertrand et al. \(2008\) \(40\), De Basti et al. \(2008\) \(41\), Lamy et al. \(2009\)](#)
1500 [\(42\), Lamy et al. \(2002\) \(43\), Szeicz et al. \(2003\) \(44\), de Porras et al. \(2012\) \(45\), de Porras et al. \(2014\) \(46\),](#)
1501 [Markgraf et al. \(2007\) \(47\), Siani et al. \(2010\) \(48\), Gilli et al. \(2001\) \(49\), Markgraf et al. \(2003\) \(50\), Stine et al.](#)
1502 [\(1990\) \(51\).](#)

1503

1504

1505 **References**

1506

1507 Abe-Ouchi, A., Saito, F., Kageyama, M., Bracannot, P., Harrison, S. P., Lambeck, K., Otto-Bliesner, B. L., Peltier,
1508 W.R., Tarasov, L., Peterschmitt, J.-Y., Takahashi, K.: Ice-sheet configuration in the CMUP5/PMIP3 Last Glacial
1509 Maximum experiments. *Geosci. Model Dev.*, 8, 3621-3637, doi:10.5194/gmd-8-3621-2015, 2015

1510 [Andersen, J., Egholm, D.L., Knudsen, M.F., Jansen, J., Nielsen, S.B., The periglacial engine of mountain erosion – Part](#)
1511 [1: Rates of frost cracking and frost creep., *Earth Surface Dynamics*, 3\(4\), 447-462, 2015.](#)

1512 Arnold, L., Breon, F.M., and Brewer, S: The earth as an extrasolar planet: the vegetation spectral signature today and
1513 during the last Quaternary climatic extrema, *Int. J. Astrobiol.*, 8, 81–94. [http://dx.doi.org/10.1017/](http://dx.doi.org/10.1017/S1473550409004406)
1514 [S1473550409004406](http://dx.doi.org/10.1017/S1473550409004406), 2009.

1515 [Asahi, K.: Equilibrium-line altitudes of the present and Last Glacial Maximum in the eastern Nepal](#)
1516 [Himalayas and their implications for SW monsoon climate, *Quatern. Int.*, 212, 26-34,](#)
1517 [doi:10.1016/j.quaint.2008.08.004, 2010.](#)

1518 Bahrenberg, G., Giese, E., and Nipper, J.: Multivariate Statistik. *Statistische Methoden in der Geographie 2*, Stuttgart,
1519 1992.

1520 [Baker, P. A., Seltzer, G. O., Fritz, S. C., Dunbar, R. B., Grove, M. J., Tapia, P. M., Cross L., Rowe H. D., and](#)
1521 [Baroda, J. P.: The History of South America Tropical Precipitation for the Past 25,000 Years, *Earth and*](#)
1522 [Atmospheric Sciences](#), 291, 640-643, doi:10.1126/science.291.5504.640, 2001.

1523 Ballantyne, A.P., Greenwood, D.R., Sinninghe Damste, J.S., Csank, A.Z., Eberle, J.J., and Ryczynski, N.:
1524 Significantly warmer Arctic surface temperatures during the Pliocene indicated by multiple independent proxies,
1525 *Geology* 38 (7), 603–606, 2010.

1526 Banfield, J.F., Barker, W.W., Welch, S.A., and Taunton, A.: Biological impact on mineral dissolution: application of
1527 the lichen model to understanding mineral weathering in the rhizosphere, *Proceedings of the National Academy of*
1528 *Sciences*, 96, 3404-3411, 1999.

1529 [Barnston, A. G. and Livezey, R. E.: Classification, Seasonality and Persistence of Low-Frequency](#)
1530 [Atmospheric Circulation Patterns, *Mon. Weather Rev.*, 115, 1083-1126, doi:10.1175/1520-](#)
1531 [0493\(1987\)1151083:CSAPOL2.0.CO;2, 1987.](#)

1532 [Barron, J.A., and Anderson, L.: Enhanced Late Holocene ENSO/PDO expression along the margins of the](#)
1533 [eastern North Pacific. *Quaternary International*, 235, 3-12, 2011.](#)

1534 Bendick, R., and Ehlers, T.A.: Extreme localized exhumation at syntaxes initiated by subduction geometry, *Geophys.*
1535 *Res. Lett.*, 41(16), 2014GL061026, doi:10.1002/2014GL061026, 2014.

1536 [Bertrand, S., Charlet F., Charlier B., Renson V., and Fagel N.: Climate variability of southern Chile since the](#)
1537 [Last Glacial Maximum: a continuous sedimentological records from Lago Puyehue \(40°S\), *J. Paleolimnol.*](#)

- 1538 | [39, 179-195, doi:10.1007/s10933-007-9117-y, 2008.](#)
- 1539 | [Betancourt, J. L., Latorre C., Rech J. A., Quade J., and Rylander K.A.: A 22,000 Year Record of Monsoonal](#)
1540 [Precipitation from Northern Chile's Atacama Desert, *Science*, 289, 1542-1546, doi: 10.1126/](#)
1541 [science.289.5484.1542, 2000.](#)
- 1542 Bigelow, N. H., Brubaker, L. B., Edwards, M. E., Harrison, S. P., Prentice, I. C., Anderson, P. M., Andreev, A. A.,
1543 Bartlein, P. J., Christensen, T. R., Cramer, W., Kaplan, J. O., Lozhkin, A. V., Matveyeva, N. V., Murray, D. V.,
1544 McGuire, A. D., Razzhivin, V. Y., Ritchie, J. C., Smith, B., Walker, D. A., Gajewski, K., Wolf, V., Holmqvist, B.
1545 H., Igarashi, Y., Kremenetskii, K., Paus, A., Pisaric, M. F. J., and Vokova, V. S.: Climate change and Arctic ecosys-
1546 tems I. Vegetation changes north of 55°N between the last glacial maximum, mid-Holocene and present. *Journal of*
1547 *Geophysical Research - Atmospheres*, 108(D19), doi: 10.1029/2002JD002558, 2003.
- 1548 | [Bird, B. W., Abbott M. B., Rodbell D. T., and Vuille M.: Holocene tropical South American hydroclimate re-](#)
1549 [vealed from a decadal resolved lake sediment \$\delta^{18}O\$ record, *Earth Planet. Sc. Lett.*, 310, 192-202,](#)
1550 [doi:10.1016/j.epsl.2011.08.040, 2011.](#)
- 1551 | [Bobst, A. L., Lowenstein, T. K., Jordan, T. E., Godfrey, L. V., Ku, T.-L., and Luo, S.: A 106 ka paleoclimate](#)
1552 [record from drill core of the Salar de Atacama, northern Chile, *Palaeogeogr. Palaeocl.*, 173, 21-42,](#)
1553 [doi:10.1016/S0031-0182\(01\)00308, 2001.](#)
- 1554 Braconnot, P., Otto-Bleisner, B., Harrison, S.P., Joussaume, S., Peterschmitt, J.-Y., Abe-Ouchi, A., Crucifix, M.,
1555 Driesschaert, E., Fichet, T., Hewitt, C.D., Kagayama, M., Kitoh, A., Loutre, M.-F., Marti, O., Merkel, U.,
1556 Ramstein, G., Valdes, P., Weber, L., Yu, Y., and Zhao, Y.: Results of PMIP2 coupled simulations of the mid-
1557 Holocene and Last Glacial maximum, part 1: experiments and large-scale features, *Clim Past* 3:261–277, 2007.
- 1558
- 1559 Bohner, J.: General climatic controls and topoclimatic variations in Central and High Asia, *Boreas*, 35(2), 279-295,
1560 2006.
- 1561 Bookhagen, B., Thiede, R.C., and Strecker, M.R.: Late Quaternary intensified monsoon phases control landscape
1562 evolution in the northwest Himalaya, *Geology*, 33(2), 149-152. doi: 10.1130/G20982.1, 2005.
- 1563 Boos, W. R., and Kuang, Z.: Dominant control of the South Asian monsoon by orographic insulation versus plateau
1564 heating, *Nature*, 463, 218-222, 2010.
- 1565 Braun, J.: A simple model for regolith formation by chemical weathering: Regolith Formation, *Journal of Geophysical*
1566 *Research Earth's Surface*, DOI: 10.1002/2016JF003914, 2016.
- 1567 | [Caves, J.: Late Miocene Uplift of the Tian Shan and Altai and Reorganization of Central Asia Climate, *GSA*](#)
1568 [Today, doi:10.1130/gsatg305a.1, 2017.](#)
- 1569 CLIMAP Project Members: Seasonal Reconstruction of the Earth's Surface at the Last Glacial Maximum.
1570 *Map and Chart Series*, Vol. 36, Geological Society of America, 18 pp., 1981.
- 1571
- 1572 Cohen, S.J.: Bringing the global warming issue close to home: The challenge of regional impact studies. *Bulletin of the*
1573 *American Meteorological Society*, 71: 520 – 526., 1990.

- 1574 | [D'Agostino, K., Seltzer, G., Baker, P., Fritz, S., and Dunbar, R.: Late-Quaternary lowstands of Lake Titicaca: evidence from high-resolution seismic data, *Palaeogeogr. Palaeoclimatol.*, 179, 97-111, doi:10.1016/S0031-0182\(01\)00411-4, 2002.](#)
- 1575 |
- 1576 |
- 1577 | [De Batist, M., Fagel N., Loutre M. F., and Chapron E.: A 17,900-year multi-proxy lacustrine record of Lago Puyehue \(Chilean Lake District\): introduction, *J. Paleolimnol.*, 39, 151-161, doi:10.1007/s10933-007-9113-2, 2007.](#)
- 1578 |
- 1579 |
- 1580 | [de Porras, M. E., Maldonado, A., Abarzua, A. M., Cardenas, M. L., Francois, J. P., Martel-Cea, A., Stern, C. R., Mendez, C., and Reyes, O.: Postglacial vegetation, fire and climate dynamics at Central Chilean Patagonia \(Lake Shaman, 44°S\), *Quaternary Sci. Rev.*, 50, 71-85, doi:10.1016/j.quascirev.2012.06.015; 2012.](#)
- 1581 |
- 1582 |
- 1583 |
- 1584 | [de Porras, M. E., Maldonado, A., Quintana, F. A., Martel-Cea, A., Reyes, O., and Méndez, C.: Environmental and climatic changes in central Chilean Patagonia since the Late Glacial \(Mallín El Embudo, 44° S\), *Clim. Past*, 10, 1063–1078, doi:10.5194/cp-10-1063-2014, 2014.](#)
- 1585 |
- 1586 |
- 1587 | Delunel, R., van der Beek, P. A., Carcaillet, J., Bourlès, D. L., and Valla, P.G.: Frost-cracking control on catchment denudation rates: Insights from in situ produced ¹⁰Be concentrations in stream sediments (Ecrins–Pelvoux massif, French Western Alps), *Earth and Planetary Science Letters*, 293(1–2), 72–83, doi:10.1016/j.epsl.2010.02.020, 2010.
- 1588 |
- 1589 |
- 1590 |
- 1591 | [DeMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Samthein, M., Baker, L., and Yarusinsky, M.: Abrupt onset and termination of the African Humid Period:: rapid climate responses to gradual insolation forcing, *Quaternary Sci. Rev.*, 19, 347-361, doi:10.1016/S0277-3791\(99\)00081-5, 2000.](#)
- 1592 |
- 1593 |
- 1594 | [Demske, D., Tarasov, P. E., Wünnemann, B., and Riedel, F.: Late glacial and Holocene vegetation, Indian monsoon and westerly circulation in the Trans-Himalaya recorded in the lacustrine pollen sequence from Tso Kar, Ladakh, NW India, *Palaeogeogr. Palaeoclimatol.*, 279, 172-185, doi:10.1016/j.palaeo.2009.05.008, 2009.](#)
- 1595 |
- 1596 |
- 1597 |
- 1598 | [Diaz, F.P., Latorre, C., Maldonado, A., Quade, J., and Betancourt, J.L.: Rodent middens reveal episodic, long-distance plant colonizations across the hyperarid Atacama Desert over the last 34,000 years, *J. Biogeogr.*, 39, 510-525, doi:10.1111/j.1365-2699.2011.02617.x, 2012.](#)
- 1599 |
- 1600 |
- 1601 | Dietrich, W. E., and Perron, J.T.: The search for a topographic signature of life, *Nature*, 439(7075), 411–418, doi:10.1038/nature04452, 2006.
- 1602 |
- 1603 | Dietrich, S., Werner, M., Spanghel, T., and Lohmann, G.: Influence of orbital forcing and solar activity on water isotopes in precipitation during the mid and late Holocene, *Clim. Past*, 9, 13-26. doi:10.5194/cp-9-13-2013, 2013.
- 1604 |
- 1605 |
- 1606 | Dettman, D.L., Fang, X.M., Garzzone, C.N., and Li, J.J.: Uplift-driven climate change at 12 Ma: a long delta O-18 record from the NE margin of the Tibetan plateau. *Earth and Planetary Science Letters*, 214(1-2), 267-277, 2003.
- 1607 |
- 1608 | Dowsett, H.J., Robinson, M., Haywood, A., Salzmann, U., Hill, D., Sohl, L., Chandler, M., Williams, M., Foley, K., and Stoll, D.: The PRISM3D paleoenvironmental reconstruction. *Stratigraphy*, 7, 123–139, 2010.
- 1609 |
- 1610 | [Egholm, D.L., Nielsen, S.B., Pedersen, V.K., Lesemann, J.: Glacial effects limiting mountain height, *Nature*, 460\(7257\).](#)

- 1611 | [884-887, 2009.](#)
- 1612 | Ehlers, T.A., and Poulsen, C.J.: Influence of Andean uplift on climate and paleoaltimetry estimates, *Earth and Planet-*
1613 *ary Science Letters*, 281(3-4), 238-248, 2009.
- 1614 Etheridge, D.M., Steele, L., Langenfelds, R., Francey, R., Barnola, J., and Morgan, V.: Natural and anthropogenic
1615 changes in atmospheric CO₂ over the last 1000 years from air in Antarctic ice and firn, *J Geophys Res* 101:4115–
1616 4128, 1996.
- 1617
- 1618 Etheridge, D.M., L. Steele, R. Francey, and R. Langenfelds (1998), Atmospheric methane between 1000 a.d. and
1619 present: evidence of anthropogenic emissions and climatic variability. *J Geophys Res*, 103:15979–15993
- 1620
- 1621 Feng, R., Poulsen, C.J., Werner, M., Chamberlain, C.P., Mix, H.T., and Mulch, A.: Early Cenozoic evolution of topo-
1622 graphy, climate, and stable isotopes in precipitation in the North American Cordillera, *American Journal of Science*,
1623 313(7), 613–648, 2013.
- 1624 | [Feng, R., Poulsen, C.J., Werner, M., 2016. Tropical circulation intensification and tectonic extension recorded](#)
1625 [by Neogene terrestrial d18O records of the western United States. *Geology* 44. doi:10.1130/G38212.1](#)
1626
- 1627 | [Feng, R., Poulsen, C.J., 2016. Refinement of Eocene lapse rates, fossil-leaf altimetry, and North American](#)
1628 [Cordilleran surface elevation estimates. *Earth Planet. Sci. Lett.* doi:10.1016/j.epsl.2015.12.022](#)
1629
- 1630 | [Fritz, S. C., Baker, P. A., Lowenstein, T. K., Seltzer, G. O., Rigsby, C. A., Dwyer, G. S., Tapia, P. M., Arnold,](#)
1631 [K. K., Ku, T. L., and Luo, S: Hydrologic variation during the last 170,000 years in the southern hemisphere](#)
1632 [tropics of South America, *Quaternary Res.*, 61, 95 – 104, doi:10.1016/j.yqres.2003.08.007, 2004.](#) Gasse, F.,
1633 [Arnold, M., Fontes, J. C., Fort, M., Gibert, E., Huc, A., Li, B., Li, Y., Liu, Q., Melleres, F., Van Campo, E.,](#)
1634 [Wang, F., and Zhang, Q.: A 13,000-year climate record from western Tibet, *Nature*, 353, 742-745,](#)
1635 [doi:10.1016/j.quaint.2006.02.001, 1991.](#)
- 1636 | [Gayo E. M., Latorre, C., Santoro, C. M., Maldonado, A., and De Pol-Holz, R.: Hydroclimate variability in the](#)
1637 [low-elevation Atacama Desert over the last 2500 yr, *Clim. Past*, 8, 287-306, doi:10.5194/cp-8-287-2012,](#)
1638 [2012.](#)
- 1639 | Gierz, P., Lohmann, G., and Wei, W.: Response of Atlantic Overturning to future warming in a coupled atmosphere-
1640 ocean-ice sheet model, *Geophysical Research Letters*, 42, 6811-6818, doi:10.1002/2015GL065276, 2015.
- 1641 | [Gilli, A., Ariztegui, D., Bradbury, J. P., Kelts, K. R., Markgraf, V., and McKenzie, J. A.: Tracking abrupt cli-](#)
1642 [mate change in the Southern Hemisphere: a seismic stratigraphic study of Lago Cardiel, Argentina](#)
1643 [\(49°S\), *Terra Nova*, 13, 443-448, doi:10.1046/j.1365-3121.2001.00377.x , 2001.](#)
- 1644 Glotzbach, C., van der Beek, P., Carcaillet, J., and Delunel, R.: Deciphering the driving forces of erosion rates on mil-
1645 lennial to million-year timescales in glacially impacted landscapes: An example from the Western Alps, *Journal of*
1646 *Geophysical Research: Earth Surface*, 118, 1491-1515, 2013.
- 1647 Gong, X., Knorr, G., Lohmann, G., and Zhang, X.: Dependence of abrupt Atlantic meridional ocean circulation changes
1648 on climate background states, *Geophysical Research Letters*, 40 (14), 3698-3704, doi:10.1002/grl.50701, 2013.
- 1649 | [Grosjean, M., Van Leeuwen, J., Van der Knaap, W., Geyh, M., Ammann, B., Tanner, W., Messerli, B.,](#)

- 1650 | [Núñez, L., Valero-Garcés, B., and Veit, H. : A 22,000 14C year BP sediment and pollen record of climate](#)
1651 | [change from Laguna Miscanti \(23°S\), northern Chile, *Global Planet. Change*, 28, 35–51, doi:10.1016/](#)
1652 | [S0921-8181\(00\)00063-1, 2001.](#)
- 1653 | Gysels, G., Poesen, J., Bochet, E., and Li, Y.: Impact of plant roots on the resistance of soils to erosion by water: a re-
1654 | view, *Progress in Physical Geography*, 29, 189–217. <http://doi.org/10.1191/0309133305pp443ra>, 2005.
- 1655 | [Hansen, B. C. S., Seltzer, G. O., and Wright Jr., H.E.: Late Quaternary vegetational change in the central](#)
1656 | [Peruvian Andes, *Palaeogeogr. Palaeocl.*, 109, 263-285, doi:10.1016/0031-0182\(94\)90179-1, 1994.](#)
- 1657 | Harris, I., Jones, P.D., Osborn, T.J., and Lister, D.H.: Updated high-resolution grids of monthly climatic observations -
1658 | the CRU TS3.10 Dataset, *International Journal of Climatology*, doi:10.1002/joc.3711, 2013.
- 1659 | Harrison, S. P., Yu, G., Takahara, H., and Prentice, I. C.: Palaeovegetation - Diversity of temperate plants in east Asia.
1660 | *Nature* 413, 129-130, 2001.
1661
- 1662 | Harrison, S.P., Bartlein, P.J., Brewer, S., Prentice, I.C., Boyd, M., Hessler, I., Holmgren, K., Izumi, K., and Willis, K.:
1663 | Climate model benchmarking with glacial and mid-Holocene climates, *Climate Dynamics*, 43, 671-688. doi
1664 | 10.1007/s00382-013-1922-6, 2013.
- 1665 | Haywood, A.M., Valdes, P.J., and Sellwood, B.W.: Global scale palaeoclimate reconstruction of the middle Pliocene
1666 | climate using the UKMO GCM: initial results. *Global and Planetary Change*, 25 (3–4), 239–256, 2000.
- 1667 | Haywood, A.M., Dowsett, H.J., Otto-Bliesner, B., Chandler, M.A., Dolan, A.M., Hill, D.J., Lunt, D.J., Robinson, M.M.,
1668 | Rosenbloom, N., Salzmann, U., and Sohl, L.E.: Pliocene Model Intercomparison Project (PlioMIP): experimental
1669 | design and boundary conditions (Experiment 1), *Geoscientific Model Development* (3), 227-242, 2010.
1670
- 1671 | Haywood, A.M., Hill, D.J., Dolan, A.M., Otto-Bliesner, B., Bragg, F., Chan, W.-L., Chandler, M.A., Contoux, C., Jost,
1672 | A., Kamae, Y., Lohmann, G., Lunt, D.J., Abe-Ouchi, A., Pickering, S.J., Ramstein, G., Rosenbloom, N.A., Sohl, L.,
1673 | Stepanek, C., Yan, Q., Ueda, H., and Zhang, Z.: Large-scale features of Pliocene climate: results from the Pliocene
1674 | Model Intercomparison Project, *Clim. Past*, 9, 191-209. doi:10.5194/cp-9-191-2013, 2013.
1675
- 1676 | Herman, F., Seward, D., Valla, P.G., Carter, A., Kohn, B., Willett, S.D., and Ehlers, T.A.: Worldwide acceleration of
1677 | mountain erosion under a cooling climate, *Nature*, 504, 423–426. doi:10.1038/nature12877, 2013.
- 1678 | [Herzschuh, U., Kuerschner, H., and Mischke, S.: Temperature variability and vertical vegetation belt shifts](#)
1679 | [during the last ~50,000 yr in the Qilian Mountains \(NE margin of the Tibetan Plateau, China\), *Quaternary*](#)
1680 | [Res., 66, 133-146, doi:10.1016/j.yqres.2006.03.001, 2006a.](#)
- 1681 | [Herzschuh, U., Winter, K., Wuennemann, B., and Li, S.: A general cooling trend on the central Tibetan](#)
1682 | [Plateau throughout the Holocene recorded by the lake Zigetang pollen spectra, *Quatern. Int.*, 154, 113-](#)
1683 | [121, doi:10.1016/j.quaint.2006.02.005, 2006b.](#)
- 1684 | [Herzschuh, U., Kramer A., Mischke S., and Zhang C.: Quantitative climate and vegetation trends since the](#)
1685 | [late glacial on the northeastern Tibetan Plateau deduced from Koucha lake pollen spectra, *Quaternary*](#)
1686 | [Res., 71, 162-171, doi:10.1016/j.yqres.2008.09.003, 2009.](#)
1687

- 1688 Hillenbrand, C.-D., and Fütterer, D.K.: Neogene to Quaternary deposition of opal on the continental rise west of the
1689 Antarctic Peninsula, ODP Leg 178, Sites 1095, 1096, and 1101. In: Barker, P.F., Camerlenghi, A., Acton, G.D.,
1690 Ramsay, A.T.S. (Eds.), Proceedings of the Ocean Drilling Programme, Scientific Results, 178. Texas A and M Uni-
1691 versity, College Station, Texas, pp. 1–40 (CD-ROM), 2002.
- 1692 [Hillyer, R., Valencia, B. G., Bush, M.B., Silman, M.R., and Steinitz-Kannan, M.: A 24,700-yr paleolimnological](#)
1693 [history from the Peruvian Andes, *Quaternary Res.*, 71, 71-82, doi:10.1016/j.yqres.2008.06.006, 2009.](#)
- 1694 Hobley, D.E., Sinclair, H.D., and Cowie, P.A.: Processes, rates, and time scales of fluvial response in an ancient post-
1695 glacial landscape of the northwest Indian Himalaya. *Geological Society of America Bulletin*, 122, 1569-1584, 2010.
- 1696 [Hodell, D. A., Brenner, M., Kanfoush, S. L., Curtis, J. H., Stoner, J. S., Song, X., Wu, Y., and Whitmore, T. J.:](#)
1697 [Paleoclimate of southwestern China for the past 50,000 yr inferred from lake sediment records, *Quatern-*](#)
1698 [ary Res., 52, 369-380, doi:10.1006/qres.1999.2072, 1999.](#)
- 1699 [Hu, G., Yi, C.-L., Zhang, J.-F., Liu, J.-L., Jiang, T., and Qin, X.: Optically stimulated luminescence dating of a](#)
1700 [moraine and a terrace in Laohugou valley, western Qilian Shan, northeastern Tibet, *Quaternary Interna-*](#)
1701 [tional 321, 37-49, doi:10.1016/j.quaint.2013.12.019, 2014.](#)
- 1702 Insel, N., Poulsen, C.J., and Ehlers, T.A.: Influence of the Andes Mountains on South American moisture transport,
1703 convection, and precipitation, *Climate Dynamics*, 35 (7-8), 1477-1492, 2010.
- 1704 Jeffery, M.L., Ehlers, T.A., Yanites, B.J., and Poulsen, C.J.: Quantifying the role of paleoclimate and Andean Plateau
1705 uplift on river incision: PALEOCLIMATE ROLE IN RIVER INCISION, *Journal of Geophysical Research: Earth*
1706 *Surface*, 118(2), 852–871, doi:10.1002/jgrf.20055, 2013.
- 1707 [Jenny, B., Valero-Garces, B. L., Urrutia, R., Kelts, K., Veit, H., and Geyh, M.: Moisture changes and](#)
1708 [fluctuations of the Westerlies in Mediterranean Central Chile during the last 2000 years: The Laguna](#)
1709 [Aculeo record \(33°50' S\), *Quatern. Int.*, 87, 3-18, doi:10.1016/S1040-6182\(01\)00058-1, 2002a.](#)
- 1710 [Jenny, B., Valero-Garcés, B.L., Villa-Martínez, R., Urrutia, R., Geyh, M. A., and Veit, H.: Early to Mid-Holo-](#)
1711 [cene Aridity in Central Chile and the Southern Westerlies: The Laguna Aculeo Record \(34°S\), *Quaternary*](#)
1712 [Res., 58, 160–170, doi:10.1006/qres.2002.2370, 2002b.](#)
- 1713 Jungclauss, J. H., Lorenz, S. J., Timmreck, C., Reick, C. H., Brovkin, V., Six, K., Segsneider, J., Giorgetta, M.A.,
1714 Crowley, T.J., Pongratz, J., Krivova, N.A., Vieira, L.E., Solanski, S.K., Klocke, D., Botzet, M., Esch, M., Gayler,
1715 V., Haak, H., Raddatz, T.J., Roeckner, E., Schnur, R., Widmann, H., Claussen, M., Stevens, B., and Marotzke, J.:
1716 Climate and carbon-cycle variability over the last millennium. *Climate of the Past*, 6, 723-737. doi:10.5194/cp-6-723-
1717 2010, 2010.
- 1718 [Junginger, A., Roller, S., Olaka, L. A., and Trauth, M. H.: The effects of solar irradiation changes on the mi-](#)
1719 [gration of the Congo Air Boundary and water levels of paleo-Lake Suguta, Northern Kenya Rift, during the](#)
1720 [African Humid Period \(15–5 ka BP\), *Palaeogeogr. Palaeoclimatol.*, 396, 1-16, doi:10.1016/](#)
1721 [j.palaeo.2013.12.007, 2014.](#)
- 1722 [Kaiser, J., Schefuss, E., Lamy, F., Mohtadi, M., and Hebbeln, D.: Glacial to Holocene changes in sea surface](#)
1723 [temperature and coastal vegetation in north central Chile : high versus low latitude forcing, *Quaternary*](#)

- 1724 | [Sci. Rev., 27, 2064–2075, doi:10.1016/j.quascirev.2008.08.025, 2008.](#)
- 1725 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen,
1726 J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K.C., Ropelewski, C., Wang, J., Leetmaa,
1727 A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project. *Bulletin of the American*
1728 *Meteorological Society*, 77(3), 437-471, 1996.
- 1729 | [Kashiwaya, K., Masuzawa, T., Morinaga, H., Yaskawa, K., Yuan, B. Y., Liu, J. Q., and Gu, Z.: Changes in](#)
1730 [hydrological conditions in the central Qing-Zang \(Tibetan\) Plateau inferred from lake bottom sediments.](#)
1731 [Earth Planet. Sc. Lett., 135, 31-39, doi:10.1016/0012-821X\(95\)00136-Z, 1995.](#)
- 1732 | [Kent-Corson, M., Sherman, L., Mulch, A. and Chamberlain, C.: Cenozoic topographic and climatic response](#)
1733 [to changing tectonic boundary conditions in Western North America, Earth Planet. Sc. Lett., 252\(3-4\),](#)
1734 [453-466, doi:10.1016/j.epsl.2006.09.049, 2006.](#)
- 1735 Kistler, R., Collins, W., Saha, S., White, G., Woollen, J., Kalnay, E., Chelliah, M., Ebisuzaki, W., Kanamitsu, M.,
1736 Kousky, V., Van den Dool, H., Jenne, R., and Fiorino, M.: The NCEP–NCAR 50–Year Reanalysis: Monthly Means
1737 CD–ROM and Documentation. *Bulletin of the American Meteorological Society*, 82(2), 247–267, 2001.
- 1738 Kirchner, J.W., Finkel, R.C., Riebe, C.S., Granger, D.E., Clayton, J.L., King, J.G., and Megahan, W.F.: Mountain
1739 erosion over 10 yr, 10 k.y., and 10 m.y. time scales, *Geology*, 29 (7), 591-594, 2001.
- 1740 Knorr, G., Butzin, M., Micheels, A., and Lohmann, G.: A Warm Miocene Climate at Low Atmospheric CO₂ levels.
1741 *Geophysical Research Letters*, 38, L20701, doi:10.1029/2011GL048873, 2011.
- 1742 Koons, P.O., Zeitler, P.K., and Hallet, B.: 5.14 Tectonic Aneurysms and Mountain Building, in *Treatise on Geomor-*
1743 *phology*, pp. 318–349, Elsevier, 2013.
- 1744 | [Kotlia, B. S., Sharma, C., Bhalla, M. S., Rajagopalan, G., Subrahmanyam, K., Bhattacharya, A., and](#)
1745 [Valdiya, K. S.: Paleoclimatic conditions in the late Pleistocene Wadda Lake, eastern Kumaun Himalaya](#)
1746 [\(India\), Palaeogeogr. Palaeocl., 162, 105-118, doi:10.1016/S0031-0182\(00\)00107-3, 2000.](#)
- 1747 | [Kramer, A., Herzs Schuh, U., Mischke, S., and Zhang, C.: Late glacial vegetation and climate oscillations on](#)
1748 [the southeastern Tibetan Plateau inferred from the lake Naleng pollen profile, Quaternary Res., 73, 324-](#)
1749 [335, doi:10.1016/j.yqres.2009.12.003, 2010.](#)
- 1750 Kraus, H.: Die Atmosphäre der Erde. Eine Einführung in die Meteorologie. Berlin, 2001.
- 1751 Kutzbach, J.E., Guetter, P.J., Ruddiman, W.F., and Prell, W.L.: Sensitivity of Climate to Late Cenozoic Uplift in
1752 Southern Asia and the American West - Numerical Experiments, *Journal of Geophysical Research-Atmospheres*,
1753 94(D15), 18393-18407, 1989.
- 1754 Kutzbach, J.E., Prell, W.L., and Ruddiman, W.F.: Sensitivity of Eurasian Climate to Surface Uplift of the Tibetan Plat-
1755 eau, *Journal of Geology*, 101(2), 177-190, 1993.
- 1756 | [Lamy, F. and Kaiser, J.: Past Climate Variability in South America and Surrounding Regions, Chapter 6:](#)

- 1757 | [Glacial to Holocene Paleoceanographic and Continental Paleoclimate Reconstructions Based on ODP Site 1233/GeoB 3313 Off southern Chile, 2009.](#)
- 1758 |
- 1759 | [Lamy, F., Hebbeln, D., and Wefer, G.: High Resolution Marine Record of Climatic Change in Mid-latitude Chile during the Last 28,000 Years Based on Terrigenous Sediment Parameters, *Quaternary Res.*, 51, 83-93, doi:10.1006/qres.1998.2010, 1999.](#)
- 1760 |
- 1761 |
- 1762 | [Lamy, F., Klump, J., Hebbeln, D., and Wefer, G.: Late Quaternary rapid climate change in northern Chile, *Terra Nova*, 12, 8-13, doi:10.1046/j.1365-3121.2000.00265.x, 2000.](#)
- 1763 |
- 1764 | [Lamy, F., Rühlemann, C., Hebbeln, D. and Wefer, G.: High- and low-latitude climate control on the position of the southern Peru-Chile Current during the Holocene, *Paleoceanography*, 17, 16-1–16-10, doi:10.1029/2001PA000727, 2002.](#)
- 1765 |
- 1766 |
- 1767 | [Latorre, C., Betancourt, J., Rylander, K., and Quade, J.: Vegetation invasions into absolute desert: A 45 000 yr rodent midden record from the Calama-Salar de Atacama basins, northern Chile \(lat 22° - 24°S\), *Geol. Soc. Am. Bull.*, 114, 349–366, doi:10.1130/0016-7606\(2002\)114<0349:VIIADA>2.0.CO;2, 2002.](#)
- 1768 |
- 1769 |
- 1770 | [Latorre, C., Betancourt, J.L., Rylander, K.A., Quade, J., and Matthei, O.: A vegetation history from the arid prepuna of northern Chile \(22-23°S\) over the last 13500 years, *Palaeogeogr. Palaeoclimatol.*, 194, 223-246, doi:10.1016/S0031-0182\(03\)00279-7, 2003.](#)
- 1771 |
- 1772 |
- 1773 | [Latorre, C., Betancourt, J.L., and Arroyo, M.T.K.: Late Quaternary vegetation and climate history of a perennial river canyon in the Río Salado basin \(22°S\) of northern Chile, *Quaternary Res.*, 65, 450-466, doi:10.1016/j.yqres.2006.02.002, 2006.](#)
- 1774 |
- 1775 |
- 1776 | [Lechler, A. and Niemi, N.: Sedimentologic and isotopic constraints on the Paleogene paleogeography and paleotopography of the southern Sierra Nevada, California, *Geology*, 39\(4\), 379-382, doi:10.1130/g31535.1, 2011.](#)
- 1777 |
- 1778 |
- 1779 | [Lechler, A., Niemi, N., Hren, M. and Lohmann, K.: Paleoelevation estimates for the northern and central proto-Basin and Range from carbonate clumped isotope thermometry, *Tectonics*, 32\(3\), 295-316, doi:10.1002/tect.20016, 2013.](#)
- 1780 |
- 1781 |
- 1782 | Lease, R.O., and Ehlers, T.A.: Incision into the Eastern Andean Plateau During Pliocene Cooling, *Science*, 341(6147), 774–776, doi:10.1126/science.1239132, 2013.
- 1783 |
- 1784 | Legates, D.R., and Willmott, C.J.: Mean Seasonal and Spatial Variability in Gauge-Corrected, Global Precipitation, *International Journal of Climatology* 10(2), 111-127. doi: 10.1002/joc.3370100202, 1990.
- 1785 |
- 1786 | Li, J., Ehlers, T.A., Werner, M., Mutz, S.G., Steger, C., Paeth, H.: Late quaternary climate, precipitation $\delta^{18}\text{O}$, and Indian monsoon variations over the Tibetan Plateau. *Earth and Planetary Science Letters* 457, 412-422, 2017.
- 1787 |
- 1788 | [Liu, X., Colman, S. M., Brown, E. T., An, Z., Zhou, W., Jull, A. J. T., Huang, Cheng, Y., P., Liu, W., and Xu, H.: A climate threshold at the eastern edge of the Tibetan plateau, *Geophys. Res. Lett.*, doi: 10.1002/2014GL060833, 2014.](#)
- 1789 |
- 1790 |
- 1791 | [Licht, A., Quade, J., Kowler, A., de los Santos, M., Hudson, A., Schauer, A., Huntington, K., Copeland, P., and Lawton, T.: Impact of the North American monsoon on isotope paleoaltimeters: Implications for the paleoaltimetry of the American southwest, *Am. J. Sci.*, 317\(1\), 1-33, doi:10.2475/01.2017.01, 2017.](#)
- 1792 |
- 1793 |

- 1794 Lohmann, G., Pfeiffer, M., Laepple, T., Leduc, G., and Kim, J.-H.: A model-data comparison of the Holocene global
1795 sea surface temperature evolution. *Clim. Past*, 9, 1807-1839, doi:10.5194/cp-9-1807-2013, 2013.
1796
- 1797 Lorenz, S.J., and Lohmann, G.: Acceleration technique for Milankovitch type forcing in a coupled atmosphere-ocean
1798 circulation model: method and application for the Holocene. *Climate Dynamics* (2004) 23: 727. doi:10.1007/
1799 s00382-004-0469-y, 2004.
1800
- 1801 [Maldonado, A. and Villagrán, C.: Climate variability over the last 9900 cal yr BP from a swamp forest pollen
1802 record along the semiardi coast of Chile. *Quaternary Res.*, 66, 146-258, doi:10.1016/j.yqres.2006.04.003,
1803 2006.](#)
- 1804 [Maldonado, A. J., Betancourt, J. L., Latorre, C., and Villagrán, C.: Pollen analyses from a 50000-yr rodent
1805 midden series in the southern Atacama Desert \(25°30' S\), *J. Quaternary Sci.*, 20, 493-507,
1806 doi:10.1002/jqs.936, 2005.](#)
- 1807 [Maldonado, A., Méndez, C., Ugalde, P., Jackson, D., Seguel, R., and Latorre, C.: Early Holocene climate
1808 change and human occupation along the semiarid coast of north-central Chile. *J. Quaternary Sci.*, 25,
1809 985–988, doi:10.1002/jqs.1385, 2010.](#)
- 1810 [Markgraf, V., Bradbury, J. P., Schwalb, A., Burns, S., Stern, C., Ariztegui, D., Gilli, D., Anselmetti, F. S.,
1811 Stine, S., and Maidana, N.: Holocene palaeoclimates of southern Patagonia: limnological and
1812 environmental history of Lago Cardiel, Argentina, *Holocene*, 13, 581–591,
1813 doi:10.1191/0959683603hl648rp, 2003.](#)
- 1814 [Markgraf, V., Whitlock, C., and Haberle, S.: Vegetation and fire history during the last 18,000 cal yr B.P. in
1815 Southern Patagonia: Mallín Pollux, Coyhaique, Province Aisén \(45°41'30" S, 71°50'30" W, 640 m
1816 elevation\), *Palaeogeogr. Palaeocl.*, 254, 492–507, doi:10.1016/j.palaeo.2007.07.008, 2007.](#)
- 1817 [Maroon, E. A., Frierson, D. M. W., and Battisti, D. S.: The tropical precipitation response to Andes
1818 topography and ocean heat fluxes in an aquaplanet model, *J. Climate*, 28, 381–398, doi:10.1175/JCLI-D-
1819 14-00188.1, 2015.](#)
- 1820 [Maroon, E. A., Frierson, D. M. W., Kang, S. M., and Scheff, J.: The precipitation response to an idealized
1821 subtropical continent, *J. Climate*, 29, 4543–4564, doi:10.1175/JCLI-D-15-0616.1, 2016.](#)
- 1822 Marshall, J.A., Roering, J.J., Bartlein, P.J., Gavin, D.G., Granger, D.E., Rempel, A.W., Praskievicz, S.J., and Hales,
1823 T.C.: Frost for the trees: Did climate increase erosion in unglaciated landscapes during the late Pleistocene? *Science
1824 Advances*, 1, 1-10, 2015.
- 1825 Marston, R.A.: Geomorphology and vegetation on hillslopes: Interactions, dependencies, and feedback loops, *Geomor-
1826 phology*, 116(3–4), 206–217. doi:http://doi.org/10.1016/j.geomorph.2009.09.028, 2010.
- 1827 Matsuoka, N., and Murton, J.: Frost weathering: Recent advances and future directions. *Permafr. Periglac. Process.* 19,
1828 195–210, 2008.
- 1829 Matsuoka, N.: Solifluction rates, processes and landforms: A global review. *Earth Science Reviews* 55, 107–134, 2001.
- 1830 Maussion, F., Scherer, D., Mölg, T., Collier, E., Curio, J., and Finkelnburg, R.: Precipitation seasonality and variability
1831 over the Tibetan Plateau as resolved by the High Asia Reanalysis, *J. Climate*, 27, 1910-1927, doi:10.1175/JCLI-D-
1832 13-00282.1, 2014
- 1833 [Méndez, C., Gil, A., Neme, G., Nuevo Delaunay, A., Cortegoso, V., Huidobro, C., Durán, and Maldano, A.:
1834 Mid Holocene radiocarbon ages in the Subtropical Andes \(~29° - 35° S\), climatic change an implicaton for
1835 human space organization, *Quatern. Int.*, 356, 15-26, doi:10.1016/j.quaint.2014.06.059, 2015.](#)

- 1836 Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P.C., Ebisuzaki, W., Jovic, D., Woollen, J., Rogers, E.,
1837 Berbery, E.H., Ek, M.B., Fan, Y., Grumbine, R., Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D., and Shi, W.:
1838 North American Regional Reanalysis, *Bulletin of the American Meteorological Society*, 87, 343–
1839 360.[doi:10.1175/BAMS-87-3-343](https://doi.org/10.1175/BAMS-87-3-343), 2006.
- 1840 [Methner, K., Fiebig, J., Wacker, U., Umhoefer, P., Chamberlain, C. and Mulch, A.: Eocene-Oligocene proto-](#)
1841 [Cascades topography revealed by clumped \(\$\Delta 47\$ \) and oxygen isotope \(\$\delta 18O\$ \) geochemistry \(Chumstick](#)
1842 [Basin, WA, USA\), *Tectonics*, 35\(3\), 546-564, \[doi:10.1002/2015tc003984\]\(https://doi.org/10.1002/2015tc003984\), 2016.](#)
- 1843 [Mischke S., Kramer M., Herzs Schuh U., Shang H., Erzinger J., and Zhang C.: Reduced early holocene](#)
1844 [moisture availability in the Bayan Har Mountains, northeastern Tibetan Plateau, inferred from a multi-](#)
1845 [proxy lake record, *Palaeogeogr. Palaeoclimatol.*, 267, 59-76, \[doi:10.1016/j.palaeo.2008.06.002\]\(https://doi.org/10.1016/j.palaeo.2008.06.002\), 2008.](#)
- 1846 Molnar, P., Boos, W.R., and Battisti, D.S.: Orographic Controls on Climate and Paleoclimate of Asia: Thermal and
1847 Mechanical Roles for the Tibetan Plateau, *Annual Review of Earth and Planetary Sciences*, 38, 77-102, 2010.
- 1848 Molnar, P., and England, P.: Late Cenozoic uplift of mountain ranges and global climate change: chicken or egg?,
1849 *Nature*, 346, 29-34, 1990.
- 1850 Montgomery, D.R.: Slope distributions, threshold hillslopes, and steady-state topography, *American Journal of Science*,
1851 301, 432-454, 2001.
- 1852 Montgomery, D.R., Balco, G., and Willett, S.D.: Climate, tectonics, and the morphology of the Andes, *Geology*, 29(7),
1853 579-582, 2001.
- 1854 Moon, S., Chamberlain, C.P., Blisniuk, K., Levine, N., Rood, D.H., Hilley, G.E.: Climatic control of denudation in the
1855 deglaciaded landscape of the Washington Cascades, *Nature Geoscience*, 4, 469-473, 2011.
- 1856 [Moreno, A., Santoro, C. M., and Latorre, C.: Climate change and human occupation in the northernmost](#)
1857 [Chilean Altiplano over the last ca. 11500 cal. a BP, *Quaternary Sci.*, 24, 373–382, \[doi:10.1002/jqs.1240\]\(https://doi.org/10.1002/jqs.1240\),](#)
1858 [2009.](#)
- 1859 [Morinaga, H., Itota, C., Isezaki, N., Goto, H., Yaskawa, K., Kusakabe, M., Liu, J., Gu, Z., Yuan, B., and Cong,](#)
1860 [S.: Oxygen-18 and carbon-13 records for the last 14,000 years from lacustrine carbonates of Siling-Co](#)
1861 [\(lake\) in the Qinghai-Tibetan Plateau, *Geophys. Res. Lett.*, 20, 2909-2912, \[doi:10.1029/93GL02982\]\(https://doi.org/10.1029/93GL02982\),](#)
1862 [1993.](#)
- 1863 [Morrill, C., Overpeck, J. T., Cole, J. E., Liu, K., Shen, C., and Tang, L.: Holocene variations in the Asian mon-](#)
1864 [soon inferred from the geochemistry of lake sediments in central Tibet, *Quaternary Res.*, 65, 232-243,](#)
1865 [doi:10.1016/j.yqres.2005.02.014, 2006.](#)
- 1866 Moulton, K.L., and Berner, R.A.: Quantification of the effect of plants on weathering: studies in Iceland, *Geology*, 26,
1867 895-898, 1998.
- 1868 [Muegler, I., Gleixner, G., Guenther, F., Maeusbacher, R., Daut, G., Schuett, B., Berking, J., Schwalb, A.,](#)
1869 [Schwark, L., Xu, B., Yao, T., Zhu, L., and Yi, C.: A multi-proxy approach to reconstruct hydrological](#)
1870 [changes and Holocene climate development of Nam Co, Central Tibet, *J. Paleolimnol.*, 43, 625-648,](#)
1871 [doi:10.1007/s10933-009-9357-0, 2010.](#)

- 1872 [Mujica, M. I., Latorre, C., Maldonado, A., González-Silvestre, L., Pinto, R., De Pol-Holz, R., and Santoro, C.](#)
1873 [M.: Late Quaternary climate change, relict populations and present-day refugia in the northern Atacama](#)
1874 [Desert: a case study from Quebrada La Higuera \(18° S\), J. Biogeogr., 42, 76–88, doi:10.1111/jbi.12383,](#)
1875 [2015.](#)
- 1876 [Mulch, A., Sarna-Wojcicki, A., Perkins, M. and Chamberlain, C.: A Miocene to Pleistocene climate and](#)
1877 [elevation record of the Sierra Nevada \(California\), P. Natl. Acad. Sci. USA, 105\(19\), 6819-6824,](#)
1878 [doi:10.1073/pnas.0708811105, 2008.](#)
- 1879
- 1880 [Mulch, A., Chamberlain, C., Cosca, M., Teyssier, C., Methner, K., Hren, M. and Graham, S.: Rapid change in](#)
1881 [high-elevation precipitation patterns of western North America during the Middle Eocene Climatic Op-](#)
1882 [timum \(MECO\), Am. J. Scie, 315\(4\), 317-336, doi:10.2475/04.2015.02, 2015.](#)
- 1883 Mutz, S.G., Ehlers, T.A., Li, J., Steger, C., Peath, H., Werner, M., Poulsen, C.J.: Precipitation $\delta^{18}\text{O}$ over the South Asia
1884 Orogen from ECHAM5-wiso Simulation: Statistical Analysis of Temperature, Topography and Precipitation.
1885 *Journal of Geophysical Research-Atmospheres*, 121(16),9278-9300, doi: 10.1002/2016JD024856, 2016.
- 1886 [Nishimura, M., Matsunaka, T., Morita, Y., Watanabe, T., Nakamura, T., Zhu, L., Nara, W. F., Imai, A., Izutsu,](#)
1887 [Y., and Hasuike, N.: Paleoclimatic changes on the southern Tibetan Plateau over the past 19,000 years](#)
1888 [recorded in lake Pumoyum Co. and their implications for the southwest monsoon evolution, Palaeogeogr.](#)
1889 [Palaeocl., 396, 75-92, doi:10.1016/j.palaeo.2013.12.015, 2014.](#)
- 1890 Otto-Bliesner, B.L., Brady, C.B., Clauzet, G., Tomas, R., Levis, S., and Kothavala, Z.: Last Glacial Maximum and
1891 Holocene Climate in CCSM3. *Journal of Climate*, 19, 2526-2544, 2006.
- 1892 Paeth, H.: Key Factors in African Climate Change Evaluated by a Regional Climate Model, *Erdkunde*, 58, 290-315,
1893 2004.
- 1894 [Peel, M.C., Finlayson, B.L., McMahon, T.A.: Updated world map of the Koppen- Geiger climate classification.](#)
1895 [Hydrol. Earth Syst. Sci. 11, 1633–1644, 2007.](#)
- 1896
- 1897 Pfeiffer, M., and Lohmann, G.: Greenland Ice Sheet influence on Last Interglacial climate: global sensitivity studies
1898 performed with an atmosphere-ocean general circulation model. *Climate of the Past*, 12, pp. 1313-1338,
1899 doi:10.5194/cp-12-1313-2016, 2016.
- 1900 Pickett, E.J., Harrison, S.P., Flenley, J., Grindrod, J., Haberle, S., Hassell, C., Kenyon, C., MacPhail, M., Martin, H.,
1901 Martin, A.H., McKenzie, M., Newsome, J.C., Penny, D., Powell, J., Raine, J.I., Southern, W., Stevenson, J., Sutra,
1902 J.-P., Thomas, I., van der Kaars, S., Ward, J.: Pollen-based reconstructions of biome distributions for Australia,
1903 South-East Asia and the Pacific (SEAPAC region) at 0, 6000 and 18,000 14C years B.P.. *Journal of Biogeography*,
1904 31, 1381–1444, doi: 10.1111/j.1365-2699.2004.01001.x, 2004.
- 1905
- 1906 [Pingel, H., Mulch, A., Alonso, R., Cottle, J., Hynek, S., Poletti, J., Rohrmann, A., Schmitt, A., Stockli, D. and](#)
1907 [Strecker, M.: Surface uplift and convective rainfall along the southern Central Andes \(Angastaco Basin,](#)
1908 [NW Argentina\), Earth Planet. Sc. Lett., 440, 33-42, doi:10.1016/j.epsl.2016.02.009, 2016.](#)
- 1909
- 1910 Prentice, I. C., Jolly, D., and BIOME 6000 Participants. (2000). Mid-Holocene and glacial-maximum vegetation geo-
1911 geography of the northern continents and Africa. *Journal of Biogeography* 27, 507-519.

- 1912
- 1913 [Pueyo, J. J., Sáez, A., Giralt, S., Valero-Garcés, B.L., Moreno, A., Bao, R., Schwalb, A., Herrera, C., Klo-](#)
- 1914 [sowska, B., and Taberner, C.: Carbonate and organic matter sedimentation and isotopic signatures in](#)
- 1915 [Lake Chungará, Chilean Altiplano, during the last 12.3 kyr, *Palaeogeogr. Palaeocl.*, 307, 339-355,](#)
- 1916 [doi:10.1016/j.palaeo.2011.05.036, 2011.](#)
- 1917 [Quade, J., Rech, J. A., Betancourt, J. L., Latorre, C., Quade, B., Rylander, K. A., and Fisher, T.: Paleowet-](#)
- 1918 [lands and regional climate change in the central Atacama Desert, northern Chile, *Quaternary Res.*, 69,](#)
- 1919 [343-360, doi:10.1016/j.yqres.2008.01.003, 2008.](#)
- 1920 Raymo, M.E., and Ruddiman, W.F.: Tectonic forcing of late cenozoic climate. *Nature*, 359(6391), 117-122, 1992.
- 1921 [Rech, J. A., Pigati, J. S., Quade, J., and Betancourt, J. L.: Re-evaluation of mid-Holocene deposits at](#)
- 1922 [Quebrada Puripica, northern Chile, *Palaeogeogr. Palaeocl.*, 194, 207-222, doi:0.1016/S0031-](#)
- 1923 [0182\(03\)00278-5, 2003.](#)
- 1924 Robinson, M.M.: New quantitative evidence of extreme warmth in the Pliocene Arctic, *Stratigraphy* 6, 265–275, 2009.
- 1925 Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Korn-
- 1926 blueh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U., and Tompkins, A.: The atmospheric general circu-
- 1927 lation model ECHAM5. Part I: Model description. Rep. 349Rep., 127 pp, Max Planck Institute for Meteorology,
- 1928 Hamburg, 2003.
- 1929 Roering, J.J., Marshall, J., Booth, A.M., Mort, M., and Jin, Q.: Evidence for biotic controls on topography and soil pro-
- 1930 duction, *Earth and Planetary Science Letters*, 298(1–2), 183–190, doi:10.1016/j.epsl.2010.07.040, 2010.
- 1931 Salzmann, U., Williams, M., Haywood, A.M., Johnson, A.L.A., Kender, S., and Zalasiewicz, J.: Climate and environ-
- 1932 ment of a Pliocene warm world. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 309 (1-8), 2011.
- 1933 [Sarnthein, M.: Sand deserts during glacial maximum and climatic optimum, *Nature*, 272, 43–46,](#)
- 1934 [doi:10.1038/272043a0, 1978.](#)
- 1935 Sarnthein, M., Gersonde, R., Niebler, S., Pflaumann, U., Spielhagen, R., Thiede, J., Wefer, G., Weinelt, M.: Overview
- 1936 of Glacial Atlantic Ocean Mapping (GLAMAP 2000), *Paleoceanography*, 18(2), doi:10.1029/2002PA000769,
- 1937 2003.
- 1938 Schäfer-Neth C., and Paul, A.: The Atlantic Ocean at the last glacial maximum: objective mapping of the GLAMAP
- 1939 sea-surface conditions. In: Wefer G, Mulitza S, Ratmeyer V (eds) *The South Atlantic in the late quaternary: recon-*
- 1940 *struction of material budgets and current systems.* Springer, Berlin, pp 531–548, 2003.
- 1941 Schaller, M., von Blanckenburg, F., Veldkamp, A., Tebbens, L.A., Hovius, N., and Kubik, P.W.: A 30 000 yr record of
- 1942 erosion rates from cosmogenic ^{10}Be in Middle European river terraces, *Earth and Planetary Science Letters*,
- 1943 204(1), 307–320, 2002.
- 1944 Schaller, M., and Ehlers T.A.: Limits to quantifying climate driven changes in denudation rates with cosmogenic radio-
- 1945 nuclides, *Earth and Planetary Science Letters*, v. 248, pp. 153-167. doi:10.1016/j.epsl.2006.05.027, 2006.

- 1946 | [Schwalb, A., Burns, S., and Kelts, K.: Holocene environments from stable isotope stratigraphy of ostracods](#)
1947 | [and authigenic carbonate in Chilean Altiplano Lakes, *Palaeogeogr. Palaeocl.*, 148, 153-168, doi:10.1016/](#)
1948 | [S0031-0182\(98\)00181-3, 1999.](#)
- 1949 | [Shen J., Liu X. Q., Wang S. M., and Matsumoto R.: Palaeoclimatic changes in the Qinghai, Lake area during](#)
1950 | [the last 18,000 years, *Quatern. Int.*, 136, 131-140, doi:10.1016/j.quaint.2004.11.014, 2005.](#)
- 1951 | [Shen C., Liu K., Tang L., and Overpeck J. T.: Quantitative relationships between modern pollen rain and cli-](#)
1952 | [mate in the Tibetan Plateau. *Rev. Palaeobot. Palyno.*, 140, 61–77, doi:10.1016/j.revpalbo.2006.03.001,](#)
1953 | [2006.](#)
- 1954 | [Siani, G., Colin, C., Michel, E., Carel, M., Richter, T., Kissel, C., and Dewilde, F.: Late Glacial to Holocene](#)
1955 | [terrigenous sediment record in the Northern Patagonian margin: Paleoclimate implications, *Palaeogeogr.*](#)
1956 | [Palaeocl., 297, 26-36, doi:10.1016/j.palaeo.2010.07.011, 2010.](#)
- 1957 | Simmons, A.J., Burrige, D.M., Jarraud, M., Girard, C., and Wergen, W.: The ECMWF Medium-Range prediction
1958 | models development of the numerical formulations and the impact of increased resolution, *Meteorology and Atmo-*
1959 | *spheric Physics*, 40(1-3), 28-60, 1989.
- 1960 | Sohl, L.E., Chandler, M.A., Schmunk, R.B., Mankoff, K., Jonas, J.A., Foley, K.M., and Dowsett, H.J.: PRISM3/GISS
1961 | topographic reconstruction, *U.S. Geological Survey Data Series*, 419, 6p., 2009.
1962
- 1963 | Sowers T., Alley, R.B., and Jubenville, J.: Ice core records of atmospheric N₂O covering the last 106,000 years. *Sci-*
1964 | *ence*, 301:945–948, 2003.
1965
- 1966 | Stepanek, C., and Lohmann, G.: Modelling mid-Pliocene climate with COSMOS , *Geosci. Model Dev.* , 5 , pp. 1221-
1967 | 1243 . doi:10.5194/gmd-5-1221-2012, 2012.
- 1968 | [Stine, S. and Stine M.: A record from Lake Cardiel of climate change in southern South America, *Nature*,](#)
1969 | [345, 705-708, doi:10.1038/345705a0,1990.](#)
- 1970 | [Szeicz, J. M., Haberle, S. G., and Bennett, K. D.: Dynamics of North Patagonian rainforests from fine-resolu-](#)
1971 | [tion pollen, charcoal and tree-ring analysis, *Chonos Archipelago, Southern Chile, Austral Ecol.*, 28, 413–](#)
1972 | [422, doi:10.1046/j.1442-9993.2003.01299.x , 2003.](#)
- 1973 | Takahashi, K., and Battisti, D.: Processes controlling the mean tropical pacific precipitation pattern. Part I: The Andes
1974 | and the eastern Pacific ITCZ. *Journal of Climate*, 20(14), 3434-3451, 2007a.
- 1975 | Takahashi, K., and Battisti, D.: Processes controlling the mean tropical pacific precipitation pattern. Part II: The SPCZ
1976 | and the southeast pacific dry zone. *Journal of Climate*, 20(23), 5696-5706, 2007b.
- 1977 | [Tang, L., Shen, S., Liu, K., and Overpeck, J. T.: Changes in south Asian monsoon: new high-resolution pa-](#)
1978 | [leoclimatic records from Tibet, China, *Chinese Sci. Bull.*, 45, 87-91, doi:10.1007/BF02884911, 2000.](#)
- 1979 | Thiede, R.C., and Ehlers, T.A.: Large spatial and temporal variations in Himalayan denudation. *Earth and Planetary*
1980 | *Science Letters*, 374, 256-257. doi:10.1016/j.epsl.2013.03.004, 2013.

- 1981 Thomas, A.: The climate of the Gongga Shan range, Sichuan Province, PR China. *Arctic and Alpine Research*, 29(2),
1982 226-232, 1997.
- 1983 Uppala, S.M., Kållberg, P.W., Simmons, A.J., Andrae, U., da Costa Bechtold, V., Fiorino, M., Gibson, J.K., Haseler, J.,
1984 Hernandez, A., Kelly, G.A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R.P., Andersson, E., Arpe, K.,
1985 Balmaseda, M.A., Beljaars, A.C.M., van de Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A.,
1986 Dragosavac, M., Fisher, M. Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B.J., Isaksen, L., Janssen, P.A.E.M.,
1987 Jenne, R., McNally, A.P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N.A., Saunders, R.W., Simon, P., Sterl, A.,
1988 Trenberth, K.E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis. *Quart. J. R.*
1989 *Meteorol. Soc.*, 131, 2961-3012, 2005.
- 1990
- 1991 von Blanckenburg, F., Bouchez, J., Ibarra, D.E., and Maher, K.: Stable runoff and weathering fluxes into the oceans
1992 over Quaternary climate cycles, *Nature Geoscience*, 8(7), 538–542, doi:10.1038/ngeo2452, 2015.
- 1993 Valla, P.G., Shuster, D.L., and van der Beek, P.A.: Significance increase in relief of European Alps during mid-Pleisto-
1994 cene glaciations. *Nature Geoscience*, 4, 688–692. doi:10.1038/ngeo1242, 2011.
- 1995 [Van Campo, E., Cour, P., and Huang, S.: Holocene environmental changes in Bangong Co basin \(Western
1996 Tibet\). Part 2: The pollen record, *Palaeogeogr. Palaeoclimatol.*, 120, 49-63, doi:10.1016/0031-0182\(95\)00033-
1997 X, 1996.](#)
- 1998
- 1999 [Villagrán, C. and Varela, J.: Palynological Evidence for Increased Aridity on the Central Chilean Coast during
2000 the Holocene, *Quaternary Res.*, 34, 198–207, doi:10.1016/0033-5894\(90\)90031-F, 1990.](#)
- 2001 [Villa-Martínez, R., Villagrán, C., and Jenny, B.: The last 7500 cal yr B.P. of westerly rainfall in Central Chile
2002 inferred from a high-resolution pollen record from Laguna Aculeo \(34°S\), *Quaternary Res.*, 60, 284-293,
2003 doi:https://doi.org/10.1016/j.yqres.2003.07.007, 2003.](#)
- 2004 Wei, W., and Lohmann, G.: Simulated Atlantic Multidecadal Oscillation during the Holocene. *Journal of Climate*, 25,
2005 6989–7002. doi: http://dx.doi.org/10.1175/JCLI-D-11-00667.1, 2012.
- 2006 [Wang, R.L., Scarpitta, S. C., Zhang, S. C., and Zheng, M. P.: Later Pleistocene/Holocene climate conditions
2007 of Qinghai-Xizhang Plateau \(Tibet\) based on carbon and oxygen stable isotopes of Zabuye Lake sedi-
2008 ments, *Earth Planet. Sc. Lett.*, 203, 461-477, doi:10.1016/S0012-821X\(02\)00829-4, 2002.](#)
- 2009 Whipple, K.X, Kirby, E., and Broecklehurst, S.H.: Geomorphic limits to climate-induced increases in topographic re-
2010 lief. *Nature*, 401, 39-43. doi:10.1038/43375, 1999.
- 2011 Whipple, K.X., and Tucker, G.E.: Dynamics of the stream-power river incision model: Implications for height limits of
2012 mountain ranges, landscape response timescales, and research needs, *Journal of Geophysical Research-Solid Earth*,
2013 104, 17661-17674, 1999.
- 2014 Whipple, K. X.: The influence of climate on the tectonic evolution of mountain belts, *Nat. Geosci.*, 2, doi:10.1038/
2015 ngeo413, 2009.
- 2016 Wilks, D.S.: Statistical methods in the atmospheric sciences - 3rd ed. Academic Press, Oxford, 2011.

- 2017 Willett, S.D., Schlunegger, F., and Picotti, V.: Messinian climate change and erosional destruction of the central
2018 European Alps. *Geology*, 34(8), 613-616, 2006.
- 2019 Wilson, G.S., Barron, J.A., Ashworth, A.C., Askin, R.A., Carter, J.A., Curren, M.G., Dalhuisen, D.H., Friedmann, E.I.,
2020 Fyodorov-Davidov, D.G., Gilichinsky, D.A., Harper, M.A., Harwood, D.M., Hiemstra, J.F., Janecek, T.R., Licht,
2021 K.J., Ostroumov, V.E., Powell, R.D., Rivkina, E.M., Rose, S.A., Stroeven, A.P., Stroeven, P., van der Meer, J.J.M.,
2022 Wizevich, M.C.: The Mount Feather Diamicton of the Sirius Group: an accumulation of indicators of Neogene Ant-
2023 arctic glacial and climatic history, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 182 (1–2), 117–131, 2002.
- 2024 [Wischnewski, J., Mischke, S., Wang, Y., and Herzsuh, U.: Reconstructing climate variability on the](#)
2025 [northeastern Tibetan Plateau since the last lateglacial – a multi-proxy, dual-site approach comparing](#)
2026 [terrestrial and aquatic signals, *Quaternary Science Reviews*, 30, 82-97,](#)
2027 [doi:10.1016/j.quascirev.2010.10.001, 2011.](#)
- 2028 [Wünnemann, B., Mischke, S., and Chen, F.: A Holocene sedimentary record from Bosten lake, China, Pa-](#)
2029 [laeogeogr. *Palaeocl.*, 234, 223-238, doi:10.1016/j.palaeo.2005.10.016, 2006.](#)
- 2030 [Yanhong, W., Lücke, A., Zhangdong, J., Sumin, W., Schleser, G.H., Battarbee, R.W., and Weilan, X.: Holo-](#)
2031 [cene climate development on the central Tibetan Plateau: A sedimentary record from Cuoe Lake, *Palaeo-*](#)
2032 [geogr. *Palaeocl.*, 234, 328-340, doi:10.1016/j.palaeo.2005.09.017, 2006](#)
- 2033 Yanites, B.J., and Ehlers, T.A.: Global climate and tectonic controls on the denudation of glaciated mountains. *Earth*
2034 *and Planetary Science Letters*, 325, 63-75, 2012.
- 2035 [Yu, L. and Lai, Z.: Holocene climate change inferred from stratigraphy and OSL chronology of Aeolian sedi-](#)
2036 [ments in the Qaidam Basin, northeastern Qinghai-Tibetan Plateau, *Quaternary Res.*, 81, 488-499,](#)
2037 [doi:10.1016/j.yqres.2013.09.006, 2014.](#)
- 2038 [Zhang, C. and Mischke, S.: A lateglacial and Holocene lake record from the Nianbaoyeze Mountains and in-](#)
2039 [ferences of lake, glacier and climate evolution on the eastern Tibetan Plateau, *Quaternary Sci. Rev.*, 28,](#)
2040 [1970-1983, doi:10.1016/j.quascirev.2009.03.007, 2009.](#)
- 2041 [Zhang, J., Chen, F., Holmes, J. A., Li, H., Guao, X., Wang, J., Li, S., Lu, Y., Zhao, Y., and Qiang, M.: Holo-](#)
2042 [cene monsoon climate documented by oxygen and carbon isotopes from lake sediments and peat bogs](#)
2043 [in China: a review and synthesis, *Quaternary Sci. Rev.*, 30, 1973-1987, doi:10.1016/](#)
2044 [j.quascirev.2011.04.023, 2011.](#)
- 2045 Zhang, X., Lohmann, G., Knorr, G., and Xu X.: Different ocean states and transient characteristics in Last Glacial Max-
2046 imum simulations and implications for deglaciation. *Clim. Past*, 9, 2319-2333, doi:10.5194/cp-9-2319-2013, 2013a.
- 2047 Zhang, R., Yan, Q., Zhang, Z.S., Jiang, D., Otto-Bliesner, B.L., Haywood, A.M., Hill, D.J., Dolan, A.M., Stepanek, C.,
2048 Lohmann, G., Contoux, C., Bragg, F., Chan, W.-L., Chandler, M.A., Jost, A., Kamae, Y., Abe-Ouchi, A., Ramstein,
2049 G., Rosenbloom, N.A., Sohl, L., and Ueda, H.: East Asian monsoon climate simulated in the PlioMIP. *Clim. Past*, 9,
2050 2085-2099, doi:10.5194/cp-9-2085-2013, 2013b.
- 2051
- 2052 Zhang, X., Lohmann, G., Knorr, G., and Purcell C.: Abrupt glacial climate shifts controlled by ice sheet changes.

2053 *Nature*, 512 (7514), 290-294, doi:10.1038/nature13592, 2014.

2054 Zhisheng, A., Kutzbach, J.E., Prell, W.L., and Porter, S.C.: Evolution of Asian monsoons and phased uplift of the South
2055 Asian plateau since Late Miocene times. *Nature*, 411(6833), 62-66, 2001.

2056 [Zhou, W. J., Lu, X. F., Wu, Z. K., Deng, L., Jull, A. J. T., Donahue, D. J., and Beck, W.: Peat record reflecting
2057 Holocene climate change in the Zoigê Plateau and AMS radiocarbon dating, *Chinese Sci. Bull.*, 47, 66-
2058 70, doi:10.1360/02tb9013, 2002.](#)

2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106

2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165




Figure 1

2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225




Figure 2

2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267
2268
2269
2270
2271
2272
2273
2274
2275
2276
2277
2278
2279
2280
2281
2282
2283
2284
2285




Figure 3




Figure 4

2289
2290
2291
2292
2293
2294
2295
2296
2297
2298
2299
2300
2301
2302
2303
2304
2305
2306
2307
2308
2309
2310
2311
2312
2313
2314
2315
2316
2317
2318
2319
2320
2321
2322
2323
2324
2325
2326
2327
2328
2329
2330
2331
2332
2333
2334
2335
2336
2337
2338
2339
2340
2341
2342
2343
2344
2345
2346
2347
2348




Figure 5

2349
 2350
 2351
 2352
 2353
 2354
 2355
 2356
 2357
 2358
 2359
 2360
 2361
 2362
 2363
 2364
 2365
 2366
 2367
 2368
 2369
 2370
 2371
 2372
 2373
 2374
 2375
 2376
 2377
 2378
 2379
 2380
 2381
 2382
 2383
 2384
 2385
 2386
 2387
 2388
 2389
 2390
 2391
 2392
 2393
 2394
 2395
 2396
 2397
 2398
 2399
 2400
 2401
 2402
 2403
 2404
 2405
 2406
 2407
 2408




Figure 6

2409
2410
2411
2412
2413
2414
2415
2416
2417
2418
2419
2420
2421
2422
2423
2424
2425
2426
2427
2428
2429
2430
2431
2432
2433
2434
2435
2436
2437
2438
2439
2440
2441
2442
2443
2444
2445
2446
2447
2448
2449
2450
2451
2452
2453
2454
2455
2456
2457
2458
2459
2460
2461
2462
2463
2464
2465
2466
2467
2468




Figure 7

2469
2470
2471
2472
2473
2474
2475
2476
2477
2478
2479
2480
2481
2482
2483
2484
2485
2486
2487
2488
2489
2490
2491
2492
2493
2494
2495
2496
2497
2498
2499
2500
2501
2502
2503
2504
2505
2506
2507
2508
2509
2510
2511
2512
2513
2514
2515
2516
2517
2518
2519
2520
2521
2522
2523
2524
2525
2526
2527
2528




Figure 8

2529
 2530
 2531
 2532
 2533
 2534
 2535
 2536
 2537
 2538
 2539
 2540
 2541
 2542
 2543
 2544
 2545
 2546
 2547
 2548
 2549
 2550
 2551
 2552
 2553
 2554
 2555
 2556
 2557
 2558
 2559
 2560
 2561
 2562
 2563
 2564
 2565
 2566
 2567
 2568
 2569
 2570
 2571
 2572
 2573
 2574
 2575
 2576
 2577
 2578
 2579
 2580
 2581
 2582
 2583
 2584
 2585
 2586
 2587
 2588




Figure 9




Figure 10

2590
2591
2592
2593
2594
2595
2596
2597
2598
2599
2600
2601
2602
2603
2604
2605
2606
2607




Figure 11

2608
 2609
 2610
 2611
 2612
 2613
 2614
 2615
 2616
 2617
 2618
 2619
 2620
 2621
 2622
 2623
 2624
 2625
 2626
 2627
 2628
 2629
 2630
 2631
 2632
 2633
 2634
 2635
 2636
 2637
 2638
 2639
 2640
 2641
 2642
 2643
 2644
 2645
 2646
 2647
 2648
 2649
 2650
 2651
 2652
 2653
 2654
 2655
 2656
 2657
 2658
 2659
 2660
 2661
 2662
 2663
 2664
 2665
 2666
 2667




Figure 12

2668
 2669
 2670
 2671
 2672
 2673
 2674
 2675
 2676
 2677
 2678
 2679
 2680
 2681
 2682
 2683
 2684
 2685
 2686
 2687
 2688
 2689
 2690
 2691
 2692
 2693
 2694
 2695
 2696
 2697
 2698
 2699
 2700
 2701
 2702
 2703
 2704
 2705
 2706
 2707
 2708
 2709
 2710
 2711
 2712
 2713
 2714
 2715
 2716
 2717
 2718
 2719
 2720
 2721
 2722
 2723
 2724
 2725
 2726
 2727




Figure 13

2728
 2729
 2730
 2731
 2732
 2733
 2734
 2735
 2736
 2737
 2738
 2739
 2740
 2741
 2742
 2743
 2744
 2745
 2746
 2747
 2748
 2749
 2750
 2751
 2752
 2753
 2754
 2755
 2756
 2757
 2758
 2759
 2760
 2761
 2762
 2763
 2764
 2765
 2766
 2767
 2768
 2769
 2770
 2771
 2772
 2773
 2774
 2775
 2776
 2777
 2778
 2779
 2780
 2781
 2782
 2783
 2784
 2785
 2786
 2787




Figure 14




Figure 15

2788
2789
2790
2791
2792
2793
2794
2795




Figure 16

2796
2797
2798
2799
2800
2801
2802
2803
2804
2805
2806
2807
2808
2809
2810
2811
2812
2813
2814
2815
2816
2817
2818
2819
2820
2821
2822
2823
2824
2825
2826
2827
2828
2829
2830
2831
2832
2833
2834
2835





Figure 17

2836
2837
2838
2839
2840
2841
2842
2843
2844
2845
2846
2847
2848
2849
2850
2851
2852
2853
2854
2855
2856
2857
2858
2859
2860
2861
2862
2863
2864
2865
2866
2867
2868
2869
2870
2871
2872
2873
2874
2875
2876
2877
2878
2879
2880
2881
2882
2883
2884



Supplemental Figure S1
(in supplement file)