

2 **Effect of changing vegetation and precipitation on denudation**
3 **(part 2): Predicted landscape response to transient climate and**
4 **vegetation cover over millennial to million year timescales**

5 Manuel Schmid ⁽¹⁾, Todd A. Ehlers^{(1)*}, Christian Werner.⁽²⁾, Thomas Hickler⁽²⁾⁽³⁾, Juan-Pablo
6 Fuentes-Espoz⁽⁴⁾

7 ⁽¹⁾ University of Tuebingen, Department of Geosciences; Wilhelmstrasse 56, 72074 Tuebingen, Germany
8 (Manuel.Schmid@Uni-Tuebingen.de, Todd.Ehlers@Uni-Tuebingen.de)

9 ⁽²⁾ Senckenberg Biodiversity and Climate Research Center (BiK-F), Senckenberganlage 25, 60325 Frankfurt/Main,
10 Germany (Christian.Werner@Senckenberg.de)

11 ⁽³⁾ Department of Physical Geography, Geosciences, Goethe-University, Frankfurt, Altenhoferallee 1, 60438
12 Frankfurt/Main, Germany (Thomas.Hickler@Senckenberg.de)

13 ⁽⁴⁾ University of Chile, Department of Silviculture and Nature Conservation, Av. Santa Rosa 11315, La Pintana, Santiago
14 RM, Chile (jufuente@uchile.cl)

15

16 * Corresponding author: (Todd.Ehlers@Uni-Tuebingen.de)

17

Abstract We present a numerical modeling investigation into the interactions between transient climate and vegetation cover with hillslope and detachment limited fluvial processes. Model simulations were designed to investigate topographic patterns and behavior resulting from changing climate and associated changes in surface vegetation cover. The Landlab surface process model was modified to evaluate the effects of temporal variations in vegetation cover on hillslope diffusion and fluvial erosion. A suite of simulations were conducted to represent present day climatic conditions and satellite derived vegetation cover at four different research areas in the Chilean Coastal Cordillera. These simulations included steady-state simulations as well as transient simulations with forcings in either climate or vegetation cover over millennial to million-year timescales. These simulations included two different transient variations in climate and vegetation cover including a step change in climate or vegetation, as well as 100 kyr oscillations over 5 Myr. We conducted eight different step-change simulations for positive and negative perturbations in either vegetation cover or climate and six simulations with oscillating transient forcings for either vegetation cover or climate, and oscillations in both vegetation cover and climate. Results indicate that the coupled influence of surface vegetation cover and mean annual precipitation shifts basin landforms towards a new steady state, with the magnitude of change highly sensitive to the initial vegetation and climate conditions of the basin. Dry, non-vegetated basins show higher magnitudes of adjustment than basins that are situated in wetter conditions with higher vegetation cover. For coupled conditions when surface vegetation cover and mean annual precipitation change simultaneously, the landscape response tends to be weaker. When vegetation cover and mean annual precipitation change independently from each other, higher magnitude shifts in topographic metrics are predicted. Changes in vegetation cover show a higher impact on topography for low initial surface cover values whereas for areas with high initial surface cover, the effect of changes in precipitation dominate the formation of landscapes. This study demonstrates a sensitivity of catchment characteristics to different transient forcings in vegetation cover and mean annual precipitation, with a crucial role for initial vegetation and climate conditions.

1. Introduction

Plants cover most of Earth's surface and interact chemically and physically with the atmosphere, lithosphere and hydrosphere. The abundance and distribution of plants throughout Earth's history is a function, amongst other things, of changing climate conditions that can impact the temporal distribution of plant functional types and vegetation cover present in an area (Hughes, 2000; Muhs et al., 2001; Walther et al. 2002). The physical feedbacks of vegetation on the Earth's near surface manifest themselves mainly through an influence of plants on weathering, erosion, transport and the deposition of sediments (Marston, 2010; Amundson et al., 2015). Although the effects of biota on surface processes has been recognized for over a 100 years (e.g., Gilbert, 1877; Langbein & Schumm, 1958), early studies focused mainly on qualitative descriptions of the underlying processes. With the rise of new techniques to quantify mass transport from the plot- to catchment-scale, and the emergence of improved computing techniques and landscape evolution models, research shifted more towards building a quantitative understanding of how biota influence both hillslope and fluvial processes (Stephan and Gutknecht, 2002; Roering et al., 2002; Marston, 2010; Curran and Hession, 2013). The previous studies motivate the companion papers presented here. In part 1 (Werner et al. 2018-this volume) a dynamic vegetation model is used to evaluate the magnitude of past (Last Glacial Maximum to present) vegetation change along the climate and ecological gradient in the Coastal Cordillera of Chile. Part 2 (this study) presents a sensitivity analysis of how transient climate and vegetation impact catchment denudation. This component is evaluated through implementation of transient vegetation effects for hillslopes and detachment limited rivers in a landscape evolution model.

Previous research in agricultural engineering has focused on plot-scale models to predict total soil loss in response to land-use change (Zhou et al. 2006; Feng et al. 2010) or general changes in plant surface cover (Gyssels et al. 2005), but do not draw conclusions about large-scale geomorphic feedbacks active over longer (millennial) timescales and larger spatial scales. However, a better understanding of how vegetation influences the large scale topographic features (e.g. relief, hillslope angles, catchment denudation) is crucial to understanding the evolution of modern landscapes. At the catchment scale, observational studies have found a correlation between higher values of mean vegetation cover and basin wide denudation rates or topographic metrics (Jeffery et al., 2014, Sangireddy et al. 2016, Acosta et al. 2015). Parallel to the previous observational studies, numerical modeling experiments of the interactions between landscape erosion and surface vegetation cover have also made progress. For example, Collins et al. (2004) were one of the first who attempted to couple vegetation dynamics with a landscape evolution model and found that the introduction of plants to their model resulted in steeper equilibrium landscapes with a higher variability in magnitude of erosional events. Following this, subsequent modeling studies built upon the previous findings with more sophisticated formulations of vegetation-erosion interactions (Istanbulluoglu and Bras, 2005) including the influence of root strength on hillslopes (Vergani et al., 2017). These studies found that not only is there a positive relationship between vegetation cover and mean catchment slope and elevation but there also exists an inverse relationship between vegetation cover and drainage density, due to the plants ability to hinder fluvial erosion and channel initiation.

The advances of the previous studies are limited mainly by their consideration of static vegetation cover or very simple formulations of dynamic vegetation cover. The exception to this is Istanbulluoglu and Bras (2005) who also considered the lag time for vegetation regrowth on hillslopes after a mass wasting event and Yetemen et al. (2015) which considered more complex hydrology in their models but on a smaller spatial scale. However, numerous studies (Ledru et al., 1997; Allen and Breshears, 1998; Bachelet et al., 2003) document that vegetation cover changes in tandem with climate change over a range of timescales (decadal to million year). Missing from previous landscape evolution studies, is consideration of not only how transient vegetation cover influences catchment denudation, but also how coeval changes in precipitation

influence catchment-wide mean denudation. While the effects of climate change over geologic timescales on denudation rates and sediment transport dynamics have been investigated by others (e.g., Schaller et al., 2002; Dosetto et al., 2010; McPhillips et al. 2013), the combined effects of vegetation and climate change on catchment denudation have not. Thus, over longer (geologic) timescales, we are left with a complicated situation of both vegetation and climate changes, and the individual contributions of these changes to catchment scale denudation are difficult to disentangle. In this study, we build upon previous works by investigating both the temporal and spatial sensitivities of landscapes to the coupled vegetation-climate system. By focusing on simplified transient forcings such as a step change, or 100 kyr oscillations in climate and vegetation cover we present a sensitivity analysis of the landscape response to each of these changes, including a better understanding of the direction, magnitude and rates of landscape change. Our model setup is motivated by four study-areas along the climate and vegetation gradient in Chile (Fig. 1a) and illuminates the transient catchment response to biotic vs. climate changes. These study areas are part of the recently initiated German priority research program *EarthShape: Earth surface shaping by biota* (www.earthshape.net). This region is used to provide a basis for our model setup for covariation in precipitation and vegetation present in a natural setting. While we present results representative of the Coastal Cordillera, Chile, it is beyond the scope of this study to provide a detailed calibration to this area and our main objective is identifying the sensitivity, and emergent behaviour, of catchment denudation to changing precipitation and vegetation cover over millennial timescales. This study also builds upon results from the companion paper (Werner et al. 2018 - this volume) by imposing temporal variations in vegetation cover, identified in that study.

2. Background to model setup

Model setup and the range of initial conditions chosen for models were based upon four study-areas located in the Coastal Cordillera of Chile (26°S to 38°S). The focus areas shown in Fig. 1a were chosen because of their similar granitic lithology and geologic and tectonic history (Andriessen and Reutter, 1994; McInnes et al., 1999; Juez-Larré et al., 2010; Makshev and Zentilli, 1999; Avdievitch et al., 2017), and the large gradient in climate and vegetation cover over the region (Fig. 1b,c). These study areas include (from north to south): Parque Nacional Pan de Azúcar; Reserva Santa Gracia; Parque Nacional La Campana, and Parque Nacional Nahuelbuta. Although this study does not explicitly present landscape evolution model results ‘calibrated’ to these specific areas, we’ve chosen the model input (e.g. precipitation, initial vegetation cover, rate of tectonic rock uplift) to represent these areas to provide simulation results that represent the non-linear relationship between precipitation and vegetation cover (e.g. Fig. 1b, c) over a large climate gradient. Topographic metrics such as mean basin slope, total basin relief, mean basin channel steepness, and mean surface vegetation cover and mean annual precipitation were extracted for the main catchments and a subset of adjacent catchments (Fig. 1; Fig. 2). Topographic metrics were extracted from 30 m resolution digital elevation model from the NASA shuttle radar topography mission (SRTM), and vegetation related datasets from the moderate resolution imaging spectroradiometer (MODIS) satellite data (https://landcover.usgs.gov/green_veg.php).

2.1 Landscape evolution modeling approach and the applicability of these results

Landscape evolution model studies can be assigned to different general approaches, which were conceptually defined by Dietrich et al. (2013). The different approaches presented in the Dietrich et al. (2013) study mostly differ in the complexity

117 of input parameters and in the resulting claim for reproducing realistic complexity in modelled landscapes. For this study
118 we have chosen the approach of essential realism, which acknowledges a system-inherent indeterminacy in the evolving
119 topography but focuses on predicting the first-order trends within a system and the differences between landscapes, based
120 on different external conditions, incorporated in the model (Howard, 1997).

121 While we do not claim to reproduce the topographic metrics of the four different focus areas in Chile on a realistic level,
122 our approach determines the general first-order effects of millennial timescale changes in precipitation and vegetation
123 cover that can impact topography. Superimposed on the effects documented in this study would be the effects of seasonal
124 changes in precipitation and vegetation cover, subcatchment variations in vegetation cover, transport limited fluvial and
125 vegetation interactions, stochastic variations in precipitation in different climate zones. Consider of the previous, more
126 detailed, aspects of precipitation-vegetation interactions on erosion could be independent studies of their own and can not
127 be covered in a single study. Thus, the modeling approach and results of this study should be considered as documenting
128 the longer (millennial) timescale climate and vegetation forcings on fluvial and surface processes.

129 **3. Methods**

130 **3.1. Model Description and governing equations**

131 For this study, we use the open-source model framework Landlab (Hobley et al., 2017). We chose a model-domain with
132 an area of 100 km² which is implemented as a rectangular grid, divided into 0.01km² spaced grid cells. For simplification
133 in the presentation of results, we present our results for the driest, northern most area (Parque Nacional Pan de Azucar)
134 and for the Parque Nacional La Campana. La Campana is situated at 32°S Latitude (Fig.1) and shows the highest values
135 in analyzed basin metrics (Fig. 2), although the general behavior and results presented here are representative of the other
136 two areas not shown. The topographic evolution of the landscape is a result of tectonic uplift and surface processes,
137 incorporating detachment limited fluvial erosion and linear diffusive transport of sediment across hillslopes (Fig.3). These
138 processes are linked to, and vary in their effectiveness due to surface vegetation density. Details of the implementation of
139 these processes into Landlab are explained in the following subsections.

140 This model setup is simplified in regards to hydrological parameters such as soil moisture and groundwater and
141 unsaturated zone flow. Also, the erosion and transport of material due to mass-wasting processes such as rockfalls and
142 landslides are not considered. We argue that those processes do not play a major role in the basins we used for model-
143 calibration and that the processes acting continuously along hillslopes and channels have the largest impact on shaping
144 our reference landscapes. The detachment-limited approach was chosen because the focus areas represent small, bedrock
145 dominated headwater catchments. Additional caveats and limitations of the modeling approach used are discussed in
146 Section 5.4. Main model parameters used in the model (and described below) are provided in Table 1.

147 **3.2. Boundary and Initial Conditions, and Model Free Parameters**

148 In an effort to keep simulations comparable, we minimized the differences in parameters between simulations. The
149 exceptions to this include the surface vegetation cover and mean annual precipitation, which were varied between
150 simulations. One of the main controls on topography is the rock uplift rate. We kept the rock uplift rate temporally and
151 spatially uniform across the domain and at 0.2 mm/yr (Table 1). Studies of the exhumation and rock uplift history of the

Coastal Cordillera, Chile, are sparse at the latitudes investigated here, but existing and in progress studies further to the north are broadly consistent with the rock uplift rate used here (Juez-Larré et al., 2010; Avdievitch et al., 2017).

The EarthShape focus sites are situated in similar granitic lithologies (Oeser et al., 2018), thereby allowing the assumption that the same critical shear-stress, baseline diffusivity, and fluvial erodibility can be used.

Vegetation cover was chosen to be spatially uniform across model domains. While vegetation can change in high-relief catchments due to precipitation and temperature changes with elevation, this simplifying assumption was made based on the low to moderate relief (500-1500m, mean ~750m) of the Coastal Cordillera areas investigated, and minimal field and MODIS observed changes in type and cover with elevation. The exception to this La Campana study area (~1,500 m relief) which has an observed change in vegetation type and cover in the upper 500 m of the catchment. Furthermore, dynamic vegetation modeling results presented in the companion paper to this (Fig. 5b in Werner et al., 2018 - this volume) indicate that although elevation gradients in plant functional types occur in the region since the last glacial maximum, the elevation range of the catchments in the Coastal Cordillera (<1500 m) exhibits only minor changes with elevation. Vegetation cover near trunk streams within catchments is observed in the field to increase, most likely due to local scale hydrology and more abundant water in these areas. However, these regions are often restricted to within 10's of meters of the trunk stream, well below the 100m grid resolution of the model, and therefore difficult to accurately resolve within the simulations presented.

The initial topography used in our simulations was a random white-noise topography with <1 m relief. To avoid unwanted transients related to the formation of this initial topography we conduct simulations to produce an equilibrium topography for each set of the different climate and vegetation scenarios (see below). These equilibrium topographies were produced by running the model for 15 Myr until a topographic steady-state is reached. The equilibrium topography after 15 Myr was used as the input topography for subsequent experiments that impose transient forcings in climate, vegetation, or both (Fig. 4). The model simulation time shown in subsequent plots is the time since completion of this initial 15 Myr steady-state topography development. In the results section, we present these results starting with differences in the initial steady-state topographies (prior to imposing transient forcings) and then add different levels of complexity by imposing either: (1) a single transient step-change for the vegetation cover (Fig. 4b); (2) a step change in the mean annual precipitation (Fig. 4d); (3) 100 kyr oscillations in the vegetation cover (Fig. 4a); (4) 100 kyr oscillations in the mean annual precipitation (Fig. 4c); or (5) 100 kyr oscillation in both the vegetation cover and mean annual precipitation (both Fig. 4a and 4c). This approach was used to produce a stepwise increase in model complexity for evaluating the individual, and then combined, effects of fluvial and hillslope processes to different forcings.

The magnitude of induced rainfall transient forcings were based upon the present-day conditions along the Coastal Cordillera study areas (Fig. 1b, c). The step change and oscillations in vegetation cover and mean annual precipitation imposed on the experiments were designed to investigate vegetation and precipitation change effects on topography over the last ~0.9 Ma, the period during which a 100 kyr orbital forcing is dominant in Earth's climate (Broecker & van Donk, 1970; Muller and MacDonald, 1997). Given this timescale of interest, we impose a 10% magnitude change in the step-increase or decrease, or the amplitude change in oscillations for the vegetation cover. This magnitude of vegetation cover change is supported by dynamic vegetation modeling of vegetation changes over glacial-interglacial cycles in Chile (see companion paper by Werner et al. 2018-this journal) and to some degree elsewhere in the world (Allen et al, 2010, Prentice et al. 2011, Huntley et al. 2013), however for the sake of simplicity we use a fixed forcing of +/-10% for all simulations and not a spatially variable forcing which would be dependent on ecosystem behaviour for each separate area. We assume

that the present-day conditions of combined vegetation cover and mean annual precipitation along the north-south gradient of the coastal cordillera are directly linked (Fig. 1b, c), and therefore follow an empirical approach based on the present day mean annual precipitation which directly links to the present-day vegetation cover in Chile (Fig. 5). We do this by associating each 10% change in vegetation cover (dV) with a corresponding change of mean annual precipitation (dP, Fig. 5) present in the study areas considered. We impose a predefined, fixed amplitude, change in surface vegetation cover as a transient forcing for simulations. For our prescribed changes in vegetation cover we then choose corresponding values of mean annual precipitation based on the relationship shown in figure 1 and 5. The simulations were parameterized in terms of changes in vegetation cover (instead of precipitation) for two reasons. First, the emphasis of this study is on advancing our knowledge of how vegetation changes impact surface processes. Given this, we wanted to present results based on reasonable changes in vegetation cover change. Second, results from the companion paper to this one (Werner et al., 2018) suggest the Chilean Coastal Cordillera experiences +/- 10% changes in vegetation covers over the last 21 kyr. We adopt this result in this study as the current best estimate for the changes in the study areas considered. Thus, the changes in precipitation and vegetation imposed in this study are empirically based on observations from the climate and ecological gradient in the Coastal Cordillera.

The boundary conditions used in the model were the same for all simulations explained above (Fig. 3). One boundary was held at a fixed elevation and open to flow outside the domain. The other three were allowed to increase in elevation and had a zero-flux condition. This design for boundary conditions is similar to previous landscape evolution modeling studies (Istanbulluoglu and Bras, 2005) and provides a means for analyzing the effects of different vegetation cover and precipitation forcings on the individual catchment and subcatchment scale.

3.3. Vegetation Cover Dependent Geomorphic Transport Laws

The governing equation used for simulating topographic change in our experiments follows the continuity of mass. Changes in elevation at different points of the model domain over time $dz(x,y,t)$ depend on

$$\frac{\delta z(x,y)}{\delta t} = U - \frac{\delta z}{\delta t}|_{hillslope} - \frac{\delta z}{\delta t}|_{fluvial} \quad (1)$$

where z is elevation, x , y are lateral distance, t is time, U is the rock uplift rate, $\frac{\delta z}{\delta t}|_{hillslope}$ is the change in elevation due to hillslope processes, $\frac{\delta z}{\delta t}|_{fluvial}$ is the change in elevation due to fluvial processes (Tucker et al., 2001a).

3.3.1 Vegetation Cover Influenced Diffusive Hillslope Transport

The change in topography in a landscape over time caused by hillslope-dependent diffusion can be characterized as:

$$\frac{\delta z}{\delta t}|_{hillslope} = -\nabla q_{sd} \quad (2)$$

Landscape evolution models characterize the flux of sediment q_s either as a linear or non-linear function of surface slope S (Culling, 1960; Fernandez and Dietrich, 1997). In order to keep the number of free parameters for the simulation to a minimum, we used the linear description of hillslope diffusion:

$$q_{sd} = K_d S \quad (3)$$

Following the approach of (; Alberts et al., 1995; Dunne, 1996; Istanbulluoglu and Bras, 2005; Dunne et al., 2010), we assign the linear diffusion coefficient K_d as a function of surface vegetation density V , an exponential coefficient α , and a baseline diffusivity K_b , such that:

$$K_d = K_b e^{-(\alpha V)} \quad (4)$$

3.3.2 Vegetation Cover Influence on Overland Flow and Fluvial Erosion

Fluvial detachment-limited erosion of material due to water is calculated in this study by the widely-used stream-power-equation (Howard and Kerby, 1983; Howard et al., 1994; Whipple and Tucker, 1999; Braun and Willet, 2013):

$$\frac{\delta z}{\delta t}|_{fluvial} = k_e(\tau - \tau_c)^p \text{ for } \tau > \tau_c \quad (5)$$

In this equation k_e represents the erodibility of the bed, τ is the bed shear stress which acts on the surface at each node, τ_c is the critical shear stress which needs to be overcome to erode the bed-material and p is a constant.

By following the approach of Istanbulluoglu and Bras (2005) and Istanbulluoglu et al. (2004), we reformulate the standard equation of shear-stress $\tau_b = \rho_w g R S$, where ρ_w is the density of water, g is the acceleration of gravity, R is the hydraulic radius and S is the local slope, to a form which incorporates Manning's roughness to quantify the effect of vegetation cover on bed shear stress (Willgoose et al., 1991, Istanbulluoglu et al., 2004):

$$\tau_v = \rho_w g (n_s + n_v)^{\frac{6}{10}} q^m S^n F_t \quad (6)$$

Here n_s and n_v represent Manning's numbers for bare soil and vegetated ground, q is the water-discharge per node which is approximated with the steady-state uniform precipitation per timestep P and the surface area per node A ($q = A * P$) and S is the local slope per node, m and n are constants. n_v for each node is calculated as a function of the local surface vegetation cover

$$n_v = n_{vr} \left(\frac{V}{V_r}\right)^w \quad (7)$$

with n_{vr} being the Manning's number for a defined reference vegetation cover, V and V_r being the vegetation cover at each node and the reference vegetation cover and w is an empirical scaling parameter.

The last variable in equation 6 represents the shear-stress partitioning ratio F_t (after Foster 1982; Istanbulluoglu and Bras, 2005), which is used to scale the shear-stress at each node to the vegetation-cover present.

$$F_t = \left(\frac{n_s}{n_s + n_v}\right)^{3/2} \quad (8)$$

By combining the formulation for shear stress out of equation 6 with the general stream-power equation 5 we formulate a new factor K_v which represent the bed erodibility per node as a function of surface vegetation cover, which leads to a new expression of fluvial erosion

$$K_v = k_e \rho_w g (n_s + n_v)^{\frac{6}{10}} F_t \quad (9)$$

$$\frac{\delta z}{\delta t}|_{fluvial} = K_v q^m S^n \quad (10)$$

3.4 Model Evaluation

Model performance was evaluated using the above equations and different initial vegetation covers and mean annual precipitation. Our focus in this study is on the general surface process response to different transient vegetation and climate conditions. Given this, topographic metrics of relief, mean slope, and normalized steepness index (K_{sn}) were computed from the model results and compared to observed values from the 30 m SRTM DEM for each of the four areas (Fig. 2). This was done to evaluate if our implementation of the governing equations in Section 3.4 produced topographies within reason of present day topographies in the four Chilean areas. A more detailed model calibration is beyond the scope of this study, and not meaningful without additional observational constraints on key parameters such as latitudinal variations in the rock uplift rate and erosivity. Our aim is not to reproduce the present day topography of the Coastal

263 Cordillera study areas but rather identify the sensitivity and emergent behaviour of vegetation-dependent surface
264 processes gradient of vegetation cover and precipitation in Chile.

265 **4. Results**

266 Our presentation of results is structured around three groups of simulations. These include: 1. steady-state simulations
267 where equilibrium topographies are calculated for different magnitudes of vegetation cover and identical precipitation
268 forcing. A second set of steady-state simulations with the same magnitudes vegetation cover as 1. but with different
269 precipitation forcings corresponding to each vegetation cover (Fig. 5, Section 4.1). 2. Simulations with a transient step-
270 change in either surface vegetation density or precipitation (Section 4.2) that is initiated after the landscape has reached
271 steady state. and 3. simulations with a transient 100 kyr oscillating time series of changing vegetation or precipitation that
272 occurs after the landscape has reached steady state (Section 4.3). For each group of transient simulations, we show the
273 topographic evolution with help of standard topographic metrics and the corresponding erosion rates after the induced
274 change.

275 **4.1 Equilibrium Topographic Metrics**

276 Topographic metrics from each of the four Chilean focus areas (Fig. 1a) were extracted for comparison to equilibrium
277 topographies predicted after 15 Myr of model simulation time. This comparison was done to document the model response
278 to changing vegetation cover (with climate held constant) and changing vegetation cover and precipitation, and also to
279 demonstrate the modeling approach employed throughout the rest of this study captures the general characteristics of
280 different topographic metrics along the Chilean Coastal Cordillera.

281 Analysis of the digital elevation model for each of our four Chilean focus areas illustrates observed changes in catchment
282 relief, slope, and channel steepness (K_{sn}) in relation to the surface vegetation (Fig. 7, red points) and latitude (Fig. 2). The
283 general trend in the observed metrics shows a non-linear increase in each metric until a maximum is reached for regions
284 with 70% vegetation cover. Following this, all observed metrics show a decline towards the area with 100% vegetation
285 cover.

286 The model predicted equilibrium topographies (Fig. 7a,b,c) from four different steady-state simulations with different
287 vegetation cover in each simulation and a constant mean annual precipitation (900 mm/yr) show a nearly linear increase
288 in all observed basin metrics with increasing vegetation cover and therefore do not reflect the overall trend observed from
289 the study areas (red line/symbols). For example, basin relief and slope are both under predicted for simulations with $V <$
290 100% (Fig.7a,b), and only the predicted maximum relief for a fully-vegetated simulation resembles the DEM maximum
291 value. For the normalized channel steepness, only two observed mean values (for $V = 10\%$ and 70%) lie within the range
292 of mean to maximum predicted K_{sn} values (Fig.7c).

293 The resulting equilibrium topographies from simulations with different mean annual precipitation and vegetation cover
294 in each simulation (Fig.7d,e,f) show an improved representation of the general trend of the DEM data. The vegetation
295 cover and precipitation values used in these simulations come from the Chilean study areas (Fig. 1b, c; Fig. 5). In these
296 simulations, the maximum in the observed basin metrics is situated at values of $V = 30\%$ with a following slight decrease
297 in the metric for $V = 30\%$ to $V = 70\%$, followed by a steeper decrease in metrics from $V = 70\%$ to $V = 100\%$. Generally
298 the model-based results tend to underestimate the basin relief and overestimate the basin channel steepness (Fig.7d,f).
299 Variations in basin slope are captured for all but the non-vegetated state (Fig.7e).

Although the above comparison between the models and observations demonstrates a range of misfits between the two, there are several key points worth noting. First, the model results shown are simplified in their setup (e.g. assuming similar rock uplift rate, identical lithology and constants), and assume the present day topography is in steady state for the comparison. Second, despite the previous simplifying assumptions, the degree of misfit between the observations and model are surprisingly small when both variable vegetation and variable precipitation, are considered (Fig. 7d,e,f). Finally (third), the general ‘humped’ shape curve observed in the Chilean areas is captured in the model predictions (Fig. 7d,e,f), with the notable exception that the maximum in observed values occurs at a higher vegetation cover ($V = 70\%$) than the model predictions ($V = 30\%$). Explanations for the possible source of these differences are revisited in the discussion section.

4.2 Transient Topography From a Step Change in Vegetation or Precipitation

The evolution of topographic metrics after a induced instantaneous disturbance (Fig. 4) of either only the surface vegetation cover (Fig. 8, green lines) or only the mean annual precipitation (Fig. 8, blue lines) is analyzed for changes in topographic metrics for either a positive disturbance (Fig. 8a,b,c) or a negative disturbance (Fig. 8d,e,f). This scenario was chosen to analyze and isolate the effects of these specific transient forcings, and are useful for understanding more complex changes in vegetation and precipitation presented later. Mean catchment erosion rates are also analyzed for their evolution after the disturbance (Fig. 9). For simplicity in presentation, results are shown for only two of the four Chilean study areas with initial vegetation (V) and precipitation (P) values for vegetation covers of 10 and 70%, and precipitation rates that correspond to these vegetation covers (i.e. $P(V=10\%)$ or $P(V=70\%)$) (Fig. 5). The results described below show a general positive correlation between all observed topographic metrics and surface vegetation cover and a negative correlation between observed topographic metrics and mean annual precipitation.

4.2.1 Positive Step Change in Vegetation Cover or Precipitation

Topographic Analysis

A positive step change in vegetation cover (V) from $V = 10\%$ to $V = 20\%$ (solid green line Fig. 8a,b,c) leads to a factor of 1.9, 1.42, and 2.1 change in mean basin relief (from 270 m to 520 m), mean basin slope (11.2° to 15.9°), and mean basin channel steepness ($108 \text{ m}^{-0.9}$ to $222 \text{ m}^{-0.9}$), respectively. The adjustment time until a new steady state in each metric is reached is 3.1 Ma. The corresponding positive change in mean annual precipitation (solid blue lines, Fig. 8a,b,c) leads to a decrease of mean basin relief to 176 m, mean basin slope to 8.6° and mean basin channel steepness to $67 \text{ m}^{-0.9}$. This corresponds to a decrease by factors of 1.5, 1.2 and 1.6, respectively. The adjustment time to new steady state conditions in this case are shorter and 1.1Ma (Fig. 8a,b,c). A second feature of these results is the brief increase and then decrease in basin average slope angles following the step change (Fig. 8b).

For simulations with $V = 70\%$ initial surface vegetation cover, a positive increase to $V = 80\%$ leads to an increase of mean basin relief from 418 m to 474 m, mean basin slope from 15.5° to 16.8° and mean basin channel steepness from $172 \text{ m}^{-0.9}$ to $199 \text{ m}^{-0.9}$. This causes an increase in each metric by factors of 1.1, 1.1 and 1.2, respectively. The adjustment time to steady-state conditions is 1.9Ma (dotted green lines, Fig 8a,b,c). The corresponding positive change in mean annual precipitation leads to a decrease of relief to 268 m, decrease in slope to 11.9° and decrease of channel steepness to $105 \text{ m}^{-0.9}$. This resembles a decrease by factors 1.5, 1.3, 1.6, respectively. Adjustment time in this case is 1.7Ma (dotted blue lines, Fig. 8a,b,c). The basin slope data shows similar behavior as the $V_{\text{ini}} = 10\%$ simulations with an initial decrease and then increase for a vegetation cover step change and an initial increase and then decrease for a step change in mean

annual precipitation. Comparison of the change in the topographic metrics for the low ($V=10\%$) and high ($V=70\%$) initial vegetation covers, the magnitude of change in each metric is larger when the step change occurs on a low, rather than higher, initial vegetation cover topography.

Erosion Rate Changes

The model results show a negative relationship between increases in vegetation cover and erosion and a positive relationship between increases in precipitation and erosion (Fig. 9). Although the response between the disturbances and changes in erosion rates are instantaneous, the maximum or minimum in the change is reached after some lag time and the magnitude and duration of non-equilibrium erosion rates varies between different simulation setups.

For initial vegetation cover of $V = 10\%$, a change in vegetation cover (dV) of $+10\%$ leads to a decrease in erosion rates from 0.2 to 0.03 mm/yr (factor of 5.7 decrease, Fig. 9a green line). The minimum erosion rate is reached 43.5 kyrs after the step change occurs. Following this minimum in erosion rates, the rates increase until the steady-state erosion rate is reached after the adjustment time. An increase in mean annual precipitation corresponding to a vegetation cover of 10% (i.e. $P(V=10\%)$ to $P(V=20\%)$; Fig. 5) leads to an increase in erosion rates to a maximum of 0.44 mm/yr after 74.8 kyrs (factor of 2.2 increase, Fig.9a, blue line). For initial vegetation of $V = 70\%$ a vegetation increase of $dV = +10\%$ results in minimum erosion rates of 0.14 mm/yr after 117.7 kyrs (factor of 1.4 decrease, Fig. 9b, green line). A corresponding increase in precipitation for these same vegetation conditions leads to maximum erosion rates of 0.44 mm/yr after 107.5 kyrs which is an increase by a factor of 2.2 (Fig.9b, blue line). The previous results for a positive step change in vegetation or precipitation demonstrate that the magnitude of change in erosion rates is larger for changes in precipitation rate than for vegetation cover changes, and in low initial vegetation cover settings ($V=10\%$) the magnitude of change in erosion rates for changing vegetation is larger (compare green lines Fig. 9a with 9b).

4.2.2 Negative Step Change in Vegetation Cover or Precipitation

Topographic Analysis

For negative step-changes in vegetation (green curves, Fig. 8d,e,f), the results show a sharp decrease in topographic metrics associated with shorter adjustment times compared to the positive step change experiments (compare Fig. 8d,e,f with a,b,c). For step changes in precipitation (blue curves, Fig. 8d,e,f), the increase of topographic metrics happens slower and therefore with longer adjustment times. A negative step change in vegetation cover from $V = 10\%$ by $dV = -10\%$ leads to a decrease of mean basin relief from 269 m to 35 m, mean basin slope from 11.2° to 2.3° and mean basin channel steepness from $108 \text{ m}^{-0.9}$ to $11 \text{ m}^{-0.9}$ which resembles decreases by factors of 7.8 , 3.8 and 9.6 , respectively. The adjustment time until a new steady-state is reached is 0.26 Ma (solid green lines, Fig. 8d,e,f). The corresponding negative change in precipitation leads to an increase in mean basin relief to 512 m, mean basin slope to 15.8° and mean basin channel steepness to $223 \text{ m}^{-0.9}$. These increases reflect changes by factors of 1.9 , 1.4 and 2.1 with an adjustment time of 4.9 Ma (dotted green lines, Fig.8d,e,f). Mean basin slope results (Fig. 8e) for a step change in vegetation illustrate a pulse-like feature of initially increasing slope values, followed by a decrease to lower slope values. In contrast, a negative step change in precipitation induce an initial decrease in slope, followed by a gradual increase in slope to a value higher than was initially observed before the change.

Simulations with initial vegetation cover $V = 70\%$ and $dV = -10\%$ show a decrease in mean basin relief from 418 m to 356 m, mean basin slope from 15.5° to 13.6° and mean basin channel steepness from $172 \text{ m}^{-0.9}$ to $144 \text{ m}^{-0.9}$ which resembles

377 changes by factors of 1.2, 1.1 and 1.2 and an adjustment time of 2.1Ma (dotted green lines, Fig.8d,e,f). Corresponding
378 negative changes in precipitation lead to increase of basin relief of 465 m, basin slope to 16.4° and channel steepness to
379 $195\text{m}^{-0.9}$ which resembles changes by factors of 1.1 for all three values. Adjustment time in this case is 2.2 Ma (dotted
380 blue lines, Fig.8d,e,f). Behavior of mean basin slope after the step-change follows the $V = 10\%$ simulations but shows
381 lower amplitudes of basin slope for both step-changes in vegetation and precipitation.

382

383 **Erosion Rates**

384 The positive step-change results (Fig. 9a, b) indicated that erosion rates reach their minimum or maximum with a lag
385 time, and show significant differences in the magnitude and duration of non-equilibrium conditions depending on if
386 vegetation or precipitation were changing. Simulations with a decrease from $V = 10\%$ to $V = 0\%$ (Fig. 9c) show a sudden
387 increase in erosion rates to a maximum value of 3.5 mm/yr which is an increase from steady state conditions by a factor
388 of 17.7 which is reached after 19.5 kyrs (green line, Fig.9c). A step decrease in precipitation for this corresponding
389 vegetation difference (i.e. $P(V=10\%)$ to $P(V=0\%)$) leads to a smaller, and protracted (longer adjustment time) decrease
390 in erosion rates to 0.03 mm/yr after 50.1 kyrs. These conditions cause a factor of 5.6 decrease (blue line, Fig. 9c).
391 Simulations of $V = 70\%$ with a vegetation change of $dV = -10\%$ show an increase in erosion rates to 0.27 mm/yr which
392 is a factor of 1.4 increase after 126.3 kyrs (Fig. 9d). For the corresponding decrease in precipitation the data show a
393 decrease in erosion rates to 0.15mm/yr after 124.5 kyrs. This resembles change by factor of 1.2 (blue line, Fig. 9d).

394 **4.3 Transient Topography – Oscillating**

395

396 **4.3.1 Oscillating Surface Vegetation Cover, Constant Precipitation**

397 **Topographic Analysis**

398 The topographic evolution in simulations with a constant precipitation (10 and 360mm/yr for $V=10\%$, and $V=70\%$,
399 respectively) and oscillating vegetation cover show a different response than the previous step change experiments. The
400 differences depend on the initial steady-state vegetation cover prior to the onset of 100kyr oscillations. All observed basin
401 metrics (Fig. 10) show an initial oscillating decrease in values until a new dynamic steady-state is reached where the
402 amplitude in oscillations is less than in the preceding initial adjustment period. Simulations with $V = 10\%$ (Fig. 10a) show
403 a factor of 2.5 decline in the basin relief (from 269 m to 107 m). For simulations with $V = 70\%$ the reaction and adjustment
404 to the new dynamic steady-state is less pronounced with a factor of 1.01 decline in relief (from 410m to 407m) with
405 positive and negative amplitudes in dynamic steady-state of 1.6 m. While these changes are unmeasurable in reality, they
406 highlight that for high initial vegetation cover settings that changes in only vegetation cover would be difficult to detect.
407 Analysis of mean basin slope for the model topographies with low ($V=10\%$) vegetation shows a similar behavior with a
408 factor of 1.6 decrease of the mean slope (from 11.2° prior to the onset of oscillations, to 6.0°, Fig. 10b). However, before
409 this new equilibrium is reached, the slopes show an increase in mean slope for the first two periods of vegetation
410 oscillation which then declines towards the new long-term stable dynamic equilibrium which is reached after
411 approximately 500 kyrs. Local maxima of mean basin slope coincide with local minima in basin relief. For the $V = 70\%$
412 simulations, the reaction is significantly smaller with no change in mean slope for the new dynamic equilibrium and
413 amplitudes of both positive and negative of 0.16°. Mean basin channel steepness (Fig. 10c) reflects the behavior of mean
414 basin elevation. For $V = 10\%$ simulations the mean channel steepness decreases by a factor of 2.7 (from $108\text{ m}^{-0.9}$ to 40

415 $m^{-0.9}$) with a positive amplitude of $3.7m^{-0.9}$ and a negative amplitude of $6.1 m^{-0.9}$. For $V = 70\%$ simulations the response is
 416 again only minor, compared to the lower initial vegetation cover simulations with a change of mean channel steepness
 417 from $186 m^{-0.9}$ to $167 m^{-0.9}$ and positive amplitudes of $1.1 m^{-0.9}$ and negative amplitudes of $0.9 m^{-0.9}$. Like the elevation
 418 data, the steepness data shows a distinct oscillating pattern with a slow increase to local maxima and rapid decreases to
 419 local minima which coincide with maxima/minima of elevation data. Taken together, the previous observations
 420 demonstrate a larger change in topography for oscillations in poorly vegetated areas compared to those with higher
 421 vegetation cover. Furthermore, the magnitude of topographic change that oscillations in vegetation impose on topography
 422 are largest in the first ~ 500 kyr after the onset of an oscillation, and diminish thereafter.

423

424

425

426 **Erosion Rates**

427 The erosion history for simulations with oscillating vegetation cover (Fig. 11) demonstrate large variations in the erosion
 428 rate that depend on the average vegetation cover of the oscillation. Furthermore, pronounced differences in the amplitude
 429 of erosion occur if the vegetation cover is above or below the mean of the oscillation (Fig. 4a). More specifically,
 430 simulations with $V = 10\%$ show a pattern of a small decrease in erosion rates (from 0.2 to 0.03 mm/yr) when vegetation
 431 cover increases above the mean cover, in contrast to a large increase in erosion rates (up to 3.3 mm/yr) when vegetation
 432 cover decreases below the mean of the oscillation (Fig. 2, Fig. 11). Maximum erosion rates decline over multiple periods
 433 of oscillation until they reach a dynamic steady-state with maximum rates (1.2mm/yr) at 760kyrs after the onset. Time
 434 periods of higher erosion rates (>0.2 mm/yr) have a mean duration of 28kyrs, whereas periods of lower erosion rates (<0.2
 435 mm/yr) have a mean duration 72kyrs. For simulations with high vegetation cover ($V = 70\%$) the maximum and minimum
 436 erosion rates are 0.28 and 0.15 mm/yr, respectively. The magnitude of maximum and minimum erosion rates are not
 437 significantly time-dependent and are therefore constant over the simulation. The mean duration of periods with higher
 438 erosion rates ($> 0.2mm/yr$) is 55 kyrs whereas the duration for periods with lower rates (< 0.2 mm/yr) is 45 kyrs. These
 439 results demonstrate that areas with low vegetation cover experience not only larger amplitudes of change in erosion rates,
 440 but also an asymmetric change whereby decreases in erosion rates are lower magnitude than the increases in erosion rates.

441

442

443 **4.3.2 Oscillating precipitation, Constant Vegetation**

444 **Topographic Analysis**

445 The evolution of topographic parameters for simulations with oscillating mean annual precipitation and two different
 446 constant surface vegetation covers ($V=10$ or 70% , Fig. 12) show a less extreme and smaller temporal change in erosion
 447 rate to variations in precipitation compared to the previous discussed effects of oscillating vegetation cover (Fig. 11). In
 448 Fig. 12a, the mean basin relief results for $V = 10\%$ and oscillating precipitation show small variations (+4.9 m to -3.8 m)
 449 in relief around a mean of 269 m, which is similar to the mean relief prior to the onset of oscillations at 5,000kyrs. For
 450 simulations with $V = 70\%$ the change in relief is slightly more pronounced with factor of 1.1 adjustment to a new mean
 451 (380 m from 418 m) in steady-state conditions. The evolution of topographic slope (Fig. 12b) for $V = 10\%$ simulations
 452 shows a factor of 1.05 adjustment to a new dynamic equilibrium (from 11.2° to 10.6°). For $V = 70\%$ the mean slope
 453 values do not significantly change from steady-state to transient conditions . Mean channel steepness (Fig. 12c) for

V=10% shows a factor of 1.01 adjustment (from $108 \text{ m}^{-0.9}$ to $110 \text{ m}^{-0.9}$), and would be difficult to measure in reality. The amplitude of oscillation is $4 \text{ m}^{-0.9}$ for both negative and positive amplitudes. For $V_{\text{ini}} = 70\%$ simulations a factor 1.1 change in channel steepness occurs (from $171 \text{ m}^{-0.9}$ to $152 \text{ m}^{-0.9}$) with amplitudes of $4.5 \text{ m}^{-0.9}$ for both positive and negative changes. Figure 12 illustrates changes in topographic metrics that result from oscillations in precipitation occurring around vegetation covers of 10 and 70%. These changes are significantly smaller than those predicted for constant precipitation, but oscillating vegetation conditions (Fig. 10).

460

461 **Erosion Rates**

462 Predicted erosion rates from simulations with constant surface vegetation cover and oscillating mean annual precipitation
463 indicate different amplitudes of change around the mean erosion rate depending on the vegetation cover. For simulations
464 with $V = 10\%$ (Fig. 13, blue solid line) erosion rates oscillate symmetrically around the steady-state erosion rate
465 (0.2 mm/yr). The maximum and minimum erosion rates of 0.42 and 0.01 mm/yr , respectively, do not lead to a shift in the
466 mean value of erosion rates over time.. In contrast, predicted rates with a higher vegetation cover of $V = 70\%$ (Fig. 13,
467 blue dotted line) demonstrate an asymmetric oscillation in rates around the mean, whereby the maximum in rates (0.43
468 mm/yr) has a larger difference above the mean rate, than do the minimums in the oscillation (0.12 mm/yr). For this higher
469 vegetation cover scenario, a gradual decrease in the mean erosion rate
470 while time progresses occurs. Furthermore, the maximum and minimum erosion rates decline over several oscillation
471 periods. Taken together, these results indicate that oscillations in precipitation impact erosion with different magnitude
472 depending on the amount of vegetation cover. Areas with low vegetation cover demonstrate the highest and symmetric
473 oscillation of erosion rates due to changes in precipitation whereas in areas with high vegetation cover the effect of
474 negative changes in precipitation is dampened by vegetation.

475 **5. Discussion**

476 The previous results highlight different sensitivities to changes in either surface vegetation cover or mean annual
477 precipitation. In the following, we synthesize the previous results and then build upon them to discuss the scenario of
478 synchronous variation in both precipitation and vegetation cover.

479 **5.1 Interpretation of Steady-State Simulations**

480 Landscapes in a topographic steady-state show distinct features in topographic metrics that are widely used to estimate
481 catchment-averaged erosion rates and therefore the leading processes of erosion within a landscape (DiBiase et al., 2010).
482 The steady-state simulations presented can reproduce (Fig. 7) variations in topographic metrics over different climate and
483 vegetation states seen in other studies (Langbein & Schumm, 1958, Walling and Webb 1983). Comparison of simulations
484 with homogeneous precipitation and changing values of vegetation cover (Fig. 7a,b,c) to simulations with both changing
485 precipitation and vegetation cover (Fig. 7d,e,f) indicates we can only reproduce a similar trend with a distinct peak in
486 topographic metrics when both variable precipitation and vegetation cover are considered. From this, we conclude that
487 modern model-based landscape evolution studies that aim to compare areas with different climates should incorporate
488 vegetation dynamics in their simulations. Misfits between the predicted and Chilean observed topographic metrics (Fig.
489 7d,e,f) present when the vegetation and precipitation both vary likely stem from the simplicity of the model setup used
490 and the likelihood of differences of the rock uplift rate and lithology's present in these areas.

5.2 Interpretation of Step-Change Experiments

Our analysis shows that changes in vegetation-cover typically have a higher magnitude of impact on topographies for lower values of initial vegetation cover, compared to simulations with high initial vegetation cover (Fig. 8, 9). In those settings the influence of vegetation cover outweighs the influence of precipitation in cases of negative and positive directions of the step change. The reason for this is due to a higher impact of changes in vegetation on erosivity and diffusivity (parameter K_v , K_d ; equation 4, 10) than changes in precipitation.

Furthermore, a negative step change in vegetation cover impacts the topographic metrics by a factor of two more than do positive step change changes (Fig. 8d,e,f). This response is interpreted to be due to the non-linear reaction of diffusivity and fluvial erodibility to changes in vegetation cover (See Fig.6). Negative changes in vegetation cover lead to a higher overall change in diffusivity and erodibility compared to positive step-changes.

Model results for the topographic metrics and erosion rates also indicate a difference in the adjustment times of the system until a new steady state is reached when either precipitation or vegetation cover changes (Figs. 8, 9). For simulations with positive step-changes (Fig. 8a,b,c) the adjustment time for changes in vegetation cover to reach a new equilibrium in topographic metrics or erosion rates is three times longer than the adjustment time for changes in precipitation. Simulations with a negative step-changes in vegetation cover show an adjustment time which is shorter by a factor of 18 compared to negative changes in precipitation. This difference in adjustment time again is a result of the non-linear behavior of erosion parameters K_d and K_v which influence how effective a signal of increasing or decreasing erosion can travel through a river basin (Perron et al., 2012). High values of K_d and K_v are associated with lower adjustment times and are a result of negative changes in vegetation cover. The influence of changing precipitation on adjustment time behaves in a more linear fashion and therefore mostly depends on the overall magnitude of change. Therefore, positive step-changes in vegetation cover decrease K_d and K_v which leads to higher adjustment times than the corresponding changes in precipitation.

An increase and then decrease, or decrease and then increase, in predicted slope and erosion rates is observed for both the positive and negative step changes experiments (Fig. 8b,e; and Fig. 9). This non-linear response in both positive and negative step changes in precipitation and vegetation cover is also manifested in the subsequent oscillation experiments, but most clearly identifiable in the step change experiments. The explanation for this behavior is as follows. A positive step change in vegetation cover (Fig. 8b) leads to a decrease in fluvial capacity because increased vegetation cover increases the Manning's roughness (parameter n_v , equation 8). The effect of changing the Manning's roughness varies with the location in the catchment and influences which processes (fluvial or hillslope) most strongly influence slopes and erosion rates. In the upper part of catchments where contributing areas (and discharge) are low, this increase in Manning's roughness causes many areas to be below threshold conditions such that fluvial erosion is less efficient, and hillslope diffusion increases in importance and reduces slopes. In the lower part of catchments, where contributing area and discharge are higher, changes in the Manning's roughness are not large enough to impact fluvial erosion because these areas remain at, or above, threshold conditions for erosion. With time, the lower regions of the catchments that are at or above threshold conditions propagate a wave of erosion up to the higher regions that are below threshold conditions. The propagating wave of erosion eventually leads to increase in slope angles, essential due to the response time of the fluvial network to adjust to the new Manning's roughness conditions.

In contrast, a positive step change in mean annual precipitation leads to an initial increase in fluvial shear stress which initially causes headward incision of river channels and leads to wave of erosion that propagates upstream and increases channel slope values (Fig. 8b, see also e.g. Bonnet and Crave, 2003). The increase in channel slopes leads to an increase

531 in the hillslope diffusive flux adjacent to the channels that then propagates upslope. Eventually, this increase in hillslope
532 flux leads to a decrease in hillslope angles, and an overall reduction in mean catchment slopes after the systems reaches
533 equilibrium.

534 Negative step-changes in vegetation cover or precipitation (Fig. 8e, green curves) shows the opposite behavior of the
535 previous positive step change description. A negative step change in vegetation cover leads to an initial increase of fluvial
536 erosion everywhere in the catchment because the Manning's roughness decreases everywhere. This catchment wide
537 decrease in Manning's roughness leads to fluvial incision everywhere in the catchment and an increase in mean slope.
538 However, eventually hillslope processes catch up with increased slopes near the channels and with time an overall
539 reduction of slope occurs. Negative changes in precipitation (Fig. 8e, blue curves) lead to an initial decrease in fluvial
540 erosion thereby leading to an increase in the significance of hillslope processes such that slope angles between channel
541 and ridge decrease as hillslope processes fill in channels. With time, the fluvial network equilibrates to lower precipitation
542 conditions by increasing slopes to maintain equilibrium between erosion and rock uplift rates.

543 Thus, the contrasting behavior of either initially increasing or decreasing slopes and erosion rates, followed by a change
544 in the opposite direction of this initial change highlight a complicated vegetation-climate induced response to changes in
545 either parameter. This non-linear behavior, and the millennial timescales over which these changes occur, suggest that
546 modern-systems that experienced past changes in climate and vegetation will likely be in a state of transience and the
547 concept of a dynamic equilibrium in hillslope angles and erosion rates may be difficult to achieve in these natural systems.
548 Previous studies have inferred relationships between mean catchment erosion rates derived from cosmogenic
549 radionuclides and topographic metrics (e.g., DiBiase 2010; DiBiase and Whipple 2011). However, the previous discussion
550 of how topographic metrics change in response to variable precipitation and vegetation suggest that empirical
551 relationships between erosion rates and topographic metrics contain a signal of climate and vegetation cover in the
552 catchment. We illustrate the effect of step changes in climate and vegetation on the new steady-state of topographic
553 metrics in figure 14. In this example, the new steady state conditions in basin relief and mean slope after a modest (+/-
554 10%) change in vegetation or precipitation (triangles) differ from the initial steady-state condition (circles). These
555 changes in topographic metrics when the new steady-state is achieved occur despite the rock uplift rate remaining
556 constant.

557 **5.3 Interpretation of Oscillation Experiments**

558 The results from the 100 kyr oscillating vegetation and precipitation conditions shows that oscillating vegetation cover
559 without the corresponding oscillations in precipitation leads to adjustments of topographic features, to a new dynamic
560 equilibrium after approximately 1.5 Ma (Figs. 10, 11). The previously described response of topographic metrics and
561 erosion rates to oscillating vegetation (see results section) are due to processes described in the previous step change
562 experiments. For example, the asymmetric oscillations in topographic metrics for $V=10\%$ (Fig. 10) are due to the
563 superposition of positive, then negative changes described in section 5.2. Variations in the imposed Manning's roughness,
564 and relative strengths of fluvial vs hillslope processes in different parts of the catchments at different times causes the
565 topographic metrics and erosion rates to have a variable amplitude and shape of response from the symmetric oscillations
566 imposed on the topography (Fig. 4a).

567 Simulations with oscillating precipitation and constant vegetation cover however show a less pronounced shift to new
568 equilibrium conditions and lower amplitudes of oscillation in both topographic metrics and erosion (Figs. 12, 13). This
569 difference in the response of the topographic metrics and erosion rates shown in figures 12 and 13, compared to the

oscillating vegetation cover experiments (Figs 10, 11), is due to a higher impact of changes in vegetation cover on parameters which guide erosion rates and therefore adjustment to topographic metrics compared to the \corresponding changes in precipitation in our model domains (Fig. 5). Especially for simulations with low initial vegetation cover the effect of changing vegetation has a larger magnitude effects because of the non-linear response of diffusivity and fluvial erodibility to changes in vegetation cover compared to the linear response to changes in precipitation.

5.4 Coupled Oscillations in Both Vegetation and Precipitation

The previous sections present a sensitivity analysis of how step changes or oscillations in either vegetation cover or precipitation influence topography. Here we present a step-wise increase towards reality by investigating the topographic response to changes in both precipitation and climate at the same time. The amplitude of change prescribed for both precipitation and vegetation is based upon the present empirical relationship observed in the Chilean study areas for initial vegetation covers of 10 and 70%, and mean annual precipitations for 10 and 360 mm/yr (Fig. 5). As with the previous experiments, oscillations in parameters were imposed upon steady-state topography that developed with the previous values, and a rock uplift rate of 0.2 mm/yr.

Figure 15 shows the evolution of topographic metrics for simulations with combined oscillations in precipitation and vegetation. The variation in topographic metrics resembles those described for simulations with constant vegetation cover and oscillating climate by showing little to no significant adjustment towards new dynamic steady-state conditions. The amplitudes of oscillation are dampened from those of previous results because of the opposing effects of changes in precipitation and vegetation cover (e.g. compare blue and green curves in Figs. 8 and 9).

However, inspection of the predicted erosion rates (Fig. 16) for the combined oscillations indicates a significant (~ 0.1 ; ~ 0.15 mm/yr), and highly non-linear response. The response between the 70% and 10% vegetation cover scenarios are very different such that for heavily vegetated areas ($P(V=70\%)$) erosion rates typically increase during an oscillation, whereas for the low vegetation cover conditions ($P(V=10\%)$) erosion rates initially show a decrease, and then an increase and decrease at a higher frequency.

To better understand this contrast in the response to combined precipitation and vegetation changes, the first 100 kyr cycle is shown in figure 17. After an oscillation starts, the 10% initial vegetation cover simulations show a decline in erosion rates with the minimum erosion rate correlated with highest values of both vegetation cover and mean annual precipitation (compare top and bottom panels). This part of the response is interpreted as resulting from the hindering effect of increased vegetation on erosion rates outweighing the impact of higher precipitation on erosion rates (Fig. 17) because vegetation decreases the effectiveness of erosion and transport of surface material. After values of vegetation cover and precipitation start to decline, erosion rates show a very rapid increase to values of ~ 0.3 mm/yr. This increase in erosion rates is due to an increase in both K_v and K_d (Fig. 3b, equations 4, 5) which outcompetes the effect of precipitation decrease.

Following this, a sudden drop in erosion rates to 0 mm/yr occurs and lasts for 3 kyrs due to the onset of hyper arid conditions at minimum precipitation. After this low in erosion rates, they increase again (to 0.3 mm/yr) as precipitation and vegetation cover increase while the effect of increased precipitation outweighs the effect of the non-linear decrease in K_v and K_d (Fig. 3b, c; equations 4, 5). Finally, at the end of this complex cycle a decrease in erosion rates occurs (Fig. 17b) while vegetation and precipitation are increasing (upper panel) because the effect of vegetation increases K_v/K_d and outweighs the effect of increasing precipitation.

608 Lastly, a clearly different behavior in erosion rates occurs for settings with higher vegetation cover (e.g. $P(V=70\%)$, Fig.
609 17) compared to the previous lower vegetation cover scenarios. As the vegetation cover and precipitation increase (Fig.
610 17A) in the first half of the 100 kyr cycle, the erosion rates increase (to 0.35 mm/yr). This is due to the increase in
611 precipitation which outcompetes the decline in vegetation influenced erosivity/diffusivity parameters K_d and K_v .
612 Following this, when vegetation cover and precipitation decrease in the second half of the cycle, little to no change occurs
613 in the erosion rates. This near static behavior in erosion rates while precipitation and vegetation cover decrease is due to
614 an equilibrium between the negative effect on erosion rates for decreasing precipitation and the positive effect on erosion
615 rates for decreasing vegetation cover.

616 In summary, the non-linear shape of the vegetation dependent erosivity (K_v) and hillslope diffusivity (K_d) in combination
617 with linear effects of mean annual precipitation on erosion rates, exert a primary control on the direction and magnitude
618 of change in catchment average erosion rates. Despite a simple oscillating behaviour in precipitation and vegetation
619 cover, a complex and non-linear response in erosion rates occurs. In Fig. 18 we depicted the conceptual end-members of
620 landscape behaviour for the different scenarios of increasing or decreasing vegetation cover and mean annual precipitation
621 for different initial landscapes. The implications of this are large for observational studies of catchment average erosion
622 rates and suggest that the direction and magnitude of response observed in a setting is highly dependent on the mean
623 vegetation and precipitation conditions of the catchment, as well as what time the observations are made within the cycle
624 of the varying vegetation and precipitation. Furthermore, these results highlight the need for future modeling studies (and
625 motivation for our ongoing work), to investigate the response of catchment topography and erosion rates to more realistic
626 climate and vegetation change scenarios, as well as a broader range of initial vegetation covers and precipitation rates
627 than those explored here such that the threshold in behaviour between the two curves shown in figure 17b can be
628 understood.

629 **5.5 Potential Observational Approaches to Test Model Predictions**

630 The behaviour discussed in the previous section matches field-data reported by Owen et al. (2010) who analysed soil
631 production rates from bedrock in different climate regimes. This data, under the assumption of steady-state soil thickness,
632 can be translated into denudation rates. They show that for low values of mean annual precipitation, soil production rates
633 vary between 0 m/Ma and 2 m/Ma due to abiotic processes controlling soil production rates. These observations resemble
634 the effect of our simulations with 10% initial vegetation cover, which shows the same variations in erosion rates with
635 intervals of zero erosion rate for hyper-arid conditions (Fig. 17). Areas with higher values of mean annual precipitation
636 show higher values in the soil production rate. These data points were not corrected for different uplift rates in the sample
637 areas so it is not possible to isolate the effect of vegetation/precipitation and tectonic uplift. In general, the observations
638 show no clear isolated trend but more of a cluster of soil production rates among a common mean, situated in a zone
639 controlled by biotic conditions. Compared to our model data for simulations with 70% initial vegetation cover, this
640 resembles the non-intuitive behaviour of an increase in erosion rate for increasing values of vegetation cover and
641 precipitation compared to a constant erosion rate for decreasing values of vegetation cover and precipitation.

642 Schaller et al. (2018) and Oeser et al. (2018) present millennial timescale (cosmogenic radionuclide derived) hillslope
643 denudation, and soil production, rates from the Chilean (EarthShape) study areas (Fig. 1A) considered in this study. They
644 find the lowest hillslope denudation rates in the arid and poorly vegetated north. Moving south towards higher
645 precipitation and vegetation cover the denudation rates increase until the southernmost location with highest rainfall and
646 vegetation cover where denudation rates decrease again. This non-linear relationship of hillslope denudation rates with

647 vegetation cover and precipitation is not directly comparable to the results presented here, but is consistent with a) the
648 notion emphasized here that interactions between precipitation and vegetation cover on denudation are non-linear, and b)
649 that the study areas considered here, although tectonically quiescent for tens of millions of years, have varying denudation
650 rates that suggest either variable rock uplift rates, and/or a persistent state of transience in hillslope denudation induced
651 by millennial timescale oscillations in climate and vegetation.

652 Beyond the previous studies, limited observations are available for comparison to the predictions shown here. The
653 millennial to million-year time scales investigated here can best be evaluated from observations of catchment wide
654 denudation over similar timescales. Cosmogenic radionuclide measurements from modern river sediments offer one
655 means to evaluate these results. Work by Acosta et al. (2015) in east Africa and Olen et al., (2016) in the Himalaya, are
656 also consistent with the results presented here for the range of vegetation cover available in each of these areas. However,
657 the integration time scales that these studies are sensitive to are shorter than what is presented here and prohibit a detailed
658 comparison. A final approach that future studies could pursue is to calculate paleo denudation rate for catchments from a
659 time series of sediments deposits preserved in either lakes (e.g. Marshall et al. 2015) or fluvial river terraces (e.g. Schaller
660 and Ehlers, 2006; Schaller et al., 2016). However, to be most effective, these studies need to target multiple study areas
661 with terrace or lake deposits that span a range of vegetation covers in the upstream catchments.

663 **5.6 Model Restrictions and Caveats**

664 Similar to previous work on this topic (Collins et al. 2004, Istanbuluoglu and Bras 2005) , the model setup used in this
665 study was simplified to document how different vegetation and climate related factors impact topography over long
666 (geologically relevant) timescales. We acknowledge that future model-studies should address some of the restrictions
667 imposed by our approach to evaluate their significance for the results presented here. Future work should consider a
668 transport-limited fluvial model or a fully-coupled alluvial sedimentation and transport model. The addition of this could
669 bring new understanding in to how vegetation not only influences detachment limited systems, but also influences
670 sedimentation and entrainment of material. This added level of complexity could however limit (due to computational
671 concerns) the temporal scales over which an investigation like this could be conducted. Future studies could improve
672 upon this work by considering a more in-depth parametrization of how vegetation related processes (e.g. changes in root
673 depth and density, plant functional type) influence topographic metrics and erosion rates. Also, although supported by
674 various publications (Dunne, 1996; Dunne et al., 2010), the assumption of a non-linear response of the effectiveness of
675 diffusional and fluvial processes to increases in vegetation cover has a major impact on the results of these study. A better
676 field-based understanding of these processes and the involved relationships could improve the accuracy of model studies
677 like this.

678 Also, due to the long-timescales considered here, mean annual precipitation rather than a stochastic distribution of
679 precipitation were implemented. Future work should evaluate how stochastic distributions in precipitation and extreme
680 events in arid, poorly vegetation settings, impact these results, however the long timescale forcings in precipitation and
681 vegetation imposed in this study will likely persist as the background template upon which high-frequency changes are
682 active.

683 Regarding the vegetation and water-budget, a more sophisticated model of evapotranspiration and infiltration as a function
684 to surface plant cover and plant functional traits such as rooting depth would improve model predictions and is a priority

685 for our future work. Improvements will come from planned coupling of surface process model with the dynamic
686 vegetation model LPJ-GUESS (Werner et al. 2018, this issue).

687 Our assumption that an increase in surface vegetation cover directly translates to an increase in Mannings roughness is
688 an additional simplification. The real value of Mannings roughness of a surface will be a function of the fractional
689 densities of different plant communities per model-patch. We argue that this simplification is however necessary because
690 it is not possible to know the composition of the plant community for specific areas in our modeled timescales. This could
691 be resolved by fully coupling a landscape evolution model to a dynamic vegetation model to resolve inter-patch
692 differences of surface vegetation cover and intra-patch plant functional types.

693 We also acknowledge that the transient forcings we have chosen for driving our model are simplistic and could be
694 improved by a higher-fidelity time-series of climate over the last millennia. We choose a 100kyr, eccentricity driven,
695 periodicity because it is widely recognized that the eccentricity cycles are a main control in driving Earth's glacial cycle
696 over the last 0.9Ma. While this approach is reasonable for a sensitivity analysis such as we've conducted, it prohibits a
697 detailed comparison to observations in specific study areas without additional refinement. Our results suggest that a
698 shorter periodicity, which would resemble other periodicities in the Milankovitch cycle (e.g., 41kyrs, 23kyrs) or shorter
699 time scale climate variations, such as Heinrich events (see Huntley et al. 2013) would lead to smaller magnitudes of
700 adjustment to new dynamic equilibria, because of short time spans in high-/low-erosive climate conditions within one
701 period. Regarding the long time-periods considered, we chose to have a steady-state climate driver in the model without
702 frequency driven modulation of rainfall events. We argue that over large time scales the occurrence of these events can
703 be integrated into a meaningful mean value but acknowledge that the incorporation of those events could alter the results
704 on a short cycle-basis. However because there is no meaningful way to test these frequency distributions against past-
705 climates, this would add additional unknowns and assumptions into our model parameterization.

706 Finally, we emphasize that a subset of our results, which resemble small magnitude changes of topographic relief (e.g.,
707 factor of change 1.01, Section 4.3.1, 4.3.2) are valid results for the predicted synthetic landscapes in our model framework
708 and not a numerical artefact. However, we acknowledge that these predicted changes are too small to be measured in a
709 real-world setting.

710 **6. Conclusions**

711 The results from our experiments show that the interactions of vegetation cover and mean annual precipitation on the
712 evolution of landscapes is a complex system with competing effects. Main conclusions which emerge from this study are:

713 (I) vegetation cover has a hindering effect on hillslope and fluvial erosion but the magnitude on which changes in
714 vegetation cover affect these processes is a function of the initial state of the system. Changes in systems with higher
715 initial values of vegetation cover have a less pronounced effect than changes in systems with lower initial vegetation
716 cover.

717 (II) In comparison to the Coastal Cordilleras of Chile, the relationship between precipitation and surface vegetation cover
718 shows a distinct shape: For a 10% increase in surface vegetation cover, the corresponding increase in mean annual
719 precipitation is smaller in areas of lower vegetation cover and increases for areas with higher vegetation cover. This has
720 an effect on transient topographies by shifting the equilibrium of vegetation and precipitation effects on erosion rates.

721 (III) Following our step-change simulations, model results show different behaviours for changes in vegetation-cover and
722 mean annual precipitation. While increases in mean annual precipitation have an increasing effect on erosion rates and

723 therefore a long-term negative effect on topographic metrics, an increase in vegetation cover hinders erosion, and leads
724 to higher topographic metrics. The magnitude of these changes is again dependent on the initial vegetation cover and
725 precipitation before the step-change.

726 (IV) Simulations with either oscillating vegetation cover or oscillating precipitation show adjustments to new dynamic
727 mean values around which the basin metrics oscillate. The magnitude of adjustment is highly sensitive to initial vegetation
728 cover, where simulations with 10% initial cover show higher magnitudes than simulations with 70% cover, for oscillating
729 vegetation. Oscillating precipitation leads to lower-/no adjustments but an oscillation of basin metrics around the initial
730 mean values with generally lower amplitudes compared to simulations with oscillating vegetation cover.

731 (V) Simulations with coupled oscillations of both vegetation cover and precipitation show only small magnitudes of
732 adjustments in topography metrics to new dynamic equilibrium similar to simulations with a oscillation in only
733 precipitation. However corresponding erosion rates show a complex pattern of rapid increases and decreases which results
734 from a interplay of competing effects of hindering of erosion by vegetation and aiding of erosion by precipitation.

735 Taken together, the above findings from this study highlight a highly variable behavior in how variations in vegetation
736 cover impact erosion and topographic properties. The complexity in how vegetation cover and precipitation changes
737 influence topography demonstrates the need for future work to consider both of these factors in tandem, rather than
738 singling out either parameter (vegetation cover or precipitation) to understand potential transients in topography.

739

740 **Acknowledgements:**

741 This study was funded as part of the German science foundation priority research program EarthShape: Earth surface
742 shaping by biota (grant EH329-14-1 to T.A.E. and T.H.). We thank Erkan Istanbuluoglu, Taylor Schildgen, and two
743 anonymous reviewers for constructive reviews that improved the manuscript. We also thank Associate Editor Rebecca
744 Hodge for thoughtful handling of the manuscript. We thank the national park service in Chile (CONAF) for providing
745 access to, and guidance through, the study areas during field trips. Daniel Hobley and the LandLab ‘slack’ community
746 are thanked for constructive suggestions during the Landlab program modifications implemented for this study. The
747 source code used in this study is freely available upon request.

748




Figure 1 Overview of the geographic location, precipitation, and vegetation cover of the Coastal Cordillera, Chile studies areas used for model setup up. A) Digital topography of the areas considered and corresponding to the EarthShape (www.earthshape.net) focus areas where ongoing related research is located. B) Observed present day mean annual precipitation from the WorldClim and CHELSA datasets used as model input. B) Present day maximum surface vegetation cover from MODIS data.




Figure 2 Normalized basin metrics for study-areas derived from 30 m SRTM digital topography from the study areas shown in Figure 1a. Colored dots represent cumulative mean values of normalized slope, relief and channel steepness calculated for all locations using 5-8 representative catchments in each area. Dotted lines represent linear interpolation between values. Note the gradual increase, then decrease in all metrics around study area at ~32°S.




Figure 3 Example model setup used in simulations in this study. Figure shows an example model predicted topography with a set drainage network, draining to the south. Boundary conditions and parameterizations used in the models are labeled. Blue colors represent low elevations, brown colors represent higher elevations. Additional details of parameters used are specified in Table 1.




Figure 4 Transient forcings in vegetation and precipitation considered in model experiments. Simulations were run for 15 Myr prior to the runtime shown in the figure. All transients imposed started a runtime of 5 Myr. A) Variations in vegetation cover imposed in the oscillating experiment conditions for initial vegetation cover of 10 and 70%. Oscillations have a 10% amplitude and a 100kyr periodicity. B) 10% positive and negative step-changes in vegetation cover imposed on simulations with 10% and 70% initial vegetation cover. C) Oscillating mean annual precipitation. Positive and negative amplitudes of oscillation resemble the magnitude of precipitation change extracted from vegetation cover/rainfall relationship from satellite data (Fig. 5, D) Positive and negative step changes in mean annual precipitation. The initial precipitation was based on values extracted from the worldclim climate dataset for respective focus areas.




Figure 5 Graphical representation of the observed precipitation – vegetation relationship in the Chilean focus areas (Fig. 1) and how precipitation amounts were selected when perturbations in vegetation cover were imposed. Black dots represent vegetation-precipitation values used in the steady-state model conditions and prior to any transients. Red dots show how vegetation cover perturbations in +/- 10% in the model simulations were used to select corresponding mean annual precipitation amounts. Note that the observed relationship between observed precipitation and vegetation cover in the Coastal Cordillera of Chile is non-linear, and is a source of the non-linear behavior in model forcing (e.g. Fig. 4) and results (Fig. 17) presented here.




Figure 6 Predicted values of hillslope diffusivity K_d (solid line) and fluvial erodibility K_v (dashed line) as a function of vegetation surface cover. Although absolute values can't be compared due to different units, the shape of the curves representing the different parameters show different sensitivities to changes in vegetation cover, and major source of the non-linearities discussed in the text. Fluvial erodibility shows the highest magnitude of change for vegetation cover values < 25% whereas hillslope diffusivity reacts in a more linearly with highest change below < 65% vegetation cover.




Figure 7 Steady-state model predicted (shaded regions) and observed (red dots) topographic metrics from the study areas shown in Figure 1 for different vegetation cover amounts. Observed topographic metrics were extracted from SRTM 90 m DEM. Model predicted values are shown for the cases of constant mean annual precipitation (a,b,c) or variable precipitation (D,E,F). Variable precipitation rates and vegetation covers were selected for these simulations using the observed values from the focus areas (Fig. 5). Note that for variable precipitation and vegetation cover simulations (d,e,f) the predicted values (similar to the observations) develop a humped shape pattern of an increase and then decrease in each parameter suggesting the changes in both precipitation and vegetation cover are needed to reproduce the general trend seen in observations. The sources of misfit between the predicted and observed values are due to the simplified (and untuned) setup of the simulations and discussed in the text.




Figure 8 Observed evolution of topographic metrics after a step-change in either vegetation (green lines) or mean annual precipitation (blue lines). Results are shown for two different initial vegetation cover amounts of V=10 and 70%. Imposed mean annual precipitation changes were done by selecting the precipitation amount corresponding to the initial and final vegetation amounts used in the simulations for vegetation cover ‘only’ change. Panels a,b,c show the reaction of model topographies to positive changes in boundary conditions, panels d,e,f show the reaction to negative changes in boundary conditions.




Figure 9 Mean catchment-wide erosion rates after a step-change disturbance in model boundary conditions. Blue lines represent erosion rates for models with changes in only precipitation, green lines represent erosion rates for models with changes in only vegetation cover. Panels a,b show the evolution after positive step-change, panels c,d for models with negative step-change. Note that the direction of change (positive or negative) from the initial state is in the opposite directions for precipitation and vegetation cover changes. This effect is manifested in the subsequent plots.




Figure 10 Evolution of topographic metrics for simulations with oscillating surface vegetation cover and constant precipitation corresponding to the initial vegetation cover prior to the transient in vegetation cover. Panels a,b,c show mean basin relief, mean basin slope and mean basin channel steepness (k_{sn}), respectively.






Figure 11 Predicted mean catchment erosion rates for simulations with oscillating surface vegetation cover and constant precipitation. Note that the magnitude of change in erosion rates for $\pm 10\%$ change in vegetation covers differs depending on the initial (or background) vegetation cover. This non-linear response is due in part to the vegetation cover effects on rock erodibility and diffusivity shown in figure 6.



830

831 **Figure 12** Evolution of topographic metrics for simulations with oscillating mean annual precipitation and constant vegetation
 832 cover. The vegetation cover was held constant at the value corresponding to the precipitation rate prior to the onset of the
 833 transient at 5000 kyrs. Panels a,b,c show mean basin relief, mean basin slope and mean basin channel steepness (k_{sn}),
 834 respectively.



835

836 **Figure 13** Mean catchment erosion rates for simulations with oscillating mean annual precipitation and constant surface
 837 vegetation cover. The amplitude of change in the erosion rates varies with the initial vegetation cover, in part due to the non-
 838 linear relationship between precipitation and vegetation cover (Fig. 4, 5).




Figure 14 Shifts in mean basin slope/mean basin relief relationship for simulations with positive and negative step-changes in either vegetation cover (green triangles) or mean annual precipitation (blue triangles). Black dots represent initial steady-state conditions prior to any imposed transient in vegetation cover or mean annual precipitation. Note that the sensitivity of topographic relief to perturbations in precipitation or vegetation cover is highest for low-vegetation cover (10%) settings.






Figure 15 Evolution of topographic metrics for coupled simulations where both changes in surface vegetation cover and a corresponding change (Fig. 5) in mean annual precipitation are simultaneously imposed. The amplitudes and frequency of the forcings that were imposed on the simulations are the same than the ones used for the simulations with isolated transient forcings. Panels a,b,c show evolution of mean basin relief, mean basin slope and mean basin channel steepness (k_{sn}) after start

850 of oscillation at 5Ma. Note the muted/damped response relative to previous simulations of oscillating vegetation cover or
 851 precipitation conditions.



852

853 **Figure 16** Mean catchment erosion rates for coupled simulations with changes in surface vegetation cover and mean annual
 854 precipitation. The first cycle in the time series is expanded in Figure 17. The variable amplitude and non-linear response shown
 855 here is due to the combined non-linear forcings in precipitation (Fig. 4, 5) and rock erodibility and diffusivity (Fig. 6) for
 856 different initial vegetation cover amounts.



857

858 **Figure 17** Mean catchment erosion rates for coupled simulations for one period of oscillation after the start of transient
 859 conditions (see also Fig. 15). Upper subplot shows conceptualized transient forcing in vegetation cover and mean annual
 860 precipitation, lower subplot shows erosion rate for simulations with low (black line) and high (dotted line) initial vegetation
 861 cover and precipitation values.

862




Figure 18 Conceptual figure showing the topographic response from simulations with coupled oscillation of mean annual precipitation and vegetation cover (see Fig. 17 for erosion rates). Upper row illustrates transient forcings, lower two rows show initial topography (yellow) and resulting transient topography (pink). Changes in topography are not to scale. Vegetation and rainfall amount is shown qualitatively on the hillslopes.

Table 1 Model parameters used for Landlab model setup.

| Model Parameter | Unit | Value |
|---|---|-----------|
| Uplift (U) | mm/yr | 0.2 |
| Fluvial Erodibility (k_e) | m/yr ($\text{Kg m}^{-1} \text{s}^{-2}$)-1 | 7.00E-06 |
| Critical Shear Stress (τ_c) | Pa | 58 |
| m, n | - | 0.6 / 0.7 |
| Base Diffusivity (K_b) | m^2/yr | 0.02 |
| Mannings Number (Vegetated, nvr) | - | 0.6 |
| Mannings Number (Soil, ns) | - | 0.01 |
| Reference Vegetation Cover (V_r) | - | 100% |
| w | - | 1 |
| alpha | - | 0.3 |
| p | - | 1 |
| Transient Vegetation Cover Amplitude (+-dV) | % | 10 |

References

Acosta, V. T., Schildgen, T. F., Clarke, B. A., Scherler, D., Bookhagen, B., Wittmann, H., von

874 Blanckenburg, F. and Strecker, M. R.: Effect of vegetation cover on millennial-scale landscape
875 denudation rates in East Africa, *Lithosphere*, 7(4), 408–420, doi:10.1130/L402.1, 2015.

876 Alberts, E. E., Nearing, M. A., Weltz, M. A., Risse, L. M., Pierson, F. B., Zhang, X. C., Laflen, J. M.
877 and Simanton, J. R.: Soil component, in USDA—Water Erosion Prediction Project, p. 47., 1995.

878 Allen, C. D. and Breshears, D. D.: Drought-induced shift of a forest–woodland ecotone: rapid
879 landscape response to climate variation, *Proceedings of the National Academy of Sciences*, 95(25),
880 14839–14842, 1998.

881 Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M.,
882 Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H. (Ted), Gonzalez, P., Fensham, R., Zhang,
883 Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W., Semerci, A. and Cobb, N.: A
884 global overview of drought and heat-induced tree mortality reveals emerging climate change risks for
885 forests, *Forest Ecology and Management*, 259(4), 660–684, doi:10.1016/j.foreco.2009.09.001, 2010.

886 Amundson, R., Heimsath, A., Owen, J., Yoo, K. and Dietrich, W. E.: Hillslope soils and vegetation,
887 *Geomorphology*, 234, 122–132, doi:10.1016/j.geomorph.2014.12.031, 2015.

888 Andriessen, P. A. and Reutter, K.-J.: K-Ar and fission track mineral age determination of igneous
889 rocks related to multiple magmatic arc systems along the 23 S latitude of Chile and NW Argentina,
890 *Tectonics of the southern central Andes: structure and evolution of an active continental margin*, 141–
891 154, doi:10.1007/978-3-642-77353-2_10, 1994.

892 Avdievitch, N. N., Ehlers, T. A. and Glotzbach, C.: Slow long-term exhumation of the West Central
893 Andean Plate Boundary, *Tectonics*, 2017.

894 Bachelet, D., Neilson, R. P., Hickler, T., Drapek, R. J., Lenihan, J. M., Sykes, M. T., Smith, B., Sitch,
895 S. and Thonicke, K.: Simulating past and future dynamics of natural ecosystems in the United States,
896 *Global Biogeochemical Cycles*, 17(2), n/a-n/a, doi:10.1029/2001GB001508, 2003.

897 Blois, J. L., Williams, J. W., Fitzpatrick, M. C., Jackson, S. T. and Ferrier, S.: Space can substitute
898 for time in predicting climate-change effects on biodiversity, *Proceedings of the National Academy*
899 *of Sciences*, 110(23), 9374–9379, doi:10.1073/pnas.1220228110, 2013.

900 Bonnet, S. and Crave, A.: Landscape response to climate change: Insights from experimental
901 modeling and implications for tectonic versus climatic uplift of topography, *Geology*, 31(2), 123–
902 126, 2003.

903 Braun, J. and Willett, S. D.: A very efficient O(n), implicit and parallel method to solve the stream
904 power equation governing fluvial incision and landscape evolution, *Geomorphology*, 180–181, 170–
905 179, doi:10/gbcb9, 2013.

906 Broecker, W. S. and van Donk, J.: Insolation changes, ice volumes, and the O¹⁸ record in deep-sea
907 cores, *Reviews of Geophysics*, 8(1), 169, doi:10/dmcjc3, 1970.

908 Broxton, P. D., Zeng, X., Scheftic, W. and Troch, P. A.: A MODIS-Based Global 1-km Maximum
909 Green Vegetation Fraction Dataset, *Journal of Applied Meteorology and Climatology*, 53(8), 1996–
910 2004, doi:10.1175/JAMC-D-13-0356.1, 2014.

911 Carretier, S., Regard, V., Vassallo, R., Aguilar, G., Martinod, J., Riquelme, R., Pepin, E., Charrier,
912 R., Herail, G., Farias, M., Guyot, J.-L., Vargas, G. and Lagane, C.: Slope and climate variability

control of erosion in the Andes of central Chile, *Geology*, 41(2), 195–198, doi:10.1130/G33735.1, 2013.

Collins, B. D., Bras, R. L. and Tucker, G. E.: Modeling the effects of vegetation-erosion coupling on landscape evolution, *Journal of Geophysical Research*, 109(F3), doi:10.1029/2003JF000028, 2004.

Culling, W. E. .: Analytical Theory of Erosion, *The Journal of Geology*, 68(3), 336–344, 1960.

Curran, J. C. and Hession, W. C.: Vegetative impacts on hydraulics and sediment processes across the fluvial system, *Journal of Hydrology*, 505, 364–376, doi:10.1016/j.jhydrol.2013.10.013, 2013.

DiBiase, R. A. and Whipple, K. X.: The influence of erosion thresholds and runoff variability on the relationships among topography, climate, and erosion rate, *Journal of Geophysical Research*, 116(F4), doi:10.1029/2011JF002095, 2011.

DiBiase, R. A., Whipple, K. X., Heimsath, A. M. and Ouimet, W. B.: Landscape form and millennial erosion rates in the San Gabriel Mountains, CA, *Earth and Planetary Science Letters*, 289(1–2), 134–144, doi:10.1016/j.epsl.2009.10.036, 2010.

Dietrich, W. E., Bellugi, D. G., Sklar, L. S., Stock, J. D., Heimsath, A. M. and Roering, J. J.: Geomorphic Transport Laws for Predicting Landscape form and Dynamics, in *Geophysical Monograph Series*, edited by P. R. Wilcock and R. M. Iverson, pp. 103–132, American Geophysical Union, Washington, D. C., 2013.

Dosseto, A., Hesse, P. P., Maher, K., Fryirs, K. and Turner, S.: Climatic and vegetation control on sediment dynamics during the last glacial cycle, *Geology*, 38(5), 395–398, doi:10.1130/G30708.1, 2010.

Dunne, T., Whipple, K. X. and Aubry, B. F.: Microtopography of hillslopes and initiation of channels by horton overland flow, in *Geophysical Monograph Series*, vol. 89, edited by J. E. Costa, A. J. Miller, K. W. Potter, and P. R. Wilcock, pp. 27–44, American Geophysical Union, Washington, D. C., 1995.

Dunne, T., Malmon, D. V. and Mudd, S. M.: A rain splash transport equation assimilating field and laboratory measurements, *Journal of Geophysical Research*, 115(F1), doi:10/dq4w5k, 2010.

Feng, X., Wang, Y., Chen, L., Fu, B. and Bai, G.: Modeling soil erosion and its response to land-use change in hilly catchments of the Chinese Loess Plateau, *Geomorphology*, 118(3–4), 239–248, doi:10.1016/j.geomorph.2010.01.004, 2010.

Fernandez, N. F. and Dietrich, W.: Hillslope evolution by diffusive processes: The timescale for equilibrium adjustments, *Water Resources Research*, 33(6), 1307–1318, 1997.

Gilbert, G. K.: Report on the Geology of the Henry Mountains, Dept of the Interior, US Govt Printing Office, Washington, DC., 1877.

Gyssels, G., Poesen, J., Bochet, E. and Li, Y.: Impact of plant roots on the resistance of soils to erosion by water: a review, *Progress in Physical Geography*, 29(2), 189–217, doi:10.1191/0309133305pp443ra, 2005.

Hobley, D. E. J., Adams, J. M., Nudurupati, S. S., Hutton, E. W. H., Gasparini, N. M., Istanbuluoglu, E. and Tucker, G. E.: Creative computing with Landlab: an open-source toolkit for building, coupling, and exploring two-dimensional numerical models of Earth-surface dynamics, *Earth Surface*

951 Dynamics, 5(1), 21–46, doi:10.5194/esurf-5-21-2017, 2017.

952 Howard, A. D.: Badland Morphology and Evolution: Interpretation Using a Simulation Model, *Earth*
953 *Surface Processes and Landforms*, 22(3), 211–227, doi:10/b4mtzw, 1997.

954 Howard, A. D. and Kerby, G.: Channel changes in badlands, *Geological Society of America Bulletin*,
955 94(6), 739–752, 1983.

956 Howard, A. D., Dietrich, W. E. and Seidl, M. A.: Modeling fluvial erosion on regional to continental
957 scales, *Journal of Geophysical Research: Solid Earth*, 99(B7), 13971–13986, doi:10/bd8k3j, 1994.

958 Hughes, L.: Biological consequences of global warming: is the signal already, *TREE*, 15(2), 2000.

959 Huntley, B., Allen, J. R. M., Collingham, Y. C., Hickler, T., Lister, A. M., Singarayer, J., Stuart, A.
960 J., Sykes, M. T. and Valdes, P. J.: Millennial Climatic Fluctuations Are Key to the Structure of Last
961 Glacial Ecosystems, edited by J. Mohan, *PLoS ONE*, 8(4), e61963,
962 doi:10.1371/journal.pone.0061963, 2013.

963 Istanbuluoglu, E. and Bras, R.: Vegetation-modulated landscape evolution: Effects of vegetation on
964 landscape processes, drainage density, and topography, *Journal of Geophysical Research*, 110(F2),
965 doi:10.1029/2004JF000249, 2005.

966 Istanbuluoglu, E., Tarboton, D. G., Pack, R. T. and Luce, C. H.: Modeling of the interactions between
967 forest vegetation, disturbances, and sediment yields, *Journal of Geophysical Research: Earth Surface*,
968 109(F1), n/a-n/a, doi:10.1029/2003JF000041, 2004.

969 Jeffery, M. L., Yanites, B. J., Poulsen, C. J. and Ehlers, T. A.: Vegetation-precipitation controls on
970 Central Andean topography, *Journal of Geophysical Research: Earth Surface*, 119(6), 1354–1375,
971 doi:10.1002/2013JF002919, 2014.

972 Juez-Larré, J., Kukowski, N., Dunai, T. J., Hartley, A. J. and Andriessen, P. A. M.: Thermal and
973 exhumation history of the Coastal Cordillera arc of northern Chile revealed by thermochronological
974 dating, *Tectonophysics*, 495(1–2), 48–66, doi:10.1016/j.tecto.2010.06.018, 2010.

975 Langbein, W. B. and Schumm, S. A.: Yield of sediment in relation to mean annual precipitation, *Eos*,
976 *Transactions American Geophysical Union*, 1958.

977 Ledru, M.-P., Labouriau, M. L. S. and Lorscheitter, M. L.: Vegetation dynamics in southern and
978 central Brazil during the last 10,000 yr B.P., *Review of Palaeobotany and Palynology*, 99, 131–142,
979 1997.

980 Maksaev, V. and Zentilli, M.: Fission Track Thermochronology of the Domeyko Cordillera, Northern
981 Chile: Implications for Andean Tectonics and Porphyry Copper Metallogenesis, *Exploration and*
982 *Mining Geology*, 8, 65–89, 1999.

983 Marshall, J. A., Roering, J. J., Bartlein, P. J., Gavin, D. G., Granger, D. E., Rempel, A. W.,
984 Praskievicz, S. J. and Hales, T. C.: Frost for the trees: Did climate increase erosion in unglaciated
985 landscapes during the late Pleistocene?, *Science Advances*, 1(10), e1500715–e1500715,
986 doi:10/gddtwz, 2015.

987 Marston, R. A.: Geomorphology and vegetation on hillslopes: Interactions, dependencies, and
988 feedback loops, *Geomorphology*, 116(3–4), 206–217, doi:10.1016/j.geomorph.2009.09.028, 2010.

989 McInnes, B. I. A., Farley, K. A., Sillitoe, R. and Kohn, B. P.: Application of Apatite (U-th)/He
990 Thermochronometry to the determination of the sense and amount of vertical fault displacement at
991 the Chuquicamata Porphyry Copper Deposit, Chile, *Economic Geology*, 94, 937–948, 1999.

992 McPhillips, D., Bierman, P. R., Crocker, T. and Rood, D. H.: Landscape response to Pleistocene-
993 Holocene precipitation change in the Western Cordillera, Peru: ^{10}Be concentrations in modern
994 sediments and terrace fills, *J. Geophys. Res. Earth Surf.*, 118(4), 2013JF002837,
995 doi:10.1002/2013JF002837, 2013.

996 Muhs, D. R., T. A. Ager and Beget, J. E.: Vegetation and paleoclimate of the last interglacial period,
997 central Alaska, *Quaternary Science Reviews*, 20, 41–61, 2001.

998 Muller, R. A. and MacDonald, G. J.: Spectrum of 100-kyr glacial cycle: Orbital inclination, not
999 eccentricity, *Proceedings of the National Academy of Sciences*, 94(16), 8329–8334, doi:10/c7bpd5,
1000 1997.

1001 Oeser, R., Stroncik, N., Moskwa, L.-M., Bernhard, N., Schaller, M., Canessa, R., Brink, L. van den,
1002 Köster, M., Brucker, E., Stock, S., Fuentes-Espoz, J. P., Godoy, R., Matus, F., Osés Pedraza, R.,
1003 Osses McIntyre, P., Paulino, L., Seguel, O., Bader, M. Y., Boy, J., Dippold, M., Ehlers, T. A., Kühn,
1004 P., Kuzyakov, Y., Leinweber, P., Scholten, T., Spielvogel, S., Spohn, M., Übernickel, K., Tielbörger,
1005 K., Wagner, D. and Blanckenburg, F. von: Chemistry and microbiology of the critical zone along a
1006 steep climate and vegetation gradient in the Chilean Coastal Cordillera, n.d.

1007 Olen, S. M., Bookhagen, B. and Strecker, M. R.: Role of climate and vegetation density in modulating
1008 denudation rates in the Himalaya, *Earth and Planetary Science Letters*, 445, 57–67,
1009 doi:10.1016/j.epsl.2016.03.047, 2016.

1010 Owen, J. J., Amundson, R., Dietrich, W. E., Nishiizumi, K., Sutter, B. and Chong, G.: The sensitivity
1011 of hillslope bedrock erosion to precipitation, *Earth Surface Processes and Landforms*, 36(1), 117–
1012 135, doi:10.1002/esp.2083, 2011.

1013 Perron, J. T., Richardson, P. W., Ferrier, K. L. and Lapôtre, M.: The root of branching river networks,
1014 *Nature*, 492(7427), 100–103, doi:10.1038/nature11672, 2012.

1015 Prentice, I. C., Harrison, S. P. and Bartlein, P. J.: Global vegetation and terrestrial carbon cycle
1016 changes after the last ice age, *New Phytologist*, 189(4), 988–998, doi:10.1111/j.1469-
1017 8137.2010.03620.x, 2011.

1018 Reinhardt, L., Jerolmack, D., Cardinale, B. J., Vanacker, V. and Wright, J.: Dynamic interactions of
1019 life and its landscape: feedbacks at the interface of geomorphology and ecology, *Earth Surface
1020 Processes and Landforms*, 35(1), 78–101, doi:10.1002/esp.1912, 2010.

1021 Roering, J. J., Almond, P., Tonkin, P. and McKean, J.: Soil transport driven by biological processes
1022 over millennial time scales, *Geology*, 30(12), 1115–1118, 2002.

1023 Sangireddy, H., Carothers, R. A., Stark, C. P. and Passalacqua, P.: Controls of climate, topography,
1024 vegetation, and lithology on drainage density extracted from high resolution topography data, *Journal
1025 of Hydrology*, 537, 271–282, doi:10.1016/j.jhydrol.2016.02.051, 2016.

1026 Schaller, M. and Ehlers, T. A.: Limits to quantifying climate driven changes in denudation rates with
1027 cosmogenic radionuclides, *Earth and Planetary Science Letters*, 248(1–2), 153–167,
1028 doi:10.1016/j.epsl.2006.05.027, 2006.

1029 Schaller, M., von Blanckenburg, F., Veldkamp, A., Tebbens, L. A., Hovius, N. and Kubik, P. W.: A
1030 30 000 yr record of erosion rates from cosmogenic ^{10}Be in Middle European river terraces, *Earth and*
1031 *Planetary Science Letters*, 204(1–2), 307–320, doi:10.1016/S0012-821X(02)00951-2, 2002.

1032 Schaller, M., Ehlers, T. A., Stor, T., Torrent, J., Lobato, L., Christl, M. and Vockenhuber, C.: Spatial
1033 and temporal variations in denudation rates derived from cosmogenic nuclides in four European
1034 fluvial terrace sequences, *Geomorphology*, doi:10.1016/j.geomorph.2016.08.018, 2016.

1035 Schaller, M., Ehlers, T. A., Lang, K. A. H., Schmid, M. and Fuentes-Espoz, J. P.: Addressing the
1036 contribution of climate and vegetation cover on hillslope denudation, Chilean Coastal Cordillera
1037 (26° – 38°S), *Earth and Planetary Science Letters*, 489, 111–122, doi:10.1016/j.epsl.2018.02.026,
1038 2018.

1039 Smith, B., Wårlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J. and Zaehle, S.: Implications
1040 of incorporating N cycling and N limitations on primary production in an individual-based dynamic
1041 vegetation model, *Biogeosciences*, 11(7), 2027–2054, doi:10.5194/bg-11-2027-2014, 2014.

1042 Stephan, U. and Gutknecht, D.: Hydraulic resistance of submerged flexible vegetation, *Journal of*
1043 *Hydrology*, 269(1), 27–43, 2002.

1044 Tucker, G., Lancaster, S., Gasparini, N. and Bras, R.: The channel-hillslope integrated landscape
1045 development model (CHILD), in *Landscape erosion and evolution modeling*, pp. 349–388, Springer.,
1046 2001.

1047 Tucker, G. E. and Hancock, G. R.: Modelling landscape evolution, *Earth Surface Processes and*
1048 *Landforms*, 35(1), 28–50, doi:10.1002/esp.1952, 2010.

1049 Vergani, C., Giadrossich, F., Buckley, P., Conedera, M., Pividori, M., Salbitano, F., Rauch, H.,
1050 Lovreglio, R. and Schwarz, M.: Root reinforcement dynamics of European coppice woodlands and
1051 their effect on shallow landslides: A review, *Earth-Science Reviews*, 167, 88–102,
1052 doi:10.1016/j.earscirev.2017.02.002, 2017.

1053 Walling, D. . and Webb, B. .: Patterns of sediment yield, *Background to Palaeohydrology*, 69–100,
1054 1983.

1055 Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J., Fromentin, J.-M.,
1056 Hoegh-Guldberg, O. and Bairlein, F.: Ecological responses to recent climate change, *Nature*,
1057 416(6879), 389–395, 2002.

1058 Werner, C., Schmid, M., Ehlers, T. A., Fuentes-Espoz, J. P., Steinkamp, J., Forrest, M., Liakka, J.,
1059 Maldonado, A. and Hickler, T.: Effect of changing vegetation on denudation (part 1): Predicted
1060 vegetation composition and cover over the last 21 thousand years along the Coastal Cordillera of
1061 Chile, *Earth Surface Dynamics*, 2018.

1062 Whipple, K. X. and Tucker, G. E.: Dynamics of the stream-power river incision model: Implications
1063 for height limits of mountain ranges, landscape response timescales, and research needs, *J. Geophys.*
1064 *Res.*, 104(B8), 17661–17674, doi:10.1029/1999JB900120, 1999.

1065 Willgoose, G., Bras, R. L. and Rodriguez-Iturbe, I.: Results from a new model of river basin
1066 evolution, *Earth Surface Processes and Landforms*, 16(3), 237–254, 1991.

1067 Yetemen, O., Istanbuluoglu, E., Flores-Cervantes, J. H., Vivoni, E. R. and Bras, R. L.: Ecohydrologic

1068 role of solar radiation on landscape evolution, *Water Resources Research*, 51(2), 1127–1157,
1069 doi:10.1002/2014WR016169, 2015.

1070 Zachos, J., Pagani, M., Sloan, L., Thomas, E. and Billups, K.: Trends, Rhythms, and Aberrations in
1071 Global Climate 65 Ma to Present, *Science*, 292(5517), 686–693, doi:10.1126/science.1059412, 2001.

1072 Zhou, Z. C., Shangguan, Z. P. and Zhao, D.: Modeling vegetation coverage and soil erosion in the
1073 Loess Plateau Area of China, *Ecological Modelling*, 198(1–2), 263–268,
1074 doi:10.1016/j.ecolmodel.2006.04.019, 2006.

1075