

New constraints to river valley evolution before and after the emplacement of the largest landslide on the exposed Earth surface: the Seymareh rockslide - debris avalanche (Zagros Mts., Iran)

Michele Delchiaro¹, Marta Della Seta¹, Salvatore Martino¹, Maryam Dehbozorgi², Reza Nozaem³

¹Department of Earth Sciences, Sapienza University of Rome, Rome, Italy

²Department of Earth Science, Kharazmi University, Tehran, Iran

³School of Geology, College of Science, University of Tehran, Iran

Correspondence to: Michele Delchiaro (michele.delchiaro@uniroma1.it)

Abstract. The Seymareh Landslide, detached ~10 ka from the northeastern flank of the Kabir-kuh fold (Zagros Mts., Iran), is recognized worldwide as the largest rock slope failure (44 Gm³) ever recorded on the exposed Earth surface. Detailed studies have been performed that have described the landslide mechanism and different scenarios have been proposed for explaining the induced changes in landscape. The purpose of this study is to provide still missing time constraints to the evolution of the Seymareh River valley, before and after the emplacement of the Seymareh Landslide, to highlight the role of geomorphic processes both as predisposing factors and as response to the landslide debris emplacement.

We used optically stimulated luminescence (OSL) to date lacustrine and fluvial terrace sediments, whose plano-altimetric distribution has been correlated to the detectable knickpoints along the Seymareh River longitudinal profile, allowing the reconstruction of the evolutionary model of the fluvial valley. We infer that the knickpoint migration along the main river and the erosion wave propagated upstream through the whole drainage network caused the stress release and the ultimate failure of the rock mass involved in the landslide. We estimated that the stress release activated a Mass Rock Creep (MRC) process with gravity-driven deformation processes occurring over an elapsed time-to-failure on the order of 10² ky. We estimated also that the Seymareh damming lake persisted for ~3500 years before starting to empty ~6.6 ka due to lake overflow. A sedimentation rate of 10 mm y⁻¹ was estimated for the lacustrine deposits, which increased up to 17 mm y⁻¹ during the early stage of lake emptying due to the increased sediment yield from the lake tributaries. We calculated an erosion rate of 1.8 cm y⁻¹ since the beginning of the landslide cut by Seymareh River, which propagated through the drainage system up to the landslide source area.

The evolutionary model of the Seymareh River valley can provide the necessary constraints for future stress-strain numerical modeling of the landslide slope to reproduce the MRC and demonstrate the possible role of seismic forcing in anticipating the time-to-failure for such an end-member case study.

1 Introduction

Tectonically active landscapes are very dynamic systems in which threshold conditions on hillslopes are often reached. Therefore, there are considerable implications for natural hazards related to seismicity and to the geomorphic coupling between hillslopes and rivers, with both fluvial control on hillslopes and landslide affecting the fluvial network. In response to rock uplift, relief and hillslope angles increase linearly in time due to erosional processes in landscapes affected by low to moderate tectonic forcing (Montgomery and Brandon, 2002; Binnie et al., 2007; Larsen and Montgomery, 2012). Nonetheless, such a linear increase in relief and hillslope angles is limited by the reaching the threshold slope conditions associated with the hillslope material strength (Schmidt and Montgomery, 1995), until the latter is exceeded by gravitational stress giving rise to bedrock landslides. This leads to a nonlinear increase in erosion rates in landscapes affected by long-lasting or high-rate tectonic forcing, where the increase in the rate of channel incision is accommodated by an increased frequency of slope failure rather than by slope steepening. These slope failures may represent the ultimate conditions of mass rock creep (MRC; Chigira, 1992) processes contributing to the development of gravity-driven deformations due to time-dependent rheology. This kind of deformations can evolve into generalized failure and slope collapse when the increased strain rate leads to an accelerating deformation. The strain limit conditions are reached when the stationary creep stage evolves into the accelerating creep stage. Failure is typically associated with mass strength reduction as an effect of rock mass damage occurring over time during the ongoing gravity-driven deformations (Eberhardt et al., 2004; Stead et al., 2006). The ultimate collapse for MRC leads to rock avalanches originating from the instantaneous fragmentation of rock masses (Hungri et al., 2001).

This work is focused on the evolution of the Seymareh River valley before and after the valley was dammed by the landslide worldwide recognized as the largest subaerial rock landslide ever observed, the Seymareh rockslide – debris avalanche that occurred in the northwestern Zagros Mts. (Iran). Different interpretations have been proposed so far by the scientific community to explain the generation of such an exceptional event and different scenarios have been hypothesized for explaining the induced changes in landscape. Harrison and Falcon (1937, 1938) provided much of the present knowledge on the rock avalanche, including the geology and structure of the source area, the general geomorphology and the basic geometry of the landslide. Oberlander (1965) included a short appendix on the landslide in his study of Zagros streams and discussed its origin in relation to the activity of Seymareh River. Later, in the 1960s, Watson and Wright (1969) characterized the geomorphology and stratigraphy of the debris, discussed the origin of the initial rockslide, and examined the debris avalanche emplacement mechanisms. Roberts (2008) and Roberts and Evans (2013) provided a detailed model of how the geological and tectonic evolution of the Kabir-kuh fold predisposed the slope to such a large-scale failure, including formation of structural/kinematic and rheological control, and inferred a seismic trigger. Specifically, Roberts and Evans (2013) obtained from a charcoal-rich layer approximately 15 m above the base of the lacustrine sequence with a ^{14}C age of 8710 years BP. Based on the interpretation of three separate radiocarbon ages provided additionally by Griffiths et al. (2001) an estimated radiocarbon bracket age of the Seymareh event was suggested between 9800–8710 ^{14}C years BP. Yamani et al.

(2012) provided some general details on the evolution of the dam lake drainage, describing a sequence of entrenched lacustrine terraces upstream of the landslide dam. Finally, Shoaie (2014) reviewed the possible mechanisms of failure and interpreted the post-failure geomorphic features through analyzing the processes responsible for the formation and erosion of the landslide dams of the Seymareh, Jaidar and Balmak (called also Chah Javal) lakes by using available annual sedimentation data and field measurements of the deposits in these lakes.

Despite the number of scenarios proposed so far, quantitative constraints on the river valley evolution before and after the emplacement of this giant landslide are still missing, which could help in quantitatively modeling the trigger scenario of this end-member case study of massive rock slope failures. In this regard, recently, some works (Bozzano et al., 2012, 2016; Della Seta et al., 2017; Martino et al., 2017) have focused on the role of landscape evolution rates on the development of MRC processes. Among these, Bozzano et al. (2016) demonstrated that erosion rates play a key role in the development of MRC processes within the rock masses and, consequently, in their possible evolution into massive rock slope failures, even without invoking transient external forcing (e.g., earthquakes). Furthermore, evidence of giant rock slides due to MRC that caused valley river damming have been recently documented by Zhao et al. (2019) in the Sichuan; these rockslides are related to the effects of river knickpoint propagation and inner gorge formation and serve as a further confirmation of the combined role played by the fluvial dynamics and the geological structural setting.

This work is aimed at filling the knowledge gap in landscape evolution associated with the emplacement of the Seymareh Landslide (hereafter referred to as SL). Detailed geomorphological mapping, correlation and dating of certain geomorphic markers (Burbank and Anderson, 2012) represented by fluvial and lacustrine terraces pre- and post-failure have been performed upstream and downstream of the landslide dam in order to provide new time constraints to the Seymareh River valley evolution and to outline the role of the geomorphic processes both as predisposing factors for MRC processes and as response to this giant gravitational instability.

2 Regional geological and geomorphological framework

The SL detached from the northeastern flank of the Kabir-kuh fold, the largest and highest anticline in the Pusht-e Kuh arc, in the northwestern part of Iran (Vergés et al., 2011). The Zagros mountain range is part of the Alpine-Himalayan orogenic system that originates from the Late-Cretaceous-Cenozoic convergence between Africa/Arabia-Eurasia (Talbot and Alavi, 1996; Stampfli and Borel, 2002; Golonka, 2004; McQuarrie, 2004; Mouthereau et al., 2012). The Zagros orogen was traditionally classified by distinctive lithological units and structural styles into four NW trending tectono-metamorphic and magmatic belts (Fig. 1). These are bounded by defects on a regional scale such as the Main Zagros Thrust (MZT), High Zagros Fault (HZF) and Mountain Front Fault (MFF) (Agard et al., 2005 and references therein). These tectonic units are from the inner to the outer sectors of the belt are: the Urumieh Dokhtar volcanic arch, the Sanandaj-Sirjan Zone, the Imbricate Zone, the Zagros (or Simply) folded belt and the continental Mesopotamian Foreland (Fig. 1).

Seismicity is distributed in a 200-300 km wide area of the Zagros mountain range (Hatzfeld et al., 2010, Paul et al., 2010, Rajabi et al., 2011), with a sharp cut along the Main Zagros Reverse Fault in the NE (e.g., Yamini-Fard et al., 2016), with recurrent earthquakes of Mw 5-6 and exceptional earthquakes of higher magnitude, i.e., up to Mw 6-8 (see supplementary material). The SL occurred in a very densely seismically active area and recurrence rate of nearby strongly felt earthquakes considerably higher than the rate of slope steepening led Roberts and Evans (2013) to hypothesize that seismic forcing may have played a primary role in triggering the landslide.

The outcropping formations in the Kabir-kuh anticline date to a time interval ranging from the Late Cretaceous to the early Miocene and are characterized by different lithological and rheological properties (Vergés et al., 2011). Since the geo-structural setting of the fold flanks represented a crucial predisposing factor for the catastrophic massive rock slope failure (Roberts and Evans, 2013), we referred to the most detailed stratigraphic column proposed by James and Wynd (1965), Alavi (2004) and to the detailed mapping of the Kabir-kuh fold conducted by the Iran Oil Operating Companies (Setudehnia and Perry 1967; Takin et al. 1970; Macleod 1970). Specifically, the investigated area includes the middle and low reaches of Seymareh River starting approximately 60 km upstream of the SL down to the SE termination of the Kabir-kuh fold. In Fig. 2, the geological map of the study area, the stratigraphic column and two geological cross-sections related to different structural sectors are reported.

It is noteworthy that, in the Kabir-kuh anticline, the Pabdeh Formation is composed of three rheologically contrasting members, which crop out in the SL scar area: i) the lower Pabdeh member (150 m thick), which is dominated by marls and shales, ii) the Taleh Zang member (50 m thick), consisting of platform limestone, and iii) the upper Pabdeh member (150 m thick), composed mainly of calcareous marl. The Asmari Formation creates a carapace originally covering the top of the Kabir-kuh fold, while in the synclinal valleys between the Kabir-kuh fold and the adjacent folds, the Asmari Formation is overlapped by a Miocene-Pliocene succession (Homke et al., 2004). Referring to the Changuleh syncline studied by Homke et al. (2004), the foreland stratigraphy includes the following: i) the Gachsaran Formation (early Miocene - 12.3 Ma, thickness approximately 400 m), composed of salt, anhydrite, marl and gypsum; ii) the Agha Jari Formation (12.3 Ma – 3 Ma., thickness approximately 1400 m); and iii) the Bakhtiari Formation (3 Ma – early Pleistocene, thickness approximately 900 m). The Agha Jari Formation consists of sandstones and conglomerates, linked to the evolution from deltaic to fluvial transitional environments (Elyasi et al., 2014), and the Bakhtiari formation consists of conglomerates characterized by coarse and mud-supported grains, sandstones, shales and silts and marks the onset of syn-orogenic fluvial environment conditions (Shafiei and Dusseault, 2008).

The reported cross-sections intersect the synclinal valley of the Seymareh River. The dip angle of the northeastern flank of the syncline considerably decreases from NW to SE from 45° (section A-A') to 18-20° (section B-B'). Therefore, along the section A-A' the Cretaceous-Paleogene bedrock (from the Sarvak Formation to the Asmari Formation) offers a greater accommodation volume to the continental and epicontinental formations (Gachsaran Formation and Agha Jari Formation), as the synclinal axis is located at a lower elevation than in the B-B' section.

The Zagros Range globally provides one of the most spectacular examples of landscape evolution in response to active tectonics (Bourne and Twidale, 2011) because its drainage network clearly adapted to the growth of the thrust-fold structures (Ramsey et al., 2008) and to the erodibility of the outcropping formations (Oberlander, 1985). Oberlander (1968) suggested that the drainage network in the NW Zagros was superimposed from structurally conformable younger horizons. In his model, the breaching of hard geological units of the antiformal ridges follows a phase of river cutting and expansion of the fold axial fold basins through the softer overlying units. In the Kabir-kuh fold, the transverse cutting of the Asmari limestone, and the exposure of the underlying more erodible Pabdeh-Gurpi marls, leads to the formation of a low-relief landscape with synformal ridges on which the new through-going drainage system can be developed. In Oberlander's hypothesis, it is the Pabdeh and Gurpi marls that facilitate the creation of a low-relief landscape across the anticline crests and are therefore integral to the story of drainage superimposition.

Tucker and Slingerland (1996) computed a numerical landscape evolution model, calibrated on the Kabir-kuh fold, to understand how the growth and propagation of the folds, the different lithologies and the drainage network can influence the sediment flux from a tectonically active belt towards the foreland basin. The authors calibrated the landscape evolution model with the current topography of the range, obtaining time constraints for landscape evolution modeling. According to the Oberlander model, Fig. 3 shows four main steps that describe the landscape evolution of the Kabir-kuh fold with the timing provided in the model by Tucker and Slingerland (1996).

- Step 1 - Approximately 4.3 Ma, in response to the initial stages of fold growth, an orthoclinal drainage develops, parallel to the main structures. The tributaries flowing along the flanks of the folds transport debris, which is deposited in the synclines. In the Kabir-kuh fold the carbonate core is still buried by the Miocene cover units.
- Step 2 - Approximately 3.8 Ma, as soon as the deformation front migrates towards the SW, new folds raise with a progressive adjustment of the drainage to these morpho-structures. The previously deposited sediments are remobilized and transported towards the depocenter of the syncline basins and partly outside; the syn-orogenic deposits are strongly eroded along the crests of the anticlines, thus exposing the underlying formations. This causes a topography characterized by resistant hogbacks that flank the inner cores.
- Step 3 - Approximately 2.4 Ma, with the ongoing deformation, the drainage develops in a "trellis" pattern. The river erosion affects the erodible units located stratigraphically between the limestone of the Asmari Formation and the inner core of the fold. At the end of this step the Miocene cover is completely removed from the ridges and the river erosion also affects the marls and evaporites of the syn-orogenic formations in the valleys, exposing the underlying limestone of the Asmari Formation.
- Step 4 - Approximately 1.6 Ma, due to the continuous uplift and exhumation of younger, more external folds, the sediment accumulation becomes negligible and the Asmari limestone is strongly eroded giving rise to syncline ridges. The following Quaternary landscape evolution is then likely driven by the evolution of the drainage network and is also influenced by climatic factors and by the slope-to-channel dynamics.

The model by Tucker and Slingerland (1996) is the unique numerical model existing on the Kabir-kuh fold and this motivates our choice of using it as a reference for the medium-to-long term evolution of the Seymareh River valley. The Seymareh River valley is arranged parallel to the Kabir-kuh fold and its evolution was inevitably influenced by the exceptional landslide event that temporarily dammed it, causing the formation of the three-lake system which includes the Seymareh, Jaidar and Balmak lakes (Fig. 4). The valley evolution before and after the event is well recorded by Quaternary landforms preserved along the valley. Yamani et al. (2012) focused on the post-failure evolution of the valley describing four levels of terraces upstream of the landslide dams as a sequence of lacustrine terraces. Shoaiei (2014), in addition to evaluating the longevity of the SL dams, identified in the merging of Seymareh River with a left tributary as the reason for strong river incision at the base of the northeastern flank of the Kabir-kuh fold and as a possible causal factor for the SL collapse. However, none of the previous studies on the Quaternary evolution of the Seymareh River valley provided absolute dating of geomorphic markers (mainly fluvial terraces) preserved upstream as well as downstream of the landslide dam or provided robust and quantitative constraints to the pre-failure valley evolution as a possible geomorphological factor for failure occurrence.

3 Methods

The geomorphological study of the area was carried out first through the analysis and interpretation of remote sensing data, such as aerial photos (National Cartographic Center of Iran, aerial photo, scale:1:20000, acquired on 24 August 1955), Google Earth satellite optical images (2018 Landsat Imagery) and vector topographic maps (National Cartographic Center of Iran, topographic map of Kuhdasht, scale: 1:25,000), which led to the first detection of possible geomorphic markers within the Seymareh River valley. Vector topographic data also allowed the construction of a 10 m Digital Elevation Model (DEM) for terrain analyses and led to the projection of the possible geomorphic markers along the river longitudinal profile (Wilson and Gallant; 2001; Burbank and Anderson, 2012). The DEM was obtained by the ArcGIS 10® software package, starting from vector topographic data (contour lines, hydrography and point elevation) and using the ANUDEM interpolation algorithm (Hutchinson et al., 2011 and references therein). To automatically extract the hydrographic network from the DEM and then to project the geomorphic markers along the longitudinal river profiles, some of the ArcGIS® 10 tools of the Hydrology toolbox were used (Flow Direction, Flow Accumulation, Reclassify, Stream Order and Stream to Feature), setting the flow accumulation threshold according to that proposed for the fluvial domain (10^{-1} km^2) by Montgomery and Foufoula-Georgiu (1993). The longitudinal profile was therefore transformed into a route along which the elevation of the top surfaces of geomorphic markers identified in the area were projected through the Linear Referencing Tools (Create Route and Locate Features along Route). A geological and geomorphological field survey was then carried out with the aims of mapping the most significant active and relict landforms for the Quaternary evolution of the Seymareh River valley and of sampling the corresponding deposits in order to date them with the OSL method (optically stimulated luminescence; Murray and Olley, 2002; Wintle and Murray, 2006 and references therein).

OSL sampling is a very delicate and quite complex technique. In fact, it is absolutely necessary to prevent the sample from being exposed to light because the luminescence signal could be reduced or even reset. In choosing the most suitable site to sample, of course, levels were identified with original sedimentary structures, avoiding bioturbations and post-depositional alterations. Once the site for sampling was identified, it was important to carefully clean off the slope and prepare, according to the state of thickening, consistency or cementation of the material, the equipment necessary for taking the sample, without it being exposed to light. Furthermore, to minimize the effects of cosmic radiation and to thereby avoid the risk of rejuvenated ages, the samples were taken at least one meter below the topographic surface (or below eventual erosional surfaces identified within the deposits). All of the samples, mainly characterized by fine-grained loose sediments (size <2 mm) were taken directly by using a hammer to insert a metal tube into the ground, which must be isolated from light and humidity immediately after collection. To maximize the uniformity of the natural radioactivity of the burial period, the sampled material was surrounded by at least 30 cm of homogeneous sediment. From the same level where it was sampled, an additional 500 - 800 g of sediment was extracted to evaluate natural radioactivity (if the annual dose rate measurement is not performed in situ), for the mineralogical and granulometric analyses, as well as to determine the moisture content.

The OSL dating was performed by the LABER OSL Laboratory, in Waterville, Ohio (U.S.). Quartz was extracted for equivalent dose (D_e) measurements. In the OSL laboratory, the sample was treated first with 10% HCl and 30% H_2O_2 to remove organic materials and carbonates, respectively. After grain-size separation, the fraction of 90-125 μm size is relatively abundant, so this fraction was chosen for D_e determination. The grains were treated with HF acid (40%) for approximately 40 minutes to remove the alpha-dosed surface, followed by 10% HCl acid to remove fluoride precipitates. Luminescence measurements were performed using an automated Risø TL/OSL-20 reader. Stimulation was carried out by a blue LED ($\lambda=470\pm 20$ nm) stimulation source for 40 s at 130°C. Irradiation was carried out using a $^{90}Sr/^{90}Y$ beta source built into the reader. The OSL signal was detected by a 9235QA photomultiplier tube through a U-340 filter with 7.5 mm thickness. For D_e determination, the SAR protocol was adopted. The preheating temperature was chosen to be 260°C for 10 s and the cut-heat was 220°C for 10 s. The final D_e is the average of the D_{es} of all aliquots, and the final D_e error is the standard error of the D_e distribution. For each sample, at least 12 aliquots were measured for D_e determination. The D_e was measured using SAR on quartz, and the aliquots that passed criteria checks were used for final D_e calculation.

Recycling ratios were between 0.90-1.1, and recuperation was relatively small. The cosmic ray dose rate was estimated for each sample as a function of depth, altitude and geomagnetic latitude. The concentration of U, Th and K was measured by neutral activation analysis (NAA). The elemental concentrations were then converted into the annual dose rate, taking into account the water content (lab measured) effect. The final OSL age is then: $D_e/Dose-rate$.

4 Results

The best geomorphic markers preserved in the study area are represented by a lacustrine terrace and two suites of fluvial terraces. The suites are distributed both upstream and downstream of the landslide debris, marking the evolutionary stages of

the valley, respectively, after and before the landslide emplacement. Fluvial and lacustrine terrace deposits mainly consist of gravel, sand, silt and clay, and conglomerates outcropping immediately upstream and downstream of the landslide pertain to inactive alluvial fans connected to a relict position of the valley floor, likely of a paleo-Seymareh river.

In the middle reach of the Seymareh River valley, the upstream geomorphic markers include: inactive, terraced conglomeratic alluvial fans (Cg_m), a terraced lacustrine deposit and a suite of four orders of fill terraces (named from Qt1_m to Qt4_m). The geometry of the terraced conglomerates (section A-A' in Fig. 2) can be associated with alluvial fans generated on the flanks of a former synclinal valley by streams likely forming the tributaries of a paleo-Seymareh river whose path was to the SW of the present one. The fill terraces are entrenched in the terraced lacustrine deposit of Seymareh Lake upstream of the landslide, in the area where Harrison and Falcon (1938), Roberts and Evans (2013) and Shoaiei (2014) hypothesized the natural damming lake could be extended (Figs. 5 and 6). Prograding lacustrine fan deltas formed by tributaries of Seymareh Lake have been recognized, which likely formed during the emptying phase of the lake. In this sector, we successfully dated 4 samples (SEY4, SEY5, SEY6, and SEY8; Table 1 and supplementary material); in particular the lacustrine deposit at two different stratigraphic levels, 560 and 590 m a.s.l., which provided OSL ages of 10.4±0.90 ka (sample SEY8) and 7.37±0.73 ka (sample SEY4), respectively. The OSL age of 17.9±1.50 ka (SEY6) obtained for an alluvial deposit at the base of the lacustrine deposits is coherent with the age of emplacement of the SL, as already inferred by Roberts and Evans (2013).

A suite of 2 strath terraces and a flood plain shaped onto the landslide debris along the Seymareh River gorge have been identified (Figs. 7a and 7b), which are important markers of the evolution of the natural dam because they formed after its cut likely due to an overflow of the damming lake. Here, we successfully dated one sample taken on a strath terrace (SEY9; Table 1 and supplementary material), which provided an age of 6.59±0.49 ka as time constrain for the initial stage of lake emptying.

In the lower reach of the Seymareh River valley, the downstream geomorphic markers include: inactive, terraced conglomeratic alluvial fans (Cg_l) and a suite of four orders of fill terraces (named from Qt1_l to Qt4_l) downstream of the SL (Figs. 7c and 8). Here, we successfully dated three samples from the fill terraces deposits (SEY3, SEY10, SEY11; Table 1 and supplementary material). The ages obtained provide useful time constraints to the main depositional events during the pre-failure valley evolution. Minimum ages of 373±34 ka and 312±45 ka have been obtained for samples SEY3 and SEY10, respectively, since these samples were saturated due to their low concentration of quartz grains, and SEY11 was dated at 60±5 ka.

The above-described geomorphic markers of the Seymareh River valley have been mapped and reported in morpho-stratigraphic profiles. The most significant landforms for the valley slope evolution are presented with a detail for the post-failure fluvial and lacustrine terrace suites upstream of the landslide dam (Fig. 6) and the pre-failure fluvial terrace suite downstream of the landslide dam (Fig. 8), respectively.

Figures 9 and 10 report the longitudinal profile of Seymareh River, along which, in addition to the geomorphic markers, were projected also: the benchmarks of the basal contact of the Quaternary deposits on the bedrock, the projection of points

corresponding to the top of the SL debris, the upstream and downstream limits of the landslide, the location of the OSL sampling, and the projection of the outcrop of the Bakhtiari Formation (Fig. 9), which is rarely preserved and marks the initial alluvial infill of the Seymareh River valley. Figure 9 shows the height distribution of the pre-failure geomorphic markers. The benchmarks along Seymareh River indicate a mostly bedrock channel, and the longitudinal profile is characterized by two knickpoints located upstream of the SL and downstream of the lowest suite of alluvial terraces, as indicated by the black arrows. The geomorphic markers downstream and upstream do not belong to the same suite of terraces, as their projections along Seymareh River do not have any topographic correlation to each other (Fig. 9 and 10). The tops of all the fluvial terraces downstream of the SL are located lower in height than the most important knickpoint located immediately upstream and sculpted in the bedrock. Figure 10 shows the height distribution of the post-failure geomorphic markers. The markers are represented by: i) the horizontal lacustrine terrace formed by the incision of the deposits pertaining to the Seymareh Lake, formed as a consequence of the landslide damming; ii) the two levels of the strath terraces and a flood plain formed on the landslide debris during the initial stages of dam cutting and emptying of the lake; and iii) the four fill terrace levels formed after the emptying of the Seymareh Lake.

The geomorphological field survey, supported by a remote survey based on optical satellite and aerial images, also allowed us to demonstrate the evidence of gravity-induced features of the downslope dipping strata, along the scar of the SL (Fig. 11).

Clear evidence of MRC driving towards stress concentration and failure has been recognized in gravity-induced folding within the thin-layered Pabdeh Formation just below the sliding surface of the SL (i.e., that cannot be ascribed to parasitic structural folding). Furthermore, impressive buckling of the downslope dipping strata, which crop on the sliding surface of the SL, have been interpreted as a release of concentrated stresses due to the post-failure rebound caused by the collapsed rock mass.

5 Discussion

5.1 Constraints to pre-failure valley evolution

The longitudinal profile of the Seymareh River and the geomorphic markers preserved mainly downstream of the landslide dam provided new constraints on the pre-failure valley evolution. The major knickpoint located immediately upstream of the SL is the most interesting to be analyzed in relation with the landslide event. Its shape in the longitudinal profile clearly let us identify it as a “slope-break knickpoint” (Kirby and Whipple, 2012; Boulton et al., 2014), thus developed as a knickpoint retreating in response to a persistent perturbation to the fluvial system (Tucker and Whipple, 2002), as frequently observed in tectonically active regions. The location of this knickpoint upstream of the SL and the outcrop of the basal contact of the landslide at the bottom of the Seymareh River gorge (Fig. 7a) suggests that this shape of the longitudinal profile was already

developed before the failure, meaning that the erosion wave which generated the knickpoint affected the SL slope foot before the failure occurrence.

The poorly preserved, well-cemented alluvial fan conglomeratic deposits outcropping upstream of the landslide lie on the Miocene Agha Jari Formation, at a higher elevation than the outcrops of the Bakhtiari Formation. Their remnants are aligned in correspondence with the axis of a relict synclinal valley, likely corresponding to a very early stage (Pliocene?) of the Seymareh valley evolution. On the other hand, the conglomerate deposits outcropping downstream of the landslide (Cg₁) are closer in height to the major knickpoint, thus suggesting that they were in equilibrium with a local base level corresponding to the early propagation of the major knickpoint. Furthermore, they must be younger than the Bakhtiari Formation, which is preserved at higher elevation.

The alluvial terraces preserved downstream of the SL likely mark the valley evolutionary stages during the major knickpoint retreat (Demoulin et al., 2017). Along the longitudinal river profile, the uppermost outcrops of each level of this terrace suite were swept away by the landslide, which unfortunately prevents estimation of the rates of knickpoint retreat. Nonetheless, according to what was observed by Bridgland et al. (2017) about river terrace development in the NE Mediterranean region, the sedimentation phases should correspond to cold periods. In particular, Bridgland et al. (2012) observed, in the valleys of the Tigris and Ceyhan in Turkey, the Kebir in Syria and the trans-border rivers Orontes and Euphrates, a regular terrace formation in synchrony with 100 ka climatic cycles that can be correlated with MIS 12, 10, 8, 6 and 4-2. Therefore, the minimum ages obtained for the SEY3 and SEY10 samples could be reasonably extended to 478 ka (MIS 12) and 374 ka (MIS 10), respectively, and the OSL age of the SEY11 fits well with the Last Glacial Period.

20 **5.2 Constraints to post-failure valley evolution**

The geomorphic markers preserved upstream of the landslide dam provided new constraints on the geomorphic response of the Seymareh valley to the 44 Gm³ natural dam. Such a response was first the formation of three lakes (Seymareh, Jaidar and Balmak; Fig. 4) whose persistence and evolution is well recorded by the deposits outcropping in the valley. In this regard, the estimation of a sedimentation rate of 10 mm y⁻¹ in the Seymareh Lake was obtained using the OSL ages of 10.4±0.90 ka and 7.37±0.73 ka for the lacustrine deposit sampled at 560 and 590 m a.s.l., respectively.

The strath terrace sculpted on the landslide deposit and dated at 6.59±0.49 ka constrains the cut of the natural dam due to overflow which caused the lake to empty. The lake overflow was likely caused by the gradual filling of the reservoir with lacustrine deposits, which progressively reduced the dam infiltration section. Nevertheless, the possible role of groundwater seepage within the pervious natural dam in balancing the Seymareh River discharge and delaying the dam overflow remains a questionable topic to be approached and solved in future studies. Despite their interpretation as progressively younger lacustrine deposits by Yamani et al. (2012), the four terrace levels entrenched in the terraced lacustrine deposit show a longitudinal downstream gradient, which, along with their sedimentological characters, identify them as fill terraces. Furthermore, the OSL age obtained for the lacustrine deposit at the base of the Qt2_m terrace (sample SEY8) is 10.4±0.90 ka, demonstrating that the suite of alluvial terraces is entrenched into the same (and unique) lacustrine deposit. The OSL age

of 4.49 ± 0.48 ka obtained for the Qt1_m terrace (sample SEY5) provides time constraints to the emptying phase of Seymareh Lake. Such time constraints are fine-tuned by the age of the strath terrace formed on the landslide debris, which corroborate the initial stage of lake emptying at 6.59 ± 0.49 ka (SEY9). As indicated by the age of the Qt1_m terrace (of 4.49 ± 0.48 ka), the Seymareh Lake likely persisted up to ~ 5 ka, much longer than the 935 years estimated by Shoaiei (2014). Since the top of the lacustrine deposit lies at 630 m a.s.l., an increased sedimentation rate of ~ 17 mm y^{-1} can be inferred for the late stage of the lake evolution, which is in agreement with an increased sediment yield from tributaries during the early stages of lake emptying (Fig. 5d).

The overflow, at 6.59 ± 0.49 ka allows us to calculate the erosion rate affecting the landslide deposit after the overflow. The ratio between the thickness of the eroded sediment (~ 120 m) and the time elapsed since the beginning of the process (~ 6.59 ky) allows estimation of an erosion rate of 1.8 cm y^{-1} for Seymareh River along the gorge. The cut of the landslide dam induced a new change in the fluvial base level, bringing the slope-to-valley floor system into disequilibrium. For this reason, a dense drainage system was set on the scar area, which, due to the high erodibility and low permeability of the less competent Pabdeh-Gurpi Formation immediately below the sliding surface on the Kabir-kuh ridge NE slope, has generated the badlands mapped in Fig. 6.

15

5.3 Evolutionary model of the Seymareh River valley

The landscape evolution of the Seymareh River valley before and after the failure occurrence can be summarized in the following six phases:

1. Setting of a paleo-Seymareh river into a synclinal valley, likely developed in the Pliocene, to the west of the present position of the Seymareh River and deposition of fan deposits (Cg_m) (Fig. 12a).
2. Development of the valley with local base level correlated to the Seymareh longitudinal profile segment upstream of the major knickpoint along the Seymareh River and coeval to the deposition of the Bakhtiari Formation (late Pliocene-early Pleistocene) (Fig. 12b).
3. Emplacement of the downstream fan deposits corresponding to the Cg_l conglomerates (early Pleistocene) and generation of the four orders of Middle-Late Pleistocene alluvial terraces (Qt1_1-Qt4_1) preserved downstream of the landslide and formed during the progressive migration of the major knickpoint, which is presently located upstream of the landslide (Fig. 12c).
4. SL event (~ 10 ka), according to the ^{14}C ages by Roberts and Evans (2013) and to the OSL ages provided in this work for the lacustrine deposits (Lac) (Fig. 12d).
5. Formation and permanence of the Seymareh Lake (~ 10 -6.6 ka), according to the ^{14}C estimated ages by Roberts and Evans (2013) and to the OSL ages provided in this study for the lacustrine deposits (Lac) (Fig. 12e). The progressive infilling of the lake reservoir progressively reduced the infiltration section on the upstream side of the landslide dam. The presence of a minor emissary on the downstream side of the landslide debris cannot be excluded.

30

6. Overflow of the lake and cut of the natural dam with formation of the first strath terrace (6.59 ± 0.49 ka), followed by a second strath terrace and a flood plain during the emptying of the lake, which upstream is associated with the sedimentation of a fluvio-lacustrine sequence at the top of the lacustrine sediments (Fig. 12f).
7. Complete emptying of the lake and generation of the suite of fill terraces entrenched in the deposits of Seymareh Lake (4.5 ka. - Present) (Fig. 12g).

5.4 Implications of the evolutionary model for future back-analysis of the SL

According to the multi-modeling approach proposed by Martino et al. (2017), Quaternary landscape evolution modeling of slope-to-valley floor systems plays a key role as a tool for chronological constraints to the creep evolution of entire slopes (Bozzano et al., 2016; Della Seta et al., 2017).

The geomorphic processes developed before the failure of the SL likely acted as predisposing factors for MRC processes in the rock mass successively collapsed. Kinematic freedom, both at the top and on the fold flank was created by the incising network of streams that dissect the Asmari Formation carbonate caprock following the major joint set in the Asmari Formation already described in Roberts and Evans (2013; and references therein). In particular, the headward erosion of streams towards the anticline's structural high described by Oberlander (1968), caused the expansion of the fold axial basins through the softer units, determining the upslope kinematic freedom. In the timing proposed by Tucker and Slingerland (1996) the latter was reached at approximately 1.6 ka. Stress release at the slope base was definitely produced by the Middle-Late Pleistocene upstream migration of the knickpoint along the Seymareh River longitudinal profile. Unfortunately, since the emplacement of the landslide swept away the uppermost outcrops of the alluvial terraces formed in response to the upstream knickpoint migration, the rate of knickpoint migration cannot be inferred. Nonetheless, an elapsed time-to-failure on the order of 10^2 ky, since the kinematic freedom at the slope base was reached, can be reasonably estimated by the age of the oldest terrace in the lower reach of the river minus the age of the landslide occurrence.

It is noteworthy that the stratigraphy of the source rock mass, also described in detail by Roberts and Evans (2013), accounts for different rheological behaviors, which could have induced differential strain rates within the slope leading to failure according to a MRC process. More particularly, the time-dependent visco-plastic behavior, more typical of clayey and marly deposits, which have lower viscosity values, can justify time-dependent (creep) strains which could have generated high stress concentration within the higher viscosity level over time (i.e., mostly characterized by elasto-plastic rheology), inducing their cracking and leading to failure. In fact, a stiffness contrast exists between the upper member of the Pabdeh Formation and the overlying Asmari Formation. The attitude of the strata is moderately dipping downslope (15° - 20°), and a reduced lateral confining effect is due to continental and epicontinental deposits ascribable to the Gachsaran and Agha Jari Formations. Moreover, the low dip angle of the strata reduces the vertical thickness of the Asmari Formation caprock, which was completely eroded by Seymareh River during its engraving, thus allowing the sliding mechanism of the Pabdeh and Asmari layered formations.

Therefore, the results of this work have implications for a future back-analysis through stress-strain numerical modeling of the SL because they can be used to constrain the elapsed time since MRC initiation and ultimate failure conditions. Such a perspective is to be regarded as a key challenge for dimensioning such an end member event in regard to both time and space distribution as well as for evaluating the possible role of impulsive triggering actions (i.e., strong to very strong earthquakes) in anticipating the time-to-failure of the slope.

6 Conclusion

In a multi-modeling approach to the study of MRC processes affecting slopes at a large space-time scale, the performed geomorphic analysis allowed us to constrain the evolution of the Seymareh River valley in the northwestern Zagros Mts., before and after the failure of the largest landslide ever recorded on the exposed Earth surface. The identification and OSL dating of different suites of lacustrine, alluvial and strath terraces constrained in time the major pre- and post-failure evolutionary steps of the river valley system.

The oldest geomorphic markers in the Seymareh River valley are represented by relict conglomerates preserved upstream of the landslide, which demonstrate the early (Pliocene?) position of a paleo-Seymareh river flowing into a synclinal valley close to the northeastern flank of the Kabir-kuh fold.

Drainage evolution associated with the growth of the Kabir-kuh fold was characterized by the deep incision of the stream network, which allowed the kinematic release of the rock mass involved in the Seymareh giant landslide. Such a stream incision was accompanied by the retreat of a major “slope-break knickpoint” along the Seymareh longitudinal profile, time-constrained by the age of a suite of river fill terraces. According to the age of pre-failure terraces, in the middle-late Pleistocene the erosion wave reached the portion of the Kabir-kuh fold that ~10 ka was affected by the SL. According to the timing of the landscape evolution model proposed by Tucker and Slingerland (1996), the upper slope underwent kinematic release about 1.6 ka. Therefore, the collapse was prepared by MRC processes acting over a time window of 10^2 ky;

The geomorphic response to the landslide dam consisted in the formation of three lakes, among which Seymareh Lake persisted for ~3500 years before its emptying phase started ~6.6 ka due to lake overflow. A sedimentation rate of 10 mm y^{-1} was estimated for the lacustrine deposits, which increased up to 17 mm y^{-1} during the early stage of lake emptying due to the increased sediment yield from the lake tributaries. Since ~4.5 ka, a suite of four alluvial terraces upstream of the landslide demonstrates the alternating erosion/deposition phases of the re-established Seymareh River.

An incision rate of 1.8 cm y^{-1} was estimated since the beginning of the landslide cut by Seymareh River, and such a strong erosion started propagating up to the landslide source area where badlands developed, eroding the marly Pabdeh-Gurpi Formation.

The results obtained here provide new constraints to the valley evolution in view of future stress-strain numerical modeling of the MRC process that involved the SL slope before its generalized collapse. Such a modeling could also be considered to

discuss the possible role of impulsive triggering (earthquakes) in anticipating the time-to-failure due to the gravity-driven deformational processes.

Acknowledgments

This work has been possible thanks to the International Agreement between Sapienza University of Rome and Kharazmi University of Tehran and to the associated mobility funds by Sapienza (Grant no. AI2615R3W4). We are particularly grateful to Mohammad Moumeni Taromsari for his efforts and fruitful scientific contributions to the field work.

References

- Alavi, M.: Regional stratigraphy of the Zagros fold-thrust belt of Iran and its proforeland evolution. *Am. J. Sci.*, 304(1), 1-20, doi: 10.2475/ajs.304.1.1, 2004.
- 10 Agard, P., Omrani, J., Jolivet, L. and Mouthereau, F.: Convergence history across Zagros (Iran): Constraints from collisional and earlier deformation, *Int. J. Earth Sci.*, 94(3), 401–419, doi:10.1007/s00531-005-0481-4, 2005.
- Binnie, S. A., Phillips, W. M., Summerfield, M. A. and Fifield, L. K.: Tectonic uplift, threshold hillslopes, and denudation rates in a developing mountain range, *Geology*, 35(8), 743–746, doi:10.1130/G23641A.1, 2007.
- Boulton, S. J., Stokes, M. and Mather, A. E.: Transient fluvial incision as an indicator of active faulting and Plio-Quaternary uplift of the Moroccan high Atlas, *Tectonophysics*, 633(1), 16–33, doi:10.1016/j.tecto.2014.06.032, 2014.
- 15 Bourne, J. A. and Twidale, C. R.: Neglected and cryptostructural effects in drainage development, *Cadernos do Laboratorio Xeolóxico de Laxe*, Vol. 36, pp. 41 - 60, 2011.
- Bozzano, F., Martino, S., Montagna, A. and Prestininzi, A.: Back analysis of a rock landslide to infer rheological parameters, *Eng. Geol.*, 131–132, 45–56, doi:10.1016/j.enggeo.2012.02.003, 2012.
- 20 Bozzano, F., Della Seta, M. and Martino, S.: Time-dependent evolution of rock slopes by a multi-modelling approach, *Geomorphology*, 263, 113–131, doi:10.1016/j.geomorph.2016.03.031, 2016.
- Bridgland, D. R., Westaway, R., Romieh, M. A., Candy, I., Daoud, M., Demir, T., Galiatsatos, N., Schreve, D. C., Seyrek, A., Shaw, A. D., White, T. S. and Whittaker, J.: The River Orontes in Syria and Turkey: Downstream variation of fluvial archives in different crustal blocks, *Geomorphology*, 165–166, 25–49, doi:10.1016/j.geomorph.2012.01.011, 2012.
- 25 Bridgland, D. R., Demir, T., Seyrek, A., Daoud, M., Abou Romieh, M. and Westaway, R.: River terrace development in the NE Mediterranean region (Syria and Turkey): Patterns in relation to crustal type, *Quat. Sci. Rev.*, 166, 307–323, doi:10.1016/j.quascirev.2016.12.015, 2017.
- Burbank, D. W. and Anderson R. S.: *Tectonic Geomorphology*. John Wiley and Sons, Second Edition, Ltd., Chichester, West Sussex, UK, 2012.


- Casciello, E., Verges, J., Saura, E., Casini, G., Fernandez, N., Blanc, E., Homke, S. and Hunt, D. W.: Fold patterns and multilayer rheology of the Lurestan Province, Zagros Simply Folded Belt (Iran), *J. Geol. Soc. London.*, 166(5), 947–959, doi:10.1144/0016-76492008-138, 2009.
- Chigira, M.: Long-term gravitational deformation of rocks by mass rock creep, *Eng. Geol.*, 32(3), 157–184, doi:10.1016/0013-7952(92)90043-X, 1992.
- 5 Della Seta, M., Esposito, C., Marmoni, G. M., Martino, S., Scarascia Mugnozza, G. and Troiani, F.: Morpho-structural evolution of the valley-slope systems and related implications on slope-scale gravitational processes: New results from the Mt. Genzana case history (Central Apennines, Italy), *Geomorphology*, 289, 60–77, doi:10.1016/j.geomorph.2016.07.003, 2017.
- 10 Demoulin, A., Mather, A. and Whittaker, A.: Fluvial archives, a valuable record of vertical crustal deformation, *Quat. Sci. Rev.*, 166, 10–37, doi:10.1016/j.quascirev.2016.11.011, 2017.
- Eberhardt, E., Stead, D. and Coggan, J. S.: Numerical analysis of initiation and progressive failure in natural rock slopes-the 1991 Randa rockslide, *Int. J. Rock Mech. Min. Sci.*, 41(1), 69–87, doi:10.1016/S1365-1609(03)00076-5, 2004.
- Golonka, J.: Plate tectonic evolution of the southern margin of Eurasia in the Mesozoic and Cenozoic, *Tectonophysics*, 15 381(1–4), 235–273, doi:10.1016/j.tecto.2002.06.004, 2004.
- Griffiths, H. I., Schwalb, A. and Stevens, L. R.: Environmental change in southwestern Iran: the Holocene ostracod fauna of Lake Mirabad. *Holocene*, 11(6), 757-764, doi:10.1191/09596830195771, 2001.
- Harrison, J. V., and Falcon, N. L.: The Saidmarreh landslip, south-west Iran, *Geogr. J.*, 89, 42-47, 1937.
- Harrison, J. V., and Falcon, N. L.: An ancient landslip at Saidmarreh in southwestern Iran, *J. Geol.*, 46, 296-309, 1938.
- 20 Hatzfeld, D., Authemayou, C., Van der Beek, P., Bellier, O., Lavé, J., Oveisi, B., Tatar, M., Tavakoli, F., Walpersdorf, A. and Yamini-Fard, F.: The kinematics of the Zagros Mountains (Iran), *Geol. Soc. London, Spec. Publ.*, 330(1), 19–42, doi:10.1144/SP330.3, 2010.
- Homke, S., Vergés, J., Garcés, M., Emami, H. and Karpuz, R.: Magnetostratigraphy of Miocene-Pliocene Zagros foreland deposits in the front of the Push-e Kush Arc (Lurestan Province, Iran), *Earth Planet. Sci. Lett.*, 225(3–4), 397–410, 25 doi:10.1016/j.epsl.2004.07.002, 2004.
- Hungr, O., Evans, S. G., and Hutchinson, J. N.: A Review of the Classification of Landslides of the Flow Type, *Environ. Eng. Geosci.*, 7(3), 221-238, doi:10.2113/gsegeosci.7.3.221, 2001.
- Hutchinson, M., Xu, T. and Stein, J.: Recent Progress in the ANUDEM Elevation Gridding Procedure, *Geomorphometry*, 19–22, doi:10.1002/osp4.29, 2011.
- 30 James, G. A. and Wynd, J. G.: Stratigraphic nomenclature of Iranian oil consortium agreement area. *AApG Bulletin*, 49(12), 2182-2245, 1965.
- Kirby, E. and Whipple, K. X.: Expression of active tectonics in erosional landscapes, *J. Struct. Geol.*, 44, 54–75, doi:10.1016/j.jsg.2012.07.009, 2012.

- Larsen, I. J. and Montgomery, D. R.: Landslide erosion coupled to tectonics and river incision, *Nat. Geosci.*, 5(7), 468–473, doi:10.1038/ngeo1479, 2012.
- Macleod, J. H.: Kabir Kuh, 1:100000 Geological Map. Iran Oil Operating Companies, Geological Exploration Division, Tehran, 1970.
- 5 Martino, S., Della Seta, M. and Esposito, C.: Back-analysis of rock landslides to infer rheological parameters, in: *Rock Mechanics and Engineering, Volume 3: Analysis, Modeling and Design*, First Edition, Xia-Ting Feng Editor, London, UK, 237-269, 2017.
- McQuarrie, N.: Crustal scale geometry of the Zagros fold-thrust belt, Iran, *J. Struct. Geol.*, 26(3), 519–535, doi:10.1016/j.jsg.2003.08.009, 2004.
- 10 Montgomery, D. R., and Brandon, M. T.: Topographic controls on erosion rates in tectonically active mountain ranges, *Earth. Planet. Sc. Lett.*, 201(3-4), 481-489, doi:10.1016/S0012-821X(02)00725-2, 2002.
- Montgomery D. R. and Fofoula-Georgiou E.: Channel Network Source Representation Using Digital Elevation Models, *Water. Resour. Res.*, Vol. 29, No. 12, pp. 3925-3934, doi:10.1029/93WR02463, 1993.
- Mouthereau, F., Lacombe, O. and Vergés, J.: Building the Zagros collisional orogen: Timing, strain distribution and the
 15 dynamics of Arabia/Eurasia plate convergence, *Tectonophysics*, 532–535, 27–60, doi:10.1016/j.tecto.2012.01.022, 2012.
- Murray, A. S., and Olley, J. M.: Precision and accuracy in the optically stimulated luminescence dating of sedimentary quartz: a status review, *Geochronometria*, 21(1), 1-16, 2002.
- Oberlander, T. M.: The Zagros streams: a new interpretation of transversedrainage in an orogenic zone, *Syracuse Geographical Series*, Syracuse University Press, New York, 1965.
- 20 Oberlander, T. M.: The origin of the Zagros defiles. *The Cambridge History of Iran*, 1: The Land of Iran, Fisher W.B (Ed.), Cambridge University Press, Cambridge, UK, 195-211, 1968.
- Oberlander, T. M.: Origin of drainage transverse to structures in orogens. In: *Tectonic Geomorphology*, Proceedings of the 15th Binghamton Geomorphology Symposium 1984. Morisawa, M. and Hack, J. T. (eds.), Allen & Unwin, Boston, MA, 155–182, 1985.
- 25 Paul, A., Hatzfeld, D., Kaviani, A., Tatar, M. and Péquegnat, C.: Seismic imaging of the lithospheric structure of the Zagros mountain belt (Iran), *Geol. Soc. London, Spec. Publ.*, 330(1), 5–18, doi:10.1144/SP330.2, 2010.
- Ramsey, L. A., Walker, R. T., and Jackson, J.: Fold evolution and drainage development in the Zagros mountains of Fars province, SE Iran, *Basin. Res.*, 20(1), 23-48, doi:10.1111/j.1365-2117.2007.00342.x, 2008.
- Rajabi, A. M., Mahdaviifar, M. R., Khamehchiyan, M. and Del Gaudio, V.: A new empirical estimator of coseismic landslide
 30 displacement for Zagros Mountain region (Iran), *Nat. Hazards*, 59(2), 1189–1203, doi:10.1007/s11069-011-9829-1, 2011.
- Roberts, N. J.: Structural and geologic controls on gigantic (1 Gm³) landslides in carbonate sequences : case studies from the Zagros Mountains, Iran and Rocky Mountains, Canada, Master's thesis, University of Waterloo, Canada, 2008.
- Roberts, N. J. and Evans, S. G.: The gigantic Seymareh (Saidmarreh) rock avalanche, Zagros Fold–Thrust Belt, Iran, *J. Geol. Soc. London.*, 170(4), 685–700, doi:10.1144/jgs2012-090, 2013.

- Schmidt, K. M. and Montgomery, D. R.: Limits to Relief, *Science*, Vol. 270, 617–621, doi:10.1126/science.270.5236.617, 1995.
- Setudehnia, A. and Perry, J. T. O.'B.: Dal Parri. 1:100000 Geological Map. Iran Oil Operating Companies, Geological Exploration Division, Tehran, 1967.
- 5 Shafiei, A. and Dusseault, M. B.: Geomechanical Properties of a Conglomerate from Iran Geomechanical Properties of a Conglomerate from Iran, In The 42nd US Rock Mechanics Symposium (USRMS), American Rock Mechanics Association, doi:10.13140/RG.2.1.1722.7684, 2008.
- Shoaei, Z.: Mechanism of the giant Seimareh Landslide, Iran, and the longevity of its landslide dams, *Environ. Earth Sci.*, 72(7), 2411–2422, doi:10.1007/s12665-014-3150-8, 2014.
- 10 Stampfli, G. M., and Borel, G. D.: A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons, *Earth. Planet. Sc. Lett.*, 196(1), 17-33, doi:10.1016/S0012-821X(01)00588-X, 2002.
- Stead, D., Eberhardt, E. and Coggan, J. S.: Developments in the characterization of complex rock slope deformation and failure using numerical modelling techniques, *Eng. Geol.*, 83(1–3), 217–235, doi:10.1016/j.enggeo.2005.06.033, 2006.
- 15 Takin, M., Akbari, Y. and Macleod, J.H.: Pul-E Dukhtar. 1:100000 Geological Map. Iran Oil Operating Companies, Geological Exploration Division, Tehran, 1970.
- Talbot, C. J. and Alavi, M.: The past of a future syntaxis across the Zagros, *Geol. Soc. London, Spec. Publ.*, 100(1), 89–109, doi:10.1144/GSL.SP.1996.100.01.08, 1996.
- Tucker, G. E. and Slingerland, R.: Predicting sediment flux from fold and thrust belts, *Basin Res.*, 8(3), 329–349, doi:10.1046/j.1365-2117.1996.00238.x, 1996.
- 20 Tucker, G. E. and Whipple, K. X.: Topographic outcomes predicted by stream erosion models: Sensitivity analysis and intermodel comparison, *J. Geophys. Res. Solid Earth*, 107(B9), 1-16, doi:10.1029/2001JB000162, 2002.
- Vergés, J., Goodarzi, M., Emami, H., Karpuz, R., Efstathiou, J. and Gillespie, P.: Multiple detachment folding in Pusht-e Kuh arc, Zagros: Role of mechanical stratigraphy, *Thrust fault-related Fold, AAPG Mem.*, 69–94, doi:10.1306/13251333M942899, 2011.
- 25 Watson R. A. and Wright H. E.: The Saidmarreh landslide, Iran, Schumm, S. A. and Bradley, W. C. (eds), *United States Contributions to Quaternary Research*, Geological Society of America, Special Papers, 123, pp. 115–139, 1969.
- Wilson, J. P., and Gallant, J. C.: *Terrain analysis: principles and applications*, John Wiley and Sons, New York, USA, 2000.
- Wintle, A. G., and Murray, A. S.: A review of quartz optically stimulated luminescence characteristics and their relevance in single-aliquot regeneration dating protocols, *Radiat. Meas.*, 41(4), 369-391, doi:10.1016/j.radmeas.2005.11.001, 2006.
- 30 Yamani, M., Goorabi, A. and Azimirad, S.: Large Seymareh Landslide and Lake Terraces Sequence, *Physical Geography Research, Quarterly*, Volume 44, Issue 4, pp. 43- 60, 2012.

Yamini-Fard, F., Hatzfeld, D., Tatar, M., and Mokhtari, M.: Microearthquake seismicity at the intersection between the Kazerun fault and the Main Recent Fault (Zagros, Iran), *Geophys. J. Int.*, 166(1), 186-196, doi:10.1111/j.1365-246X.2006.02891.x, 2006.

Zhao, S., Chigira, M., Wu, X.: Gigantic rockslides induced by fluvial incision in the Diexi area along the eastern margin of the Tibetan Plateau, *Geomorphology*, <https://doi.org/10.1016/j.geomorph.2019.04.008>, 2019.



10 **Figure 1: Simplified structural map of the Zagros mountain range with location of the. MZT, Main Zagros Thrust; HZF, High Zagros Fault; MFF, Mountain Front Fault; BR, Bala Rud fault zone (modified from Casciello et al., 2009).**




Figure 2: Geological Map, stratigraphic column and cross sections of the study area.




Figure 3: Evolution of the drainage network in the Zagros chain sector of the Kabir-kuh fold, according to the Oberlander's model and with the timing provided in the landscape evolution model by Tucker and Slingerland (1996). See the text for explanation of the four steps. (modified from Oberlander, 1985).





Figure 4: Overview and focus (in the red box) of the SL, Zagros Fold–thrust Belt, NW Iran. (modified from Google Earth®).



5 **Figure 5: Geomorphic markers upstream of the SL, represented by a suite of four orders of alluvial terraces entrenched in the lacustrine deposits (Lac) of Seymareh Lake upstream of the landslide, in the areas where Harrison and Falcon (1938), Roberts and Evans (2013) and Shoaie (2014) hypothesized the natural damming lake could be extended. a) Overall view of the suite of terraces; b) example of fluvial terrace deposit; c) example of lacustrine deposit; d) evidence of a prograding lacustrine fan delta formed by one of the right tributaries of Seymareh Lake during its early emptying phase.**





Figure 6: Map of the lacustrine and alluvial terrace suite and of the most significant landforms for the valley slope evolution upstream of the SL.



5 **Figure 7: Geomorphic markers upstream of the SL. a) A strath terrace and a flood plain developed over the landslide debris, which are important markers of the evolution of the natural dam since they testify to the moment of the overcoming of the damming lake and the overflowing of the river onto the landslide debris, respectively; b) detail of the strath terrace deposit sampled for OSL dating; c) The suite of fluvial terraces downstream of the SL; the Qt1 level is poorly preserved and not visible in this photo.**




Figure 8: Map of the alluvial terrace suite and of the most significant landforms for the valley slopes evolution downstream of the

5 SL.




Figure 9: Projection of the pre-failure geomorphic markers (upstream and downstream conglomerates; downstream fluvial terrace suite) along the longitudinal profile of Seymareh River. The obtained OSL ages are indicated.





Figure 10: Projection along the longitudinal profile of Seymarche River of the post-failure geomorphic markers (fluvial terraces and lacustrine deposits relative to the Seymarche Lake suite). The obtained OSL ages are indicated.

5



5 **Figure 11: Scar area of the SL. a) front view of the scar area with the location of sites where evidence of buckling has been recognized; b) ductile buckling deformations within the layers of the Upper Pabdeh Member which cannot be referred to parasitic structural folding due to their localisation just below the SL sliding surface and to their reduced persistency (both lateral and vertical) within the rock mass; c) brittle buckling deformation of the Taleh Zang Member along the scar area.**




Figure 12: Evolutionary model of the Seymareh River valley. See text for explanation. Traces and legend of geological cross-sections are reported in Fig. 2.

Table 1: OSL ages obtained for the geomorphic markers recognized in the Seymareh River valley. Detailed information and photos of sampling sites are available in the supplementary material.

SAMPLE	DESCRITPION	COORDINATES	ELEVATION (m a.s.l.)	OSL AGE (ka)	ERROR (ka)
SEY4	lacustrine deposit	33° 13.197'N 47° 18.382'E	590	7.37	±0.73
SEY5	alluvial terrace deposit (Qt1_m)	33° 13.437'N 47° 18.219'E	607	4.49	±0.48
SEY6	alluvial deposit beneath the lacustrine deposit	33° 13.291'N 47° 18.358'E	580	17.9	±1.50
SEY8	lacustrine deposit at the base of Qt2_m	33° 7.402'N 47° 28.795'E	560	10.4	±0.90
SEY9	deposit of a strath terrace on landslide debris	33° 4.462'N 47° 34.197'E	570	6.59	±0.49
SEY3	alluvial terrace deposit (Qt2_l)	32° 59.591'N 47° 46.144'E	485	≥373*	±34
SEY10	alluvial terrace deposit (Qt3_l)	32° 59.335'N 47° 46.071'E	436	≥312*	±45
SEY11	alluvial terrace deposit (Qt4_l)	32° 59.265'N 47° 45.869'E	400	60	±5

5