

Landslides as geological hotspots of CO₂ emission: clues from the instrumented Séchilienne landslide, Western European Alps.

Pierre Nevers¹, Julien Bouchez², Jérôme Gaillardet^{2,4}, Christophe Thomazo³, Delphine Charpentier¹,
5 Laëticia Faure² and Catherine Bertrand¹

¹UMR Chrono-Environnement 16 route de Gray, 25000 Besançon, France

²Université de Paris, Institut de physique du globe de Paris, CNRS, F-75005 Paris, France

³UMR CNRS/uB6282 Biogéosciences 6 Boulevard Gabriel, 21000 Dijon, France

10 ⁴Institut Universitaire de France, 75231 Paris, France

Correspondence to: Pierre Nevers (pierre.nevers@univ-fcomte.fr)

Abstract. This study makes use of a highly instrumented active landslide observatory (9 years of data) in the French Alps, the Séchilienne slope. Here we use a combination of major element chemistry and isotopes ratios (⁸⁷Sr/⁸⁶Sr, δ³⁴S) measured in different water types of the stable and unstable part of the Séchilienne instability to assess the contribution of the different
15 lithologies of the slope and the chemical weathering mechanisms. Chemical and isotopic ratios are used to characterize weathering processes and the origin of waters and their flowpaths through the massif. A mixing model allows us to allocate the different major elements to different sources, to identify secondary carbonate formation as a major process affecting solutes in the subsurface waters of the instability, and to quantify the involvement of sulfuric and carbonic acids as a source of protons.

20 We show that the instability creates favorable and sustained conditions for the production of sulfuric acid by pyrite oxidation, by opening new fractures and supplying fresh reactive surfaces. We clearly identify the contribution of the dissolution of each mineral phase to the chemistry of the waters, with a clear role of remote gypsum dissolution to the sulfate budget in the sampled waters. We are also able to refine the pre-existing hydrogeological views on the local water circulation and water flow paths in the instability by showing the hydrological connectivity of the different zones. Overall,
25 our results show that the Séchilienne landslide, despite its role in accelerating rock chemical and physical weathering, acts as a geological source of CO₂ to the atmosphere. If generalizable to other large instabilities in mountain ranges, this study illustrates the complex coupling between physical and chemical erosion and their impact on the carbon cycle and global climate. The study also highlights the importance of distinguishing between sulfite oxidation and gypsum dissolution as a source of sulfate ions to rivers, particularly in mountain ranges.

30 1 Introduction

The weathering of rocks plays a key role in the chemical and climatic evolution of the Earth surface and is one of the geological processes that impacts atmospheric CO₂ concentration. When carbonic acid is the proton supplier, silicate weathering removes carbon dioxide from the atmosphere (Lerman et al., 2007, Berner and Berner, 2012). However, the oxidative dissolution of sulfides (*e.g.* pyrite FeS₂) produces sulfuric acid that can act as an alternative proton supplier to
35 chemical weathering reactions. Although not directly influencing atmospheric CO₂, silicate weathering by sulfuric acid does reduce the potential of rock weathering for CO₂ sequestration by "removing" silicates from the Earth surface and thus limiting their weathering by carbonic acid. When sulfuric acid reacts with carbonate minerals, dissolved inorganic carbon is added to ambient waters which leads on the long term to CO₂ release towards the atmosphere (Lerman et al., 2007, Calmels et al., 2007, S.-L. Li et al., 2008, Torres et al., 2014). The relevance of this process for the global carbon cycle is two-fold.

40 First, even though carbonate rocks do not constitute the major fraction of the rock types exposed at the Earth surface, the dissolved products of carbonate dissolution dominate global weathering fluxes (Gaillardet et al., 1999) as carbonate minerals dissolve several orders of magnitude faster than silicates (Lasaga 1984). Second, because weathering by sulfuric acid is mainly limited by the supply of sulfide minerals to the Earth surface, it is particularly prominent in active mountain belts characterized by high erosion rates (Calmels et al., 2007; Torres et al., 2016; Blattmann et al., 2019). Within these
45 tectonically active environments, landslides are likely to be hotspots of sulfuric acid production, carbonate weathering, and CO₂ release (Emberson et al., 2015, 2018). Indeed, slope instability leads to sustained grain comminution and fractures opening, thereby providing a continuous supply of contact surfaces between water, air, and minerals that can in particular allow for sulfuric acid production and carbonate mineral weathering (Binet et al., 2009; Bertrand et al., 2014).

Here we explore the hypothesis that slope instability can constitute a mechanism promoting coupled sulfide oxidation and
50 carbonate weathering, in a contribution to the study of the role active mountain ranges play on the global carbon cycle (Raymo and Ruddiman, 1992; Hilton and West, 2020). We focus on the S echilienne slope instability located in the French Alps. This site of active, slow landsliding serves as an observatory for landslide processes and has been the subject of previous hydrogeological and geophysical investigation (Vengeon, 1998; Guglielmi et al., 2002; Meric et al., 2005; LeRoux et al., 2011; Vallet et al., 2015; Lajaunie et al., 2019). We combine measurements of the concentration of major elements and
55 of the isotope composition of strontium and sulfur (⁸⁷Sr/⁸⁶Sr, δ³⁴S) dissolved in groundwater and springs to estimate the contribution of different rock types to the dissolved species produced by weathering reactions in the landslide. In particular, we estimate the relative role of different acid types (carbonic vs. sulfuric) and of two rock types (silicates vs. carbonates), and evaluate the role played by secondary carbonate formation on the solute budget of percolating waters. Besides shedding light on the global impact of landsliding on atmospheric CO₂, Sr and S isotopes coupled to water chemistry allow for a
60 quantitative analysis of solute sources in natural waters and of the chemical evolution of natural waters, which in turn opens the possibility to improve existing hydrogeological model in complex environments such as landslides.

2 Study area


2.1 Geological setting

The "S echilienne" site hosts a highly-instrumented, continuously-monitored landslide, part of the French National Landslide
65 Observatory (OMIV, <http://www.ano-omiv.cnrs.fr/>). The S echilienne massif is located at the SW border of the Paleozoic crystalline Belledonne mountain range in the French Alps, 20 km southeast of Grenoble (Is ere, France; Fig. 1). The active zone of the site is a gravitational instability affecting 60.10⁶ m³ of material, with a maximum depth of about 150 m, located on a south-facing slope of the massif (Le Roux et al. 2011). The most active part of the landslide, referred to as "les Ruines", is located on the eastern border of the unstable zone. Long-term monitoring (extensometers, geodetic measurements,
70 tacheometers and microwave radar) shows that the displacement velocity is around 300 cm y⁻¹, while the less active parts of the site are moving at a mean of 10 cm y⁻¹ (Le Roux et al. 2011; Dubois et al. 2014).

Geological and structural information are provided by the geological map and by two boreholes drilled in 2010 in the unstable area at depths down to 150 m. Baudement et al. (2013) has integrated these information in a GOCAD® 3D model, recently used by Lajaunie et al. (2019) to propose a new vision of the S echilienne slope based on a 3D resistivity model. The
75 basement of the massif is mainly composed of micaschists showing a N-S trending sub-vertical foliation. Stratigraphically discordant deposits dating from the Carboniferous to the Liassic periods cover the micaschists on the top north – northeast of the massif (Mont Sec) and along the Sabot Fault (Fig. 1). The slope is locally covered by Quaternary (W urm) glacio-fluvial deposits made of material reworked from the surrounding formations (Vengeon 1998, Vallet 2014). The micaschists consist primarily of quartz, biotite, phengite, and chlorite with occurrence of carbonate veins and pyrite in fractures. Carboniferous
80 deposits are made of black shales, sandstones, and conglomerates with quartz and serpentine pebbles. Triassic rocks

correspond to sandstone, quartzite, dolomite and locally of black shales, argilites, and gypsum. Liassic deposits are limestones with intercalation of layers rich in breccia consisting of micaschist, dolomite, and coal (Barf  ty et al. 1972, Vengeon 1998, Vallet 2014). Strong local heterogeneities exist in terms of lithology and fracturation and are induced by the gravitational deformation (Lajaunie et al. 2019).

- 85 The part of the slope affected by the landslide extends from 400 m to 1,100 m above sea level (a.s.l.; Le Roux et al. 2011; Fig. 1d). Above the elevation of 1,100 m a.s.l., the morphology of the Mont Sec corresponds to a plateau of glacial origin underlain by moraine deposits concentrated in small topographic depressions. The landslide is delimited at its northern border by a major head scarp of about 10 meters high and several hundreds of meters wide, which separates the glacial plateau of Mont Sec from the unstable zone. Eastward, N-S faults scarps limit the landslide whereas the western and southern parts are not well defined by geomorphological evidence. The motion of the landslide consists in a deeply-rooted, toppling movement with N50-70 slabs toward the valley, coupled with the sagging of the upper zone of the slope near the Mont Sec (Vengeon 1998). The S  chilienne instability is assumed to originate from the decompression of the basement rocks after the Romanche glacier retreated at the last glaciation (15 kyr ago). Decompression caused the opening of fractures and then the collapse of the summit of the Mont Sec (Montjuvent and Winistorfer 1980, Vengeon et al. 1999, Potherat and Alfonsi, 2001.). The S  chilienne slope is thus affected by a dense network of near-vertical, open fractures trending N70 and N110/120, controlling the deformation of the S  chilienne landslide which is characterized by a deep progressive deformation (about 100-150 m) and the absence of a well-defined basal sliding surface. Two N20 major fractures are also crossing the S  chilienne massif and the Sabot and S  chilienne faults. Open fractures are locally filled with detrital material resulting from the erosion of the massif (Vallet et al. 2015a).
- 90
- 95
- 100 Borehole logs available within the instability (Lajaunie et al. 2019) show that the rock formations below the slope are relatively unstructured, and that pyrite is heterogeneously distributed therein. Rock samples along the boreholes seem to have been subjected to oxidizing conditions, albeit with no clear sulfide reaction front at the scale of the instability. In addition, petrological observations on thin sections from these boreholes, combined with mineralogical analyses obtained from X-ray diffraction (XRD; supplementary materials, section 1) show that pyrite is disseminated within the rocks, with no particular association with calcite. Gypsum was not detected from XRD analyses in the sampled rocks, consistent with results from inverse modeling by Vallet et al. (2015) suggesting that sulfate in waters from the unstable zone (UZ) essentially originates from pyrite oxidative weathering.
- 105



110

Figure 1: Map of the Séchilienne site a. Location of the Séchilienne massif in the French Alps, b. Simplified geological map of the Séchilienne massif and sampling locations, c. schematic hydrogeological model of the Séchilienne massif, d. hydrogeological cross-section of the instability, modified after Vallet et al. (2015).

115 2.2 Hydrogeological setting

The high degree of fracturation and heterogeneity of the Séchilienne massif leads to distinct and complex patterns of hydrological flow paths. At Séchilienne, water pathways are characterized by different transit times related to a dual permeability behavior that is typical of fractured rock aquifers where conductive fractures play a major role in the drainage (Fig 1.c): rapid transit of infiltration waters through fractures reflects the functioning of a so-called "reactive" hydrological component, whereas slower transit of water through the micro-fissured, less permeable rock matrix, resulting in a smeared response of flow rate to rainfall, typifies an "inertial" circulation (Maréchal 1998, Cappa et al. 2004, Vallet et al. 2015a). In particular, the Sabot and Séchilienne faults play an important role on fluid flow through the massif by draining waters from the sedimentary cover at a fast rate (0.7 km day^{-1}), and bypassing the less pervious and more inertial micro-fissured matrix characterized by lower flow velocity (0.08 km day^{-1}) (Mudry et Etievant 2007; Vallet et al. 2015a). Local perched aquifers develop during high-flow periods and discharge downwards to the main aquifer, due to the contrast of permeability between the decompressed zone at the surface and the unaltered rock (Lajaunie et al., 2019). An underground tunnel for the production of electricity in a local hydropower plant, named "Galerie EDF", built by Electricité de France (EDF) and located at the base of the slope, acts as a major westward drain for groundwater (Vallet et al. 2015a).

Difference in hydraulic conductivity between the highly fractured unstable zone (thickness about 150-200 m, Le Roux et al. 2011) and the basement situated under the landslide (Fig 1.d) led to the build-up of a two-layer aquifer system. Those two

130

layers are connected to one another through major fractures (Vengeon 1998, Meric et al. 2005; Le Roux et al. 2011, Guglielmi et al. 2002, Vallet et al. 2015a). A temporary and discontinuous shallow perched aquifer is present in the landslide with extension and connectivity varying according to short-term recharge variations. This aquifer is almost dry during the low flow periods, with numerous disconnected saturated pockets (such as open fractures filled by colluvial deposit and altered material) linked to the heterogeneity of the landslide (Guglielmi 2002, Cappa et al. 2004, Vallet et al. 2015a). The recharge of this aquifer is mainly local (through trenches and counterscarps, limiting the runoff) with a contribution of remote groundwater through near-surface drainage at high-flow periods from the sedimentary cover above the landslide (near the Mont Sec summit) (Guglielmi 2002, Vallet et al. 2015a). The deep aquifer, which extends all over the massif (altitude around 550 m asl), corresponds to a saturated layer hosted by the fractured metamorphic bedrock and to an overlying, 100-m thick vadose layer (Vallet et al. 2015a). The deep aquifer level is controlled by the constant water heads of the Romanche alluvium in the valley, and of the Galerie EDF (425 m a.s.l.).

3. Samples and analytical methods

Nine outflows draining the whole massif were investigated for physico-chemical parameters and dissolved load chemistry (Fig 1.a). Two of these outflows (G1 and G2) are located within the Unstable Zone (UZ) and correspond to seep water collected in a tunnel excavated to monitor the landslide at 710 m a.s.l. (G710). The remaining outflows (S10, S12, S13, S15, S18, S20, S21) correspond to springs draining the Stable Zone (Fig 1.a). These outflows can be differentiated regarding the dominant local lithology: S10 is located in the Bedrock Stable Zone (BSZ), S12 in the Carbonate Stable Zone (CSZ), and S13, S15, S18, S20, and S21 are located in an area of the Stable Zone characterized by mixed lithology (Mixed Stable Zone, MSZ). Samples were collected every three months over the period 2010-2019. Between 2014 and 2017, waters were sampled once a year for Sr isotopes, while samples for S isotope were collected in 2019. In total, 360 water samples were collected and analyzed for this study. Four local rocks samples were taken, reflecting the main lithological types encountered at Séchilienne: basement micaschist, carbonate (both calcite-rich and dolomite-rich) from the sedimentary cover, and a recrystallized vein in micaschist.

Field measurements of water temperature, pH, and electrical conductivity (EC) were made with a WTW pH/Cond 340i (Xylem Inc.) sensor, with a precision of 0.1 unit and $0.1 \mu\text{S cm}^{-1}$ for pH and EC, respectively. Water samples were collected in polyethylene bottles and filtered with a $0.45 \mu\text{m}$ pore diameter nylon filter, before being preserved in cold conditions for measurements of major element concentration and Sr and S isotopes. Analyses of dissolved major elements were all carried out at the research laboratory Chrono-Environnement at the University of Franche-Comté. Dissolved major cation concentrations were measured by atomic absorption spectrometry (AA 100 Perkin-Elmer) with detection limits of 0.5, 0.1, 0.01, and 0.1 mg L^{-1} for Ca^{2+} , Mg^{2+} , Na^{+} and K^{+} , respectively. Dissolved anion concentrations were determined using high-pressure ion chromatography (Dionex DX 100) with detection limits of 0.1, 0.1, and 0.05 mg L^{-1} for Cl^{-} , SO_4^{2-} and NO_3^{-} , respectively. The concentration of HCO_3^{-} was measured by acid titration ($\text{N}/50 \text{ H}_2\text{SO}_4$) within 48 hours after sampling, with 1% accuracy. Dissolved silica concentration was analyzed with a spectrophotometer (Spectroquant, Pharo 300, Merck) using a silica-test kit (Merck) with 3% accuracy. Only analyses with a charge balance better than 10% were taken into account.

Strontium isotope analyses were carried out at the High-Resolution Analytical Platform (PARI) of the Institut de Physique du Globe de Paris (IPGP). For water samples, dissolved Sr was first isolated from the water sample matrix by automated ion chromatography following the method of Meynadier et al. (2006). For rocks, in addition to bulk samples analysis after digestion in concentrated HF and HNO_3 , a three-step sequential leaching procedure was conducted using H_2O , 1M acetic acid, and 1M HCl. The first step was designed to recover the exchangeable fraction adsorbed onto the solid surface; step 2 was for extracting Sr from carbonates, amorphous hydroxides, and phosphate minerals (Tessier et al., 1979); step 3 was to dissolve any high-order Fe-Mn oxide/oxyhydroxide phases that might be present after HCl leaching (Tessier et al., 1979).

The leachate solutions and residual samples were measured for major and trace elements by Quadrupole ICP-MS (Agilent 7900) with a precision better than 5% and processed for $^{87}\text{Sr}/^{86}\text{Sr}$ ratio analysis following the same procedure used for bulk samples. To that effect, 3M HNO_3 aliquots of digestion solutions were loaded on columns loaded with 0.2 mL of Sr-SPEC resin (Eichrom). Then, 3M HNO_3 was used to elute the sample matrix before Sr was eluted in H_2O . Strontium isotope ratios were then measured using a Multi-Collector Inductively Coupled Plasma Mass Spectrometer (MC-ICP-MS; Thermo-Fisher Neptune) in low resolution mode (Hajj et al., 2017). Purified Sr solutions were introduced using an APEX desolvation unit and a PFA nebulizer at a rate of 50 to 100 $\mu\text{L min}^{-1}$ depending on the measurement session, and at Sr concentrations between 50 and 150 ppb. The accuracy and reproducibility of the $^{87}\text{Sr}/^{86}\text{Sr}$ analysis was assessed using repeated measurements of the international isotope Sr carbonate standard (SRM987, NIST). Over three sessions of measurements, the average value for SRM987 standard NIST was 0.710249 ± 0.000025 , in agreement with the accepted value.

Sulfur isotopes measurements were performed at the Biogéosciences Laboratory, University of Bourgogne, Dijon, France on both sulfates from water samples and sulfides from basement micaschist. Nine samples were treated with an excess of 250 g l^{-1} BaCl_2 solution to precipitate BaSO_4 . After centrifugation, the BaSO_4 precipitate was washed several times with deionized distilled water and dried at 60°C for 24 hours in an oven. Five hundred micrograms of purified barite samples were poured into tin capsules and homogeneously mixed with 1/3 of vanadium pentoxide before isotopic measurements (^{34}S , ^{32}S) using a Vario PYRO cube (Elementar GmbH) connected online *via* an open-split device to an IsoPrime IRMS system (Isoprime, Manchester, UK). Sulfur isotope data are expressed in delta notation and reported in units per mille (‰). The $\delta^{34}\text{S}$ data are reported with respect to the international standard Vienna Cañon Diablo Troilite (VCDT). Analytical errors are $\pm 0.3\text{‰}$ (1σ) based on replicate analyses of the international barite standard NBS-127, which was used for data correction assuming a $\delta^{34}\text{S}$ value of +20.3 on the VCDT scale.

Sulfur contained in sulfides was extracted from eight rock samples of the basement micaschist formation on aliquots of 3 grams (4 sub-samples on each of two rock samples, including unaltered pyrite and iron oxides) following the method described by Canfield et al. (1986). Dried and rinsed Ag_2S precipitates recovered after wet chemistry sulfides extraction are weighted for gravimetric quantification of sample sulfur content. Five hundred micrograms of silver sulfides precipitates were then mixed with an equivalent weight of tungsten trioxide in tin capsules before combustion in a Vario pyro cube (Elementar GmbHTM). Sulfur isotope compositions ($\delta^{34}\text{S}$) were measured using an IsoPrime IRMS device (Isoprime, Manchester, UK). International standards (IAEA-S-1, IAEA-S-2, IAEA-S-3) were used for calibration and results are reported in the δ -notation relative to the Vienna Canyon Diablo Troilite (VCDT) standard. Reproducibility (1σ) is better than 0.2‰ based on duplicate analyses of standard materials and samples.

4. Results

The concentration of major and trace elements, as well as Sr and S isotope composition of rock samples are given in Table 1 and 2 (data available at [10.5281/zenodo.4606732](https://zenodo.org/record/4606732)). The concentration of major elements, EC, pH, temperature, and Sr and S isotope composition of water samples are given in Table 3.

4.1 Major elements

Rock samples

The two limestone samples show distinctive response to the leaching procedure, with most of the Ca of the "Laffrey" limestone located in the acetic acid leachate (67%) and the rest in the HCl fraction (11%), which is indicative of the calcitic nature of this rock sample (Table 1). By contrast, most of the Ca of the "Lias" sample is hosted in the residue, while a significant fraction (37%) is HCl-soluble, suggesting that the carbonate fraction of this sample is dolomitic. The "micaschist"

and "vein" samples have higher bulk Sr/Ca, Mg/Ca, Al/Ca and Na/Ca ratios than the two limestone samples, confirming that they are mostly made of silicate minerals.

215 *Spring water samples*

The chemical composition of waters sampled on the Séchilienne site is very diverse, reflecting heterogeneity in rock types and the existence of various groundwater flow paths. Water pH at Séchilienne is relatively high and varies from 6.5 to 9.4 with a mean value of 7.9. Electrical conductivity range between $79 \mu\text{S cm}^{-1}$ and $1114 \mu\text{S cm}^{-1}$ (Table 3). From the major ion perspective, samples can be grouped into 4 main water types (Fig. 2). These water types correspond to those identified previously by Vallet et al. (2015a). Type 1 corresponds to Ca-HCO₃ waters, typical of water draining carbonate formations, typified by the S12 spring draining the carbonate cover at the top of the Séchilienne slope (CSZ). S12 has low EC values ranging from $79 \mu\text{S cm}^{-1}$ to $147 \mu\text{S cm}^{-1}$ with a mean of $117 \mu\text{S cm}^{-1}$. The second group corresponds to Mg-Ca-HCO₃ rich waters, which have circulated through the sedimentary cover (carbonate and dolomite) and the micaschists bedrock and is represented by the S10 spring (BSZ). All S10 samples have higher electrical conductivities ranging from to $308 \mu\text{S cm}^{-1}$ to $509 \mu\text{S cm}^{-1}$ with a mean of $443 \mu\text{S cm}^{-1}$. Waters sampled in the unstable part of the slope (UZ), include the underground outflows G1 and G2 and show a chemical composition that vary from Mg-Ca-HCO₃-SO₄ waters to Mg-Ca-SO₄ waters and constitute the third hydrogeochemical group. The highest EC values of this study are observed for the G1 outflow with electrical resistivities ranging from $613 \mu\text{S cm}^{-1}$ to $1114 \mu\text{S cm}^{-1}$ with a mean value of $824 \mu\text{S cm}^{-1}$. The other outflow of the unstable zone (outflow G2), in contrast to the previous one, shows a mean electrical conductivity value around $391 \mu\text{S cm}^{-1}$, with a minimum value of $313 \mu\text{S cm}^{-1}$ (and a maximum value of $470 \mu\text{S cm}^{-1}$). The fourth and last type of waters include the S13, S15, S18, S20, S21 outflows, sampled in the stable part of the slope (MSZ) along the Sabot fault, and show Ca-Mg-HCO₃-SO₄ type waters. The MSZ group exhibits EC values ranging from $357 \mu\text{S cm}^{-1}$ to $567 \mu\text{S cm}^{-1}$, with a mean of $479 \mu\text{S cm}^{-1}$. Waters of the Unstable Zone group (G1, G2) are characterized by the highest concentrations in SO₄²⁻ (from 1.32 to 3.90 mmol L⁻¹) compared to the other outflows sampled which have values ranging from 0.57 to 1.34 mmol L⁻¹ for the MSZ outflows (S13, S15, S18, S20, S21) and from 0.48 to 0.67 for the S10 (BSZ) outflow (Fig. 2; Tab. 3). Fig. 2 clearly shows that SO₄²⁻ ions are significantly contributing to the electrical balance of the analyzed waters. Dissolved Cl⁻ concentrations are lower than $50 \mu\text{mol L}^{-1}$ in springs S10, S12, S13, S18, but can reach values above $100 \mu\text{mol L}^{-1}$ in springs S15, S20, and S21. Dissolved NO₃⁻ concentrations are typically below $20 \mu\text{mol L}^{-1}$ in springs G1, G2, and S10, but are higher in springs S12, S13, S18, S15, S20, and S21, with concentrations above $100 \mu\text{mol L}^{-1}$ observed in the latter (Tab. 3).

240 Rainwater samples (Tab. 3) show very low EC values with a mean of $26 \mu\text{S cm}^{-1}$ and low concentrations for all elements analyzed. In particular, chloride concentrations range from $3.3 \mu\text{mol L}^{-1}$ to $20.3 \mu\text{mol L}^{-1}$.




Figure 2: Piper diagram representing the major water types samples in the outflows of the Séchilienne massif

245

4.2 Strontium isotopes

Rock samples

The two carbonate rock samples have the lowest Sr isotopic ratios (Table 1), with the lowest value being 0.7095 slightly higher than the Sr isotopic composition of lower Jurassic seawater (from 0.7065 to 0.7076; Koepnick et al., 1990). The acetic-acid and HCl soluble fractions of the limestone samples, as well as the bulk analysis of the dolomitic "Lias" sample and its H₂O leachate, are characterized ⁸⁷Sr/⁸⁶Sr ratios of ~ 0.7105. The ⁸⁷Sr/⁸⁶Sr ratio of all the leachates of the calcitic "Laffrey" sample, as well as of the bulk sample, show a wider range of variation, in the range 0.7104-0.7179. By contrast, the highest ⁸⁷Sr/⁸⁶Sr ratios were found in the micaschists samples with a value of 0.7351 (Table 1), typical of silicate rocks (0.73±0.01; Négrel et al., 1993). The veins contained in the micaschists also show high Sr isotopic ratios (0.7277). Such high ⁸⁷Sr/⁸⁶Sr ratios are particularly encountered in the residues and bulk samples.

Spring water samples

The Sr isotope ratios measured in the spring samples in and around the Séchilienne instability range from 0.7093 to 0.7231 (Table 3). The four main groups of waters have contrasted ⁸⁷Sr/⁸⁶Sr isotopic ratios. The highest ⁸⁷Sr/⁸⁶Sr values are found in the UZ underground outflows samples (G1, G2) with an average of 0.7210 ± 0.0006 (one standard deviation). The lowest ⁸⁷Sr/⁸⁶Sr values correspond to the waters of the MSZ (springs S13, S15, S18, S20, and S21), and average at 0.7095 ±

0.00012, *i.e.* at the value measured in the carbonate rock. The S12 outflow (CSZ) is characterized by $^{87}\text{Sr}/^{86}\text{Sr}$ ratios of around 0.7095 ± 0.00005 , *i.e.* close to those of the MSZ group. Intermediate values of Sr isotopic ratios are found for the samples of the BSZ group (S10), with an average of 0.7148 ± 0.00019 .

265 4.3 Sulfur isotopes

Rock samples

Sulfur isotope composition of unaltered rock samples (4 samples) range from -7.9‰ to 17.8‰ with an average value of $1.2‰ \pm 11.8‰$, whereas weathered micaschists (4 samples) exhibit $\delta^{34}\text{S}$ values ranging from -13.1 to 9.9‰ and an average of $-1.4‰ \pm 9.5‰$; (Table 2). These numbers show the extremely large range of possible sulfur isotope signals co-existing in
270 the various rock types present in the landslide.

Spring water samples

Waters show a much narrower range of $\delta^{34}\text{S}$ values, ranging from -5.5‰ to 6.5‰ (mean $0.43‰ \pm 5.12‰$). The highest $\delta^{34}\text{S}$ values are observed for water of the MSZ group (outflows S13, S15, S20, S21) with an average of $6.28‰ \pm 0.34‰$. Samples
275 of the S12 outflow (CZS group) also exhibits a high $\delta^{34}\text{S}$ value of 6.03‰. The G1 and G2 outflows (UZ group) exhibit negative $\delta^{34}\text{S}$ values with an average of $-3.74‰ \pm 1.75‰$, with lower $\delta^{34}\text{S}$ values for G2, (averaging at $-5.33‰ \pm 0.22‰$) than for G1 ($2.2‰ \pm 0.06‰$). The BSZ group (outflow S10) is characterized by $\delta^{34}\text{S}$ values that are intermediate between those of the MSZ and UZ groups (2.4‰).

5. Discussion

280 5.1 Identification of sources to dissolved species

5.1.1 Atmospheric and anthropogenic sources

Rainwater is potentially a significant source of elements to the water sampled in the different springs at Séchilienne. To assess the importance of atmospheric inputs, we use Cl^- concentrations. Chloride is not significantly involved in chemical reactions at the Earth surface and its presence in waters has three main origins: rainwater (through the dissolution of seasalt
285 aerosols), dissolution of saline rocks or inclusions, and anthropogenic inputs. On the other hand, at Séchilienne, NO_3^- is most likely to be derived from human activity, through fertilizer input and/or domestic waste. Therefore, the correlation between Cl^- and NO_3^- concentrations in springs S15, S20, and S21 ($R^2 = 0.65$; Fig. S8; supplementary materials) suggests that beyond rainwater, anthropogenic inputs are a significant Cl^- source to these springs. This inference is consistent with the presence of villages upslope from these springs. High Cl^- concentrations ($> 100 \mu\text{mol L}^{-1}$) are also found in some samples from
290 underground outflows G1 and G2 (Table 3). However, in the case of these two springs such high Cl^- concentrations are not accompanied with high NO_3^- concentrations but correlate to some extent with dissolved Na^+ and K^+ concentrations (Fig. S8). This observation could be indicative of the dissolution of salts (NaCl and KCl) as a significant process delivering Cl^- and cations to the springs. Although the origin of these salts is unclear, we note that Zn-Pb ore deposits are reported discovered
295 in the bedrock of Séchilienne landslide exploitation (Barnes, 1997). The presence of fluid inclusions containing alkali elements and Cl in these ores is likely, and leaching of such fluid inclusions could have occurred because of the exposure of new mineral surface during ore. To summarize, the excess of Cl in Séchilienne springs is most probably due to a combination of human activity (road salts and agriculture) and to dissolution of fluid inclusions of hydrothermal origin.

The expected concentration of Cl^- derived from precipitation in spring waters (hereafter called $[\text{Cl}]_{\text{crit}}$ for "critical chloride", Stallard et al., 1983) of the Séchilienne massif can be estimated by multiplying the mean Cl^- concentration found in
300 rainwater by the mean evapotranspiration factor (P/ETP: 4.02 with P: precipitation, ETP: evapotranspiration calculated from

temperature and latitude of the study site, according to Oudin et al. 2005). Alternatively, $[Cl]_{crit}$ can be estimated as being equal to the lowest Cl^- concentrations in the sample set (S10, S12 and S21 springs). Both methods concur to fix the atmospheric contribution of Cl^- to the S echilienne waters at a maximum of $30 \mu mol l^{-1}$. Above this concentration, additional sources must be involved. Once $[Cl]_{crit}$ is known, it is possible to correct all cation concentrations from the atmospheric seasalt input by using:

$$[X]^* = [X] - [Cl]_{crit} \times \left(\frac{X}{Cl}\right)_{seawater} \quad (1)$$

In this equation, $[X]^*$ denotes the concentration of an element X in the water sample, corrected from the atmospheric input, and $(X/Cl)_{seawater}$ is the seawater elemental ratio. This correction is only significant for Na^+ , due to the relatively high concentrations found in the S echilienne waters.

As explained above, significant excess of Cl^- ($[Cl] > [Cl]_{crit}$) is found for the S15, S20, and S21 springs where this excess is the highest (about 30 to 60 $\mu mol l^{-1}$) and due to domestic and/or agricultural activity; and in the G1 and G2 underground outflows (between 15-20 $\mu mol l^{-1}$ of excess), where this excess can be attributed to salt dissolution. For the first group of springs, it is most likely that input of Cl^- is associated with input of K^+ , through the use of fertilizers. For the second group of springs each mole of Cl^- released by salt dissolution can be associated with one mole of K^+ or one mole of Na^+ . Because of the challenge associated to assessing the exact cause for the observed Cl^- excess in these springs, in the quantitative source apportionment (section 5.1.4) we use a stochastic approach to reflect the uncertainty linked to the nature of the cations delivered to the springs by these Cl^- sources. However, we emphasize that for the springs where silicate weathering is the most prominent process in terms of cation production (G1 and G2; see section 5.1.2 below), the correction for the solute sources causing the Cl^- excess (human activity and salt dissolution) is relatively minor.




Figure 3: Na vs Cl concentrations measured in the different groups of water outflow from the S echilienne massif. Rainwater data points are aligned along the seawater composition (seasalts).

5.1.2 The importance of silicate weathering

The concentration of Na^+ , once corrected from atmospheric and anthropogenic inputs, can be used as a proxy of silicate weathering reactions, if the dissolution of silicate minerals is a dominant source of Na to surface waters compared to salt dissolution and human activity. The underground outflows G1 and G2 have the highest Na^* concentrations (mean values 200±30 $\mu\text{mol L}^{-1}$ and 158±21 $\mu\text{mol L}^{-1}$ for G1 and G2, respectively) (Fig. 3). This observation suggests the importance of silicate weathering reactions in the Unstable Zone, made of fractured micaschists. Although as explained above the source of excess chloride could also be contributing Na (were this additional source NaCl inclusions), the Na^* concentrations remain the highest found in the S chilienne landslide area assuming that all Cl^- release to waters is associated to an equivalent Na^+ release (in moles). Despite their excess of Cl^- , the MSZ outflows chemistry also reveals that silicate weathering reactions are releasing Na^+ to those waters (Fig. 3). However, the Na^* concentrations of the Mixed Stable Zone (60 $\mu\text{mol L}^{-1}$ on average) are equal to around half of those encountered in the Unstable Zone. This contrast between the stable (MSZ) and unstable zone (UZ) illustrates that the intensity of silicate weathering is linked to the degree of fracturation at S chilienne. Finally, the low Na^* concentration in the S12 outflow (Fig. 3) can be attributed to the fact that it mainly drains the carbonate cover. The most plausible explanation for the non-zero Na^* concentration in S12 is the release of Na from silicate material disseminated in the carbonate rocks, and/or to anthropogenic inputs.

5.1.3 Identifying sources to solutes in the springs of the S chilienne massif

In the following we use dissolved elemental and isotopic ratios to quantitatively constrain the contribution of various rock sources (silicates, carbonates, and gypsum) for solutes in the S chilienne springs. As shown above, strong contrasts exist in $^{87}\text{Sr}/^{86}\text{Sr}$ ratios between the sedimentary carbonate cover and the crystalline rocks of the basement. The isotopic ratio of dissolved Sr released by water-rock interaction reflects that of the minerals undergoing dissolution and is not affected by the reincorporation of Sr in secondary minerals (*e.g.* Negrel et al. 1993). Sr isotopes can thus be used to trace the provenance of dissolved Sr, and by extension of the different cations, in the waters of S chilienne. By contrast, elemental ratios such as Ca/Sr and Mg/Sr may be affected by the precipitation of secondary minerals and in particular by the formation of secondary carbonates (Bickle et al., 2015) and should be used more carefully to identify the provenance of cations.

A series of plots using $^{87}\text{Sr}/^{86}\text{Sr}$ as a common Y-axis are shown in Fig. 4. In $^{87}\text{Sr}/^{86}\text{Sr}$ vs. Ca/Sr or Na/Sr plots (Fig. 4a and 4b) conservative mixing between reservoirs is indicated by straight lines joining the end members. In this set of figures, the Ca/Sr ratios for the carbonate, silicate, and evaporite end members are taken from Negrel et al. (1993) and Gaillardet et al. (1997). The corresponding Sr isotopic ratios are those measured in the rock samples from the S chilienne massif (Table 3).

The position of the data points corresponding to springs S13, S15, S18, S20, and S21 (MSZ group) in Fig. 4a and 4b shows that their relatively low Sr isotopic composition cannot only be derived from the dissolution of carbonates. Another unradiogenic end member with low Ca/Sr and Na/Sr ratios needs to be invoked. As indicated by Fig. 4c this end member is enriched in sulfate as shown by its high SO_4/Na ratio. Although gypsum outcrops are not visible at S chilienne, gypsum is known to exist in the local Triassic formations present in the upper part of the slope as indicated by the regional geological map (Fig.1). More generally, the presence of gypsum is well documented in the Triassic strata of the "external Alps" where it plays a major role in large-scale deformation and thrusting (Barf ty et al. 1972). The occurrence of gypsum and carbonate dissolution inferred from the chemistry of the S chilienne springs indicates that the Sabot Fault, which lies at the North-East of the MSZ outflows (Barf ty et al. 1972), plays a major role in draining aquifers hosted by sedimentary rocks to the MSZ and BSZ outflows. The Sr and S isotope composition of Triassic seawater (between 0.7075 and 0.708 and 15±3 ‰, respectively; Burke et al. 1982, Fanlo and Aroya 1998, Kampschutte and Strauss 2004) and the typical Ca/Sr ratio of waters draining gypsum (Gaillardet et al., 1997; Meybeck et al., 1986) are consistent with the contribution of gypsum dissolution.

Fig. 4 also shows first that the S12 spring, reported by Vallet et al. (2015a) to be supplied by rapid flowpaths through the sedimentary cover, in addition to being solute-poor compared to springs of the MSZ group, is not influenced by gypsum

dissolution despite its geographical position on the sedimentary part of the slope. The relatively high Na/Sr ratios observed in the S12 spring is probably due to anthropogenic influence, as revealed by the high nitrate concentrations measured in this spring (section 5.1.1 and Fig. S8c).

By contrast, $^{87}\text{Sr}/^{86}\text{Sr}$ and chemical ratios of waters from the UZ outflows (G1, G2) are clearly influenced by a silicate end member. However, their Sr isotopic signature is lower than those of the local micaschist, indicating the additional contribution of Sr from a carbonate and / or evaporitic source to the G1 and G2 spring (Fig. 4a). The $^{87}\text{Sr}/^{86}\text{Sr}$ ratios of waters of the S10 outflow also exhibit intermediate values between the silicate and carbonate-gypsum mixing line, but with $^{87}\text{Sr}/^{86}\text{Sr}$ ratio lower than those of the UZ outflows, supporting the idea that water-silicate interaction in the BSZ are less intense than in the UZ. These inferences based on Sr isotopes are in full agreement with those made above based on Na^* concentrations and can be interpreted as reflecting the lesser degree of fracturation of the stable zone compared to the unstable zone. Fig. 4c and Fig. 4d show that the higher $^{87}\text{Sr}/^{86}\text{Sr}$ ratios observed in springs of the UZ (G1, G2), and to a lesser extent of the BSZ (S10) are associated with sulfate enrichment. However, unlike for samples of the MSZ (S13-S21), dissolved sulfate in UZ and BSZ samples has a relatively low S isotope composition (Fig. 4d). This observation is compatible with a significant influence of sulfide oxidation, despite the very wide range of $\delta^{34}\text{S}$ values measured in the bedrock micaschists (between -13,14‰ and 17,77‰, average -0,10‰, S.D: 10,05; Table 2).

The presence of pyrite has been reported in the unstable zone of Séchilienne (Bertrand et al. 2014, Vallet et al. 2015a). The concomitant increase in SO_4^{2-} and radiogenic Sr (Fig. 4c), combined with the decrease in $\delta^{34}\text{S}$ (Fig. 4d) suggests a coupling in the unstable zone between sulfide oxidation and silicate weathering. Indeed, the oxidative weathering of pyrite, possibly by O_2 or water, leads to the release of sulfate to waters (e.g. Spence and Telmer, 2005):



The oxidation of sulfide to intermediate sulfur species or to sulfate appears to produce only small isotope effects (Fry et al., 1986, 1988; Zerkle et al., 2009; Balci et al., 2012). The significance of these reactions in the unstable zone of Séchilienne can be related to the role of fracturation and grain comminution in favoring the contact between water, air, and minerals, which is the rate-limiting factor for a fast chemical reaction such as pyrite oxidation.

Altogether, our analyses shows that the composition of the water outflows from the Séchilienne site can be interpreted by a variable contribution of waters having interacted with micaschists and sedimentary rocks, and by a dual origin (sulfide oxidation vs. gypsum dissolution) of sulfate ions. The chemical and isotopic characteristics of the MSZ and UZ waters show that these waters have percolated through the sedimentary cover before reaching their outlet in the massif through the Sabot fault.

400




Figure 4: Sr isotopic composition of the different groups of water outflows from the S echilienne massif as a function of Sr-normalized ratios (a, b and c) and S isotopic composition (d). Mixing end members are discussed in the text. Straight lines indicate a mixing process in Fig. 4a, 4b and 4c. In Fig. 4d, the dashed line is a mixing hyperbola calculated based on the composition of the end members ($^{87}\text{Sr}/^{86}\text{Sr}$, $\delta^{34}\text{S}$, and SO_4/Sr ratios of the silicate and carbonate end members).

5.1.4 Quantitative apportionment

Spring dissolved Sr isotopes and major element chemistry make it possible to estimate the relative contribution of each identified end member to the different cations. The details of these calculations and results are given in supplementary materials. Following the above discussion, mixing equations can be written for the conservative trace element Sr:

$$\left(\frac{^{87}\text{Sr}}{^{86}\text{Sr}}\right)_{\text{mix}} = X_{\text{sil}}^{\text{Sr}} \cdot \left(\frac{^{87}\text{Sr}}{^{86}\text{Sr}}\right)_{\text{sil}} + X_{\text{carb}}^{\text{Sr}} \cdot \left(\frac{^{87}\text{Sr}}{^{86}\text{Sr}}\right)_{\text{carb}} + X_{\text{gyyps}}^{\text{Sr}} \cdot \left(\frac{^{87}\text{Sr}}{^{86}\text{Sr}}\right)_{\text{gyyps}} \quad (4)$$

$$\left(\frac{\text{Na}}{\text{Sr}}\right)_{\text{mix}} = X_{\text{sil}}^{\text{Sr}} \cdot \left(\frac{\text{Na}}{\text{Sr}}\right)_{\text{sil}} + X_{\text{carb}}^{\text{Sr}} \cdot \left(\frac{\text{Na}}{\text{Sr}}\right)_{\text{carb}} + X_{\text{gyyps}}^{\text{Sr}} \cdot \left(\frac{\text{Na}}{\text{Sr}}\right)_{\text{gyyps}} \quad (5)$$

$$1 = X_{\text{sil}}^{\text{Sr}} + X_{\text{carb}}^{\text{Sr}} + X_{\text{gyyps}}^{\text{Sr}} \quad (6)$$

Where the subscripts *mix*, *sil*, *carb*, and *gyyps* denote the mixture (water), the silicate, carbonate, and gypsum end members respectively. Proportions of Sr derived from each of those end members *i* are denoted X_i^{Sr} . All ratios are corrected from atmospheric and salt inputs according to the above method. Because the carbonate and gypsum end member add relatively few Na compared to Na* (Na from silicates), equation 5 simplifies into:

$$\left(\frac{Na}{Sr}\right)_{mix} = X_{sil}^{Sr} \cdot \left(\frac{Na}{Sr}\right)_{sil} \quad (7)$$

This assumption is supported by the positions of the different springs in Fig. 4a, which indicates that the low-⁸⁷Sr/⁸⁶Sr component of the springs - encompassing both carbonate and gypsum weathering - has a negligible Na content. The proportions of Sr in the different mixing reservoirs can then be estimated and the contribution of each of these end members to the load of the dissolved major species SO₄²⁻ and Mg²⁺ then calculated following:

$$X_i^E = X_i^{Sr} \left(\frac{E}{Sr}\right)_i / \left(\frac{E}{Sr}\right)_{spring} \quad (8)$$

with $i = sil, carb, \text{ or } gyp$, and $E = SO_4$, or Mg (corrected from rain inputs). Full discussion is given in supplementary materials (section 3) on the choice of the $(Na/Sr)_{sil}$ (eq. 7) and more generally of the $(E/Sr)_i$ ratios (eq. 8), based on regression of the spring hydrochemical data and independent constraints from our geochemical analyses of the rock samples.

In carbonate-rich contexts like that of S echilienne, dissolved Ca²⁺ concentrations can be affected by precipitation of secondary carbonates which tend to scavenge significant amounts of dissolved Ca relative to Mg and Sr (Bickle et al., 2015). For this reason, in principle eq. 8 cannot be applied for $E = Ca$ and $i = carb$. The relatively high Mg/Ca ratios (around 0.9 mol/mol in springs G1, G2, and S10; and 0.2-0.4 mol/mol in springs S10 to S21; Table 3) and Sr/Ca ratios (around 2 mmol/mol in springs G1, G2, S10, S12, and S18; and above 6 mmol/mol for springs S13, S15, S20, and S21) compared to the estimated Mg/Ca and Sr/Ca ratios of the calcite end member at S echilienne (below 0.1 mol/mol and 1 mmol/mol, respectively) determined from our geochemical analyses of rock samples (Table 1) are indeed suggestive of a significant role of secondary carbonate formation. We quantify the role of secondary carbonate formation using the method proposed by Bickle et al. (2015), which is based on the comparison in the Sr-Ca-Mg-Na compositional space between the measurements made in springs and the prediction from conservative mixing between the rock end members (supplementary materials; section 3). In this analysis we contend that secondary carbonate formation affects waters containing solutes derived from the three rock end members identified for the S echilienne springs (silicates, carbonates, and gypsum) after they mix. We estimate that along the water flowpath secondary carbonate formation scavenges around 60% of the Ca initially released to solution by the combined dissolution of silicates, carbonates, and gypsum for springs G1, G2, and S10, whereas the effect of secondary carbonate precipitation is negligible for the other springs (supplementary materials; section 3). These results highlight the potential role of lithological diversity, a characteristic of the bedrock material drained by springs G1, G2, and S10 in the UZ and BSZ (compared to other springs mostly influenced by the carbonate cover), in promoting secondary carbonate formation through mixing of compositionally different waters.

Another challenge in using eq. 8 at S echilienne is the fact that both calcite and dolomite are reported to occur as carbonate minerals at S echilienne - as confirmed by our own chemical analyses of rock samples (Tab. 1). Therefore, we take into account the presence of dolomite in our quantitative source apportionment, in particular regarding the $(Mg/Sr)_{carb}$ ratio used in eq. (8) (supplementary materials; section 3). Based on arguments linked to the extent of secondary carbonate precipitation needed to explain the spring data (see above), we estimate that the contribution of dolomite dissolution to the overall Ca released by carbonate weathering at S echilienne is about 10 to 20% (supplementary materials).

In order to quantify the uncertainty associated to our mixing model, a Monte Carlo approach was used with 10,000 simulations. Results are given in Tab. S2 (tables available in supplementary materials) and represented in Fig. 5 as a stacked bar plot. In the following text, results on mixing proportions X_i^E are reported as $D_{50-D_{16}}^{+D_{84}}$ (D_n is the n^{th} percentile of the output distribution over the 10,000 simulations; D_{50} is thus the median).

Carbonate dissolution appears as the major contributor to dissolved Ca, and to about half of dissolved Mg in all the springs sampled on the different parts of the studied zone (UZ, BSZ, MSZ), making this process a major supplier of cationic charges

to waters at Séchilienne (Tab. S2; Fig. 5). In the most active part of the landslide (G1, G2), despite the silicate-dominated lithology, carbonate contribution is significant (about 40%), indicating the waters percolating through the unstable zone acquired part of their chemical and isotopic composition from above the hillslope. Calculations of the proportions of sulfate derived from the different end members show a minor but non negligible contribution of gypsum dissolution (reaching 88% of the total sulfate in spring S15) and a very clear contribution of pyrite oxidative weathering particularly important in the fractured zone. In the G1-G2-S10 group of springs, most of the anionic charge (> 80%) is provided by the oxidative weathering of pyrite. Springs from the BSZ exhibit a lower proportion from silicate end member with a median of $0.48_{0.24}^{0.76}$ against $0.61_{0.38}^{0.84}$, $0.56_{0.31}^{0.81}$ for G1 and G2 (UZ), respectively. This contrast can be attributed to the unstable context of G1 and G2 compared to that of the stable part of the slope at the BSZ outflow.

Gypsum dissolution is a major process at some sites (with fraction of SO_4 from gypsum ranging between $0.57_{0.43}^{0.79}$ and $0.88_{0.80}^{0.97}$ in waters of the MSZ S13-S21, for example), but nowhere does this process dominates the overall solute production (Fig. 5). Therefore, and although significant uncertainty exists regarding the S isotope composition of the pyrite endmember, S isotope data lend support to our inference from the mixing model that waters are derived essentially from pyrite oxidation in the unstable zone, and from gypsum dissolution the MSZ waters (S13, S15, S18, S20, S21). The relative contribution of carbonates is also significant in MSZ waters with proportions ranging from $0.32_{0.09}^{0.58}$ for S18 to $0.33_{0.13}^{0.56}$ for S15.

Based on results of the mixing model, we can estimate a value for $\delta^{34}\text{S}$ of the pyrite endmember. Indeed, a significant linear negative relationship ($R^2 = 0.8$) exists between the $\delta^{34}\text{S}$ measured in springs across the Séchilienne massif and the modal estimates of their $X_{\text{sil}}^{\text{SO}_4}$ (Fig. S12; supplementary materials). The intercept of this relationship at $X_{\text{sil}}^{\text{SO}_4} = 1$ (equivalently at $X_{\text{gyp}}^{\text{SO}_4} = 0$) gives an estimate for $\delta^{34}\text{S}_{\text{sulfur}}$ of -3.1‰. Such estimates are consistent with the range of measurements of solid sulfur reported in this study (ranging between -13.1‰ to 17.8‰) and reflect an average value of the S isotope composition of sulfides for the Séchilienne unstable zone.




Figure 5: Concentration (in charge equivalents) calculated for major dissolved species and for each end member from the mixing model presented in the text and in the supplementary materials (Table S2). Silicates contribute for Na+K+Ca+Mg (Na+K not indicated), carbonates for Ca+Mg, gypsum dissolution for Ca+Mg (not distinguished). Note that ultimately the hydrogenocarbonate ion originates from respiration in soils. Sulfuric acid is generated by the oxidation of sulfide minerals, a process that occurs preferentially in the fractured zone. Relative contributions of different end members were obtained by solving a set of mixing equations using a Monte Carlo approach (section 5.1.3). Note that the relative contribution of each rock end member to the Ca^{2+} load here refers to that calculated for the Ca "initially" released into solution, that is before secondary carbonate precipitation (supplementary materials).

5.2 Implications for hydrogeological processes at the Séchilienne site

495 Water plays an important role in the dynamics of slope instabilities, first as a physical (hydrogeological) agent that can lead to aggravation of the instability. Secondly, water is a geochemical agent, which weathers rocks and makes them less cohesive (Rutqvist and Stephansson, 2003; Binet, 2006; Cappa et al., 2004). These two categories of processes interact with each other in time as weathering leads to modifications in subsurface permeability and porosity, and thus in water flowpaths through the massif (Gu et al. 2020). Hydrological triggering is the most usual mechanism of initiation and reactivation of

500 landslides but water flows in the subsurface have also been shown to have a major impact on the destabilization of a slope (de Montety et al. 2007; Guglielmi et al. 2002, Vallet et al. 2015b). However, landslides constitute very heterogeneous media due to their intense fracturation, which makes hydrogeological investigation complicated. The use of hydrochemistry and isotopic investigation can provide new insight to classical investigation. For example, groundwater dissolved $^{87}\text{Sr}/^{86}\text{Sr}$ ratios have proven to be useful in determining the sources of solutes in natural waters (Négreil and Deschamps, 1996, Négreil et al., 2001, Dotsika et al., 2010), investigating mineral weathering reactions (Brass, 1975; Åberg et al., 1989; Bullen et al., 1996; Clow and Drever, 1996; Bullen and Kendall, 1998), and identifying mixing processes involving groundwaters of different sources (Woods et al., 2000; Frost and Toner, 2004; Singleton et al., 2006) also inside an unstable context (Deiana et al., 2018). Values of groundwater dissolved $\delta^{34}\text{S}-\text{SO}_4^{2-}$ have also been used in aquifer studies to identify sulfate sources (Moncaster et al., 2000; Cortecci et al., 2002; Gammons et al., 2013). In particular, the Séchilienne hydrogeological model

510 proposed by Vallet et al. (2015a) use sulfates as a tracer of waters flowing through the instability with the assumption that all SO_4^{2-} measured in groundwaters is sourced from pyrite oxidation. High sulfate concentrations in MSZ waters were indeed inferred by Vallet et al. (2015a) to be derived from a mixture of 30% of waters from the UZ (drained through the micro-fissured matrix) and 70% from the sedimentary cover (drained through both micro-fissured matrix and larger fractures), thereby establishing a hydraulic connection between the UZ and the MSZ waters (Fig. 1c).

515 Results from the present study partly support the hydrogeological model established by Vallet et al. (2015a) but allow us to refine this model through the identification of the contribution of another, unexpected end member corresponding to the dissolution of gypsum with a remote origin. Based on the local geological map (Barfély et al. 1972) gypsum occurrence has been reported but outside of the study zone, upstream along the Sabot fault which lies at the North-East of the MSZ outflows. As the fault is a major flowpath (Lajaunie et al., 2019), draining aquifers hosted by the sedimentary cover to the

520 MSZ outflows, it contributes to the enrichment in SO_4^{2-} of those waters (Fig. 6). Our study therefore indicates a significant evaporitic origin for the sulfates in the MSZ waters, challenging the interpretation of Vallet et al. (2015a) of a hydrogeological connection between waters of the unstable and stable zones. Sulfate in outflows draining the UZ and BSZ is not strongly sourced from evaporites, but the part of those sulfates with evaporitic origin can be explained by a contribution of water flows through the Sabot fault towards the sedimentary cover and the basement formations (Fig. 6).

525 In addition, the systematic differences in elemental concentrations observed between the UZ and BSZ outflows (Tab. 3) can be linked to the structure of the slope and water flowpaths in the subsurface. Indeed, the S10 outflow drains a stable area (BSZ) just above the lowly-weathered, only slightly fractured basement. By contrast, the G1 and G2 outflows drain the unstable part of the slope (UZ), where the basement is highly fractured. This leads in turn to a stronger weathering degree of rocks and minerals there, and in particular of pyrite which is a major contributor to dissolved sulfate in the G1 and G2

530 outflows - more so than in the S10 outflow (stable and lowly weathered) characterized by lower sulfate contents. Improving numerical and predictive models requires the incorporation of hydrological processes such as the dynamics of water circulation within a slope (directly dependent on fracturation, volumes of water involved, etc.). This study shows that isotopic proxies such as Sr and S isotopes ($^{87}\text{Sr}/^{86}\text{Sr}$, $\delta^{34}\text{S}$) coupled to water chemistry can be a very powerful tool to constrain groundwater origin and flowpaths in landslides and can substitute to tracer surveys, and constitute an alternative

535 for hydrogeological investigations in logistically-challenging field environments such as unstable slopes.




Figure 6: Sketch of the Séchilienne groundwater conceptual model, modified after Vallet et al., (2015).

540 5.3 Role of landslides on silicate weathering and CO₂ consumption

Recent studies have shown the importance of pyrite oxidation, sulfuric acid production, and associated chemical weathering in active landslides (Emberson et al. 2015, Emberson et al. 2016). The novelty of the present study is the use of waters sampled within an active, slow landslide, rather than percolation waters sampled at the base of the colluvium of recent landslides, of river waters draining landslide-affected watersheds. In the following paragraph we examine the potential implications of the present study of the Séchilienne landslide for the global carbon cycle.

Silicate weathering by carbonic acid consumes atmospheric CO₂ and, combined with the precipitation of carbonates in the ocean, is the mechanism that has allowed for the sequestration of atmospheric CO₂ and consequently lower the Earth's surface temperature on geological timescales (Bernier and Bernier 2012). Rock forming-minerals uplifted to the Earth surface react with oxygen, carbonic acid produced by soil respiration, and sulfuric acid produced by the oxidation of sulfide minerals. The following reactions describe how carbonic (equations 9 and 11) and sulfuric (equations 10 and 12) acids interact with silicate (here wollastonite CaSiO₃) and carbonate minerals, and lead to the production of alkalinity (here HCO₃⁻):



It is usually considered that when Ca^{2+} reaches the ocean, over a time period longer than 0.1 to 1 Myr, the precipitation of CaCO_3 releases CO_2 into the ocean-atmosphere system according to the reaction:




The influence of the above reactions (equations 9-13) on atmospheric CO_2 partial pressure depends on the time scale considered (Torres et al., 2016). At short timescales (typically $< 10^5$ yrs), the chemistry of river discharge is able to influence the carbonate system in the ocean. Indeed, the delivery of alkalinity and Dissolved Inorganic Carbon (DIC) to the ocean to a ratio lower than that of the modern seawater ratio ($\text{Alk}/\text{DIC} \sim 1$) leads to increased dissolved CO_2 concentration, and consequently higher CO_2 content in the atmosphere through re-equilibration (Zeebe and Wolf-Gladrow, 2001). If the Alk/DIC ratio is higher than 1 but lower than 2, at time scales longer than that typical of carbonate precipitation in the ocean (10^5 to 10^6 yrs) but shorter than that of marine sulfate reduction to sulfide in sea bottom sediments (several 10^6 yrs), atmospheric CO_2 will increase because the precipitation of carbonates releases CO_2 to the ocean-atmosphere system that was not consumed on land by weathering reactions (combination of equations. 12 and 13). This mechanism should lead to global warming (Calmels et al., 2007) and has been invoked by Torres et al. (2014) for maintaining atmospheric CO_2 levels during the Himalayan orogeny, which otherwise should have led to a rapid atmospheric CO_2 depletion by enhanced silicate weathering. Finally, at timescales longer than that typical of sedimentary burial of sulfide in the ocean, only silicate weathering by carbonic acid leads to net C sequestration (Bernier et Bernier, 1996; Calmels et al. 2007).

However, our analysis demonstrates that at S echilienne reaction (13) also occurs directly on the continent through the formation of secondary carbonates, favored by the addition of calcium and alkalinity derived from silicate weathering, which results in a "short cut" of the carbon cycle. This short-term CO_2 release has to be taken into account when evaluating the overall CO_2 effect of weathering reactions at S echilienne, and more generally in lithologically complex weathering systems where secondary carbonate formation is likely to involve solutes produced by a variety of processes, and in particular by carbonate weathering by sulfuric acid.

To this aim, we use the stoichiometry of reactions (9) to (13), together with the results of our quantitative source apportionment (section 5.1.4) to calculate the impact of weathering reactions at S echilienne on atmospheric CO_2 over two scales (convoluted spatially and temporally), referred to in the following as "on site" or "local" (*i.e.*, immediately when weathering processes take place in the unstable zone) and "long term" or "global" (*i.e.*, taking into account marine carbonate precipitation in the ocean ensuing solute delivery to the ocean).

Fig. 7 shows that waters produced in the Unstable Zone of S echilienne (G1 and G2) act as CO_2 sources on the "long term", whereas waters produced in the bedrock stable zone (S10) or the mixed stable zone (S15, S18) are CO_2 sinks or CO_2 -neutral within uncertainty. Our study thus shows that instabilities such as the S echilienne landslide can act as hotspots of long-term CO_2 release to the atmosphere depending on the types of mineral-fluid interactions and also on the flow paths followed by the water drained in the landslide. We suggest that chemical weathering in similar landslides throughout the Alps (*i.e.* Clapi ere, Super Sauze, and Valabres in the French Alps, Rosone in Italy; Barla et Chiriotti, 1995; Follacci, 1999; Binet, 2006) have a similar impact on global biogeochemical cycles and climate. Although it is beyond the scope of the present study to quantify the CO_2 fluxes linked to weathering in the S echilienne landslide - let alone to attempt an extrapolation of such local results to the scale of the Alpine range - our work clearly shows that silicate and carbonate weathering by sulfuric acid generated in landslide zones of active mountain ranges have a climatic impact though a complex set of entangled short-term and long-term effects. Furthermore, this impact contradicts the textbook view that silicate weathering in mountain ranges consumes CO_2 from the atmosphere and cool the global climate (Raymo, 1991; Hilton and West 2020), and motivates more detailed studies associating hydrogeological and mineralogical approaches to build a more realistic understanding of the impact of mountains on climate change.



605 Figure 7: Evaluation of the effect of weathering processes at Séchilienne on atmospheric CO₂. The hydrochemistry of the springs G1 and
 610 G2 draining the instability ("Unstable Zone") show that the weathering results in long-term CO₂ production to the atmosphere because the
 cations are preferentially released in spring waters by the action of sulphuric acid and not carbonic acid from the soils (Table S3;
 supplementary materials). In the Bedrock Stable Zone (spring S10) and in the Mixed Stable Zone (springs S15 and S18), weathering
 processes act as CO₂ sinks or are CO₂-neutral within uncertainty. Secondary carbonate precipitation returns CO₂ to the atmosphere. In
 particular, in the spring G2 of the Unstable Zone, this process results in a net "on site" CO₂ release because it involves precipitation of Ca
 and alkalinity derived from carbonate weathering by sulfuric acid.

6. Conclusion

We use measurements of dissolved major element chemistry coupled to Sr and S isotopic ratios in spring waters of the
 Séchilienne active landslide site in order to identify the chemical processes at play in the subsurface of the landslide area.
 615 Among these tracers, strontium isotopes allow us to allocate cations to different sources, circumventing issues affecting
 elemental ratios related to the precipitation of secondary. Silicate, carbonate, and evaporite weathering all appear to
 contribute to the cation load of the Séchilienne waters. Scavenging of dissolved calcium by secondary carbonate formation is
 identified as a major process affecting solutes in the subsurface waters of the Séchilienne instability and favored by the
 mixing of different solution having interacted with a heterogeneous set of minerals. Sulfur isotopes bring a unique
 620 qualitative constraint on the origin of the sulfate ion, which is abundant in the Séchilienne groundwaters, showing the
 contribution of not only pyrite oxidation but also of gypsum dissolution.

The provenance of dissolved species at Séchilienne also reveals the complex water flow paths there. In particular, waters
 percolating through the landslide have acquired part of their hydrochemical characteristics far away from the unstable zone
 itself. For example, sulfur isotopes clearly indicate an unexpected contribution from Triassic sedimentary gypsum

625 dissolution, that can only occur in the sedimentary layers capping the upper part of the massif and pointing out the importance of water drainage by a major fault of the massif.

The comparison between the stable and unstable parts of the site suggests that silicate weathering is enhanced in the fractured, unstable zone, where the landslide is active. Sulfur isotopes indicate that the production of acidity by the oxidation of magmatic sulfides enhances rock alteration in the unstable zone, this leads us to suggest the following feedback. By favoring the penetration of oxic waters and allowing contact with silicate minerals, fracturation and grain comminution controls the oxidation of pyrite that in turn rapidly generates sulfuric acid. The weathering of silicate minerals by sulfuric acid weakens the rock structure what in turn favors fracturation in response to the gravitational stress. For example, Fletcher et al. (2006) and Behrens et al. (2015) have shown that opening of porosity at the rock-soil interface in soil profiles can be initiated by oxidation of Fe(II) minerals inducing a positive volume budget leading to the production of micro-cracks, inducing further weathering, at the origin of the opening of fractures, provided that enough carbonate and pyrite is present in the bedrock. At a larger scale, the feedback we propose here exemplifies a similar process of coupling between physical and chemical processes sustaining mass wasting in mountain ranges.

635
640
645 Finally, we demonstrate that the Séchilienne landslide is a hotspot of CO₂ release to the atmosphere over the long term. Although it remains difficult to upscale the results of the present study to the entire Alpine range, or to a global scale, landslides developed on sulfide-hosting sedimentary rocks appear to have a climatic impact opposite to the conventional view that rock weathering in mountains ranges consumes CO₂ from the atmosphere, and thus contributes to global cooling. In addition, our study shows a strong control of weathering processes and rates by local hydrogeological features, such as the complexity of flow paths setting the chemistry of the groundwaters within the unstable zone. More work is needed to assess the importance of landslides as hotspots of chemical alteration and geological CO₂ emissions, in particular to investigate their hydrological and hydrochemical response to weather and climate change. More generally, landslides epitomize the coupling between landscape evolution, tectonics and climate-weather. For this reason - as well as that of their societal impact in terms of natural hazard - monitoring landslides over a range of time scales and frequency should become a priority.

Team list

650 Pierre Nevers
Julien Bouchez
Jérôme Gaillardet
Thomazo Christophe
Charpentier Delphine
655 Laëticia Faure
Catherine Bertrand

Author contribution

Pierre Nevers measured the Sr isotopic ratios, worked on the interpretation and wrote text. Julien Bouchez and Jérôme Gaillardet helped with the isotopic measurements, in the inversion calculations and in writing text. Christophe Thomazo
660 measured the sulfur isotopic ratios. Delphine Charpentier worked on XRD and mineralogic analyses presented in supplementary materials. Laëticia Faure helped with the rock sample leaching procedure and Sr isotope measurements. Catherine Bertrand organized the sampling strategy and is in charge of the Séchilienne Observatory (SNO OMIV) and helped in writing text.

Competing interests

665 The authors declare that they have no conflict of interest.

Acknowledgments

Part of this work was supported by IPGP multidisciplinary program PARI and by Region île-de-France SESAME Grant no. 12015908. Caroline Gorge, Christophe Loup and Caroline Amiot are acknowledged for analytical support, and Vanessa
670 Stefani is acknowledged for support on the field. The authors warmly thank: the PEA²t platform (Chrono environment, University of Bourgogne Franche-Comté, UMR CNRS 6249, France), which manages and maintains the analytical equipment used in this study; and OMIV and OSU Theta for supporting the project through financial contributions (SRO_2015).

References

675 Barféty, J., Bordet, P., Carme, F., Debelmas, J., Meloux, M., Montjuvent, G., Sarrot Reynauld, J.: Carte géologique détaillée de la France (1/50000) n°797 Vizille. Editions du BRGM, p. 38, 1972

Barla, G. et Chiriotti, E., 1995. Insights into the behaviours of the large deep seat
gravitational slope deformation of Rosone, in the Piemont region (Italy). Felsbau, 13:
680 425-432

Baudement, C., Bertrand, C., Guglielmi, Y., Viseur, S., Vallet, A., & Cappa, F.: Quantification de la dégradation mécanique et chimique d'un versant instable : approche géologique, hydromécanique et hydrochimique Etude du versant instable de Séchilienne, Isère (38). 1–6, 2013

Behrens R., Bouchez J., Schuessler J. A., Dultz S., Hewawasam T. and von Blanckenburg F. (2015) Mineralogical transformations set slow weathering rates in low-porosity metamorphic bedrock on mountain slopes in a tropical climate. *Chem. Geol.* 411, 283–298, <https://doi.org/10.1016/j.chemgeo.2015.07.008>, 2015

690 Berner, K.E. and Berner, R.A.: *Global Environment: Water, Air and Geochemical Cycles*, Princeton Univ. Press, 369–382, 2012

Bertrand, C., Vallet, A., Mudry, J.: Hydrochemical Approach of Mechanical Degradation of the S echilienne Unstable Slope, in *Engineering Geology for Society and Territory – Volume 2*, edited by: Lollino, G., Giordan, Crosta, G. B., Corominas, J.,
695 Azzam, R., Wasowski, J., Sciarra, N., Springer International Publishing Switzerland, chapter: 383, https://doi.org/10.1007/978-3-319-09057-3_383, 2015

Bickle, M. J., Tipper, E. T., Galy, A., Chapman, H., and Harris, N.: On Discrimination Between Carbonate and Silicate Inputs to Himalayan Rivers, *Am. J. Sci.*, 315, 120–166, <https://doi.org/10.2475/02.2015.02>, 2015

700

Binet, S., Spadini, L., Bertrand, C., Guglielmi, Y., Mudry, J., and Scavia, C.: Variability of the groundwater sulfate concentration in fractured rock slopes: a tool to identify active unstable areas, *Hydrol. Earth Syst. Sci.*, 13, 2315–2327, <https://doi.org/10.5194/hess-13-2315-2009>, 2009

705 Blattmann T. M., Wang, S. L., Lupker, M., M arki, L., Haghypour, N., Wacker, L., Chung, L. H., Bernasconi, S. M, Pl tze, M., Eglinton, T. I.: Sulphuric acid-mediated weathering on Taiwan buffers geological atmospheric carbon sinks, *Sci Rep*, 2945, <https://doi.org/10.1038/s41598-019-39272-5>, 2019

Brass, G. W.: The effect of weathering on the distribution of strontium isotopes in weathering profiles. *Geochimica et Cosmochimica Acta*, 39, 1647–1653, [https://doi.org/10.1016/0016-7037\(75\)90086-1](https://doi.org/10.1016/0016-7037(75)90086-1), 1975

710

Bullen, T. D., Krabbenhoft, D. P., Kendall, C.: Kinetic and mineralogic controls on the evolution of groundwater chemistry and $^{87}\text{Sr}/^{86}\text{Sr}$ in a sandy silicate aquifer, northern Wisconsin, USA. *Geochimica et Cosmochimica Acta*, 60(10), 1807–1821, [https://doi.org/10.1016/0016-7037\(96\)00052-X](https://doi.org/10.1016/0016-7037(96)00052-X), 1996

715 Bullen, T. D. and Kendall, C.: Tracing of Weathering Reactions and Water Flowpaths: A Multi-isotope Approach, in: *Isotope Tracers in Catchment Hydrology*, edited by Kendall, C., McDonnell, J. J, Elsevier Science B.V., Amsterdam, pp. 611-646, <https://doi.org/10.1016/B978-0-444-81546-0.50025-2>, 1998

Burke, W.H., Denison, R.E., Hetherington, E.A., Koepnick, R.B., Nelson, H.F. and Otto, J.B.: Variation of seawater $^{87}\text{Sr}/^{86}\text{Sr}$ throughout Phanerozoic time. *Geology* 10: 516-519., [https://doi.org/10.1130/0091-7613\(1982\)10<516:VOSSTP>2.0.CO;2](https://doi.org/10.1130/0091-7613(1982)10<516:VOSSTP>2.0.CO;2), 1982

720

Calmels, D., Gaillardet, J., Brenot, A., & France-Lanord, C.: Sustained sulfide oxidation by physical erosion processes in the Mackenzie River basin: Climatic perspectives, *Geology*, 35(11), 1003–1006, <https://doi.org/10.1130/G24132A.1>, 2007

725

- Canfield, D. E., Raiswell, R., Westrich, J. T., Reaves, C. M., Berner, R. A.: the use of chromium reduction in the analysis of reduced inorganic sulfur in sediments and shales, *Chemical Geology*, 54(1-2), 149–155, [https://doi.org/10.1016/0009-2541\(86\)90078-1](https://doi.org/10.1016/0009-2541(86)90078-1), 1986
- 730 Cappa, F., Guglielmi, Y., Soukatchoff, V., Mudry, J., Bertrand, C., Charmoille, A.: Hydromechanical modeling of a large moving rock slope inferred from slope levelling coupled to spring long-term hydrochemical monitoring: example of the La Clapière landslide (Southern Alps, France), *Journal of Hydrology*, 291(1–2), 67–90, <https://doi.org/10.1016/J.JHYDROL.2003.12.013>, 2004
- 735 Cervi, F., Ronchetti, F., Martinelli, G., Bogaard, T. A., and Corsini, A.: Origin and assessment of deep groundwater inflow in the Ca' Lita landslide using hydrochemistry and in situ monitoring, *Hydrol. Earth Syst. Sci.*, 16, 4205–4221, <https://doi.org/10.5194/hess-16-4205-2012>, 2012
- Clow, D. W and Drever, J. I.: Weathering rates as a function of flow through an alpine soil, *Chemical Geology*, 132(1–4),
 740 131–141, [https://doi.org/10.1016/S0009-2541\(96\)00048-4](https://doi.org/10.1016/S0009-2541(96)00048-4), 1996
- Cortecchi, G., Dinelli, E., Bencini, A., Adorni-Braccesi, A., La Ruffa, G.: Natural and anthropogenic SO₄ sources in the Arno river catchment, northern Tuscany, Italy: a chemical and isotopic reconnaissance, *Applied Geochemistry*, 17(2), 79–92, [https://doi.org/10.1016/S0883-2927\(01\)00100-7](https://doi.org/10.1016/S0883-2927(01)00100-7), 2002
- 745 Deiana, M., Cervi, F., Pennisi, M., Mussi, M., Bertrand, C., Tazioli, A., Ronchetti, F.: Chemical and isotopic investigations ($\delta^{18}\text{O}$, $\delta^2\text{H}$, ^3H , $^{87}\text{Sr}/^{86}\text{Sr}$) to define groundwater processes occurring in a deep-seated landslide in flysch. *Hydrogeology Journal*, 26(8), 2669–2691, <https://doi.org/10.1007/s10040-018-1807-1>, 2018
- 750 Dotsika, E., Poutoukis, D., Kloppmann, W., Guerrot, C., Voutsas, D., Kouimtzis, T. H.: The use of O, H, B, Sr and S isotopes for tracing the origin of dissolved boron in groundwater in Central Macedonia, Greece. *Applied Geochemistry*, 25(11), 1783–1796. <https://doi.org/10.1016/J.APGEOCHEM.2010.09.006>, 2010
- Dubois, L., Chanut, M.-A., Duranthon, J. P.: Amélioration continue des dispositifs d'auscultation et de surveillance intégrés
 755 dans le suivi du versant instable des Ruines de Séchilienne in : *Géologues*, 182, edited by : Editions du BRGM, 50–55, 2014
- Emberson, R., Hovius, N., Galy, A. Marc, O.: Chemical weathering in active mountain belts controlled by stochastic bedrock landsliding, *Nature Geoscience*, 9, 42–45, <https://doi.org/10.1038/ngeo2600>, 2015
- 760 Emberson, R., Hovius, N., Galy, A., Marc, O.: Oxidation of sulfides and rapid weathering in recent landslides, *Earth Surface Dynamics*, 4(3), 727–742, <https://doi.org/10.5194/esurf-4-727-2016>, 2016b
- Emberson, R., Galy, A., Hovius, N.: Weathering of Reactive Mineral Phases in Landslides Acts as a Source of Carbon Dioxide in Mountain Belts, *Journal of Geophysical Research: Earth Surface*, 123(10), 2695–2713.
 765 <https://doi.org/10.1029/2018JF004672>, 2018

- Fanlo, I. and Ayora, C.: The evolution of the Lorraine evaporitic basin: implications for the chemical and isotope composition of the Triassic ocean, *Chemical Geology*, vol. 146 n° 3-4, p. 135–154, <https://doi.org/10.1016/S0009-25419800007-2>, 1998
- 770
- Fletcher, R. C., Buss, H. L., and Brantley, S. L.: A spheroidal weathering model coupling porewater chemistry to soil thicknesses during steady-state denudation, *Earth Planet. Sci Lett.*, 244, 444–457, <https://doi.org/10.1016/j.epsl.2006.01.055>, 2006
- Follacci, J.: Seize ans de surveillance du glissement de la Clapière (Alpes maritimes). 220, Rapport, *Bulletin du Laboratoire des Ponts et Chaussées*. 35-51, 1999
- 775
- Frost, C. D., Toner, R. N.: Strontium isotopic identification of water-rock interaction and ground water mixing, *Ground Water*, 42(3), 418–432, <https://doi.org/10.1111/j.1745-6584.2004.tb02689.x>, 2004
- 780
- Gaillardet, J., Dupre, B., Allegre C. J., Négrel, P.: Chemical and physical denudation in the Amazon River Basin, *Chemical Geology*, vol. 142 n° 3–4, p. 141–173, [https://doi.org/10.1016/S0009-2541\(97\)00074-0](https://doi.org/10.1016/S0009-2541(97)00074-0), 1997
- Gaillardet, J., Dupre, B., Louvat, P., Allegre, C. J.: Global silicate weathering and CO₂ consumption rates deduced from the chemistry of large rivers, *Chemical Geology*, 159, 3–30, [https://doi.org/10.1016/S0009-2541\(99\)00031-5](https://doi.org/10.1016/S0009-2541(99)00031-5), 1999
- 785
- Gammons, C. H., Brown, A., Poulson, S. R., Henderson, T. H.: Using stable isotopes (S, O) of sulfate to track local contamination of the Madison karst aquifer, Montana, from abandoned coal mine drainage, *Applied Geochemistry*, 31, 228–238, <https://doi.org/10.1016/J.APGEOCHEM.2013.01.008>, 2013
- 790
- Gu, X., Rempe, D., Dietrich, W., West, A., Lin, T.C., Jin, L., Brantley, S.: Chemical reactions, porosity, and microfracturing in shale during weathering: The effect of erosion rate. *Geochimica et Cosmochimica Acta*. 269, 2019
- Guglielmi, Y., Vengeon, J., Bertrand, C., Mudry, J., Follacci, J., Giraud, A.: Hydrogeochemistry: an investigation tool to evaluate infiltration into large moving rock masses (case study of La Clapière and Séchilienne alpine landslides), *Bulletin of Engineering Geology and the Environment*, 61(4), 311–324. <https://doi.org/10.1007/s10064-001-0144-z>, 2002
- 795
- Hajj, F., Poszwa, A., Bouchez, J., Guérol, F.: Radiogenic and “stable” strontium isotopes in provenance studies: A review and first results on archaeological wood from shipwrecks, *Journal of Archaeological Science*, 86, 24–49, <https://doi.org/10.1016/j.jas.2017.09.005>, 2017
- 800
- Hilton, R.G. and West, A.J.: Mountains, erosion and the carbon cycle, *Nature reviews earth & environment*, 1 (6). pp. 284–299, <https://doi.org/10.1038/s43017-020-0058-6>, 2020
- Kampschulte, A., Strauss, H.: The sulfur isotopic evolution of Phanerozoic seawater based on the analysis of structurally substituted sulfate in carbonates, *Chemical Geology*, 204(3–4), 255–286, <https://doi.org/10.1016/j.chemgeo.2003.11.013>, 2004
- 805

- 810 Koepnick, R. B., Denison, R. E., Burke, W. H., Hetherington, E. A., Dahl, D. A.: Construction of the Triassic and Jurassic portion of the Phanerozoic curve of seawater $87\text{Sr}/86\text{Sr}$, *Chemical Geology: Isotope Geoscience Section*, 80(4), 327–349, [https://doi.org/10.1016/0168-9622\(90\)90014-4](https://doi.org/10.1016/0168-9622(90)90014-4), 1990
- 815 Lajaunie, M., Gance, J., Nevers, P., Malet, J.-P., Bertrand, C., Garin, T., and Ferhat, G.: Structure of the S echilienne unstable slope from large-scale 3d electrical tomography using a resistivity distributed automated system (r-das). *Geophysical Journal International*, <https://doi.org/10.1093/gji/ggz259>, 2019
- Lasaga, A. C.: Chemical kinetics of water-rock interactions, *Journal of Geophysical Research: Solid Earth*, 89(B6), 4009–4025, <https://doi.org/10.1029/jb089ib06p04009>, 1984
- 820 Lerman, A., Wu, L., Mackenzie, F. T.: CO_2 and H_2SO_4 consumption in weathering and material transport to the ocean, and their role in the global carbon balance, *Marine Chemistry*, 106(1-2 SPEC. ISS.), 326–350, <https://doi.org/10.1016/j.marchem.2006.04.004>, 2007
- 825 Le Roux, O., Jongmans, D., Kasperski, J., Schwartz, S., Potherat, P., Lebrouc, V., Meric, O.: Deep geophysical investigation of the large S echilienne landslide (Western Alps, France) and calibration with geological data, *Engineering Geology*, 120(1–4), 18–3, <https://doi.org/10.1016/J.ENGGEOL.2011.03.004>, 2011
- Li, S. L., Calmels, D., Hana, G., Gaillardet, J., Liu, C. Q.: Sulfuric acid as an agent of carbonate weathering constrained by $\delta^{13}\text{CDIC}$: Examples from Southwest China, *Earth and Planetary Science Letters*, 270, 3–4, 30, 189–199, <https://doi.org/10.1016/j.epsl.2008.02.039>, 2008
- 830 Mar echal, J.-C., 1998. Les circulations d’eau dans les massifs cristallins alpins et leurs relations avec les ouvrages souterrains, *Ecole Polytechnique F ed erale de Lausanne (EPFL)*, 1998
- 835 Meric, O., Garambois, S., Jongmans, D., Wathelet, M., Chatelain, J. L., Vengeon, J. M.: Application of geophysical methods for the investigation of the large gravitational mass movement of S echilienne, France, *Canadian Geotechnical Journal*, 42(4), 1105–1115, <https://doi.org/10.1139/t05-034>, 2005
- 840 Meybeck Michel. Composition chimique des ruisseaux non pollu es en France. Chemical composition of headwater streams in France. In: *Sciences G eologiques*, tome 39, n o 1, pp. 3-77, <https://doi.org/10.3406/sgeol.1986.1719>, 1986
- 845 Meynadier, L., Gorge, C., Birck, J., All egre, C.: Automated separation of Sr from natural water samples or carbonate rocks by high performance ion chromatography, *Chemical Geology*, 227(1–2), <https://doi.org/10.1016/j.chemgeo.2005.05.012>, 2006
- 845 Moncaster, S. J., Bottrell, S. H., Tellam, J. H., Lloyd, J. W., Konhauser, K. O.: Migration and attenuation of agrochemical pollutants: Insights from isotopic analysis of groundwater sulphate, *Journal of Contaminant Hydrology*, 43(2), 147–163, [https://doi.org/10.1016/S0169-7722\(99\)00104-7](https://doi.org/10.1016/S0169-7722(99)00104-7), 2000

- Montety, V. de, Marc, V., Emblanch, C., Malet, J.-P., Bertrand, C., Maquaire, O., Bogaard, T. A.: Identifying the origin of groundwater and flow processes in complex landslides affecting black marls: insights from a hydrochemical survey, *Earth Surface Processes and Landforms*, 32(1), 32–48, <https://doi.org/10.1002/esp.1370>, 2007
- 850
- Montjuvent, G. and Winistorfer, J.: Glaciations quaternaires dans les Alpes franco-suissees et leur piedmont, in *Géologie Alpine*, 56, 251–282, 1980
- 855
- Mudry, J., & Etievant, K., 2007. Synthèse hydrogéologique du versant instable des Ruines de Séchilienne (Unpublished report), UMR Chrono-Environnement, University of Franche-Comté, Besançon, 2007
- Négrel, P., Allègre, C. J., Dupré, B., Lewin, E.: Erosion sources determined by inversion of major and trace element ratios and strontium isotopic ratios in river: The Congo Basin case, *Earth and Planetary Science Letters*, 120(1–2), 59–76, [https://doi.org/10.1016/0012-821X\(93\)90023-3](https://doi.org/10.1016/0012-821X(93)90023-3), 1993
- 860
- Négrel, P. and Deschamps, P.: Natural and anthropogenic budgets of a small watershed in the massif central (France): Chemical and strontium isotopic characterization of water and sediments, *Aquatic Geochemistry*, 2(1), 1–27, <https://doi.org/10.1007/BF00240851>, 1996
- 865
- Négrel, P., Casanova, J., Aranyossy, J.-F.: Strontium isotope systematics used to decipher the origin of groundwaters sampled from granitoids: the Vienne Case (France), *Chemical Geology*, 177(3–4), 287–308, [https://doi.org/10.1016/S0009-2541\(00\)00414-9](https://doi.org/10.1016/S0009-2541(00)00414-9), 2001
- 870
- Oudin, L., Hervieu, F., Michel, C., Perrin, C., Andréassian, V., Anctil, F., and Loumagne, C.: Which potential evapotranspiration input for a lumped rainfall–runoff model?: Part 2—towards a simple and efficient potential evapotranspiration model for rainfall–runoff modelling, *Journal of hydrology*, 303(1–4):290–306, 2005
- 875
- Raymo, M. E.: Geochemical evidence supporting T. C. Chamberlin's theory of glaciation, *Geology*, 19, 344–347, [https://doi.org/10.1130/0091-7613\(1991\)019<0344:GESTCC>2.3.CO;2](https://doi.org/10.1130/0091-7613(1991)019<0344:GESTCC>2.3.CO;2), 1991
- Raymo, M., Ruddiman, W.: Tectonic forcing of late Cenozoic climate, *Nature*, 359, 117–122, <https://doi.org/10.1038/359117a0>, 1992
- 880
- Russell, W. A. and Papanastassiou, D. A.: Calcium isotope fractionation in ion-exchange chromatography, *Analytical Chemistry*, 50(8), 1151–1154, <https://doi.org/10.1021/ac50030a036>, 1978
- Singleton, M. J., Maher, K., DePaolo, D. J., Conrad, M. E., Evan Dresel, P.: Dissolution rates and vadose zone drainage from strontium isotope measurements of groundwater in the Pasco Basin, WA unconfined aquifer, *Journal of Hydrology*, 321(1–4), 39–58, <https://doi.org/10.1016/J.JHYDROL.2005.07.044>, 2006
- 885
- Spence, J., Telmer, K.: The role of sulfur in chemical weathering and atmospheric CO₂ fluxes: evidence from major ions, $\delta^{13}\text{CDIC}$, and $\delta^{34}\text{S SO}_4$ in rivers of the Canadian Cordillera, *Geochimica Cosmochemica*, 69, 5441–5458, <https://doi.org/10.1016/j.gca.2005.07.011>, 2005
- 890

- Stallard, R. F. and Edmond, J. M.: Geochemistry of the Amazon: 2. The influence of geology and weathering environment on the dissolved load, *Journal of Geophysical Research: Oceans*, 88, 9671–9688, <https://doi.org/10.1029/jc088ic14p09671>, 1983
- 895
- Torres, M. A., West, A. J., Li, G.: Sulphide oxidation and carbonate dissolution as a source of CO₂ over geological timescales, *Nature*, 507(7492), 346–349, <https://doi.org/10.1038/nature13030>, 2014
- Torres, M. A., West, A. J., Clark, K. E., Paris, G., Bouchez, J., Ponton, C., Adkins, J. F.: The acid and alkalinity budgets of weathering in the Andes–Amazon system: Insights into the erosional control of global biogeochemical cycles. *Earth and Planetary Science Letters*, 450, 381–391, <https://doi.org/10.1016/j.epsl.2016.06.012>, 2016
- 900
- Vallet, A. *Mouvements de fluides et processus de déstabilisation des versants alpins : Apport de l'étude de l'instabilité de Séchilienne*, Université de Franche-Comté, 2014
- 905
- Vallet, A., Bertrand, C., Mudry, J., Bogaard, T., Fabbri, O., Baudement, C., Régent, B.: Contribution of time-related environmental tracing combined with tracer tests for characterization of a groundwater conceptual model: a case study at the Séchilienne landslide, western Alps (France), *Hydrogeology Journal*, 23(8), 1761–1779, <https://doi.org/10.1007/s10040-015-1298-2>, 2015a
- 910
- Vallet, A., Charlier, J. B., Fabbri, O., Bertrand, C., Carry, N., Mudry, J.: Functioning and precipitation-displacement modelling of rainfall-induced deep-seated landslides subject to creep deformation, *Landslides*, 13, 653–670, <https://doi.org/10.1007/s10346-015-0592-3>, 2015b
- 915
- Vengeon, J. M.: *Déformation et rupture des versants en terrain métamorphique anisotrope : Apport de l'étude des Ruines de Séchilienne*, Université Joseph-Fourier - Grenoble I, 1998
- Vengeon, J.-M., Couturier, B., Antoine, P.: Déformations gravitaires post glaciaires en terrains métamorphiques. Comparaison des indices de déformation du versant sud de la Toura (Saint-Christophe-en-Oisans, France) avec le phénomène de rupture interne du versant sud du Mont Sec (Séchilienne, France), *Bulletin of Engineering Geology and the Environment*, 57(4), 387–395, <https://doi.org/10.1007/s100640050062>, 1999
- 920
- Woods, T. L., Fullagar, P. D., Spruill, R. K., Sutton, L. C.: Strontium isotopes and major elements as tracers of ground water evolution: example from the Upper Castle Hayne Aquifer of North Carolina, *Ground Water*, 38(5), 762–771, <https://doi.org/10.1111/j.1745-6584.2000.tb02712.x>, 2000
- 925
- Zeebe, R. E. and Wolf-Gladrow, D. A.: *CO₂ in Seawater: Equilibrium, Kinetics, Isotopes*, edited by Elsevier Oceanography Series, Amsterdam, 65, 2001.
- 930