

Sediment size on talus slopes correlates with fracture spacing on bedrock cliffs: Implications for predicting initial sediment size distributions on hillslopes

Joseph P. Verdian¹, Leonard S. Sklar^{1,2}, Clifford S. Riebe³, Jeffrey R. Moore⁴

¹Department of Earth and Climate Sciences, San Francisco State University, San Francisco, 94302, USA

²Department of Geography, Planning and Environment, Concordia University, Montreal, H3G1M8, Canada

³Department of Geology and Geophysics, University of Wyoming, Laramie, 82071, USA

⁴Department of Geology and Geophysics, University of Utah, Salt Lake City, 84112, USA

Correspondence to: Leonard S. Sklar (leonard.sklar@concordia.ca)

Abstract. The detachment of rock fragments from fractured bedrock on hillslopes creates sediment with an initial size distribution that sets the upper limits on particle size for all subsequent stages in the **evolution** of sediment in landscapes. We hypothesize that the initial size distribution should depend on the size distribution of latent sediment (i.e., **fracture-bound blocks in unweathered bedrock**) and weathering of sediment **both before and during** detachment (e.g., disintegration along crystal grain boundaries). However, the initial size distribution is difficult to measure, because the interface across which sediment is produced is often shielded from view by overlying soil. Here we overcome this limitation by comparing fracture spacings measured from exposed bedrock on cliff faces with particle size distributions in adjacent talus deposits at 15 talus-cliff pairs spanning a wide range of climates and lithologies in California. Median fracture spacing and particle size vary by more than tenfold and correlate strongly with lithology. Fracture spacing and talus size distributions are also closely correlated in central tendency, spread, and shape, with b-axis diameters showing the closest correspondence with fracture spacing at most sites. This suggests that weathering has not modified latent sediment either before or during detachment from the cliff face. In addition, talus has not undergone much weathering after deposition and is slightly coarser than the latent sizes, **because it contains unexploited fractures inherited from bedrock**. We introduce a new conceptual framework for understanding the relative importance of latent size and weathering in setting initial sediment size distributions in mountain landscapes. In this framework, hillslopes exist on a spectrum defined by the ratio of two characteristic timescales: the residence time in saprolite and weathered bedrock, and the time required to detach a particle of a characteristic size. At one end of the spectrum, where weathering residence times are negligible, the latent size distribution can be used to predict the initial size distribution. At the other end of the spectrum, where weathering residence times are long, the latent size distribution can be erased by weathering in the critical zone.

Style Definition: Normal

Style Definition: Heading 1: Complex Script Font: Times New Roman, 10 pt, Not Bold, (Asian) Chinese (China), (Other) English (US)

Style Definition: Heading 2: Complex Script Font: Times New Roman, 10 pt, Not Bold, Not Italic, (Asian) Chinese (China), (Other) English (US)

Style Definition: Heading 3: Complex Script Font: Times New Roman, 10 pt, Not Bold, (Asian) Chinese (China), (Other) English (US)

Style Definition: Heading 4: Complex Script Font: 10 pt, Not Bold, (Asian) Chinese (China), (Other) English (US)

Style Definition: Bullets

Formatted: Font: 17 pt, Bold, Font color: Black

Formatted: From text: 0", Numbering: Restart each page

Formatted: Normal, Justified, Space Before: 18 pt, Line spacing: Multiple 1.83 li, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Deleted: ¶

Formatted: Font color: Black

Formatted: Font: 10 pt, Font color: Black

Deleted: life

Deleted: defined by through-going fractures

Deleted: or

Deleted: suggesting that

Deleted: some

Deleted: ¶

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

1 Introduction

The detachment of rock fragments from fractured or weathered bedrock creates sediment particles of various sizes that evolve due to chemical and physical weathering during transport and storage on slopes and in rivers. At all stages in the evolution of sediment, its size distribution influences local chemical, physical, and biological processes, including throughflow of reactive fluids in soil (Maher, 2010; Brantley et al., 2011), river incision into bedrock (Sklar and Dietrich, 2004; Turowski et al., 2015; Shobe et al., 2016), and the reproductive potential of aquatic habitat (Riebe et al., 2014; Overstreet et al., 2016). The first stage begins when particles are initially detached from saprolite (in soil-mantled landscapes) or fractured bedrock (when soil is absent). At any specific source location, whether it's at the surface or at the base of soil, a set of these newly detached sediment particles have an "initial size distribution," where initial is defined by the moment when particles are first disturbed by processes that drive particle motion on hillslopes (Sklar et al., 2017; Roda-Boluda et al., 2018; Neely and DiBiase, 2020). The initial size distribution sets the initial condition for subsequent evolution of the size distribution by abrasion and fragmentation of particles during transport, both on hillslopes near where sediment is produced and in river channels downstream. The initial size distribution is therefore a fundamental control on sediment size distributions across landscapes. It sets the upper limit for sediment size in a catchment; in the absence of flocculation or cementation, sediment particles can only become smaller. In addition, the initial size sets the scale for particle size reduction by comminution during transport away from its source. The downstream evolution of sediment size is commonly modeled as a function of three factors: the initial size, the travel distance downstream from the source, and a rate constant that depends primarily on rock durability (Sklar et al., 2006; Attal and Lavé, 2009; Dingle et al., 2017). Thus, a coarser initial size will result in a coarser downstream size, for a given transport distance and rock durability. Particle shape also evolves during transport downstream and can be used to estimate the distance traveled from the particle's source when initial shape is known (Miller et al., 2014; Szabo et al., 2015). However, the factors that regulate variability in the initial size and shape of sediment are poorly understood.

The initial size distribution and initial particle shape should depend on three factors: the size distribution of discrete blocks latent within bedrock; the weathering processes that modify rock in situ; and the weathering processes that detach and mobilize the sediment. Where effects of weathering on size are minimal, the initial size distribution should strongly reflect the size distribution of latent blocks (i.e. the "latent size distribution"), which is set by the spacing and orientation of fractures, foliations, and bedding planes (Sklar et al., 2017). These represent preexisting planes of weakness and determine the volume and shape of newly created sediment particles, which can be quantified by measuring the distributions of the major-, intermediate-, and minor-axis particle diameters. The three-dimensional template for latent particles should depend on conditions experienced during formation of the rock at depth, including rate of cooling for igneous rocks (Lore et al., 2001), pressure and temperature for metamorphic rocks (Manda et al., 2008), and deposition and diagenesis in sedimentary rocks (Narr and Suppe, 1991). These factors are overprinted by fracturing induced by the evolving stress field experienced by the rock as it is exhumed from deep in the crust (Molnar et al., 2007; Leith et al., 2014; Moon et al., 2017). Thus, the latent size distribution reflects everything that has happened to the rock before it arrives at the base of the critical zone, where it is first

Deleted: due to chemical and physical weathering.

Deleted: life

Deleted: 2015

Deleted: The resulting

Moved (insertion) [1]

Deleted: the starting point

Deleted: the

Deleted: on the hillslope where the

Deleted: sets the upper limits

Deleted: particle

Deleted: at each successive stage

Deleted: sediment's life (Sklar et al.,

Moved up [1]: 2017; Roda-Boluda et al., 2018

Deleted: .

Deleted: particle shape and

Deleted: size distribution

Deleted: both

Deleted: blocks in

Deleted: and on

Deleted: characteristic length scales of subsequent

Deleted: . The "latent"

Deleted: , and mineral grain boundaries.

Deleted: experiencing weathering in

Deleted: 2

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

exposed to near-surface conditions that promote weathering (Riebe et al., 2017). Where near-surface residence times are long or weathering is intense, the initial size distribution can be affected by in-situ weathering during exhumation through the critical zone (Sklar et al., 2017). For example, chemical reactions such as mineral dissolution and physical processes such as mineral expansion can create new planes of weakness and thus influence the initial size distribution created during detachment (Fletcher and Brantley, 2010; Brantley et al., 2011; Shen et al., 2019). In-situ chemical and physical weathering can also modify rock strength, density, and other properties that might affect initial size without altering the apparent fracture spacing (Goodfellow et al., 2016; Riebe et al., 2021). In addition to these in-situ weathering processes, the initial size distribution should also depend on the characteristic length scales of weathering processes that detach and mobilize sediment, such as segregation ice growth, root wedging, and animal burrowing (Sklar et al., 2017; Messenzehl et al., 2018), particularly on slopes where preexisting planes of weakness in bedrock are widely spaced (Marshall and Roering, 2014). Although these hypotheses are both intuitive and mechanistic, to our knowledge, the relative importance of latent sizes, in-situ weathering, and detachment processes in initial size distributions has not been systematically explored. The initial detachment of rock fragments is generally hidden from view under soil, and particle sizes can therefore evolve during mixing, transport, and storage, before they are measured. This makes it difficult to isolate the influence of the latent size distribution from the effects of weathering processes.

To overcome this limitation, previous studies have compared fracture spacings measured from exposed bedrock on cliff faces with particle size distributions in adjacent sediment deposits. For example, at Inyo Creek, on the east side of the Sierra Nevada, California, where hillslope sediment size distributions are bimodal, in-situ measurements from bedrock cliff faces show that fracture spacing distributions do not vary with elevation and closely correspond with the likewise spatially invariant coarse mode of the hillslope particle size distribution (Sklar et al., 2020). Similarly, in California's San Gabriel and San Jacinto mountains, two regions with differing fracture spacing distributions, particle sizes in stream sediment correlate with fracture spacings measured in adjacent bedrock cliff faces at locations where the sediment contributing area is dominated by bare bedrock (DiBiase et al., 2018; Neely et al., 2019; Neely and DiBiase, 2020). Results from both of these studies are consistent with latent sizes in bedrock dominating over weathering on slopes in regulating the initial size distribution of coarse sediment. In contrast, results from paired cliffs and talus slopes in the Swiss Alps suggest that weathering by frost cracking can impose a characteristic scale upon talus particle sizes (Messenzehl et al., 2018). At these sites, talus size is roughly uniform across a suite of locations where fracture spacing differs by over an order of magnitude. Measured talus size is always either larger than or approximately equal to the mean fracture spacing, consistent with the hypothesis that talus production by frost-cracking can preferentially exploit a subset of pre-existing fractures with a characteristic spacing, leaving other fractures intact within the blocks detached from the cliff wall. The results from California and the Swiss Alps show that both the latent size distribution and sediment detachment mechanisms can strongly influence initial sediment size, when in-situ weathering is minimized. Quantifying the relative importance of these factors at additional sites is vital to predicting initial sediment sizes across a range of conditions.

Deleted: environment. As

Deleted: commences, but before particles are detached, physical and chemical processes can create new surfaces that can be exploited during detachment. Hence

Deleted: should also depend on the characteristic length scales of

Deleted: processes, such as mineral expansion, segregation ice growth, root wedging, and animal burrowing, particularly on slopes where preexisting planes of weakness in bedrock are widely spaced

Moved down [2]: 2017; Messenzehl et al., 2018

Deleted:). In addition to these physical weathering processes

Moved (insertion) [2]

Deleted: Goodfellow et al., 2016).

Deleted: and

Deleted: saprolite,

Deleted: weathered rock, and

Deleted: can be accessed and

Deleted: Elsewhere in California, at two sites

Deleted: DiBiase

Deleted: a suite of sites

Deleted: , leading to poor correlations with fracture spacing distributions on adjacent bedrock cliffs (Messenzehl et al., 2018).

Deleted: 3

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

Here we quantify correlations between initial sediment size and fracture spacing distributions across 15 talus-cliff pairs spanning a wide range of settings, including rock types and climatic conditions not investigated in previous work. Our sites span a 3-km range in elevation across granodiorite, andesite, basalt, metasedimentary, and chert lithologies in California. All cliffs are sufficiently steep to prevent any accumulation of regolith, such that surface fracture spacing can be assumed to reflect the latent size distribution with minimal influence of weathering. Thus our study design allows us to test the null hypothesis that the initial sizes closely match latent sizes across a wide, geologically driven range in latent size distributions. An alternate hypothesis is that talus size distributions and the latent size distributions exposed in cliff faces are not strongly correlated for one or more of the following reasons: (i) blocks are detached along only a subset of preexisting fractures, because of characteristic scales of detachment processes such as frost cracking (Messenzehl et al., 2018), or due to unequal fracture persistence (Kim et al., 2007); (ii) talus particles are created by detachment along newly formed fractures, as in the case of grus production from granite with low fracture density (Wahrhaftig, 1965); (iii) physical or chemical weathering reduces particle sizes after the talus is detached from the cliff, for example when falling particles impact the slope below or as they sit in the talus deposit. None of these alternatives to the null hypothesis are consistent with our results. Measurements of central tendency, spread, and shape of the talus size and fracture spacing distributions all correlate strongly across a 40-fold variation in median fracture spacing. We also found statistically significant differences in mean talus shape among rock types, contrary to the null hypothesis that initial particle shape is invariant for blocks produced from bedrock by mechanical weathering (Domokos et al., 2015). Together these results confirm that initial sediment size distributions can be predicted from fracture spacing distributions at sites where the latent size distribution dominates over weathering. They also imply that lithologic and tectonic controls on latent size distributions can have a strong influence on the initial size and shape of individual particles and thus on the evolution of particle size distributions across landscapes. To generalize our findings beyond the talus-cliff pairs studied here, we introduce a conceptual framework for quantifying the relative importance of latent sizes and weathering using the timescale of detachment of latent particles and the timescale of weathering that occurs before the particle is detached.

Deleted: 3000-m

Deleted:

Deleted: cones

Deleted: distribution

Deleted: both

Deleted: for example

Deleted:), or

Deleted:

Deleted: ii

Deleted: hit

Deleted: adjacent

Deleted: Neither

2 Methods

2.1 Site selection

To test the null hypothesis that latent size, set by fracture spacing, should control the initial size of sediment produced on bare bedrock slopes, we selected 15 cliff faces and adjacent talus slopes at five sites in California, USA (Fig. 1). The talus-cliff pairs in the Sierra Nevada are a subset of the bedrock slopes used by Moore et al. (2009) to quantify the influence of rock-mass strength on cliff retreat rates. These include ten slopes in the vicinity of Conness Basin, Mount Tallac, and Ebbetts Pass (Table 1), three sites where differences in lithology correspond to differences in average fracture spacing. To diversify the range of conditions that might contribute to differences in weathering and thereby produce deviations from the null hypothesis, we selected five additional talus-cliff pairs at Grizzly Peak and Twin Peaks, two sites in the San Francisco Bay Area (Table 1).

Deleted:

Deleted: about strong connections between

Deleted: and

Deleted: and shape

Deleted: represent

Deleted: featured in previous work on connections between

Deleted: and

Deleted: (Moore et al., 2009).

Deleted: a series of

Deleted: 4

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

The Sierra Nevada sites include exposures of the three major classes of rock comprising the high elevation terrain of the range (Bateman et al., 1966). The youngest rocks are volcanic basalts and andesites, exposed at Mount Tallac and Ebbetts Pass, which erupted during the Miocene Epoch (Armin et al., 1984). These volcanic rocks overlie intrusive granitic rocks characteristic of the Sierra Nevada batholith, emplaced during the Cretaceous Period, including the granodiorites exposed at Conness Basin (Bateman, 1983) and Mount Tallac (Saucedo, 2005). Also exposed at Mount Tallac are the oldest rocks of our study sites, Paleozoic-age metasedimentary roof pendants of the country rock intruded by the Cretaceous granitics (Saucedo, 2005). Bedrock at the San Francisco Bay Area sites is characteristic of the Franciscan Formation, and includes Mesozoic pillow basalt and overlying chert exposed at Twin Peaks, as well as Miocene subaerial basalt flows exposed at Grizzly Peak (Graymer et al., 2006).

This spectrum of rock types and geologic settings provides a potentially large range of latent size distributions, from the widely-spaced fractures in the massive granodiorite at Conness Basin to the closely-spaced bedding planes of the Twin Peaks ribbon chert. Similarly, fractures in bedrock at our sites are produced by a range of mechanisms, from cooling of subaerial lava at Grizzly Peak, Ebbetts Pass, and Mount Tallac to the crystallization and exhumation of granodiorite through ~10 km of crust at Conness Basin (Ague, 1997). In addition, the sites span a range in weathering conditions, due to the 3-km range in elevation, the 10°C range in mean annual temperature, and the twofold variation in mean annual precipitation (Table 1). Collectively, this combination of rock types, fracturing mechanisms, and weathering conditions can be expected to produce talus spanning much of the range in sizes found at the bases of cliffs in mountains around the world.

2.2 Fracture spacing distributions on cliff faces

To quantify fracture spacing at each site, we used a horizontal scan line (Priest and Hudson, 1976; Moore et al., 2009; Messenzehl et al., 2018) consisting of a survey tape stretched across the cliff face at a constant elevation (e.g., Fig. 2). The height of the scan line above the top of the adjacent talus cone ranged from 0.3 to 1.5 meters in our study, depending mostly on ease of sampling. Our approach assumes that fracture spacing along a single horizontal line is representative of the contributing area of the talus, including unreachable sections above the scan line. We set the length of each scan line equal to the width of contact between the cliff face and its adjacent talus cone, which ranged between 5- and 15-meters long across our sites. Thus we limited our measurements of fracture spacing to the width of the talus source area. Along each scan line, we measured the position of every fracture that crossed the tape, irrespective of orientation. This yields a distribution of fracture spacings measured as the distance between successive fractures. Our goal was to sample the spacing between fractures that could produce a particle via rockfall. Therefore, we quantified the distance between fractures that were unambiguous partings in the rock (with aperture generally larger than 0.5 mm) that were also through-going (i.e., persistent enough to intersect other fractures on the cliff face). We ignored fractures with spacings less than 2 mm, and variations in surface roughness and other

Deleted: 1

Deleted: Thus

Deleted: focused on open,

Deleted: greater

Deleted: , ignoring changes

Deleted: 5

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

rock defects. This approach does not account for the potential role of micro-fracturing at the scale of mineral grains (Eppes and Keanini, 2017; Eppes et al., 2018) in generating detachable particles on the cliff faces.

Deleted: that did not extend far enough to intersect other fractures on the cliff face.

Deleted: ignores

Deleted: microfracturing

Deleted: 2

2.3 Particle size distributions in talus cones

To quantify surface particle size distributions in talus at each of the 10 Sierra Nevada talus-cliff pairs, we measured the a-, b-, and c-axis diameters of particles at evenly-spaced points along three slope-parallel transects extending from the base of the cliff to the toe of the talus slope (e.g. Fig. 2). At each talus-cliff pair, we measured a total of 100 particles. To obtain an equal spacing (and thus obtain a random sample), we divided the sum of the three transect lengths by 100 particles, yielding spacings that varied from 1.0 to 2.5 m/particle across the sites, with larger spacings at talus-cliff pairs that have longer talus cones. At each sampling point, we used a ruler to measure sizes of particles with diameters less than 300 mm and stadia rods to measure sizes for larger particles. In some cases, the a-, b-, and/or c-axis could not be readily measured because the particle was too heavy to move and thus to fully expose it for identification of long-, intermediate-, and short-axis orientations. In those cases, we assumed that the c axis was perpendicular to the surface slope and estimated the a- and b-axis diameters using the two longest exposed axes. We then estimated a minimum value for the c axis as the height of the particle normal to the slope.

Deleted: ,

Deleted: To measure sizes

Deleted: number of particles on each slope, we

Deleted: tape

Deleted: , and used the result to define the sample spacing, which

Deleted: to 1

Deleted: everything else. We matched the precision of the fracture spacing measurements by rounding particle diameters to the nearest millimeter and lumping diameters less than 2 mm into a <2 mm bin.

Deleted: ,

Deleted: ,

Deleted: ,

The even spacing in our talus sampling approach should yield a representative particle size distribution, even if size-selective transport leads to downslope coarsening, which is commonly observed on angle-of-repose slopes (Kirkby and Stratham, 1975). This coarsening arises because finer particles encounter relatively larger friction angles and therefore travel shorter distances before coming to rest on the talus cone. Because size-selective disentrainment occurs across the entire slope, the talus surface can be treated as a single population whose grain size distribution can be quantified representatively by uniformly spaced sampling (Bunte and Abt, 2001).

Deleted: have

Deleted: in

Deleted: .

We addressed the potential for bias due to kinetic sieving (a vertical sorting process) by supplementing the surface-based measurements with bulk samples of relatively fine subsurface sediment accessed through openings between particles at the surface at five of the Sierra Nevada talus-cliff pairs. The particle size distribution of each ~2 kg sample was measured in the lab by standard mechanical sieving. In the absence of substantial vertical sorting, these subsurface distributions should overlap with and continuously extend the fine tail of the size distribution from the surface, rather than represent a distinct population with its own mode (Bunte and Abt, 2001).

Deleted: our

Deleted: These

Deleted: match

Deleted: latent

Deleted: measured in the scan lines if

Deleted: null hypothesis is correct (i.e., the initial size distribution is strongly controlled by the latent size distribution).

Deleted: two

Deleted: 6'

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

At the three San Francisco Bay Area sites, where the talus cones are relatively small, we did not sample along linear transects. Instead, we measured surface particle size distributions using standard random point counting methods (Bunte and Abt, 2001) to sample 100 particles from each talus cone. At the Grizzly Peak site, we used rulers to quantify a-, b-, and c-axis diameters of sampled particles. At the Twin Peaks site, where talus produced from both pillow basalt and ribbon chert were small compared to talus produced at the other sites, we used a mix of calipers and rulers to quantify just the b-axis diameters.

3 Results and Interpretations

Spacings between individual fractures on cliff faces range from 2 to 5000 mm across the suite of sites, with median spacings at individual sites ranging from 10 to 390 mm (Table 2). Particle sizes span a similar range, with a-axis diameters as large as 5450 mm, c-axis diameters as small as 2 mm, and median b-axis diameters ranging from 10 to 575 mm. Both fracture spacing and particle size vary systematically with lithology: Granodiorite sites have the largest fracture spacings and particle sizes while the pillow basalt site has the smallest (Fig. 3).

Across all 15 sites, the distributions of particle sizes in talus generally correspond to the distribution of fracture spacings on adjacent cliffs. This is evident in both the similar shape and overlap of the size and spacing distributions (Fig. 3). For example, at most of the sites, the empirical cumulative distribution function (ECDF) of fracture spacing is parallel to the ECDFs of particle size, which are also generally parallel to each other where the a-, b-, and c-axis diameters were measured together (Fig. 3). In many cases, the ECDFs of size and spacing also overlap for at least one of the particle diameters. For example, at CB-1, the fracture spacing distribution closely overlaps with the size distribution of the a-axis particle diameters (Fig. 3e). In contrast, the overlap is closest with the b-axis diameters at EP-26, TP-1, and TP-3, and with the c-axis diameters at both MT-38 and CB-5. Only two of the sites (CB-2 and CB-3) have particle diameter ECDFs that do not closely parallel the fracture spacing ECDF, and only one (MT-39) has a fracture spacing ECDF that plots outside the envelope defined by the a- and c-axis diameters.

The close correspondence between distributions at each talus-cliff pair is reflected in cross-site correlations between the central tendency of fracture spacing and particle size distributions for each of the three particle axes (Fig. 4). To test for a systematic correspondence between fracture spacing and talus size, unlike at the Swiss Alps sites (Messenzehl et al., 2018), we used linear regression to fit trends to the log-transformed medians of the fracture spacing and particle diameter distributions measured at each site. We then compared the slopes of the best-fit trend lines to a slope of 1.0, which corresponds to the null hypothesis that there is a 1:1 relationship between median talus size and median fracture spacing. We found that for each of the three particle axes, the increase in median particle diameter with increasing fracture spacing (Fig. 4a-c) follows a trend with a slope that is statistically indistinguishable from 1.0 (two-tailed t-test, $p > 0.45$). This result suggests that the correspondence between talus size and fracture spacing is independent of both scale and rock type across the full range of measured sizes (up to two orders of magnitude for the b-axis diameters).

Although the site-to-site variation in talus size scales with the site-to-site variation in fracture spacing, each of the three talus-diameter trend lines is offset vertically from an exact 1:1 relationship. Of the three axes, the a-axis diameters have the largest offsets, with median diameters that are systematically greater than the median fracture spacing by a factor of ~2.5 on average (Fig. 4a). The c-axis diameters have a smaller offset (52%) and plot below the 1:1 line (Fig. 4c). The offset is smaller still and positive for the b-axis diameters (Fig. 4b), which are 42% larger on average than the fracture spacing. This offset is equivalent

7

- Deleted: At each site
- Deleted: distribution
- Deleted: closely corresponds
- Deleted: empirical
- Deleted: EDF
- Deleted: EDFs
- Deleted: EDFs
- Deleted: EDFs
- Deleted: EDF
- Deleted: EDF
- Deleted: In
- Deleted: case
- Deleted: (
- Deleted:) from a 1:1 relationship in log-log space (Fig. 4a-c).
- Deleted: is scale-
- Deleted: —
- Deleted: also independent of
- Deleted: —
- Deleted: The vertical
- Deleted: between the trend and the 1:1 line is smallest for the b-axis diameters (Fig.
- Moved down [3]: 4b), which are 42% larger on average than the fracture spacing.
- Deleted: The c-axis diameters have a slightly larger offset, plotting below the
- Deleted: line (Fig. 4c), while
- Deleted: offset
- Moved (insertion) [4]
- Moved (insertion) [3]
- Deleted: (Fig.
- Deleted: 7
- Formatted: Font: 10 pt, Font color: Black
- Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

to a $\Phi/2$ interval, using the common log transformation $\Phi = \log_2(\text{size})$. All but 2 of the 15 site-specific b-axis medians plot above the 1:1 line, suggesting that the vertical offset is not due to random variation. A binomial test indicates that 2 or fewer negative offsets would arise by chance just 0.64% of the time when negative and positive offsets are equally likely. This confirms the systematically higher b axis diameters relative to fracture spacings is likely due to a systematic measurement bias or a natural process such as preferential detachment along a subset of the measured fractures.

The link between talus and fracture-bound blocks exposed on adjacent cliff faces is further supported by the close site-to-site correspondence between the spread in the b-axis diameter distribution and the spread in the fracture spacing distribution (Fig. 4d). Because our measurements of fracture spacing and talus b-axis diameter vary over several orders of magnitude, we used the geometric standard deviation, a non-dimensional metric, to quantify and compare the spreads in the distributions of both variables. To test the null hypothesis that site-to-site variation in the spread in b-axis distribution scales directly with the site-to-site variation in fracture spacing distribution, we compared the slope of a linear trend through the data with a 1:1 slope. We found that the relationship between the geometric standard deviation of the b-axis diameters and fracture spacing across all sites is statistically indistinguishable from a 1:1 relationship (two-tailed t-test, $p > 0.67$), a quantitative reflection of the fact that the ECDFs of the b-axis diameters and fracture spacings are parallel at many of the sites in Fig. 3. Thus, both the central tendency and spread in the size and spacing distributions are closely coupled across the range of rock types and climates spanned by our study sites.

A third aspect of the particle size and fracture spacing distributions that we explored is distribution shape, which can be quantified using probability density functions. To determine if the fracture spacing and b-axis size distributions share similar shapes, we first fit exponential, log-normal, and power distribution functions to the data, recognizing that fracture spacing distributions in rock commonly have shapes that follow one of these distributions (Gillespie et al., 1993). For both the fracture spacing and talus size distributions, we found that the Weibull form of the exponential distribution (Weibull, 1951) yielded the best fit to the data in most cases. The degree to which the data follow a Weibull distribution can be evaluated graphically using a standard linearization of the Weibull equation (e.g. Nakamura et al., 2007; Litwin et al., 2012), as illustrated in Figure 5. Data sampled from a population that follows a Weibull distribution will plot on a straight line in the Weibull probability space, which is defined by the plot axes of log-transformed cumulative probability (P) and log-transformed size. In addition, because we normalized the particle size and fracture spacing measurements by their respective medians, cumulative distributions that coincide in the plotting space (i.e. have the same slopes and heights) are indicative of population distributions that have the same shape.

For most sites, both the particle size and fracture spacing data fall along straight lines and often closely coincide, as in the case of MT-38, CB-1, CB-5, and TP-1 (Fig. 5a, 5e, 5f and 5o), indicating that they share roughly the same Weibull distributions. In some cases, the slope of the b-axis distribution is steeper than the fracture spacing distribution, as in the case of MT-39 and

8

Moved up [4]: 4a).

Deleted: Given that the b-axis is the characteristic dimension that scales most closely with volume, the close correspondence between the b-axis diameters and fracture spacings (Fig. 4b) is consistent with talus production from cliffs by spallation of blocks defined by intersecting fractures without substantial size reduction during deposition...

Deleted: connection

Deleted: particles

Moved (insertion) [5]

Deleted: (Fig.

Moved up [5]: 4d).

Deleted: The

Deleted: reflecting

Deleted: parallel EDFs

Deleted: evident

Deleted: .

Deleted: there was a match in shape between

Deleted: distributions

Deleted: at each site is

Deleted: that

Deleted: are indicative of a population sampled from a Weibull distribution...

Deleted: are normalized to

Deleted: .

Deleted: 8

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

CB-3 (Fig. 5b, 5c), indicative of narrower particle size distributions and consistent with the systematically lower geometric standard deviations at these sites (Fig. 4d). In some cases, the lower tails of the distributions follow a steeper trend than the rest of the data, as in the case of CB-3, GP-1, and EP-26 (Fig. 5c, 5j, and 5k), potentially reflecting an undersampling of the smallest fractures that could result from the limited sample size and our emphasis on quantifying spacings of throughgoing fractures in the scan lines. The one rock type with data that deviate substantially from the Weibull distribution is the chert: at TP-3 and especially at TP-4 (Fig. 5l, 5m), the data show curvature in the Weibull space, and the particle size and fracture spacing distributions do not closely match. Aside from these exceptions, the fracture spacing and talus size distributions have similar shapes (Figs. 3 and 5) and are closely correlated in their central tendencies and spreads across all six lithologies (Fig. 4), consistent with our hypothesis that fracture spacing distributions can be used to predict initial particle size distributions in sediment.

Our talus size measurements do not, in contrast, support the null hypothesis that initial particle shape is invariant for blocks produced by mechanical weathering. At the 11 sites where we measured the a-, b-, and c-axes diameters, we quantified shape by calculating b:a and c:a ratios, which can be plotted together on a ternary diagram (Fig. 6) that displays rods, slabs, and equisided blocks at the vertices (Sneed and Folk, 1958; Graham and Midgley, 2000). At many of the sites, individual particles span nearly the full range of shapes represented in the diagram. Within each rock type there is little site-to-site variability in mean particle shape, suggesting that we can group sites together by rock type. When we do, we find statistically significant differences in mean particle shape among some rock types, despite substantial overlap in the distributions of individual shapes among the lithologies (Fig. 6). For example, talus produced from the metasediment have b:a and c:a ratios of 0.57 and 0.26, respectively, on average, and is therefore more elongate on average than talus produced from the granodiorite, which have corresponding ratios of 0.64 and 0.35 (Fig. 6a). In addition, andesite particles are more slab-like than basalt on average, with a lower mean c:a ratio of 0.26, compared to 0.36 (Fig. 6b). In both of these comparisons, many of the talus deposits have similar elevation and therefore similar climatic conditions (Table 1), suggesting that the differences in shape among the rock types are due to intrinsic differences in bedrock rather than differences in weathering conditions. Our results show that different rock types have different initial b:a and c:a ratios and thus conflict with the theoretical expectation that b:a and c:a ratios should have universal values of 0.67 and 0.45, respectively, for particles produced by mechanical weathering (Domokos et al., 2015). This suggests that lithology-specific values for initial shape may be needed when using shape to infer distance traveled from sediment sources (Miller et al., 2014; Szabo et al., 2015; Novak-Szabo et al., 2018), particularly for lithologies that have foliation and other anisotropic properties.

4 Discussion

The close correlations between talus size and fracture spacing distributions at our sites (Figs. 3–5) suggest that particles are detached from nearly the full network of fractures exposed on the cliff faces and do not undergo much size reduction due to

Deleted: ,

Deleted: ,

Deleted: ,

Deleted: We quantified shape at

Deleted:

Deleted: Fig. 6, after

Deleted: has lower mean

Deleted: is

Deleted: indicating

Deleted: environment. Of the six possible comparisons, only one—between the granodiorite and basalt—had no statistically significant differences in either axis ratio. In the three remaining comparisons, metasediment has a lower mean b:a ratio than andesite; metasediment also has a lower mean b:a and a:c than the basalt; and finally the andesite has a lower mean c:a ratio than the granodiorite. Given the overlap in mean shape between the granodiorite and the basalt, which differ in mean particle size by up to a factor of 10 (Fig. 4), there is no evidence in our data that initial particle shape varies with size, contrary to the prediction from previous work that smaller particles should be more block-like on average (Domokos et al., 2015)....

Deleted: prediction

Deleted: the

Deleted: ratio

Deleted: a single average value

Deleted: ,

Deleted: correlation

Deleted: suggests

Deleted: 9

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

physical or chemical weathering in talus deposits. This finding, while limited to our sites, is robust across a wide range of lithologies and weathering conditions, suggesting that it spans a range of processes that could lead to particle detachment and subsequent weathering in talus deposits, including subcritical cracking and segregation ice growth.

4.1 Predicting initial sizes from fracture spacing

Of the three particle dimensions measured here, the distribution of b-axis diameters most closely matches the fracture spacing distributions (Figs. 3–5). This suggests that fracture spacing measurements can be used to predict the initial size distribution of intermediate particle diameters. This is useful because the b-axis diameter is the most characteristic linear measure of particle volume and is therefore commonly used to represent particle mass in sediment transport theory and applications (Bagnold, 1966). Nevertheless, across our sites the b-axis is systematically $\sim\frac{1}{2}\phi$ interval larger on average than the median fracture spacing (Fig 4b), raising the question of whether this reflects a bias in our methods or incomplete exploitation of fractures during detachment from the cliff faces.

There are three potential sources of bias in our methods. First, the systematically coarser b-axis diameters may be partly driven by vertical sorting that causes fine particles to be underrepresented in point counts conducted on talus slope surfaces. However, we find no evidence of this in our measurements of sediment extracted from openings between surface particles: at each site where we made these measurements, the size distributions of the bulk samples overlap with the fine tail of the talus distribution enough that they could be combined using established techniques (Bunte and Abt, 2001) into a single continuous distribution in which the median is equal to the median of the distribution of talus at the surface. This suggests that analysis of the talus at the surface provides unbiased estimates of the size distributions of material shed from cliff faces at our sites.

Another possible explanation for the systematic offset in b-axis size may stem from limitations inherent in one-dimensional measures of size to characterize three-dimensional objects. Fracture spacing is an indirect measure of latent block volume, and axis diameter is an indirect measure for talus volume. One alternate approach, used by Messenzehl et al. (2018), is to estimate talus particle volumes from linear measurements of talus axes and characterize fracture density in joints per cubic meter using scanline measurements of fracture spacing. However, this approach does not eliminate the offset when we apply it to our data. Thus, we conclude that using linear rather than volumetric metrics is an unlikely source of the difference between median b-axis and fracture-spacing data.

A third possible methodological explanation stems from the random orientation of the scan lines relative to the joint sets exposed on the rock face. If a scan line traces diagonal transects across a set of prismatic rectangular blocks, where the latent a- and b- axes are exposed and the c-axes extend into the rock mass, then the measured fracture spacings would be systematically larger than the typical b-axis, by a factor that depends on the angle between the scan line and the joint set making up the b-axis fractures. However, if conversely the a-axes extend into the rock mass, then the scan line measurements would

Deleted: phi

Deleted:); only two in 15 sites have a median b-axis diameter less than

Deleted: median fracture spacing, which is unlikely to arise by chance ($p = 0.0064$). The positive deviations shown in Figure 4 contrast with previous measurements

Deleted: other mountain landscapes in California, where sediment sizes also correlate strongly with—but

Deleted: systematically finer than—fracture spacings in the source bedrock (Neely and DiBiasi, 2020; Sklar et al., 2020). At our sites

Deleted: , in part,

Deleted: sufficiently

Deleted: can

Deleted: Hence, we interpret the positive offset in b-axis diameters (Fig. 4b) to reflect incomplete use of exposed fractures during detachment of talus from cliff faces. These unused fractures are presumably contained within the talus blocks in the deposit and may be exploited during later size reduction by physical and chemical weathering....

Deleted: 10*

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

instead underestimate the b-axis spacings. In the most general case, blocks are formed by three intersecting joint sets with non-perpendicular orientations, and thus are not rectangular prisms. In this case, the inter-fracture distance measured along any given scan line crossing a block could possibly range from near zero (near the tip of an acute-angled point) to greater than the a-axis length (for example if spanning the longest possible linear distance across a rectangular block face). An essential assumption in using the scan-line technique is that this variability can be overcome by a large sample size, resulting in an accurate if imprecise estimate of the central tendency and spread in the underlying population of fracture spacings (Priest and Hudson, 1976; Roy et al., 2014). The close correspondence being particle sizes and spacings in all aspects of the distributions except axis offset suggests that our sample sizes are large enough to overcome any scanline biases.

The fourth explanation is not related to measurement technique, but to the mechanics of block detachment. At many sites we observe talus blocks that contain fractures that were not exploited by the detachment process (Fig. 2d, 2e). This could at least partly explain the offset between measured median b-axis diameters and median fracture spacing, with detachment favoring longer, more persistent fractures exposed on the cliff face. The effect appears to be most prominent at MT39, where even the c axes are coarser than the fracture spacing overall (Fig. 3b). We cannot rule out the possibility that some of the unexploited fractures were created or extended by stresses during or after detachment from the cliff and deposition on the talus slope. In any case, these unused fractures can be exploited during later size reduction by physical and chemical weathering. Based on our observations, we conclude that incomplete exploitation of fractures provides the best explanation for why b-axis diameters are systematically greater than fracture spacings across our sites.

The finding that the b-axes are systematically greater than the fracture spacings contrasts with previous measurements from other mountain landscapes in California, where sediment sizes also correlate strongly with—but are systematically finer than—fracture spacings in the source bedrock (Neely and DiBiase, 2020; Sklar et al., 2020). In the Inyo Creek catchment, for example, the fine mode of the bimodal sediment size distribution on hillslopes is controlled by weathering processes that vary independently of fracture spacings (Sklar et al., 2020). The coarse mode is also smaller than the fracture spacings, potentially reflecting limited resolution of the photo-based fracture spacing measurements. This may also help explain why particles are also smaller on average than fracture spacings at the San Gabriel and San Jacinto mountains sites, where photos were used to quantify fracture spacings (Neely and DiBiase, 2020). However, variations in particle size across these sites can also be explained by differences in weathering that are driven by variations in the fractional coverage of regolith. Hence, the Inyo Creek, San Gabriel, and San Jacinto sites are all subject to substantial weathering not experienced at the talus-cliff pairs of our sites.

4.2 Latent size versus weathering

Our analysis suggests that there are offsets but no systematic site-to-site deviations in slope from the 1:1 trend between median b-axis diameters and median fracture spacings, despite the large differences in climate and thus weathering environment (Table

Deleted: 11

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

1). Moreover, there is no significant trend in residuals relative to the 1:1 trend with either mean annual temperature or average annual precipitation. This implies that the latent size distribution (embedded within the fractures exposed on the cliff faces) dominates over in-situ weathering as the main control on the particle sizes in talus cones across our sites.

Deleted: produced on slopes

The dominance of latent size over weathering is also supported by previous analyses of correlations between fracture density and erosion rates at the Sierra Nevada sites, where talus deposit volumes have accumulated since deglaciation ~13,000 years ago (Gillespie and Zehfuss, 2004) were used to quantify cliff retreat rates (Moore et al., 2009). Higher fracture density (and thus lower fracture spacing) corresponds to faster cliff retreat rates (Table 2), because denser fractures contribute to lower rock mass strength, and thus make bedrock cliffs more susceptible to erosion (Howard and Selby, 2009; Moore et al., 2009; Glade et al., 2017; Neely et al., 2019). Thus, at these sites, where weathering is minimal and cliffs are still responding to deglaciation, fracture spacing controls both initial size and the production rate of sediment through its effects on rock mass strength. In soil mantled landscapes, in contrast, where hillslope erosion rates are set by stream incision rates, theory and observations suggest that faster erosion should generally lead to larger particle sizes due to lower regolith residence times (T_R) and thus less-extensive weathering in the critical zone (Sklar et al., 2017; Callahan et al., 2019). At our Sierra Nevada sites, and in general at other sites where $T_R \sim 0$, latent size should commonly dominate over weathering in setting initial particle size distributions.

4.3 Sediment production timescales

Our analysis of cliff retreat rates and fracture spacings from the Sierra Nevada points to another potentially insightful timescale: T_P (T), the time required to liberate a latent particle having the characteristic, median size, calculated following Eq. (1):

$$T_P = F_{50}/E \quad (1)$$

Here, F_{50} is the median fracture spacing (L), a proxy for the characteristic latent particle size, and E is the erosion rate of the fractured bedrock surface (L/T), equal to the cliff retreat rate at the Sierra Nevada sites. Application of Equation 1 to data from our sites indicates that T_P is as short as 88 years for production of one layer of 60-mm diameter latent particles at EP-26, the most rapidly eroding cliff face. At the most slowly eroding cliff face, CB-1, it took 16,500 years to produce one layer of 330-mm diameter particles, indicating that the entire post-glacial accumulation time and more was needed to detach a single layer of latent particles with the characteristic median size (Table 2). The calculated T_P at the remaining talus-cliff pairs in the Sierra Nevada sites is less than 13,000 years, consistent with the assumption in the erosion rate calculations (Moore et al., 2009) that all of the sediment was produced after the glaciers retreated.

Deleted: ranges from

Deleted: to produce a

Deleted: , to as long as

Deleted: at CB-1

Deleted: at the most slowly eroding cliff face

The relative importance of latent size and weathering can be evaluated by quantifying the ratio of T_R to T_P , as in Eq. (2):

$$T_R/T_P = \frac{H}{E} / \frac{F_{50}}{E} = H/F_{50} \quad (2)$$

Here, H is the thickness of saprolite and weathered rock (L), and the erosion rate at the top of fractured rock is assumed to be equal to the erosion rate at the base of soil (such that the thickness of saprolite and weathered rock is in steady state). At our sites, which represent an extreme end member with no saprolite or weathered bedrock (i.e., $H=0$), $T_R/T_P \ll 1$, and the latent

Deleted: 12

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

size distribution dominates over weathering in setting initial particle size. Although uncommon, this extreme may occur in diverse settings, including arid landscapes, glacial valleys, steep rapidly eroding mountains, and over the scale of individual hillslopes in soil mantled landscapes where bedrock exposures (and thus areas with $T_R = 0$) are patchy. In contrast, where soil, saprolite, and/or weathered bedrock is both thick and spatially extensive, weathering may dominate over latent size distributions. This can produce the other extreme, $T_R/T_P \gg 1$, particularly where erosion is also slow and fractures are closely spaced.

In between the two extremes, we envision a spectrum in the relative importance of weathering and latent size as a function of T_R/T_P . This spectrum is illustrated conceptually in Figure 7 for three cases with the same slope and erosion rate: Increasing saprolite and weathered bedrock thickness and decreasing fracture spacing should lead to higher T_R/T_P ratios (from left to right in Fig. 7), which in turn would correspond to finer initial sediment size distributions produced at the top of fractured bedrock or saprolite (cf. Fig. 7a, b, and c). Fig. 7a depicts a case at the transition from bare bedrock (e.g., the cliff faces studied here), to slopes with patchy soil cover, such as those observed at the other California sites where fracture spacing and sediment size have been quantified (Neely and DiBiase, 2020; Sklar et al., 2020). Such sites should have enhanced potential for weathering relative to our sites. This might help explain why the median b-axis diameters plot higher than median fracture spacings at our sites and vice versa at the other sites in California. In landscapes that are completely covered with regolith and weathered rock (Fig. 7b–c), the signal of the latent size distribution (and also of initial shape; Fig. 6) may fade before liberation of sediment into the soil. Thus, initial size should be dominated by weathering as residence times increase and thereby increase in exposure to chemical and physical weathering (Fig. 7c).

4.4 Implications for future work

Our results from talus-cliff pairs show that, where bedrock is exposed, the latent sediment size observed in fracture spacing distributions can be used to predict initial sediment size. Latent size should also influence initial sediment size in soil-mantled landscapes, in combination with the influences of in-situ physical and chemical weathering, except in the extreme condition of long and/or intense exposure to weathering. This suggests that more widespread fracture spacing measurements would contribute improved understanding of controls on the evolution of sediment size in landscapes. However, these measurements are difficult to obtain. In soil-mantled landscapes, direct measurements of fractures in bedrock can often be made where relatively unweathered bedrock is exposed in roadcuts or in outcrops, but these measurements may be biased if outcrops are not representative of nearby soil-mantled rock. Fracture spacing has also frequently been quantified from boreholes and cores, but have only rarely been used in critical zone studies (e.g., Holbrook et al., 2019) because they have mostly been obtained for different purposes (e.g., oil and gas exploration). Geophysical techniques can also be used to characterize subsurface fracture density (e.g., St Clair et al., 2015), and when calibrated by direct observations from cores and outcrops (Flinchum et al., 2018; Callahan et al., 2020), may provide estimates of absolute fracture spacing (Parsekian et al., 2015). Fracture density can also be estimated from rock mechanical models (Shen et al., 2019) and based on analysis of topographic and regional stresses (Slim

Deleted: when

Deleted: at the other endmember,

Deleted: , saprolite and/or weathered bedrock is thick,

Deleted: endmembers

Deleted: DiBiasei

Deleted: 13*

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

et al., 2015; Moon et al., 2020). These techniques will be central to quantifying the interplay of fracturing and weathering in determining initial sediment size using our time-scale-based conceptual framework.

Our finding of a strong association between rock type and fracture spacing, and thus initial sediment size, also suggests a potentially fruitful avenue for future work. Our study was not designed to systematically test for the influence of rock type on fracture spacing, but the topic has spawned a large literature that might be mined for general relationships relevant for predicting hillslope sediment size (Narr and Suppe, 1991; Lore et al., 2001; Manda et al., 2008). Overall, our finding of a strong link between fracture spacing and initial sediment size for the end-member case of talus-cliff pairs strengthens the foundation for future field and modeling studies of the influence of lithology, tectonics, and climate on landscape-scale variations in hillslope sediment size.

5 Conclusions

The detachment of rock fragments from fractured bedrock on hillslopes creates sediment with initial size distributions that set the upper limits on particle size for all subsequent stages in the evolution of sediment as it is exposed to chemical and physical weathering during transport from source to sink. The initial size distribution should depend on three main factors: the size distribution of latent sediment (i.e., blocks defined by throughgoing fractures); weathering that occurs in-situ in fractured bedrock before the sediment is detached; and weathering and during the detachment process (e.g., disintegration along crystal grain boundaries). However, the initial size distribution is difficult to measure, because the interface across which sediment is produced is often shielded from view by overlying soil. Talus deposits that have accumulated beneath cliff faces offer opportunities to test the hypothesis that, when in-situ weathering is minimal, the initial size distribution should strongly reflect the latent size distribution defined by fractures on the cliff faces.

Here, we presented measurements of fracture spacing and particle size distributions from talus-cliff pairs spanning a wide range of climates and lithologies in California. Median fracture spacing varies by a factor of 40, median particle size varies by a factor of 60, and both of these variables correlate strongly with lithology. In addition, fracture spacing and talus size distributions are closely correlated with each other in central tendency, spread, and distribution shape, with b-axis diameters showing the closest correspondence with the fracture spacing at most sites. This suggests that weathering has not modified latent sediment, either before or during detachment from the cliff face. In addition, talus has not undergone much weathering after deposition and is slightly coarser than the latent sizes implied by the fractures, because the talus contains unexploited fractures inherited from the cliff face. Where detachment processes leave many of these unexploited fractures, initial sediment size and fracture spacing may not correlate as well as they do across our sites, even where in-situ weathering is minimal (e.g., Messenzehl et al., 2018). The positive offset in b-axis diameter relative to fracture spacing at all but 2 of our talus-cliff pairs differs from previous work elsewhere in California, where b-axis diameters are systematically finer than bedrock fracture

Deleted: life

Deleted: We hypothesize that the

Deleted: two

Deleted: and

Deleted: both

Deleted: an opportunity

Deleted: suggesting that

Deleted: These results differ

Deleted: 14

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

spacings (Neely and DiBiase, 2020; Sklar et al., 2020), likely due to post-detachment weathering and possible biases in photo-based fracture spacing measurements. Together, these observations support a new conceptual framework illustrating the relative importance of latent size distributions and weathering on the initial sediment size distribution in mountain landscapes. In this framework, hillslopes occupy a spectrum defined by the ratio of two characteristic timescales: the residence time in saprolite and weathered bedrock, and the time required to detach the characteristic particle size. Where weathering residence times are negligible, as at our 15 talus-cliff pairs, the latent size distribution can be used to predict the initial size distribution. At the other end of the spectrum, where weathering residence times are long, the latent size distribution will provide limited predictive information about initial sediment size distributions.

Deleted: ,

Deleted: in the patchy soil deposits where particles were sampled (Neely and DiBiase, 2020; Sklar et al., 2020).

Deleted: setting

Author Contributions

JV, LS, and JM designed the field investigation and JV and LS carried it out. JV, LS, and CR analyzed the data. LS and CR prepared the manuscript with contributions from all co-authors.

Competing interests

The authors declare that they have no conflict of interest

Acknowledgments

The authors thank Nolen Brown, Navek Ceja, Van Jackson-Weaver, Bradley Penner, Paul Vawter, and Dan Rosenburg for assistance in the field; Russell Callahan for assistance with graphic design in Figure 1; John Caskey and Zan Stine for helpful discussions; and two anonymous reviewers whose comments helped improve the manuscript. Funded by the National Science Foundation grants EAR 1324830 to Sklar and EAR 2012357 to Riebe. Additional support for Verdian and Sklar provided by the Doris and David Dawdy Fund for Hydrological Science at San Francisco State University.

Deleted: and

Deleted: .

Deleted: 1325033

References

Ague, J. J.: Thermodynamic calculation of emplacement pressures for batholithic rocks, California: Implications for the aluminum-in-hornblende barometer. *Geology*, 25, 6, 563-566, 1997.

Armin, R. A., John D. A., Moore, W. J. and Dohrenwend, J. C.: Geologic map of the Markleeville 15-minute Quadrangle, Alpine County, California, U.S. Geological Survey, Miscellaneous Investigations Series Map I-1474, doi: 10.3133/i1474, 1984.

Deleted: 15

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

Attal, M. and Lavé, J.: Pebble abrasion during fluvial transport: Experimental results and implications for the evolution of the sediment load along rivers, Journal of Geophysical Research, Earth Surface, 114, F04023, doi:10.1029/2009JF001328, 2009.

Bagnold, R. A.: An approach to the sediment transport problem from general physics, USGS Professional Paper 422-I, US government printing office, Washington DC, 1966.

Bateman, P. C.: Geologic map of the Tuolumne Meadows quadrangle, Yosemite National Park, California, U.S. Geological Survey, Miscellaneous Investigations Series Map I-1474, doi: 10.3133/gq1570, 1983.

Bateman, P. C., Wahrhaftig, C. and Bailey, E. H.: Geology of the Sierra Nevada, California Division of Mines and Geology Bulletin 190 (1966): 107-172.

Brantley, S. L., Buss, H., Lebedeva, M., Fletcher, R. C., and Ma, L.: Investigating the complex interface where bedrock transforms to regolith, Applied Geochemistry, 26, S12-S15, doi:10.1016/j.apgeochem.2011.03.017, 2011.

Bunte, K., and Abt, S. R.: Sampling surface and subsurface particle-size distributions in wadable gravel-and cobble-bed streams for analyses in sediment transport, hydraulics, and streambed monitoring, Gen. Tech. Rep. RMRS-GTR-74. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research Station. 428 p., 2001.

Callahan, R. P., Ferrier, K. L., Dixon, J., Dosseto, A., Hahn, W. J., Jessup, B. S., Miller, S. N., Hunsaker, C. T., Johnson, D. W., Sklar, L. S., and Riebe, C. S.: Arrested development: Erosional equilibrium in the southern Sierra Nevada, California, maintained by feedbacks between channel incision and hillslope sediment production, GSA Bulletin, 131, 7-8, 1179-1202, doi:10.1130/B35006.1, 2019.

Callahan, R. P., Riebe, C. S., Pasquet, S., Ferrier, K. L., Grana, D., Sklar, L. S., Taylor, N. J., Flinchum, B. A., Hayes, J. L., Carr, B. J. and Hartsough, P. C.: Subsurface weathering revealed in hillslope-integrated porosity distributions, Geophysical Research Letters, 47, 15, doi:10.1029/2020GL088322, 2020.

DiBiase, R. A., Rossi, M. W. and Neely, A. B.: Fracture density and grain size controls on the relief structure of bedrock landscapes, Geology, 46, 5, 399-402, doi:10.1130/G40006.1, 2018.

Dingle, E. H., Attal, M., and Sinclair, H. D.: Abrasion-set limits on Himalayan gravel flux: Nature, 544, 471-474, doi:10.1038/nature22039n, 2017.

Domokos, G., Kun, F., Sipos, A. A. and Szabó, T.: Universality of fragment shapes, Scientific Reports, 5, 9147, doi:10.1038/srep09147, 2015.

Eppes, M. C., and Keanini, R.: Mechanical weathering and rock erosion by climate-dependent subcritical cracking, Reviews of Geophysics, 55, 2, 470-508, doi:10.1002/2017RG000557, 2017.

Eppes, M. C., Hancock, G. S., Chen, X., Arey, J., Dewers, T., Huettnermoser, J., Kiessling, S., Moser, F., Tannu, N., Weiserbs, B. and Whitten, J.: Rates of subcritical cracking and long-term rock erosion, Geology, 46, 11, 951-954, doi:10.1130/G45256.1, 2018.

Fletcher, R. C., and Brantley, S. L.: Reduction of bedrock blocks as corestones in the weathering profile: observations and model, American Journal of Science, 310, 3, 131-164, doi:10.2475/03.2010.01, 2010.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Font color: Black

Deleted: doi:10.1016/j.apgeochem.2011.03.017

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Deleted: reports

Formatted: Font: Times New Roman, Font color: Black

Deleted: p.

Formatted: Font: Times New Roman, Font color: Black

Deleted: doi:10.1038/srep09147

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Field Code Changed

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman

Deleted: 16

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

Flinchum, B. A., Holbrook, W. S., Rempe, D., Moon, S., Riebe, C. S., Carr, B. C., Hayes, J. L., St. Clair, J., and Peters, M. P.: Critical zone structure under a granite ridge inferred from drilling and three-dimensional seismic refraction data, *Journal of Geophysical Research: Earth Surface* 123, 6, 1317-1343, doi:10.1029/2017JF004280, 2018.

Gillespie, P. A., Howard, C. B., Walsh, J. J., and Watterson, J.: Measurement and characterisation of spatial distributions of fractures, *Tectonophysics*, 226, 1-4, 113-141, doi:10.1016/0040-1951(93)90114-Y, 1993.

Gillespie, A. R., and Zehfuss, P. H.: Glaciations of the Sierra Nevada, California, USA, in *Quaternary Glaciations - Extent and Chronology*, Ehlers, J. and Gibbard, P. L., eds, *Developments in Quaternary Sciences*, 2, 51-62, Elsevier, 2004.

Goodfellow, B. W., Hilley, G. E., Webb, S. M., Sklar, L. S., Moon, S., and Olson, C. A.: The chemical, mechanical, and hydrological evolution of weathering granitoid, *Journal of Geophysical Research: Earth Surface*, 121, 8, 1410-1435, doi:10.1002/2016JF003822, 2016.

Glade, R. C., Anderson, R. S., and Tucker, G. E.: Block-controlled hillslope form and persistence of topography in rocky landscapes, *Geology*, 45, 4, 311-314, doi:10.1130/G38665.1, 2017.

Graham, D. J. and Midgley, N. G.: Graphical representation of particle shape using triangular diagrams: an Excel spreadsheet method, *Earth Surface Processes and Landforms*, 25, 13, 1473-1477, doi:10.1002/1096-9837(200012)25:13<1473::AID-ESP158>3.0.CO;2-C, 2000.

Graymer, R. W., Moring, B. C., Saucedo, G. J., Wentworth, C. M., Brabb, E. E. and Knudsen, K. L.: *Geologic map of the San Francisco Bay region*, US Department of the Interior, US Geological Survey, 2006.

Holbrook, W. S., Marcon, V., Bacon, A. R., Brantley, S. L., Carr, B. J., Flinchum, B. A., Richter, D. D. and Riebe, C. S.: Links between physical and chemical weathering inferred from a 65-m-deep borehole through Earth's critical zone, *Scientific Reports*, 9, 1, 1-11, doi:10.1038/s41598-019-40819-9, 2019.

Howard, A. D., and Selby, M. J.: Rock slopes, In: Parsons A. J., Abrahams A. D. (eds) *Geomorphology of Desert Environments* (pp. 189-232), Springer, Dordrecht, doi:10.1007/978-1-4020-5719-9_8, 2009.

Kim, B. H., Cai, M., Kaiser, P. K., and Yang, H. S.: Estimation of block sizes for rock masses with non-persistent joints, *Rock Mechanics and Rock Engineering*, 40, 2, 169, doi:10.1007/s00603-006-0093-8, 2007.

Kirkby, M. J. and Statham, I.: Surface stone movement and scree formation, *The Journal of Geology*, 83, 3, 349-362, doi:10.1086/628097, 1975.

Leith, K., Moore, J. R., Amann, F., and Loew, S.: In situ stress control on micro-crack generation and macroscopic extensional fracture in exhuming bedrock, *Journal of Geophysical Research*, 119, 1-22, doi: 10.1002/2012JB009801, 2014.

Litwin, K. L., Zyguelbaum, B. R., Polito, P. J., Sklar, L. S., and Collins, G. C.: Influence of temperature, composition and grain size on tensile failure of water ice: implications for erosion on Titan, *Journal of Geophysical Research – Planets*; doi:10.1029/2012JE004101, 2012.

Lore, J., Aydin, A. and Goodson, K.: A deterministic methodology for prediction of fracture distribution in basaltic multiflows, *Journal of Geophysical Research: Solid Earth*, 106, B4, 6447-6459, doi:10.1029/2000JB900407, 2001.

- Formatted: Font: Times New Roman
- Formatted: Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman, Font color: Black
- Field Code Changed
- Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman
- Field Code Changed
- Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman
- Formatted: Font: Times New Roman, Font color: Black
- Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black
- Field Code Changed
- Formatted: Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman
- Formatted: Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman
- Field Code Changed
- Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman, Font color: Black
- Formatted: Font: Times New Roman
- Formatted: Font: Times New Roman, Font color: Black
- Deleted: doi:10.1029/2000JB900407
- Formatted: Font: Times New Roman, Font color: Black
- Deleted: 17
- Formatted: Font: 10 pt, Font color: Black
- Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

Maher, K.: The dependence of chemical weathering rates on fluid residence time, *Earth and Planetary Science Letters*, 294, 1-2, 101-110, doi:10.1016/j.epsl.2010.03.010, 2010.

Manda, A. K., Mabee, S. B., and Wise, D. U.: Influence of rock fabric on fracture attribute distribution and implications for groundwater flow in the Nashoba Terrane, eastern Massachusetts, *Journal of Structural Geology*, 30, 4, 464-477, doi:10.1016/j.jsg.2007.12.006, 2008.

Marshall, J. A. and Roering, J. J.: Diagenetic variation in the Oregon Coast Range: Implications for rock strength, soil production, hillslope form, and landscape evolution, *Journal of Geophysical Research Earth Surface*, 19(6), 1395-1417, doi:10.1002/2013JF003004, 2014.

Messenzehl, K., Viles, H., Otto, J. C., Ewald, A., and Dikau, R.: Linking rock weathering, rockwall instability and rockfall supply on talus slopes in glaciated hanging valleys (Swiss Alps), *Permafrost and Periglacial Processes*, 29, 3, 135-151, doi:10.1002/ppp.1976, 2018.

Miller, K. L., Szabó, T., Jerolmack, D. J., and Domokos, G.: Quantifying the significance of abrasion and selective transport for downstream fluvial grain size evolution, *Journal of Geophysical Research: Earth Surface*, 119, 11, 2412-2429, doi:10.1002/2014JF003156, 2014.

Molnar, P., Anderson, R. S., and Anderson, S. P.: Tectonics, fracturing of rock, and erosion, *Journal of Geophysical Research: Earth Surface*, 112, F3, doi:10.1029/2005JF000433, 2007.

Moon, S., Perron, J. T., Martel, S. J., Holbrook, W. S., and St. Clair, J.: A model of three-dimensional topographic stresses with implications for bedrock fractures, surface processes, and landscape evolution, *Journal of Geophysical Research: Earth Surface*, 122, 4, 823-846, doi:10.1002/2016JF004155, 2017.

Moon, S., Perron, J. T., Martel, S. J., Goodfellow, B. W., Ivars, D. M., Hall, A., Heyman, Munier, R., Näslund, J. O., Simeonov, A. and Stroeven, A. P.: Present-day stress field influences bedrock fracture openness deep into the subsurface, *Geophysical Research Letters* 47, 23, doi:10.1029/2020GL090581, 2020.

Moore, J. R., Sanders, J. W., Dietrich, W. E. and Glaser, S. D.: Influence of rock mass strength on the erosion rate of alpine cliffs, *Earth Surface Processes and Landforms*, 34, 10, 1339-1352, doi:10.1002/esp.1821, 2009.

Nakamura, A. M., Michel, P. and Setoh, M.: Weibull parameters of Yakuno basalt targets used in documented high-velocity impact experiments, *J. Geophys. Res.*, 112, E02001, doi:10.1029/2006JE002757, (2007).

Narr, W., and Suppe, J.: Joint spacing in sedimentary rocks, *Journal of Structural Geology*, 13, 9, 1037-1048, doi:10.1016/0191-8141(91)90055-N, 1991.

Neely, A. B., DiBiase, R. A., Corbett, L. B., Bierman, P. R. and Caffee, M. W.: Bedrock fracture density controls on hillslope erodibility in steep, rocky landscapes with patchy soil cover, southern California, USA, *Earth and Planetary Science Letters*, 522, 186-197, doi:10.1016/j.epsl.2019.06.011, 2019.

Neely, A. B., and DiBiase, R. A.: Drainage area, bedrock fracture spacing, and weathering controls on landscape-scale patterns in surface sediment grain size, *Journal of Geophysical Research: Earth Surface* 125, 10, doi:10.1029/2020JF005560, 2020.

Formatted ... [1]

Deleted: doi:10.1016/j.epsl.2010.03.010

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman

Field Code Changed

Formatted ... [2]

Formatted: Font: Times New Roman

Deleted: doi:10.1002/ppp.1976

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman

Deleted: doi:10.1002/2014JF003156

Formatted: Font: Times New Roman, Font color: Black

Formatted ... [3]

Deleted: doi:10.1029/2005JF000433

Formatted: Font: Times New Roman, Font color: Black

Formatted ... [4]

Field Code Changed

Formatted ... [5]

Formatted ... [6]

Field Code Changed

Formatted ... [7]

Formatted: Font: Times New Roman

Field Code Changed

Formatted ... [8]

Formatted: Font: Times New Roman

Deleted: . and

Formatted: Font: Times New Roman

Deleted: -

Formatted: Font: Times New Roman

Formatted ... [9]

Deleted: ESSOAr [preprint], doi:10.1002/essoar.10502617.1, 11 June

Formatted ... [10]

Deleted: 18*

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

Novak-Szabo, T., Sipos, A. Á., Shaw, S., Bertoni, D., Pozzebon, A., Grotoli, E., Sarti, G., Ciavola, P., Domokos, G. and Jerolmack, D. J.: Universal characteristics of particle shape evolution by bed-load chipping, *Science advances*, 4,3, doi:10.1126/sciadv.aao4946, 2018.

Formatted: Font: Times New Roman

Overstreet, B. T., Riebe, C. S., Wooster, J. K., Sklar, L. S., and Bellugi, D.: Tools for gauging the capacity of salmon spawning substrates, *Earth Surface Processes and Landforms*, 41, 1, 130-142, doi:10.1002/esp.3831, 2016.

Formatted: Font: Times New Roman, Font color: Black

Parsekian, A. D., Singha, K., Minsley, B. J., Holbrook, W. S., and Slater, L.: Multiscale geophysical imaging of the critical zone, *Reviews of Geophysics*, 53, 1, 1-26, doi:10.1002/2014RG000465, 2015.

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Field Code Changed

Priest, S. D., and Hudson, J. A.: Discontinuity spacings in rock, *International Journal of Rock Mechanics and Mining Sciences & Geomechanics Abstracts*, 13, 5, 135-148, doi:10.1016/0148-9062(76)90818-4, 1976.

Formatted: Font: Times New Roman

Prism Climate Group, Oregon State University, <http://prism.oregonstate.edu>, last access: 11 December, 2019.

Formatted: Font: Times New Roman, Font color: Black

Riebe, C. S., Sklar, L. S., Overstreet, B. T., and Wooster, J. K.: Optimal reproduction in salmon spawning substrates linked to grain size and fish length, *Water Resources Research*, 50, 2, 898-918, doi:10.1002/2013WR014231, 2014.

Field Code Changed

Riebe, C. S., Hahm, W. J., and Brantley, S. L.: Controls on deep critical zone architecture: A historical review and four testable hypotheses, *Earth Surface Processes and Landforms*, 42, 1, 128-156, doi:10.1002/esp.4052, 2017.

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Riebe, C. S., Callahan, R. P., Granke, S. B.-M., Carr, B. J., Hayes, J. L., Schell, M. S., and Sklar, L. S.: Anisovolumetric weathering in granitic saprolite controlled by climate and erosion rates, *Geology*, 49, doi:10.1130/G48191.1, 2021.

Roda-Boluda, D. C., D'Arcy, M., McDonald, J., and Whittaker, A. C.: Lithological controls on hillslope sediment supply: insights from landslide activity and grain size distributions, *Earth Surface Processes and Landforms*, 43, 5, 956-977, <https://doi.org/10.1002/esp.4281>, 2018.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Field Code Changed

Roy, A., Perfect, E., Dunne, W. M., and McKay, L. D.: A technique for revealing scale-dependent patterns in fracture spacing data, *Journal of Geophysical Research: Solid Earth*, 119, 7, 5979-5986, doi:10.1002/2013JB010647, 2014.

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Saucedo, G. J.: Geologic map of the Lake Tahoe basin, California and Nevada, California Geological Survey, Regional Geological Map Series, Map 4, 2005.

Shen, X., Arson, C., Ferrier, K. L., West, N. and Dai, S., Mineral weathering and bedrock weakening: Modeling microscale bedrock damage under biotite weathering, *Journal of Geophysical Research: Earth Surface*, 124, 11, 2623-2646, doi:10.1029/2019JF005068, 2019.

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Font color: Black

Field Code Changed

Shobe, C. M., Tucker, G. E., and Anderson, R. S.: Hillslope-derived blocks retard river incision, *Geophysical Research Letters*, 43, 10, 5070-5078, doi:10.1002/2016GL069262, 2016.

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Sklar, L. S., and Dietrich, W. E.: A mechanistic model for river incision into bedrock by saltating bed load, *Water Resources Research*, 40, 6, doi:10.1029/2003WR002496, 2004.

Deleted: 19

Formatted: Font: 10 pt, Font color: Black

Sklar, L. S., Dietrich, W. E., Fofoula-Georgiou, E., Lashermes, B. and Bellugi, D.: Do gravel bed river size distributions record channel network structure? *Water Resources Research*, 42, 6, W06D18, doi:10.1029/2006WR005035, 2006.

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

Sklar, L. S., Riebe, C. S., Marshall, J. A., Genetti, J., Leclere, S., Lukens, C. L. and Mercers, V.: The problem of predicting the size distribution of sediment supplied by hillslopes to rivers, *Geomorphology*, 277, 31-49, doi:10.1016/j.geomorph.2016.05.005, 2017.

Sklar, L. S., Riebe, C. S., Genetti, J., Leclere, S. and Lukens, C. E.: Downvalley fining of hillslope sediment in an alpine catchment: implications for downstream fining of sediment flux in mountain rivers, *Earth Surface Processes and Landforms*, 45, 8, 1828-1845, doi:10.1002/esp.4849, 2020.

Slim, M., Perron, J. T., Martel, S. J., and Singha, K.: Topographic stress and rock fracture: A two-dimensional numerical model for arbitrary topography and preliminary comparison with borehole observations, *Earth Surface Processes and Landforms*, 40, 4, 512-529, doi:10.1002/esp.3646, 2015.

Sneed, E. D. and Folk, R. L.: Pebbles in the lower Colorado River, Texas a study in particle morphogenesis, *The Journal of Geology*, 66, 2, 114-150, doi:10.1086/626490, 1958.

Clair, J. S., Moon, S., Holbrook, W. S., Perron, J. T., Riebe, C. S., Martel, S. J., Carr, B., Harman, C., Singha, K. D. and Richter, D. D.: Geophysical imaging reveals topographic stress control of bedrock weathering, *Science*, 350, 6260, 534-538, doi:10.1126/science.aab2210, 2015.

Szabó, T., Domokos, G., Grotzinger, J. P., and Jerolmack, D. J.: Reconstructing the transport history of pebbles on Mars, *Nature Communications*, 6, 1, doi:10.1038/ncomms9366, 2015.

Turowski, J. M., Wyss, C. R., and Beer, A. R.: Grain size effects on energy delivery to the streambed and links to bedrock erosion, *Geophysical Research Letters*, 42, 6, 1775-1780, doi:10.1002/2015GL063159, 2015.

Wahrhaftig, C.: Stepped topography of the southern Sierra Nevada, California, *Geological Society of America Bulletin*, 76, 10, 1165-1190, doi:10.1130/0016-7606(1965)76[1165:STOTSS]2.0.CO;2, 1965.

Weibull, W.: A statistical distribution function of wide applicability, *Journal of Applied Mechanics*, 103, 33, 1951.

Formatted: Font: Times New Roman

Deleted: doi:10.1016/j.geomorph.2016.05.005

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Field Code Changed

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Font color: Black

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Field Code Changed

Formatted: Font: Times New Roman

Deleted: communications

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Font color: Black

Deleted: doi:10.1002/2015GL063159

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Font color: Black

Field Code Changed

Formatted: Default Paragraph Font, Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman, Font color: Black

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman, Not Bold, Font color: Auto

Formatted: Font: Times New Roman, 12 pt, Not Bold, No underline

Formatted: Font: Times New Roman

Deleted: 20

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right

Table 1. Study sites in Sierra Nevada and San Francisco Bay Area

Location	Pair	Lithology	Latitude (°N)	Longitude (°W)	Elevation (m)	MAT ¹ (°C)	MAP ² (mm)
Conness Basin	CB-1	Granodiorite	37.9763	119.3082	3262	1.3	1190
Conness Basin	CB-2	Granodiorite	37.9750	119.3033	3293	1.1	1226
Conness Basin	CB-3	Granodiorite	37.9797	119.3006	3171	1.5	1179
Conness Basin	CB-5	Metasediment	37.9928	119.2870	3140	1.3	1152
Ebbetts Pass	EP-24	Andesite	38.5655	119.8084	2530	4.8	1343
Ebbetts Pass	EP-25	Andesite	38.5665	119.8114	2549	4.8	1343
Ebbetts Pass	EP-26	Basalt	38.5473	119.8136	2732	3.9	1455
Ebbetts Pass	EP-28	Basalt	38.5483	119.8144	2744	3.9	1455
Mount Tallac	MT-38	Granodiorite	38.9430	120.1235	2134	6.5	1481
Mount Tallac	MT-39	Granodiorite	38.9420	120.1247	2195	6.5	1481
Grizzly Peak	GP-1	Basalt	37.8903	122.2346	393	13.8	727
Twin Peaks	TP-1	Pillow basalt	37.7504	122.4483	260	13.6	705
Twin Peaks	TP-2	Pillow basalt	37.7502	122.4476	252	13.6	705
Twin Peaks	TP-3	Chert	37.7533	122.4480	280	13.6	705
Twin Peaks	TP-4	Chert	37.7533	122.4480	280	13.6	705

¹Mean annual temperature (Prism Climate Group, 2019)

²Mean annual precipitation (Prism Climate Group, 2019)

Formatted: Border: Bottom: (Single solid line, 0.5 pt Line width), Tab stops: 5.56", Left + Not at 5.51"

Formatted: Tab stops: 5.69", Decimal aligned + Not at 5.63"

Formatted: Font: 12 pt

Deleted: 21

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

Table 2. Results

Pair	Fracture spacing ¹ (mm)	a-axis diameter ¹ (mm)	b-axis diameter ¹ (mm)	c-axis diameter ¹ (mm)	Fracture geometric stdev	b-axis geometric stdev	Erosion rate (mm/yr)	T_p (yrs)	Layers removed ³
CB-1	330	375	250	130	0.346	0.357	0.02	125,000	0.8
CB-2	200	500	270	205	0.458	0.612	0.09	3,000	5.9
CB-3	280	720	420	190	0.551	0.477	0.05	8,400	2.3
CB-5	120	335	220	95	0.309	0.306	0.25	880	27
EP-24	80	225	150	55	0.283	0.255	0.31	484	50
EP-25	155	320	200	70	0.401	0.294	0.12	1,667	10
EP-26	60	95	55	25	0.285	0.201	0.68	81	150
EP-28	70	160	100	65	0.358	0.338	0.26	385	48
MT-38	390	1010	575	310	0.387	0.333	0.09	6,389	3.0
MT-39	200	850	570	280	0.411	0.352	0.14	4,071	9.1
GP-1	77	130	82	48	0.286	0.203	-	-	-
TP-1	10	-	10	-	0.251	0.328	-	-	-
TP-2	10	-	14	-	0.314	0.215	-	-	-
TP-3	24	-	27	-	0.214	0.14	-	-	-
TP-4	19	-	26	-	0.257	0.184	-	-	-

¹Fracture spacings and particle diameters are reported as medians of distributions measured in field

²Cliff retreat rates were measured by Moore et al. (2009).

³Layers removed is the number of layers of thickness equal to the median fracture spacing that have been removed since the glacier retreated and is calculated as $13,000/T_p$, where T_p is calculated according to equation 1.

Formatted: Font: Times New Roman

Formatted: Tab stops: 1.5", Decimal aligned + 2.75", Decimal aligned + 3.31", Decimal aligned + 4", Decimal aligned + 5.56", Decimal aligned + 6.06", Decimal aligned + Not at 1.56" + 2.81" + 3.38" + 4.06" + 5.63" + 6.13"

Deleted: ¶ Page Break

... [11]

Deleted: 22¶

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right



Figure 1. Study site map (upper left) and representative talus-cliff pairs from each site, with label designating lithology (red B squares = basalt; pink G circles = granodiorite; purple M diamond = metasediment; orange A triangles = andesite; purple P triangles = pillow basalt; green C triangles = chert). Scale varies between images. See Table 1 and text for site descriptions.

Deleted: 23

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right





Figure 2. Field survey methods, showing: (a) an example of transect line and scan line layout at EP-26 site in the Sierra Nevada; (b) detail of scan line at EP24 illustrating measurement of distances between fractures along scan line; (c) establishment of transect 2 at CB-5; and talus boulders containing unexploited fractures at MT-39 (d) and EP-28 (e). 2 m scale in (a) highlights a person near transect 3.



Deleted:

Deleted: (top) with

Deleted: (lower left) and Transect

Deleted: (lower right). Scale

Deleted: the top image

Deleted: Transect

Deleted: Tape in lower left is labeled in 10 cm increments.

Deleted: 24

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right




Figure 3. Particle size and fracture spacing distributions for each study site, sorted by median b-axis diameter in descending order from left to right and top to bottom. Colored lines show fracture spacing distributions, solid black lines show b-axis diameter distributions, dashed lines show a- and c-axis diameter distributions. (a-axis and c-axis diameters plot to the right of left the b-axis diameters respectively). Color codes and labels for lithology are as in Fig. 1; see Table 1 for site abbreviations. At the chert and pillow basalt sites, only b-axis diameters were measured (see text).

- Deleted: Major
- Deleted: of
- Deleted: intermediate
- Formatted: Font: 12 pt
- Deleted: 25
- Formatted: Font: 10 pt, Font color: Black
- Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right





Figure 4. Central tendency and spread of particle size and fracture spacing distributions for median a-axis diameter (a), median b-axis diameter (b), median c-axis diameter (c), and the geometric standard deviation of the b-axis diameters (d). In each case, across the wide range in particle sizes and fracture spacings represented by the different lithologies sampled here, there are strong correlations between the particle size and the fracture spacing distributions that are statistically indistinguishable from a 1:1 relationship (a–d). The correspondence is closest for the b-axis diameters (b), though they are systematically 42% larger on average than the fracture spacings.



Deleted:

Deleted: roughly parallel with...statistically indistinguishable from a 1:1 relationship (a–d). The correspondence is especially close...losest for the b-axis diameters (b), though they are systematically 42% larger on average than the fracture spacings. Symbol type and color represent lithology as in Fig. 1. ... [13]

Deleted: 26

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right





Figure 5. Comparison of distribution shape for the b-axis diameter (open symbols) and fracture spacing (colored symbols) distributions in Weibull probability space (see text). Points fall along a straight line in these plots when the sample is drawn from a population having a Weibull distribution. Best-fit linear regressions for b-axis diameters and fracture spacings are shown as dashed and colored lines, respectively. Colors correspond to lithologies following conventions in Fig. 1. For most sites, most points plot along a straight line, implying that their population distributions are consistent with a Weibull distribution. In addition, the data commonly overlap, consistent with a close match between the shape of the particle size and fracture spacing distributions at many of the sites. Examples and exceptions are highlighted in the text.



Deleted:

Deleted: ... (see text). Points fall along a straight line in these plots when the sample is drawn from a population having a Weibull distribution. Best-fit linear regressions for b-axis diameters and fracture spacings are shown as dashed and colored lines, respectively. Colors correspond to lithologies following conventions in Fig. 1. For most sites, most points plot along a straight line, implying that their population distributions are consistent with a Weibull-like

[14]

Deleted: 27

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between: (No border), Tab stops: 3.13", Centered + 6.27", Right





Figure 6. Initial particle shape at sites where all three particle diameters were measured, as revealed in ternary diagrams with blocks, rods, and slabs at vertices (inset). Although data within each site and within each lithology are widely scattered in shape, the central tendencies for samples grouped by lithology yield several statistically significant differences. For example, granodiorite has a higher b:a and c:a ratio than the metasediment (left), indicating that metasediment is more rod-like on average. Symbols and colors represent lithology following conventions of Fig. 1.



Deleted:

Deleted: Particle

Formatted: Font: 12 pt

Deleted: 28*

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right




Figure 7. Conceptual framework illustrating how initial sediment size is influenced by latent sediment size and weathering. Panels depict idealized vertical profiles of subsurface weathering where initial sediment is produced by detachment at the top of intact bedrock (a) or saprolite (b,c). Erosion rate is the same in each panel. Where fracture spacing is wide and where the weathering zone is thin, latent size should dominate over weathering (a), and vice versa where fractures are closely spaced and the weathering zone is comparatively thick (c). These examples lie on a spectrum of outcomes that correlate with the ratio of two characteristic timescales: the timescale of weathering (T_w) and the time required to detach a layer of characteristic (median) particles at the base of mobile regolith (T_d). Higher ratios correspond to a greater influence of weathering and a lesser influence of latent size on initial sediment size distributions.

Formatted: Font: 9 pt, Bold, Font color: Black

Formatted: Normal, Justified, Space After: 10 pt, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Deleted: 29

Formatted: Font: 10 pt, Font color: Black

Formatted: Normal, Justified, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border), Tab stops: 3.13", Centered + 6.27", Right

Page 18: [1] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [1] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [2] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [2] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [2] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [3] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [3] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [4] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [4] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [4] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [5] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [5] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [6] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [6] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [7] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [7] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [8] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [8] Formatted Revisions 4/30/21 3:02:00 PM

Default Paragraph Font, Font: Times New Roman, Font color: Black

Page 18: [9] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman

Page 18: [10] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman, Font color: Auto

Page 18: [10] Formatted Revisions 4/30/21 3:02:00 PM

Font: Times New Roman, Font color: Auto

Page 22: [11] Deleted Revisions 4/30/21 3:02:00 PM

Page 25: [12] Deleted Revisions 4/30/21 3:02:00 PM

Page 26: [13] Deleted Revisions 4/30/21 3:02:00 PM

Page 26: [13] Deleted Revisions 4/30/21 3:02:00 PM

Page 26: [13] Deleted Revisions 4/30/21 3:02:00 PM

Page 27: [14] Deleted Revisions 4/30/21 3:02:00 PM

Page 27: [14] Deleted Revisions 4/30/21 3:02:00 PM

Page 27: [14] Deleted Revisions 4/30/21 3:02:00 PM