

Interactions between deforestation, landscape rejuvenation, and shallow landslides in the North Tanganyika - Kivu Rift region, Africa

Arthur Depicker¹, Gerard Govers¹, Liesbet Jacobs¹, Benjamin Campforts², Judith Uwihirwe^{3,4}, and Olivier Dewitte⁵

¹KU Leuven, Department of Earth and Environmental Sciences, Division of Geography and Tourism (Celestijnenlaan 200E, 3001 Heverlee, Belgium)

²CSDMS, Institute for Arctic and Alpine Research, University of Colorado at Boulder (Boulder, CO, USA)

³University of Rwanda, Department of Soil and Water Management, Faculty of Agricultural Engineering (Street KK 737, Kigali, Rwanda)

⁴Delft University of Technology, Faculty of Civil Engineering and Geosciences, Department of Water Management (Stevinweg 1, 2628 Delft, The Netherlands)

⁵Royal Museum for Central Africa, Department of Earth Sciences (Leuvensesteenweg 13, 3080 Tervuren, Belgium)

Correspondence: Arthur Depicker (arthur.depicker@kuleuven.be)

Abstract. Deforestation is associated with a decrease in slope stability through the alteration of hydrological and geotechnical conditions. As such, deforestation increases landslide activity over short, ~~contemporary-decadal~~ timescales. However, over longer timescales (0.1-10 Ma) the location and timing of landsliding is controlled by the interaction between uplift and fluvial incision. Yet, the interaction between (human-induced) deforestation and landscape evolution has hitherto not been explicitly considered. We address this issue in the North Tanganyika-Kivu Rift region (East African Rift). In recent decades, the regional population has grown exponentially and the associated expansion of cultivated and urban land has resulted in widespread deforestation. ~~On a much longer time scale, tectonic uplift has~~ In the past 11 Ma, active continental rifting and tectonic processes have forged two parallel mountainous Rift shoulders that are continuously rejuvenated (i.e. actively incised) through knickpoint retreat, enforcing topographic steepening. In order to link deforestation and rejuvenation to landslide erosion, we compiled an inventory of nearly 8,000 recent shallow landslides in © Google Earth imagery from 2000-2019. To accurately calculate landslide erosion rates, we developed a new methodology to remediate inventory biases linked to the spatial and temporal inconsistency of this satellite imagery. Moreover, to account for the impact of rock strength on both landslide occurrence and knickpoint retreat, we limit our analysis to rock types with threshold angles of 24-28°. Rejuvenated landscapes were defined as the areas draining towards Lake Kivu or Lake Tanganyika, and downstream of retreating knickpoints. We find that erosion rates in these rejuvenated landscapes are roughly 40 % higher than in the surrounding relict landscapes. ~~This difference is due to the generally steeper relief~~ In contrast, we find that slope exerts a stronger control on landslide erosion in relict landscapes. These two results are reconciled by the observation that landslide erosion generally increases with slope gradient and that the relief is on average steeper in rejuvenated landscapes ~~which more than compensates for the fact that rejuvenated slopes, when compared to similarly angled slopes in relict zones, often display a somewhat lower landslide erosion rate. These lower-~~ The weaker effect of slope steepness on landslide erosion rates in the rejuvenated landscapes could be the result of ~~a drier climate,~~

three factors: the absence of earthquake-induced landslide events in our landslide inventory, ~~and potentially a smaller regolith stock~~ a thinner regolith mantle, and a drier climate. More frequent extreme rainfall events in the relict ~~zones~~ landscapes, and the presence of a thicker regolith, may explain a stronger landslide response to deforestation compared to rejuvenated landscapes. Overall, deforestation initiates a landslide peak that lasts approximately 15 years and increases landslide erosion by a factor 2 to 8. Eventually, landslide erosion in deforested land falls back to a level similar to that observed under forest conditions, most likely due to the depletion of the most unstable regolith. Landslides are not only more abundant in rejuvenated landscapes but are also smaller in size, which may again be a consequence of ~~the~~ a thinner regolith mantle and/or seismic activity that fractures the bedrock and reduces the minimal critical area for slope failure. With this paper, we highlight the importance of considering the geomorphological context when studying the impact of recent land use changes on landslide activity.

30 *Copyright statement.* TEXT

1 Introduction

On steep terrain, ~~shallow landslide erosion~~ the erosion caused by shallow landslides (with a maximal depth of a couple of meters) increases significantly as a result of deforestation (e.g. Montgomery et al., 2000; Mugagga et al., 2012). The removal of trees, due to either human or natural causes, decreases the slope stability through the alteration of hydrological and geotechnical conditions, such as the loss in soil cohesion due to tree root decay (Sidle et al., 2006; Sidle and Bogaard, 2016). After 10-20 years, depending on the climate and vegetation regeneration rate, this effect starts wearing off (Sidle and Bogaard, 2016). However, when forests are permanently converted to grassland or cropland, the consequences of deforestation for landsliding can last much longer or even be permanent (Sidle et al., 2006).

While these general principles are ~~now~~ well described, we do not yet fully understand the extent to which ~~deforestation increases landslide erosion in different types of landscapes~~ the response to deforestation is modulated by tectonic forcing, which typically occurs over timescales of 0.1-10 Ma (e.g. Whipple and Meade, 2006). A key distinction can be made here between actively incising, rejuvenating landscapes, in which landslides are a prime slope-limiting mechanism and ‘old’, so-called relict landscapes, where hillslopes have had a long time to adapt to river incision (Burbank et al., 1996; Larsen and Montgomery, 2012). These two landscape types can be expected to respond differently to deforestation: in rejuvenating landscapes, hillslopes are already continuously adapting to river incision through landsliding (Egholm et al., 2013). In relict landscapes, on the other hand, hillslopes will slowly become less steep and landslides occur more sporadically, potentially allowing for a thick regolith ~~cover~~ mantle to develop (Schoenbohm et al., 2004; Bennett et al., 2016). Climatic variations can also drive differences in landscape response to deforestation (Crozier, 2010), and in the context of lithologically diverse landscapes, the effect of rock strength on both knickpoint retreat and landsliding must also be acknowledged (Parker et al., 2016; Baynes et al., 2018; Campforts et al., 2020).




Figure 1. Overview of the NTK Rift and its active faults (Delvaux and Barth, 2010). Overview of the NTK Rift. a) Extent of the studied area and active faults (Delvaux and Barth, 2010). LANDSAT-8 imagery is used as background (USGS, 2018). b) The transects of the elevation at different latitudes illustrate the elevated Rift shoulders, being the result of tectonic uplift. The four most populous cities of the NTK Rift (Goma, Bukavu, Uvira, Bujumbura) are located in between the Rift shoulders.

Here, we aim to explore the interplay of deforestation and uplift-driven landscape rejuvenation on shallow landslide erosion. We focus our research on the North Tanganyika-Kivu Rift region (hereafter referred to as ‘the NTK Rift’, **Fig. 1**), part of the western branch of the East African Rift. ~~Deforestation, mainly driven by agriculture, is widespread and a result of the fast-growing population (from 89 inhabitants per km² in 1975 to 241 inhabitants per km² in 2015) (Hansen et al., 2013; JRC and CIESIN, 2006).~~ Furthermore, the The area is characterized by frequent landsliding, mostly triggered by rainfall (Monsieurs et al., 2018a; Depicker et al., 2018). ~~Furthermore, widespread deforestation and active continental rifting (Hansen et al., 2013; Saria et al., 2014; Monsieurs et al., 2018a; Depicker et al., 2018).~~

. The study area is therefore an ideal setting to evaluate how deforestation affects landslide erosion in different landscape settings.

2 The North Tanganyika - Kivu Rift region

60 Active continental rifting in our study area is driven by the divergence of the Victoria and Nubia plates that started ca. 11 Ma ago and currently continues at a rate of ca. 2 mm/yr (~~Saria et al., 2014~~)(Saria et al., 2014; Pouclet et al., 2016). Due to this setting, there is widespread seismic activity, active volcanism, and uplift initiating landscape rejuvenation through knickpoint retreat (Smets et al., 2015; Delvaux et al., 2017; Dewitte et al., 2021). Adding to the geological complexity of the NTK Rift is the wide variability in age and strength of rock formations. The majority of rocks in the northern and eastern parts of the study

65 area are of Mesoproterozoic age (1600-720 Ma), being mostly quartzites, granites, or pelites. The southwest is largely covered by either weathering-resistant quartzites or weathering-prone gneiss and micaschists of Paleoproterozoic age (2500-1600 Ma). Within the Rift shoulders, the same pattern of Meso- and Paleoproterozoic rocks is observed, save for the occurrence of much younger lithologies such as the river and lake sediments in the Ruzizi floodplain and the volcanic deposits (12 Ma - present) found around Bukavu and north of Goma (Delvaux et al., 2017; Laghmouch et al., 2018).

70 ~~Until recently, the entire study area was covered in forests~~In the natural context, prior to widespread human activity, forests covered most of the DRC and the mountainous Rift shoulders in Rwanda and Burundi, while the vegetation transitioned to woodland savanna towards the east of our study area (Ellis et al., 2010; Aleman et al., 2018; Roche and Nzabandora, 2020). Only since the beginning of the 20th century, ~~has~~ large scale deforestation has taken place, especially ~~between~~ along the Rift shoulders and in Rwanda and Burundi (Ellis et al., 2010; Aleman et al., 2018). In 2000, the study area (ca. 88,500 km²)

75 had an estimated forest coverage of 73.1 %. Between 2001 and 2018, 4.5 % of these forests were cleared, mainly for the purpose of agriculture (Hansen et al., 2013; Tyukavina et al., 2018; Musumba Teso et al., 2019). ~~The study area~~ Deforestation is therefore an ~~ideal setting to evaluate how deforestation affects landslide erosion in different landscape settings~~indirect result of the fast-growing population, which increased from 89 inhabitants per km² in 1975 to 241 inhabitants per km² in 2015 (Hansen et al., 2013; JRC and CIESIN, 2015).

80 3 Methods

In the sections below, we first focus on the landscape characteristics of the NTK Rift that can exert a control on landslide erosion: forests, rejuvenation, rainfall, and rock strength. Next, we elaborate on the different aspects of the landslide erosion assessment: the compilation of an inventory and the calculation of shallow landslide erosion rates (whether or not in the context of the previously determined landscape characteristics).




Figure 2. Environmental characterization of the NTK Rift. **a)** Tree cover for the year 2000 and deforestation from 2001-2017 (Hansen et al., 2013). **b)** Elevation and some renowned mountain peaks (USGS, 2006). **c)** Average annual rainfall between 2005-2015 (Van de Walle et al., 2020). **d)** Lithostratigraphy (Table 1) (Laghmouch et al., 2018).

85 3.1 Regional controls on landslide erosion

3.1.1 Forest cover and deforestation

We characterize the NTK Rift in terms of forest dynamics by means of the global forest data presented by Hansen et al. (2013) (Fig. 2a, the data was updated in 2018). This dataset contains a tree cover map for the year 2000 and forest loss data for the period 2001-2017, both provided at a resolution of one arc-second (ca. 30 m). The tree cover data shows the percentage of tree coverage per pixel in 2000 and the forest loss data displays discrete values between 1 and 17, indicating the year in which deforestation took place. Based on these data, we distinguish three land cover classes: i) forest, having >25 % tree cover (as suggested by Hansen et al. (2013)), ii) deforested land, and iii) non-forest land, with ≤25 % tree cover. Both deforested and non-forest land encompass land use classes such as bare land, cropland, grassland, and urban land. ~~There is no~~ Historically, current 'non-forest' land used to be either savanna grassland or forest (Roche and Nzabandora, 2020). The difference between 'non-forest' land that used to be forested in the past and 'deforested' land ,except for is the elapsed time since deforestation. ~~'Non-forest' land has suffered~~ Thus, the 'non-forest' land either underwent deforestation before the year




Figure 3. Field-validated knickpoints in Rwanda. The red arrow indicates the location of the knickpoint. **a)** East of Mabanze, Rwanda (29.469678° E, 2.047852° S). **b)** East of Kanama, Rwanda (29.391763° E, 1.705683° S). **c)** Southwest of Kibilira, Rwanda (29.61586° E, 1.980443° S; image ©2019 Google Earth).

2000, ~~while ‘deforested~~or was never forest in the first place. ‘Deforested’ land experienced deforestation over the last two decades.

3.1.2 Landscape rejuvenation

100 To ~~delineate the rejuvenated and relict landscapes~~distinguish the rejuvenated landscapes within the Rift shoulders from the surrounding landscapes (hereafter referred to as the relict landscapes), we use the spatial pattern of knickpoints ~~migrating retreating~~ upstream towards the Rift shoulders, away from the active faults. ~~Anchored-Stationary~~ knickpoints, here defined as knickpoints at a distance shorter than 1 km from a geological contact, are considered unrelated to the rejuvenation process and removed from the analysis (Kirby and Whipple, 2012; Bennett et al., 2016). Two criteria are applied to identify the rejuvenated
105 Rift: (i) the area must drain towards Lake Kivu or Lake Tanganyika, and (ii) the area must be located downstream of any

non-stationary knickpoint, unless there is no knickpoint observed in the area. In the latter case, we assume the knickpoint reached the Rift shoulder and the landscape is completely rejuvenated.

We use the KNICKPOINTFINDER function in TopoToolbox to identify knickpoints. This function requires a drainage network and tolerance value, reflecting the maximum expected error in the true river profiles (Schwanghart and Scherler, 2017). The drainage network for this purpose is modeled with the 1 arc second SRTM DEM data (USGS, 2006) and a threshold catchment area of $2 \times 10^6 \text{ m}^2$. The tolerance value is used to distinguish knickpoints from discrepancies in the longitudinal river profile that are caused by errors in the DEM data. The tolerance value is calculated as the maximal difference between the 90th and 10th quantile of the smoothed river profiles (Schwanghart and Scherler, 2017), and subsequently lowered until the algorithm identifies the three knickpoints we validated in the field (Fig. 3).

3.1.3 Rainfall

Active rifting does not only trigger landscape rejuvenation but also impacts local rainfall patterns (Van de Walle et al., 2020). ~~The majority of observed recent shallow landslides~~ Depicker et al. (2020) showed a significant link between landslide occurrence and the frequency of extreme rainfall events in the NTK Rift. At the same time, field observations and local reports confirm that the majority of recent shallow landslides are rainfall-triggered (Monsieurs et al., 2018a; Depicker et al., 2020; Dewitte et al., 2021). To explore any relationship between rejuvenation and rainfall, we analyze two metrics: the average annual rainfall and the number of times where accumulated rainfall was sufficiently large to trigger landsliding. ~~Based on the landslide initiation model of Guzzetti et al. (2008), we assume this criterion is met when there is more than 15 mm rainfall over a period of two days~~ As a proxy of the latter criterion, we use a 2 day-15 mm threshold as it is a conservative estimation for global thresholds set by Guzzetti et al. (2008). Our intent is not to approximate an actual in-situ threshold, but rather to reflect spatial patterns in intense rainfall capable of triggering landslide events. For the comparison of rainfall patterns between rejuvenated and relict landscapes, we apply the non-parametric Mann-Whitney U test, whereby each observation is the average metric (rainfall or threshold exceedance) in fifth order catchments. These units are derived from the 30 m DEM using a river catchment threshold of 10^5 m^2 .

The rainfall pattern is derived from ~~the COSMO-CLM regional climate model proposed by Van de Walle et al. (2020)~~ which a regional climate simulation with COSMO-CLM, a physical model, for the period 2005-2015 and using the ERA5 reanalysis product for the initial and boundary conditions of the atmosphere (Van de Walle et al., 2020; Hersbach et al., 2020). Due to in-situ data scarcity, evaluation of the simulated precipitation amounts is restricted to a comparison with a set of satellite products. Generally, the satellite data suggests lower amounts of rainfall compared to the simulated model output (Van de Walle et al., 2020), yet this was expected as satellite products tend to underestimate the actual precipitation (Dinku et al., 2011; Mor The final model output has a spatial resolution of 2.8 km and a temporal resolution of 1 hour (Fig. 2c). Note that the spatial resolution of the rainfall products might be too low to fully capture the impact of orographic controls and the local convective storm patterns (Monsieurs et al., 2018b).

Age	Chronostratigraphy	Lithostratigraphy	Main lithological constitution
10 Ka - present	Late Quaternary	Recent volcanics	lava, tuff, and ash, deposited in the past decades and centuries, a result of eruptions of the Nyiragongo and Nyamulagira.
2–1 Ma	Early Quaternary	Young volcanics	Relatively fresh basalts, deposited ± 2 Ma ago.
12–6 Ma	Neogene	Old volcanics	Highly weathered basalt, deposited 11–4 Ma years ago.
23 Ma - present	Late Cenozoic	Rift sediments	Sand along the lake or swamps more inland.
360–201 Ma	Karoo		Black shales, tillite, not metamorphosed.
1000–540 Ma	Neoproterozoic	Itombwe	Black shales, tillite, silicified tillite, weakly methamorphosed.
		Malagarasi	Black shales, tillite, silicified tillite, weakly methamorphosed. Presence of dolomites and volcanic rocks (basalts).
820–720 Ma	Mesoproterozoic	Alcaline complexes	Granitic rocks, intrusive volcanic rocks (rhyolite).
1375–980 Ma		Granites	two-mica and leucogranites.
1600–1000 Ma		Kivu	Pelites, quartzopelites, and quartzites at different degrees of weathering. Moderately metamorphosed.
2500–1600 Ma	Paleoproterozoic	Ruzizi and ante-Ruzizi	Gneiss and micaschists, prone to chemical weathering, and quartzites, resistant to weathering. Strongly methamorphosed.
4000–2500 Ma		Archaen	Gneiss and micaschists, prone to chemical weathering, and quartzites, resistant to weathering.

Table 1. Lithostratigraphical units in the NTK Rift, as presented by (Depicker et al., 2020) (Depicker et al., 2020), and based on the work of Laghmouch et al. (2018).

3.1.4 Rock strength and threshold slopes

In order to account for the control of lithology on the hillslope response to uplift and incision (Schmidt and Montgomery, 1995; Korup, 2008; Korup and Weidinger, 2011; Bennett et al., 2016), we classify the 12 lithostratigraphical units present in the NTK Rift (**Fig. 2d** and **Table 1**) into major categories based upon the analysis of their threshold slope, a proxy for rock strength (Korup and Weidinger, 2011). ~~We determine this value~~ Rock strength is a factor that must be taken into account when investigating landslide characteristics; equal slopes with different rock strength properties are expected to display different behavior in terms of landsliding and knickpoint retreat (Parker et al., 2016; Baynes et al., 2018; Campforts et al., 2020). We determine the rock strength by analyzing the dependency of the average hillslope gradient, S , on the normalized steepness index, k_{sn} , averaged on a catchment scale (~~DiBiase et al., 2010; Bennett et al., 2016~~) (Safran et al., 2005; DiBiase et al., 2010; Bennett et al., 2016). We analyze first order catchments, whereby a drainage network was derived from the 1 arc second SRTM DEM data and a threshold catchment area of 10^5 m^2 . For each lithostratigraphical unit, we only consider watersheds where more than 50 % of the area is covered with the dedicated lithostratigraphy.

The k_{sn} values of a river segment is a proxy for the river incision rate and is calculated using the following equation (Wobus et al., 2006):

$$k_{sn} = S_{chan} A^{\theta_{ref}}, \quad (1)$$

with S_{chan} the local channel slope, A the upstream catchment area, and θ_{ref} the reference concavity index, set at 0.45 (See e.g. DiBiase and Whipple, 2011).

155 Theory suggests a positive linear relationship between S and k_{sn} in catchments with relatively low river incision rates. For catchments with high river incision rates, an increase in k_{sn} will not lead to further hillslope steepening, but to slope failure (DiBiase et al., 2010; Korup and Weidinger, 2011; Larsen and Montgomery, 2012), so that the S becomes independent of the k_{sn} . To capture this non-linear dependency of average basin slope on channel steepness, we introduce a new empirical relationship to describe the response of S to k_{sn} :

160
$$S = TA \left(1 - \exp\left(-\frac{a}{TA} k_{sn}\right) \right), \quad (2)$$

where parameter a is the slope of the curve at $k_{sn} = 0$. Thus, for low incision rates, a approximates the slope of the linear relationship between S and k_{sn} . Parameter TA is the slope angle to which the terrain converges for high k_{sn} values. Hence, TA can be considered equivalent to the threshold slope. However, when there is a linear relationship for $S = f(k_{sn})$ in the entire k_{sn} range (when the $R^2 > 0$ for a linear fit), we do not consider the threshold estimate reliable.




Figure 4. Examples of deforestation followed by landsliding. a) Landslide event North-north of Butezi, DRC (28.296984° E, 2.843201° S; image ©2019 Google Earth). **b)** Landslide event in Matala, DRC (28.360656° E, 2.645874° S; image ©2019 Google Earth).

3.2.1 Inventory

The assessment of the shallow landslide erosion is based on a © Google Earth landslide inventory which is an update from the dataset presented by Depicker et al. (2020). Only recent landslides, for which we can estimate the time of occurrence, are considered in our inventory. In other words, the moment of failure must be situated between the timing of two images. ~~Thus, landslides that are detected in the oldest image of an imagery sequence are not incorporated in the database. Since~~ Moreover, since deforestation mainly affects the stability of the first few meters of the regolith (Sidle and Bogaard, 2016), we only consider shallow landsliding in this study, ~~we~~. Deep-seated and bedrock landslides are excluded from the inventory. We apply a maximum depth of a couple of meters for landslides to be inventoried. We estimate the relative depth of the landslides (shallow or deep-seated) through *in situ* field observations and/or by visually analyzing the shape and size of the landslide scarp and deposits in © Google Earth imagery (Depicker et al., 2020). All images used in the analysis are of very-high spatial resolution, ranging from 30 to 60 cm. The images in © Google Earth are provided by either © DigitalGlobe or © CNES/© Airbus and they were captured between 2000 and 2019. Each landslide is manually assigned a polygon delineating the source area so that the total source area LS_S can be calculated ($\text{m}^2 \text{ km}^{-2} \text{ year}^{-1}$; Section 3.2.2). The LS_S is the area over which regolith material has been removed by landsliding on an annual basis and serves as a proxy for landslide erosion. To each landslide, we also manually assign a point of initiation used to calculate the landslide frequency LS_F ($\# \text{LS km}^{-2} \text{ year}^{-1}$; Section 3.2.2). In order to calculate the LS_F as accurately as possible and avoid amalgamation, we differentiate between separate source areas (Li et al., 2014; Marc et al., 2015; Roback et al., 2018). We also pay attention to not inventory landslides linked to mining and quarrying. Such sites were identified either during fieldwork, or in © Google Earth imagery through characteristics such as a gradual growth of the affected area over a time span of several years and the presence of mining infrastructure (road tracks, trucks, buildings, spoil tips) within the affected area.

~~Apart from amalgamation, our inventory can be biased due to the temporal and spatial inconsistency of © Google Earth. The longer the observed time range, defined as the age difference between the oldest and youngest image, the more landslides that are expected to have occurred. Similarly, we expect the landslide inventory to be less complete in areas of lower imagery density (being the total number of images), as vegetation regrowth might erase the spectral signature of landslides before they are captured in imagery. In order to quantify the imagery range and density, we manually identify all imagery footprints in © Google Earth. All images used in the analysis are of very-high spatial resolution, ranging from 30 to 60 cm. Areas with no available imagery range or density are excluded from the analysis. We also pay attention to not inventory landslides linked to mining and quarrying.~~

~~We investigate the impact of rejuvenation and forest cover on the size of the individual landslides in our inventory.~~ The one-sided Mann-Whitney U test is applied to statistically quantify any differences between different landslide populations (McDonald, 2014). Furthermore, we illustrate the potential differences between the landslide areas of rejuvenated and relict landscapes by comparing the frequency density of the landslide areas. The frequency density curves are fitted to the inverse Γ distribution (Malamud et al., 2004).

3.2.2 Calculating landslide erosion rates from a biased © Google Earth inventory

200 ~~In areas with no spatial or temporal bias in the available imagery, i.e. the imagery range and density are constant~~

Generally, the LS_F is calculated as:

$$LS_F = \frac{n}{rA} \quad (3)$$

with n the total number of shallow landslides, A the total area ~~covered by imagery (km²)~~, and r the imagery ~~time range that is equal for all observed landslides~~ range (years) in © Google Earth, i.e. the age difference between the oldest and youngest

205 ~~image. The imagery range is thus equal to the period of observation. However, © Google Earth imagery is biased and the imagery range is highly variable throughout the study area due to differences in the availability of © Google Earth imagery (Fig. 5a). Since Eq. (3) is valid is valid when r is constant within the study area r has no constant value throughout the entire study area. Hence we have to subdivide, we divide~~ our study area in subareas j ~~of equal time that each have a constant imagery range r^j . The landslide frequency LS_F in each subarea LS_F^j is then:~~

$$210 \quad LS_F^j = \frac{n^j}{r^j A^j}, \quad (4)$$

with n^j the number of landslides in subarea j , A^j the surface area of j , and r^j the constant ~~time imagery~~ range in j . To calculate the frequency for the entire study area, the frequencies LS_F^j are averaged out using weights proportional to their corresponding area A^j :

$$LS_F = \sum_{j=1}^N \frac{A^j}{A} LS_F^j, \quad (5)$$

215 with N the number of subareas j . Substituting **Eq. (4)**, **Eq. (5)** becomes:

$$LS_F = \frac{1}{A} \sum_{j=1}^N \frac{n^j}{r^j A^j} \frac{n^j}{r^j}. \quad (6)$$

~~which is equivalent to~~ Hence, we do not require the size A^j of each subarea j for the calculation of the total LS_F . Instead of aggregating the LS_F over all subareas, we can aggregate the LS_F over the individual landslides. **Equation (6)** then becomes:

$$LS_F = \frac{1}{A} \sum_{i=1}^n \frac{1}{r^i}, \quad (7)$$

220 with r^i the time range observed in landslide i . ~~To compensate for differences in imagery density, The landslide inventory is expected to be biased due to spatial differences in the imagery density d (Fig. 5b), defined as the total number of available images at each location, as vegetation regrowth might erase the spectral signature of landslides before they are captured in imagery. Hence, we expect to detect more landslides in areas with higher imagery density. To compensate for this bias necessary when comparing regions of different imagery coverage,~~ we assume that the probability of identifying ~~landslides a landslide in a certain region~~ increases linearly with imagery density. ~~The equation in that specific region. Equation (7)~~ then becomes:

$$LS_F = \frac{1}{A} \sum_{i=1}^n \frac{1}{r^i d^i}, \quad (8)$$

with d^i the imagery density observed ~~in~~ at the location of landslide i . Note that there can be a saturation of the information provided by the imagery: when the imagery density is high, the availability of one extra image will have no to little effect on the observed number of landslides. We ~~address the issues~~ validate our assumptions of linearity and saturation by visually
 230 assessing the dependency of landslide density (# landslides km^{-2}) on imagery density. If the assumption of linearity does not hold, we have to apply a non-linear transformation on the d^i values. If saturation is problematic to our inventory, we have to set a maximum value for d^i .

Deriving the LS_S equations is analogous to deriving the ones for the LS_F . We only have to slightly modify **Eq. (7)** and **Eq. (8)**:

$$235 \quad LS_S = \frac{1}{A} \sum_{i=1}^n \frac{a_{src}^i}{r^i}, \quad (9)$$

$$LS_S = \frac{1}{A} \sum_{i=1}^n \frac{a_{src}^i}{r^i d^i}, \quad (10)$$

whereby a_{src}^i is the source area of landslide i . Note that the calculation of the LS_S will be less accurate than for the LS_F due to biases in the delineation of the landslide source area. These biases are caused by the time lag between the landslide
 240 occurrence and the landslide detection in © Google Earth, whereby part of the source area might already have recovered. To avoid biases linked to the interpretation of the source area, all landslides were delineated by the same person. In order to statistically verify a difference in landslide activity between regions (for example rejuvenated versus relict landscapes), we use the one-sided non-parametric Mann-Whitney U test to compare the different landslide activity measures in fifth order water catchments (calculated with **Eq. (8)** and **Eq. (10)** to compensate for imagery density differences).

245 3.2.3 Impact of slope on landslide erosion

~~We~~ In order to assess the impact of slope steepness on the LS_S (a proxy for landslide erosion), we first reclassify the slope values between 0-50° into 10 classes of equal width. ~~For~~ and subsequently apply Eq. (10) to each slope class and the landslides therein. Similarly, to assess the impact of slope steepness on LS_F , we apply Eq. (8) and (10) separately. The uncertainty on the to each slope class and its landslides. Furthermore, we estimate the degree to which our LS_S and LS_F ~~is assessed by~~
 250 ~~dividing~~ calculations are affected by outliers and/or extreme landslide events. First, we divide the study area in 50 ~~East-West~~ east-west bands of equal width. ~~Subsequently~~ Second, we calculate the LS_S and LS_F for each slope class 50 times, each time leaving ~~the out the slope and landslide~~ data for a single ~~band out of the analysis~~ east-west band. In other words, for each run we slightly perturbate the landslide inventory. ~~As such, we can assess the degree to which our LS_S and LS_F calculations are affected by outliers and/or extreme landslide events.~~

255 3.2.4 Linking forest cover and deforestation to landslide erosion

In order to link forest dynamics to landslide erosion, we must distinguish between landslides that followed deforestation (**Fig. 4**) and landslides that caused deforestation. To identify the correct causality, we reconstructed the timeline of every landslide that




Figure 5. Visualization of the imagery bias in © Google Earth. a) Imagery range. b) Imagery density. The range and density were calculated by manually identifying 932 imagery footprints. The highest imagery density is available for the major cities in the study area (Goma, Bukavu, and Bujumbura), whereas the northwest and southwest regions have fewer observations. For some areas (in black) there is no available image.

occurred on deforested land (**Fig. 6b**). Landslides following deforestation are defined as those that happened within the post-deforestation time range, being the period between the first image after the year of deforestation and the most recent image.

260

Determining the LS_S in function of the time elapsed since deforestation (t_{def}) is necessary to characterize the post-deforestation landslide wave. ~~Yet, this analysis is far from straightforward (as~~ t_{def} is temporally dynamic ~~) and this analysis~~ requires two components: i) the total ~~affected area of landslides that happened t_{def} years after deforestation, and ii) the total area~~ $A_{t_{def}}$ in which we can observe land that was deforested t_{def} years ago. ~~For the first, and ii) the total affected area of landslides that happened t_{def} years after deforestation. The first component, $A_{t_{def}}$, entails all areas where the sum of t_{def} and the year of deforestation lies in the time range between the age of the oldest and newest post-deforestation image in © Google Earth. For the second component, we only include landslides for which the time between deforestation and~~ $t_{def \rightarrow LS}$ is equal to t_{def} .

265

There is a considerable degree of uncertainty associated to $t_{def \rightarrow LS}$. ~~Firstly, for each landslide, the exact moment of failure must be assumed to be distributed uniformly since we do not know the exact timing of the landslides (the occurrence is situated between the capture times of the image where it was initially observed and the preceding image. Secondly).~~ Similarly, we

270




Figure 6. Schematic overview of the three considered forest cover scenarios in © Google Earth. The satellite icons signal the availability of a © Google Earth image and the colored circles indicate whether we can observe recent landslides in the concerned image. **a)** Forest scenario: each landslide observed in these areas is linked to forest cover. **b)** Deforestation scenario: only landslides observed in-starting from the second © Google Earth image after the year of deforestation are considered to be linked to deforestation (in other words, we can only observe deforestation-induced landslides in imagery that is encircled in red on the figure). The red bar indicates Note that we do not know the year-exact moment of deforestation, only the year (indicated with the red bar) is reported. **c)** Non-forest scenario: every landslide observed in these regions is linked to non-forest.

know the year of deforestation but not the exact date. ~~Here, we assume the deforestation time to be uniformly distributed in the reported year. To visualize these uncertainties, we recalculate the $t_{def \rightarrow LS}$~~ To assess the uncertainty on the timing of deforestation and landslide occurrence, we calculate the LS_S 100 times for each landslide, each time randomly sampling the moments of deforestation and landsliding. The second component, $A_{t_{def}}$, entails all areas where the sum of t_{def} and the year of deforestation lies in the range between the oldest and newest post-deforestation image in © Google Earth sampling a new $t_{def \rightarrow LS}$ for each landslide. For each sample of $t_{def \rightarrow LS}$, we make two assumptions: i) the exact moment of deforestation (the lower limit of $t_{def \rightarrow LS}$) is assumed to be distributed uniformly in the reported deforestation year. ii) The timing of landslide occurrence (the upper limit of $t_{def \rightarrow LS}$), is assumed to be distributed uniformly between the capture times of the image where it was initially observed and the preceding image.




Figure 7. Landslide and knickpoint inventory for the NTK Rift. **a)** We identified ~~7994~~ 7,994 shallow recent landslides and ~~834~~ knickpoints that occurred either in forest, of which ~~non-forest~~ or after deforestation, and 673 ~~were considered~~ non-stationary knickpoints. These ~~knickpoint~~ knickpoints were used to separate the rejuvenated landscapes between the Rift shoulders from the surrounding relict landscapes (~~black-and-white line~~). **b)** Example of shallow landslides in Rwanda (29.7909 °E, -1.7151 °S) and the delineation of their total area (red) and source area (green). **c)** Example of the Rift shoulder west of Lake Tanganyika. The method for delineating the rejuvenated landscapes is specified in Section 3.1.2.

4 Results

4.1 Regional controls on landslide erosion

We identified ~~834~~ 673 non-stationary knickpoints using a tolerance value of 100 m. ~~From these, 673 knickpoints were considered non-stationary and were~~ These knickpoints were used to demarcate the rejuvenated ~~Rift~~ (landscapes inside the Rift (Fig. 7a)).

285 The rejuvenated landscapes encompass 15,526 km², or 18 % of the entire study area.

The average annual rainfall in the rejuvenated landscapes is significantly lower than in the relict landscapes (1,905 mm/year versus 2,297 mm/year, $p < 0.01$). Similarly, we find that within the rejuvenated landscapes the 2 day 15 mm threshold is exceeded ~~significantly~~ less often compared to the relict landscapes (17 % difference, $p < 0.01$), indicating that intense (potentially landslide-triggering) rainfall events occur less frequently.

290 Based on the analysis of 234,840 first-order catchments, we identify three major lithological categories (Fig. 8). Category I comprises units that do not display clear threshold angles. These are lithostratigraphies of relatively young age such as recent and young volcanic basalts and lake and river sediments (all < 23 Ma) except for the Malagarasi rock formations. The latter ~~formation is~~ formations are of old age (1000-540 Ma) and ~~covers~~ cover only a small area in the ~~Southeast~~ southeast of the study area. The lack of threshold landscapes in Category I could be related to the fact that the duration of exposure to weathering
295 for these rocks has been too short. Category II, consisting of old volcanic basalts and Karoo lithostratigraphy (both younger than 210 Ma), has threshold slopes of roughly 17°. Rocks of Category III, with observed threshold slopes ranging between 24-28°, are generally of older age (> 540 Ma) and display a high resistance to slope failure. The lithostratigraphy of Category III includes the following formations: Itombwe, Alkaline complexes, Granites, Kivu, Ruzizi, and Archaen.

4.2 Shallow landslide erosion in the NTK Rift: impacts of deforestation and rejuvenation

300 We inventoried 7,944 recent shallow landslides (Fig. 7a). Following the classification of Hungr et al. (2014), the observed landslides were mostly debris slides ~~(often evolving into avalanches), debris,~~ caused by the sliding of regolith on a planar surface parallel to the ground. These debris slides, once initiated, often transform into avalanches, characterized by the flow of (at least partially) saturated debris on a steep slope. Another commonly observed landslide type was the debris/mud flows, and flows ~~slides~~ flow, defined as the rapid flow of saturated debris in a steep channel. In total, we found 873 landslides in deforested
305 land, yet for only 378 of those landslides we could be certain that they were preceded by deforestation (Section 3.2.4). Furthermore, 3,155 landslides were associated with forest, and 4,411 with non-forest. Rocks of Category I and II combined contained only 344 instances, hampering a robust analysis. We therefore focus our further analysis on the ~~7600~~ 7,600 landslides in areas with rocks of Category III.

~~A total of 932 © Google Earth images, dating between 2000 and 2019, were identified. There is a great spatial variety in~~
310 ~~imagery range and density (Fig. 9a-b). The highest imagery density is available for the major cities in the study area (Goma, Bukavu, and Bujumbura), whereas the northwest and southwest regions have fewer observations. The~~ The number of observed landslides ~~increases in~~ © Google Earth appears to increase linearly with the available imagery up to a density of 12 images (Fig. 9a). ~~At more~~ The proportion of the study area with a higher density than 12 images ~~, the imagery density is 'saturated':~~




Figure 8. Threshold slope analysis for the different lithostratigraphical units in the NTK Rift. Based on Each point represents a first order river catchment over which we averaged the trends of the landscape-slope gradient S versus the and normalized river steepness index k_{sn} . The black curves represent the $S = f(k_{sn})$ relationship is presented by the black curve on each subplot; fitted to Eq. (2), both averaged over first-order river catchments. a)→d) **Category I** Young lithostratigraphy for which no clear threshold angle is observed. e)→f) **Category II** Young lithostratigraphy with a low threshold angle of ca. 17° . g)→l) **Category III** Older rocks with higher observed threshold slopes of $24-28^\circ$.

315 additional images offer no added value for landslide detection. In our inventory, nearly 99 % of landslides occur in areas with imagery density ≤ 12 images is negligible (1.5 %) and contains merely 1 % of all landslides in the inventory. Hence, the assumption that landslide density is linearly dependent on imagery density is not violated valid within our study area and we take no measures to correct for saturation (Section 3.2.2). The annual extent of imagery made available in © Google Earth increases with time, especially after 2010 (Fig. 9db).

320 After accounting for differences in imagery density, the overall LS_S in rejuvenated landscapes is roughly 40 % higher than in relict landscapes ($p=0.034$, Fig. 10a). The difference becomes even larger when looking at the LS_F (160 %, $p=0.014$, Fig. 10b), which implies that landslides are on average smaller in rejuvenated landscapes. This difference in landslide size




Figure 9. Visualization of the imagery bias in © Google Earth. a) Imagery-rangeThe impact of imagery density (the number of available images in © Google Earth) on the number of observed landslides. We only show the results for the rocks of Category III (Section 4.1).**b) Imagery**Therefore, major cities that are characterized by a high imagery density like Bukavu, Bujumbura, and Goma are excluded from this figure.**c) a) Impact of the imagery density on the number of observed landslides.**d)** The number of landslides seems to increase linearly with imagery density up to 12 images. The cumulative landslide proportion for a certain value shows the % of the landslide inventory contained in areas with an imagery density equal or lower than that value.**b)** The evolution of imagery availability between 2000 and 2018.**

between rejuvenated and relict landscapes is confirmed in all three land cover types: forests (114 versus 308 m², $p < 0.01$, [Fig. 11a](#)), non-forests (111 versus 138 m², $p < 0.01$, [Fig. 11b](#)), and deforested land (94 versus 239 m², $p < 0.01$). Similar to the rejuvenation status, forest cover also influences the landslide size. The average source area for forests (223 m²) decreases non-significantly after deforestation (165 m², $p = 0.06$). In non-forest lands, the landslide size (126 m²) is significantly smaller than in recently deforested lands ($p < 0.01$).

The LS_S and LS_F increase with slope gradient (**Fig. 12a-b,d-e**). A decrease is observed for forested slopes $> 45^\circ$, which could be linked to limitations on regolith formation, whereby weathering and sediment deposition are outpaced by erosion ([Montgomery, 2001](#)) ([Montgomery, 2001](#); [Dykes, 2002](#); [Prancevic et al., 2020](#)). When comparing slopes of equal steepness, we observe that the LS_S is generally higher in relict landscapes than in rejuvenated landscapes (**Fig. 12c**). Nevertheless, the overall LS_S is higher in rejuvenated landscapes, because the overall predominance of steeper relief (**Fig. 12g-h**) compensates for the fact that, comparing similarly angled individual slopes in rejuvenated and relict zones, rejuvenated slopes are shown to have a lower or equal rate of shallow landslide erosion (**Fig. 12c**).




Figure 10. Overall landslide activity in the NTK Rift, adjusted for imagery density. a) Overall landslide source area (LS_S), a proxy for landslide erosion. **b)** Landslide frequency (LS_F).

Recently deforested slopes are up to eight times more sensitive to shallow landsliding compared to forested slopes (**Fig. 12a-b**). The deforestation effect lasts approximately 15 years (**Fig. 13**). However, deforestation increases LS_S much more in relict landscapes compared to rejuvenated areas (**Fig. 12c**). The LS_S in the ‘non-forested’ areas (blue lines in **Fig. 12**) corresponds to the situation that prevails once the deforestation-induced landslide wave has passed. In this situation, the LS_S drops back to a level similar to that observed under forest (green line) (**Fig. 12a-b**).

5 Discussion

340 5.1 Interactions between deforestation, rejuvenation, and landslide erosion




Figure 11. Frequency density in function of the landslide source area. a) The area frequency density of shallow landslides in forest, separated for rejuvenated and relict landscapes. **b)** The area frequency density of shallow landslides in non-forest, separated for rejuvenated and relict landscapes. There were not enough landslide observations in deforested land to fit to their area frequency density to the inverse Γ distribution. The general frequency density distributions for inventories of different magnitudes (the black lines on the curves) are derived from Malamud et al. (2004). Note that the frequency is skewed towards smaller landslide sizes due to the omission of deep-seated (and generally larger) landslides from our inventory.




Figure 12. The effect of slope steepness and rejuvenation on landslide activity, corrected for imagery density. We only show results for slope classes in which we observed more than 20 landslides. **a)→c)** Landslide source area (LS_S) in function of slope. **d)→f)** Landslide frequency (LS_F) in function of slope. **g)→h)** Slope distribution for the terrain and landslides in the rejuvenated and relict landscapes. The blue and green arrows indicate the median slope in non-forest and forest landscapes. The slopes in rejuvenated landscapes are clearly steeper, both in forest and non-forest land.

We While the landslide erosion rate (approximated by the LS_S) is higher in rejuvenated landscapes due to a steeper relief, the relative effect of slope steepness on landslide erosion appears to be weaker in rejuvenated landscapes: we found that steep ($>35^\circ$) forested slopes of equal steepness display higher shallow landslide erosion rates (approximated by the LS_S) in relict landscapes than in rejuvenated landscapes (Fig. 12c). A first explanation for this difference could be the drier climate in the rejuvenated landscapes. Based on the global rainfall threshold proposed by Guzzetti et al. (2008), it is likely that landslide-triggering rainfall events are on average more frequent in relict areas compared to rejuvenated landscapes. However, it should be noted

345




Figure 13. Deforestation-induced landslide wave. Overall landslide source area (LS_S , $\text{m}^2 \text{ km}^{-2} \text{ year}^{-1}$) in function of time elapsed since deforestation, based on the analysis of 374 post-deforestation landslides in rocks of category III (Section 4.1). The grey area is the 90 % confidence interval, derived from 100 iterations of LS_S calculations (Section 3.2.4). The dashed and dotted line represent the overall erosion rates in rejuvenated and relict landscapes. There are not enough observations to make two separate consistent plots for rejuvenated and relict landscapes (Fig. A1)

~~that the difference in landslide erosion rates between~~ We evaluate three mechanisms that could explain this difference: seismic activity, regolith availability, and climate.

Seismic activity is a first factor that could explain why slope has a different impact on landslide erosion in rejuvenated and relict landscapes under forest is not due to a higher frequency of landsliding events in relict areas. On the contrary, LS_F is much higher in rejuvenated landscapes (Fig. 12f). If rainfall explains the difference in erosion between rejuvenated and relict landscapes, this could be due to the triggering of larger, rather than more, landslides. However, disentangling the relationship between rainfall and landslide frequency/erosion is complicated by the influence of other regional controls, such as seismicity and regolith availability.

A second factor that needs to be considered when evaluating differences in landslide erosion is the difference in seismic activity between the rejuvenated and relict landscapes (Delvaux and Barth, 2010). Generally, there is more and stronger seismic activity within the rejuvenated landscapes (Delvaux et al., 2017). We hypothesize that the higher seismic activity in rejuvenated landscapes would result in elevated landslide erosion rates on longer timescales due to the occurrence of major landslide events triggered by large earthquakes (Mare et al., 2015) (Delvaux and Barth, 2010; Marc et al., 2015). However, in our observed period, no landslide-triggering earthquakes occurred chances of earthquake-triggered landsliding were very limited (Dewitte et al., 2021). The absence of such events lack of such observations suggests that our window of observation is was too short to capture the direct impact of large earthquakes on landslide erosion within the rejuvenated NTK-Rift earthquakes that were large enough to trigger landsliding. Over the long term, the contribution of earthquake-induced landsliding to regolith mobilization in the rejuvenated landscapes may nevertheless be important. Weak seismic activity also contributes to the smaller size and higher frequency of observed landslides in the rejuvenated part of the study area; earthquakes Earthquakes fracture and weaken the

~~bedrock-hillslope material~~ and hence reduce the minimum critical area for landslide initiation (~~Delvaux and Barth, 2010; Milledge et al., 20~~
(Delvaux et al., 2017; Milledge et al., 2014; Vanmaercke et al., 2017). As such, seismic activity may also contribute to a smaller
average landslide size in rejuvenated landscapes. A previous study in the NTK Rift established an indirect link between spatial
patterns of seismic activity (approximated by a modelled PGA product by Delvaux et al. (2017)) and the spatial pattern of the
370 landslide occurrence, though this study did not differentiate between deep-seated and shallow landsliding (Depicker et al., 2020)

~
A ~~third~~^{second} factor potentially contributing to the difference in ~~erosion-on-equal-slopes~~^{slope impact on erosion rates in}
~~rejuvenated and relict landscapes~~ (Fig. 12c) is that the regolith ~~cover~~^{mantle} on rejuvenated slopes is expected to be thinner
and less continuous due to the drier climate, the younger age of the landscape, the continuous adaptation to river incision, and
375 sporadic earthquake-triggered landslide events (Schoenbohm et al., 2004; Egholm et al., 2013; Marc et al., 2015; Braun et al.,
2016), thereby inducing a supply-limited landsliding regime. This can also partly explain why the rejuvenated landscapes have
more, but smaller landslides in comparison to relict landscapes, as the size of the shallow landslides is constraint by the regolith
availability (Prancevic et al., 2020). However, we do not have direct evidence supporting this hypothesis: ~~The~~^{the} collection of
field data on regolith thickness is hampered by limited access to the field, especially in the eastern DRC. Alternatively, regolith
380 depth could be derived from landslide scars observed on a high-resolution DEM, but such product is currently not available.

A difference in the frequency of landslide-triggering rainfall events could be a third explanation for the lower impact of slope
steepness on landslide erosion in rejuvenated landscapes. Based on the global rainfall threshold proposed by Guzzetti et al. (2008)
, we observe that the rainfall threshold for landsliding is exceeded more often in relict landscapes. However, due to these
differences in rainfall, we would not only expect a higher erosion rate in relict landscapes, but also a higher LS_F . The latter
385 is not the case: slope steepness appears to have a lower effect on LS_F in relict landscapes than in rejuvenated landscapes
(Fig. 12f). This discrepancy between erosion and frequency could be linked to two factors: differences in regolith thickness
which allow for larger landslides in the relict landscapes, and seismic fracturing allowing for smaller (and more) landslides in
the rejuvenated landscapes.

Deforestation drastically increases the landslide frequency and landslide erosion rate. The observed landslide erosion and
390 frequency increases two- to eight-fold after deforestation (Fig. 12a-b), which is the same order of magnitude as what has been
reported in literature (Jakob, 2000; Guthrie, 2002; Glade, 2003). The effect of deforestation on landslide erosion and frequency
is temporary, lasting approximately 15 years (Fig. 13; Sidle et al., 2006; Sidle and Bogaard, 2016). After the wave has passed,
landslide erosion rates decrease to a level similar to the average erosion rate in the region (Fig. 13). However, a longer period
of observation would be useful to confirm if or this effect persists after 15 years.

395 We find that the landslide ~~erosion~~^{response} to deforestation is much more pronounced in relict landscapes than in rejuvenated
landscapes (Fig. 12c). This observation may also be linked to the drier climate and ~~seismic fracturing in the rejuvenated~~
~~landscape, as we observe more but smaller landslides~~ (Fig. 12e and 12f) ~~higher seismic activity in the rejuvenated landscape:~~
~~there are less landslide triggering rainfall events and earthquakes may induce a higher average landslide frequency thereby~~
~~removing sensitive pockets of regolith on a semi-regular basis.~~ Differences in regolith availability can also be invoked to explain
400 the different response of these landscapes to deforestation. Assuming that rejuvenated areas are indeed devoid of regolith (in

comparison to relict areas), one may expect that deforestation will lead to a less important response in rejuvenated areas, simply because the stock of material that can be mobilized through landsliding is smaller. ~~Overall, the deforestation wave persists over a time-scale of about 15 years (Fig. 13), which is consistent with reports in other studies (e.g. Sidle and Bogaard, 2016). After the wave has passed, landslide erosion rates decrease to a level similar to that of forest conditions.~~

405 The landslide erosion rates in non-forest land are much lower than in deforested areas and, in fact, similar to or lower than what has been observed in forests (Fig. 12a-b). Thus, ~~even when there is no regrowth of forest vegetation, the fading-out of the deforestation-induced landslide wave in non-forest land cannot be explained by the recovery of forest vegetation.~~ Moreover, ~~the landslide erosion rate returns to normal levels some time after deforestation.~~ ~~results imply that soils are~~ A possible hypothesis that might explain this result is that once the effect of deforestation on landsliding has worn out, the ~~regolith mantle is~~ protected as efficiently by forest cover as by grassland and crops, despite the presence of human practices such as terracing that could promote landsliding (Sidle and Ochiai, 2006). However, this is extremely unlikely given the much smaller rooting depths of both grasses and crops (Holdo et al., 2018). A more probable explanation for the fading-out of the deforestation-induced landslide wave is therefore that ~~the~~ landslide frequency and erosion rate return to lower levels once the most landslide-sensitive regolith pockets have been removed. For those slopes that are stripped of their regolith ~~cover mantle~~ after deforestation, the rainfall threshold for slope failure is temporarily increased and it may take thousands of years to redevelop a regolith depth that matches the pre-failure conditions ~~(Hufschmidt and Crozier, 2008; Parker et al., 2016)~~ (Dykes, 2002; Hufschmidt and Crozier, 2008; Parker et al., 2016). Additionally, depending on the properties of the landslide deposits (e.g. fine-grained or rock debris), slopes might also experience depositional hardening due to an increase in bulk density and cohesion of the slope material (Crozier and Preston, 1999; Brooks et al., 2002), yet field data is required to test this hypothesis.

420 Despite the fact that equal slopes in non-forest and forest land display similar landslide erosion rates, the average source area is significantly smaller in non-forest landscapes. The smaller size is likely due to ~~a the absence of trees and the associated lower overall root cohesion. Regolith with a lower root cohesion exhibits a~~ smaller minimum critical ~~landslide area linked to the absence of tree cover area needed to initiate landsliding~~ (Milledge et al., 2014; Sidle and Bogaard, 2016). Hence, landslides in non-forest and forest land have different size characteristics (Fig. 11), but the total erosion rate in function of slope remains similar (Fig. 12).

5.2 A new approach for calculating landslides erosion rates?

Using Eq. (9) which deals with the biases in the © Google Earth imagery range, we obtain an overall LS_S of $4.86 \text{ m}^2 \text{ km}^{-2} \text{ year}^{-1}$ in rejuvenated landscapes. ~~Adopting an estimated average landslide depth of 2.5 m, this figure corresponds to a denudation rate of 0.012~~ Using the volume~source area relationships presented by Larsen et al. (2010) for soil landslides in Uganda, we obtain a rough estimate of the landslide volumes. As such, we find that the LS_S in rejuvenated landscapes corresponds to an erosion rate of ca. $0.006 \text{ mm year}^{-1}$. This rate can be compared to the regional uplift rates to estimate the importance of shallow landsliding in the overall evolution of the NTK Rift. There are no accurate estimates of the uplift rates in the study area, but the maximal estimation in the Rwenzori Mountains, a particularly tectonically active re-

gion located 150 km North of our study area, is 2 mm year^{-1} (Kaufmann et al., 2016). If we consider similar rates in the NTK Rift, shallow landslide erosion compensates merely ~~0.6~~ 0.3 % of the uplift in the rejuvenated landscapes, assuming a steady state between uplift and denudation. ~~If we assume a much~~ Based on a global relationship between mean local relief and erosion rate, formulated by Montgomery and Brandon (2002), we obtain a more conservative value of ~~0.2~~ 0.6 mm year^{-1} ~~(Montgomery and Brandon, 2002, e.g.)~~ for the overall erosion rate in landscapes with a similar mean local relief as the rejuvenated landscapes in the NTK Rift (ca. 1300 m). In this scenario, shallow landslide erosion ~~compensates 6.1~~ accounts for 1.0 % of the ~~uplift rate~~ total erosion. Both the upper and lower estimate suggest that, while shallow landslides are highly visible in the landscapes we studied, their geomorphic effect is somewhat limited. However, it must be noted that the estimated erosion rate due to shallow landsliding is most likely an underestimation. First, we did not observe earthquake-induced landslide events, which are rare but may lead to catastrophic landslide erosion (Marc et al., 2015; Dewitte et al., 2021). Second, the landslide inventory used to calculate the erosion rate is incomplete due to limitations in © Google Earth coverage. Furthermore, we focused on shallow landsliding but other ~~erosion~~ erosion-processes such as deep-seated landsliding ~~may~~ may also contribute significantly to ~~sediment mobilization erosion~~ sediment mobilization erosion (Depicker et al., 2020; Dewitte et al., 2021). Nevertheless, it is to be expected that overall erosion rates are lower than uplift rates: this is the basic explanation as to why mountainous topography is formed.

6 Conclusions

We studied shallow landsliding along the NTK rift in order to understand how the interplay of landscape rejuvenation and deforestation affects landslide erosion rates. Rejuvenated landscapes display a higher shallow landslide erosion rate than relict landscapes. ~~This difference is due to the generally steeper relief~~ Contrarily, the relative effect of slope steepness on landslide erosion rates is smaller in rejuvenated landscapes ~~which more than compensates for the fact that rejuvenated slopes, when compared to similarly angled slopes in relict zones, often display a somewhat lower landslide erosion rate. These lower rates~~ . These two seemingly contradicting results are reconciled by the observations that erosion generally increases with slope gradient, and that the average slope is much steeper in the rejuvenated landscapes. The lower impact of slope steepness on landslide erosion in the rejuvenated landscapes could be the result of ~~a drier climate,~~ three factors: the omission of earthquake-induced landslide events in our inventory, ~~and potentially a smaller regolith stock~~ a thinner regolith mantle, and a drier climate. The hypothesis is consistent with our observations that deforestation initiates a much larger landslide peak in relict landscapes and that landslides are, on average, much smaller in rejuvenated landscapes. Thus, the response of a landscape to deforestation does not only depend on local topography and climate but also on the geomorphic status of the landscape. Understanding this differential response is also important to assess the risk for the local population. Our study shows that such understanding is only possible if (i) inventory biases linked to © Google Earth imagery are properly eliminated, (ii) landscape status (rejuvenated versus relict) is accounted for, and (iii) a sufficiently long time frame is considered to capture the transient nature of the deforestation-induced landslide wave.

Author contributions. **Arthur Depicker** was responsible for the compilation of the inventory data, the conceptualization of the paper storyline, the development and execution of the statistical analyses, the conduction of fieldwork, and the writing of the manuscript. **Gerard Govers** was involved in conceptualising the paper storyline, shaping the discussion, writing of the manuscript, and obtaining funding for this work. **Liesbet Jacobs** helped to fine-tune the methodology and statistical analysis, to conceptualize the paper storyline, and to write the manuscript. **Benjamin Campforts** provided the know-how to calculate drainage networks, knickpoint locations, and watershed statistics in the TopoToolbox. He contributed to the paper storyline and writing of the manuscript. **Judith Uwihirwe** was a key figure for the completion of fieldwork in Rwanda that lead to the identification of knickpoints and helped in improving our inventory. She provided feedback for the manuscript and helped to better understand landslide processes in the study area. **Olivier Dewitte** was involved in compiling the inventory, conducting fieldwork, conceptualizing the paper storyline, shaping the discussion, writing the manuscript, and obtaining funding for this work.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study was supported by the Belgium Science Policy (BELSPO) through the PASTeCA project (BR/165/A3/PASTECA) entitled 'Historical Aerial Photographs and Archives to Assess Environmental Changes in Central Africa' (<http://pasteca.africamuseum.be/>). We would like to thank Jonas Van de Walle for the provision of the rainfall dataset used in this work.

- Aleman, J. C., Jarzyna, M. A., and Staver, A. C.: Forest extent and deforestation in tropical Africa since 1900, *Nature Ecology and Evolution*, 2, 26–33, <https://doi.org/10.1038/s41559-017-0406-1>, 2018.
- Baynes, E. R., Lague, D., Attal, M., Gangloff, A., Kirstein, L. A., and Dugmore, A. J.: River self-organisation inhibits discharge control on waterfall migration, *Scientific Reports*, 8, 1–8, <https://doi.org/10.1038/s41598-018-20767-6>, 2018.
- 485 Bennett, G. L., Miller, S. R., Roering, J. J., and Schmidt, D. A.: Landslides, threshold slopes, and the survival of relict terrain in the wake of the Mendocino Triple Junction, *Geology*, 44, 363–366, <https://doi.org/10.1130/G37530.1>, 2016.
- Braun, J., Mercier, J., Guillocheau, F., and Robin, C.: A simple model for regolith formation by chemical weathering, *Journal of Geophysical Research: Earth Surface*, 121, 2140–2171, <https://doi.org/10.1002/2016JF003914>, 2016.
- Brooks, S. M., Crozier, M. J., Preston, N. J., and Anderson, M. G.: Regolith stripping and the control of shallow translational hillslope failure: Application of a two-dimensional coupled soil hydrology-slope stability model, Hawke’s Bay, New Zealand, *Geomorphology*, 45, 165–179, [https://doi.org/10.1016/S0169-555X\(01\)00153-2](https://doi.org/10.1016/S0169-555X(01)00153-2), 2002.
- 490 Burbank, D. W., Leland, J., Fielding, E., Anderson, R. S., Brozovic, N., Reid, M. R., and Duncan, C.: Bedrock incision, rock uplift and threshold hillslopes in the northwestern Himalayas, *Nature*, 379, 505–510, <https://doi.org/10.1038/379505a0>, 1996.
- Campforts, B., Vanacker, V., Herman, F., Vanmaercke, M., Schwanghart, W., Tenorio, G. E., Willems, P., and Govers, G.: Parameterization of river incision models requires accounting for environmental heterogeneity: insights from the tropical Andes, *Earth Surface Dynamics*, 8, 447–470, <https://doi.org/10.5194/esurf-8-447-2020>, <https://www.earth-surf-dynam.net/8/447/2020/>, 2020.
- 495 Crozier, M. J.: Deciphering the effect of climate change on landslide activity: a review, *Geomorphology*, 124, 260–267, <https://doi.org/10.1016/j.geomorph.2009.09.010>, 2010.
- Crozier, M. J. and Preston, N. J.: Modelling Changes in Terrain Resistance as a Component of Landform Evolution in Unstable Hill Country, in: *Process Modelling and Landform Evolution. Lecture Notes in Earth Sciences*, vol. 78, edited by Hergarten, S. and Neugebauer, H. J., pp. 267–284, Springer-Verlag Berlin Heidelberg, 1999.
- 500 Delvaux, D. and Barth, A.: African stress pattern from formal inversion of focal mechanism data, *Tectonophysics*, 482, 105–128, <https://doi.org/10.1016/j.tecto.2009.05.009>, 2010.
- Delvaux, D., Mulumba, J.-L., Sebagenzi, M. N. S., Bondo, S. F., Kervyn, F., and Havenith, H.-B.: Seismic hazard assessment of the Kivu rift segment based on a new seismotectonic zonation model (western branch, East African Rift system), *Journal of African Earth Sciences*, 134, 831–855, <https://doi.org/10.1016/j.jafrearsci.2016.10.004>, 2017.
- 505 Depicker, A., Jacobs, L., Delvaux, D., Havenith, H.-B., Maki Mateso, J.-C., Govers, G., and Dewitte, O.: The added value of a regional landslide susceptibility assessment: The western branch of the East African Rift, *Geomorphology*, 353, 106886, <https://doi.org/10.1016/j.geomorph.2019.106886>, 2020.
- 510 Dewitte, O., Dille, A., Depicker, A., Kubwimana, D., Maki-Mateso, J.-C., Mugaruka Bibentyo, T., Uwihirwe, J., and Monsieurs, E.: Constraining landslide timing in a data-scarce context: from recent to very old processes in the tropical environment of the North Tanganyika-Kivu Rift region, *Landslides*, 18, 161–177, <https://doi.org/10.1007/s10346-020-01452-0>, 2021.
- DiBiase, R. A. and Whipple, K. X.: The influence of erosion thresholds and runoff variability on the relationships among topography, climate, and erosion rate, *Journal of Geophysical Research: Earth Surface*, 116, 1–17, <https://doi.org/10.1029/2011JF002095>, 2011.

- 515 DiBiase, R. A., Whipple, K. X., Heimsath, A. M., and Ouimet, W. B.: Landscape form and millennial erosion rates in the San Gabriel Mountains, CA, *Earth and Planetary Science Letters*, 289, 134–144, <https://doi.org/10.1016/j.epsl.2009.10.036>, <http://dx.doi.org/10.1016/j.epsl.2009.10.036>, 2010.
- Dinku, T., Ceccato, P., and Connor, S. J.: Challenges of satellite rainfall estimation over mountainous and arid parts of east africa, *International Journal of Remote Sensing*, 32, 5965–5979, <https://doi.org/10.1080/01431161.2010.499381>, 2011.
- 520 Dykes, A. P.: Weathering-limited rainfall-triggered shallow mass movements in undisturbed steep-land tropical rainforest, *Geomorphology*, 46, 73–93, [https://doi.org/10.1016/S0169-555X\(02\)00055-7](https://doi.org/10.1016/S0169-555X(02)00055-7), 2002.
- Egholm, D. L., Knudsen, M. F., and Sandiford, M.: Lifespan of mountain ranges scaled by feedbacks between landsliding and erosion by rivers, *Nature*, 498, 475–478, <https://doi.org/10.1038/nature12218>, 2013.
- Ellis, E. C., Goldewijk, K. K., Siebert, S., Lightman, D., and Ramankutty, N.: Anthropogenic transformation of the biomes, 1700 to 2000, *Global Ecology and Biogeography*, 19, 589–606, <https://doi.org/10.1111/j.1466-8238.2010.00540.x>, 2010.
- 525 Glade, T.: Landslide occurrence as a response to land use change: a review of evidence from New Zealand, *Catena*, 51, 297–314, <https://doi.org/10.1016/j.cageo.2015.04.007>, 2003.
- Guthrie, R. H.: The effects of logging on frequency and distribution of landslides in three watersheds on Vancouver Island, British Columbia, *Geomorphology*, 43, 273–292, [https://doi.org/10.1016/S0169-555X\(01\)00138-6](https://doi.org/10.1016/S0169-555X(01)00138-6), 2002.
- 530 Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: The rainfall intensity-duration control of shallow landslides and debris flows: An update, *Landslides*, 5, 3–17, <https://doi.org/10.1007/s10346-007-0112-1>, 2008.
- Hansen, M. C., Potapov, P., Moore, R., Hancher, M., Turubanova, S., Tyukavina, A., Thau, D., Stehman, S., Goetz, S., Loveland, T., Komareddy, A., Egorov, A., Chini, L., Justice, C., and Townshend, J.: High-Resolution Global Maps of 21st-Century Forest Cover Change, *Science*, 342, 850–853, <https://doi.org/10.1126/science.1244693>, 2013.
- 535 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 540 Holdo, R. M., Nippert, J. B., and Mack, M. C.: Rooting depth varies differentially in trees and grasses as a function of mean annual rainfall in an African savanna, *Oecologia*, 186, 269–280, <https://doi.org/10.1007/s00442-017-4011-4>, 2018.
- Hufschmidt, G. and Crozier, M. J.: Evolution of natural risk: Analysing changing landslide hazard in Wellington, Aotearoa/New Zealand, *Natural Hazards*, 45, 255–276, <https://doi.org/10.1007/s11069-007-9158-6>, 2008.
- 545 Hungr, O., Lerouel, S., and Picarelli, L.: The Varnes classification of landslide types, an update, *Landslides*, 11, 167–194, <https://doi.org/10.1007/s10346-013-0436-y>, 2014.
- Jakob, M.: The Impacts of Logging on landslide activity at Clayoquot Sound, British Columbia, *Catena*, 38, 279–300, [https://doi.org/10.1016/S0341-8162\(99\)00078-8](https://doi.org/10.1016/S0341-8162(99)00078-8), 2000.
- JRC and CIESIN: GHS population grid, derived from GPW4, multitemporal (1975, 1990, 2000, 2015). Accessed August 12, 2020 (http://data.europa.eu/89h/jrc-ghsl-ghs_pop_gpw4_globe_r2015a), 2015.
- 550 Kaufmann, G., Hinderer, M., and Romanov, D.: Shaping the Rwenzoris: balancing uplift, erosion, and glaciation, *International Journal of Earth Sciences*, 105, 1761–1778, <https://doi.org/10.1007/s00531-015-1174-2>, <http://dx.doi.org/10.1007/s00531-015-1174-2>, 2016.

- Kirby, E. and Whipple, K. X.: Expression of active tectonics in erosional landscapes, *Journal of Structural Geology*, 44, 54–75, <https://doi.org/10.1016/j.jsg.2012.07.009>, <http://dx.doi.org/10.1016/j.jsg.2012.07.009>, 2012.
- 555 Korup, O.: Rock type leaves topographic signature in landslide-dominated mountain ranges, *Geophysical Research Letters*, 35, 1–5, <https://doi.org/10.1029/2008GL034157>, 2008.
- Korup, O. and Weidinger, J. T.: Rock type, precipitation, and the steepness of Himalayan threshold hillslopes, *Geological Society, London, Special Publications*, 353, 235–249, <https://doi.org/10.1144/SP353.12>, <http://sp.lyellcollection.org/lookup/doi/10.1144/SP353.12>, 2011.
- Laghmouch, M., Kalikone, C., Ilombe, G., Ganza, G., Delvaux, D., Safari, E., Bachinyaga, J., Wazi, N., Nzolang, C., Fernandez, M., Nimpagaritse, G., Tack, L., Dewaele, S., and Kervyn, F.: Carte géologique du Kivu au 1/500 000, *Africamuseum (Tervuren, Belgique), Université Officielle de Bukavu (Bukavu, DRC)*, 2018.
- 560 Larsen, I. J. and Montgomery, D. R.: Landslide erosion coupled to tectonics and river incision, *Nature Geoscience*, 5, 468–473, <https://doi.org/10.1038/NGEO1479>, 2012.
- Larsen, I. J., Montgomery, D. R., and Korup, O.: Landslide erosion controlled by hillslope material, *Nature Geoscience*, 3, 247–251, <https://doi.org/10.1038/NGEO776>, 2010.
- 565 Li, G., West, A. J., Densmore, A. L., Jin, Z., Parker, R. N., and Hilton, R. G.: Seismic mountain building: landslides associated with the 2008 Wenchuan earthquake in the context of a generalized model for earthquake volume balance, *Geochemistry, Geophysics, Geosystems*, 15, 833–844, 2014.
- Malamud, B. D., Turcotte, D. L., Guzzetti, F., and Reichenbach, P.: Landslide inventories and their statistical properties, *Earth Surface Processes and Landforms*, 29, 687–711, <https://doi.org/10.1002/esq.1064>, 2004.
- 570 Marc, O., Hovius, N., Meunier, P., Uchida, T., and Hayashi, S.: Transient changes of landslide rates after earthquakes, *Geology*, 43, 883–886, <https://doi.org/10.1130/G36961.1>, 2015.
- McDonald, J.: *Handbook of Biological Statistics*, Sparky House Publishing, Baltimore, MD, 3rd edn., 2014.
- Milledge, D. G., Bellugi, D., Mckean, J. A., Densmore, A. L., and Dietrich, W. E.: A multidimensional stability model for predicting shallow landslide size and shape across landscapes, *Journal of Geophysical Research: Earth Surface*, 119, 2481–2504, <https://doi.org/10.1002/2014JF003135>, 2014.
- 575 Monsieurs, E., Jacobs, L., Michellier, C., Basimike, J., Bamulezi Ganza, G., Kervyn, F., Maki Mateso, J.-C., Mugaruka Bibentyo., T., Kalikone Buzera, C., Nahimana, L., Ndayisenga, A., Nkurunziza, P., Thiery, W., Demoulin, A., Kervyn, M., and Dewitte, O.: Landslide inventory for hazard assessment in a data- poor context: a regional-scale approach in a tropical African environment, *Landslides*, 15, 2195–2209, <https://doi.org/10.1007/s10346-018-1008-y>, 2018a.
- 580 Monsieurs, E., Kirschbaum, D. B., Tan, J., Maki Mateso, J. C., Jacobs, L., Plisnier, P. D., Thiery, W., Umutoni, A., Musoni, D., Bibentyo, T. M., Ganza, G. B., Mawe, G. I., Bagalwa, L., Kankurize, C., Michellier, C., Stanley, T., Kervyn, F., Kervyn, M., Demoulin, A., and Dewitte, O.: Evaluating TMPA rainfall over the sparsely gauged East African Rift, *Journal of Hydrometeorology*, 19, 1507–1528, <https://doi.org/10.1175/JHM-D-18-0103.1>, 2018b.
- 585 Montgomery, D., Schmidt, K., Greenberg, H., and Dietrich, W.: Forest clearing and regional landsliding, *Geology*, 28, 311–314, [https://doi.org/10.1130/0091-7613\(2000\)28<311:FCARL>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<311:FCARL>2.0.CO;2), 2000.
- Montgomery, D. R.: Slope distributions, threshold hillslopes, and steady-state topography, *American Journal of Science*, 301, 432–454, <https://doi.org/10.2475/ajs.301.4-5.432>, 2001.
- Montgomery, D. R. and Brandon, M. T.: Topographic controls on erosion rates in tectonically active mountain ranges, *Earth and Planetary Science Letters*, 201, 481–489, [https://doi.org/10.1016/S0012-821X\(02\)00725-2](https://doi.org/10.1016/S0012-821X(02)00725-2), 2002.
- 590

- Mugagga, F., Kakembo, V., and Buyinza, M.: Land use changes on the slopes of Mount Elgon and the implications for the occurrence of landslides, *Catena*, 90, 39–46, <https://doi.org/10.1016/j.catena.2011.11.004>, 2012.
- Musumba Teso, P., Kavira, M., and Katcho, K.: Key Factors Driving Deforestation in North-Kivu Province, Eastern DR-Congo Using GIS and Remote Sensing, *American Journal of Geographic Information System*, 8, 11–25, <https://doi.org/10.5923/j.ajgis.20190801.02>,
595 <http://article.sapub.org/10.5923.j.ajgis.20190801.02.html>, 2019.
- Parker, R. N., Hales, T. C., Mudd, S. M., Grieve, S. W. D., and Constantine, J. A.: Colluvium supply in humid regions limits the frequency of storm-triggered landslides, *Scientific Reports*, 6, 1, 2016.
- Poucllet, A., Bellon, H., and Bram, K.: The Cenozoic volcanism in the Kivu rift: Assessment of the tectonic setting, geochemistry, and geochronology of the volcanic activity in the South-Kivu and Virunga regions, *Journal of African Earth Sciences*, 121, 219–246,
600 <https://doi.org/10.1016/j.jafrearsci.2016.05.026>, 2016.
- Prancevic, J. P., Lamb, M. P., McArdell, B. W., Rickli, C., and Kirchner, J. W.: Decreasing Landslide Erosion on Steeper Slopes in Soil-Mantled Landscapes, *Geophysical Research Letters*, 47, 1–9, <https://doi.org/10.1029/2020GL087505>, 2020.
- Roback, K., Clark, M. K., West, A. J., Zekkos, D., Li, G., Gallen, S. F., Chamlagain, D., and Godt, J. W.: The size, distribution, and mobility of landslides caused by the 2015 Mw7.8 Gorkha earthquake, Nepal, *Geomorphology*, 301, 121–138,
605 <https://doi.org/10.1016/j.geomorph.2017.01.030>, 2018.
- Roche, E. and Nzabandora, C. K.: From the congo-nile ridge to the hills area : 2500 years of environmental evolution in southern rwanda, *Geo-Eco-Trop*, 44, 43–63, 2020.
- Safran, E. B., Bierman, P. R., Aalto, R., Dunne, T., Whipple, K. X., and Caffee, M.: Erosion rates driven by channel network incision in the Bolivian Andes, *Earth Surface Processes and Landforms*, 30, 1007–1024, <https://doi.org/10.1002/esp.1259>, 2005.
- 610 Saria, E., Calais, E., Stamps, D. S., Delvaux, D., and Hartnady, C. J. H.: Present-day kinematics of the East African rift, *Journal of Geophysical Research: Solid Earth*, 119, 3584–3600, <https://doi.org/DOI: 10.1002/2013JB010901>, 2014.
- Schmidt, K. M. and Montgomery, D. R.: Limits to Relief, *Science*, 270, 617–620, <https://doi.org/10.1126/science.270.5236.617>, 1995.
- Schoenbohm, L. M., Whipple, K. X., Burchfiel, B. C., and Chen, L.: Geomorphic constraints on surface uplift, exhumation, and plateau growth in the Red River region, Yunnan Province, China, *Geological Society of America Bulletin*, 116, 895–909,
615 <https://doi.org/10.1130/B25364.1>, 2004.
- Schwanghart, W. and Scherler, D.: Bumps in river profiles: Uncertainty assessment and smoothing using quantile regression techniques, *Earth Surface Dynamics*, 5, 821–839, <https://doi.org/10.5194/esurf-5-821-2017>, 2017.
- Sidle, R. C. and Bogaard, T. A.: Dynamic earth system and ecological controls of rainfall-initiated landslides, *Earth-Science Reviews*, 159, 275–291, <https://doi.org/10.1016/j.earscirev.2016.05.013>, 2016.
- 620 Sidle, R. C. and Ochiai, H.: Landslides: Processes, Prediction and Land Use, *American geophysical union*, <https://doi.org/10.1029/WM018>, 2006.
- Sidle, R. C., Ziegler, A. D., Negishi, J. N., Nik, A. R., Siew, R., and Turkelboom, F.: Erosion processes in steep terrain—Truths, myths, and uncertainties related to forest management in Southeast Asia, *Forest Ecology and Management*, 224, 199–225, 2006.
- Smets, B., Kervyn, M., D’Oreye, N., and Kervyn, F.: Spatio-temporal dynamics of eruptions in a youthful extensional setting: Insights
625 from Nyamulagira Volcano (D.R. Congo), in the western branch of the East African Rift, *Earth-Science Reviews*, 150, 305–328, <https://doi.org/10.1016/j.earscirev.2015.08.008>, <http://dx.doi.org/10.1016/j.earscirev.2015.08.008>, 2015.
- Tyukavina, A., Hansen, M. C., Potapov, P., Parker, D., Okpa, C., Stehman, S. V., Kommareddy, I., and Turubanova, S.: Congo Basin forest loss dominated by increasing smallholder clearing, *Science Advances*, 4, <https://doi.org/10.1126/sciadv.aat2993>, 2018.

- USGS: Shuttle Radar Topography Mission, Global Land Cover Facility. 1 Arc-Second. University of Maryland, College Park, Maryland, 2006.
- USGS: LANDSAT-8, ETM+SLC-on. 60 m resolution., 2018.
- Van de Walle, J., Thiery, W., Brousse, O., Souverijns, N., Demuzere, M., and van Lipzig, N. P.: A convection-permitting model for the Lake Victoria Basin: evaluation and insight into the mesoscale versus synoptic atmospheric dynamics, *Climate Dynamics*, 54, 1779–1799, <https://doi.org/10.1007/s00382-019-05088-2>, <https://doi.org/10.1007/s00382-019-05088-2>, 2020.
- Vanmaercke, M., Ardizzone, A., Rossi, M., and Guzzetti, F.: Exploring the effects of seismicity on landslides and catchment sediment yield: An Italian case study, *Geomorphology*, 278, 171–183, <https://doi.org/10.1016/j.geomorph.2016.11.010>, 2017.
- Whipple, K. X. and Meade, B. J.: Orogen response to changes in climatic and tectonic forcing, 243, 218–228, <https://doi.org/10.1016/j.epsl.2005.12.022>, 2006.
- Wobus, C., Whipple, K. X., Kirby, E., Snyder, N., Johnson, J., Spyropolou, K., Crosby, B., and Sheehan, D.: Tectonics from topography: Procedures, promise, and pitfalls, *Special Paper of the Geological Society of America*, 398, 55–74, [https://doi.org/10.1130/2006.2398\(04\)](https://doi.org/10.1130/2006.2398(04)), 2006.

Appendix A: Deforestation wave in rejuvenated and relict landscapes




Figure A1. Deforestation-induced landslide wave in different geomorphic contexts. Overall landslide source area (LS_s , $\text{m}^2 \text{ km}^{-2} \text{ year}^{-1}$) in function of time elapsed since deforestation. The grey area represents the 90 % confidence interval, derived from 100 iterations of LS_s calculations (Section 3.2.4). The dotted line shows the overall erosion rates in the different contexts. **a) Rejuvenated**—Landslide response to deforestation in rejuvenated landscapes, characterized with 193 landslide instances. The irregular curve suggests that ~~we need more landslide observations~~, in order to better characterize the deforestation-induced landslide wave, ~~we need more landslide and deforestation observations and over a larger area~~. **b) Relict**—Landslide response to deforestation in relict landscapes, characterized with 181 landslide instances. The dotted line represents the background erosion rate in relict landscapes.