

1 **Time-Of-Flight monitoring reveals higher sediment redistribution rates related to burrowing animals**  
2 **than previously assumed**

3  
4 *Paulina Grigusova*<sup>1</sup>, *Annegret Larsen*<sup>2</sup>, *Sebastian Achilles*<sup>1</sup>, *Roland Brandl*<sup>3</sup>, *Camilo del Río*<sup>4,5</sup>, *Nina Farwig*<sup>6</sup>,  
5 *Diana Kraus*<sup>6</sup>, *Leandro Paulino*<sup>7</sup>, *Patricio Plischoff*<sup>4,8,9</sup>, *Kirstin Übernickel*<sup>10</sup>, *Jörg Bendix*<sup>1</sup>

6  
7  
8 <sup>1</sup> Laboratory for Climatology and Remote Sensing, Department of Geography, University of Marburg, 35037  
9 Marburg, Germany; paulina.grigusova@staff.uni-marburg.de (P.G.); bendix@geo.uni-marburg.de (J.B.)

10 <sup>2</sup> Soil Geography and Landscape, Department of Environmental Sciences,  
11 Wageningen University & Research, 6700 AA Wageningen, The Netherlands; annegret.larsen@wur.nl

12 <sup>3</sup> Animal Ecology, Department of Biology, University of Marburg, 35032 Marburg, Germany;  
13 brandlr@biologie.uni-marburg.de

14 <sup>4</sup> Facultad de Historia, Geografía y Ciencia Política, Instituto de Geografía, Pontificia Universidad Católica de  
15 Chile, 782-0436 Santiago, Chile; plischoff@uc.cl; cdelriol@uc.cl

16 <sup>5</sup> Centro UC Desierto de Atacama, Pontificia Universidad Católica de Chile, 782-0436 Santiago, Chile;  
17 cdelriol@uc.cl

18 <sup>6</sup> Conservation Ecology, Department of Biology, University of Marburg, 35047 Marburg, Germany;  
19 diana.kraus@biologie.uni-marburg.de (D.K.); nina.farwig@biologie.uni-marburg.de (N.F.)

20 <sup>7</sup> Facultad de Agronomía, Universidad de Concepción, 3780000 Chillán, Chile; lpaulino@udec.cl

21 <sup>8</sup> Facultad de Ciencias Biológicas, Departamento de Ecología, Pontificia Universidad Católica de Chile, 8331150  
22 Santiago, Chile; plischoff@uc.cl

23 <sup>9</sup> Center of Applied Ecology and Sustainability (CAPES), Pontificia Universidad Católica de Chile, 8331150  
24 Santiago, Chile; plischoff@uc.cl

25 <sup>10</sup> Earth System Dynamics, Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany;  
26 kirstin.uebernickel@uni-tuebingen.de

27  
28 *Corresponding author:*

29 Paulina Grigusova  
30 paulina.grigusova@staff.uni-marburg.de

31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 **Abstract**

42 Burrowing animals influence surface microtopography and hillslope sediment redistribution, but changes often  
43 remain undetected due to a lack of automated high resolution field monitoring techniques. In this study, we  
44 present a new approach to quantify microtopographic variations and surface changes caused by burrowing  
45 animals and rainfall-driven erosional processes applied to remote field plots in arid and mediterranean Chile.  
46 We compared the mass balance of redistributed sediment between burrow and burrow embedding area,  
47 quantified the cumulative sediment redistribution caused by animals and rainfall, and upscaled the results to a  
48 hillslope scale. The newly developed instrument, a Time-of-Flight camera, showed a very good detection  
49 accuracy. The animal-caused cumulative sediment redistribution was  $8.52 \text{ cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$  in the  
50 mediterranean and  $9.57 \text{ cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$  in the arid climate zone. The rainfall-caused cumulative sediment  
51 redistribution within burrow was higher ( $-6.09 \text{ cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ ) in the mediterranean than the arid climate  
52 zone ( $-0.82 \text{ cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ ). Daily sediment redistribution during rainfall within burrows were up to 350% /  
53 40% higher in the mediterranean / arid zone compared to burrow embedding area, and much higher than  
54 previously reported in studies not based on continuous microtopographic monitoring. Furthermore, 38% of the  
55 sediment eroding from the burrows accumulated within the burrow entrance while 62% was incorporated into  
56 overall hillslope sediment flux. The animals burrowed between on average 1.2 – 2.3 times a month and the  
57 burrowing intensity increased after rainfall. Our findings can be implemented into long-term soil erosion models  
58 that rely on soil processes but do not yet include animal-induced surface processes on microtopographical  
59 scales in their algorithms.

60

61 **Keywords:** Biogeomorphology, bioturbation, sediment transport, burrowing animals, rainfall, Time-of-Flight  
62 camera, Chile

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77


78

79

80

81

82 **Graphical abstract**



83

84

## 85 1. Introduction

86 Animal burrowing activity affects surface microtopography (Kinlaw and Grasmueck, 2012; Reichman  
87 and Seabloom, 2002), surface roughness (Hancock and Lowry, 2021; Jones et al., 2010; Yair, 1995) and soil  
88 physical properties (Coombes, 2016; Corenblit et al., 2021; Hall et al., 1999; Hancock and Lowry, 2021; Larsen  
89 et al., 2021; Reichman and Seabloom, 2002; Ridd, 1996; Yair, 1995). Previous studies estimated both positive  
90 as well as negative impacts of burrowing animals on sediment redistribution rates. The results were obtained  
91 by applying tests under laboratory conditions using rainfall simulators, conducting several field campaigns  
92 weeks to months apart, or by measuring the volume of excavated or eroded sediment in the field using methods  
93 such as erosion pins, splash boards, or simple rulers (Chen et al., 2021; Imeson and Kwaad, 1976; Le Hir et  
94 al., 2007; Li et al., 2019b; Li et al., 2019a; Li et al., 2019c; Li et al., 2018; Reichman and Seabloom, 2002;  
95 Übernicketl et al., 2021b; Voiculescu et al., 2019; Wei et al., 2007). Although burrowing animals are generally  
96 seen as ecosystem engineers (Gabet et al., 2003; Wilkinson et al., 2009), their role in soil erosion, in general,  
97 and for numerical soil erosion models, in particular, is, to date, limited to predictions of the burrow locations  
98 and particle mixing at these locations (Black and Montgomery, 1991; Meysman et al., 2003; Schiffers et al.,  
99 2011; Yoo et al., 2005). The complex interaction of sediment excavation and accumulation, and erosion  
100 processes at the burrow and hillslope scales are not yet included in the modelling, as for this, a suitable method  
101 capable of measuring all occurred redistribution processes is needed.

102 The reason for this knowledge gap is that previous studies have not provided data on low magnitude  
103 but frequently occurring sediment redistribution due to the specific limitations of their approaches. Field  
104 experiments with, for example, rainfall simulators can unveil processes but cannot cover the time-dependant  
105 natural dynamics of sediment redistribution. For data samplings that used methods such as erosion pins or  
106 splash boards, the sites had to be revisited each time and the data were thus obtained only sporadically  
107 (Hazelhoff et al., 1981; Imeson and Kwaad, 1976; Richards and Humphreys, 2010). Similarly, estimations of  
108 the excavated sediment volume are currently limited to one-time measurements or studies conducted several  
109 months apart (Black and Montgomery, 1991; Hall et al., 1999; Yoo et al., 2005). We expect that non-  
110 continuously conducted measurements do not include all frequently occurring excavation and erosion  
111 processes. For this, a spatio-temporally high-resolution and continuous monitoring of sediment redistribution  
112 is needed.

113 High-resolution, ground-based imaging sensing techniques might overcome such aforementioned  
114 problems. Terrestrial laser scanner systems have shown to be a suitable tool for estimation of sediment  
115 redistribution and erosion processes (Afana et al., 2010; Eltner et al., 2016a; Eltner et al., 2016b; Longoni et  
116 al., 2016; Nasermoaddeli and Pasche, 2008). However, they are expensive and labour-intensive. A  
117 continuous, automated monitoring of many mound areas in parallel is for this reason not possible. An already  
118 applied low-cost (up to 5000 USD) topographic monitoring technique is time-lapse photogrammetry which can  
119 be applied at variable observation distances and scales (e.g. (Blanch et al., 2021; Eltner et al., 2017; Galland  
120 et al., 2016; James and Robson, 2014; Kromer et al., 2019; MALLALIEU et al., 2017). For this technique, the  
121 surface has to be monitored under various angles for which several devices are needed to be installed in the  
122 field.

123 In contrast, The Time-of-Flight (ToF) technology exhibits lower spatial resolution and aerial coverage  
124 compared to time-lapse photogrammetry. However, as an active remote sensing tool it can also be used at  
125 night. Additionally, the processing is less complex compared to photogrammetry because the distance values  
126 are immediately received in a local coordinate system. ToF offers here a new possibility for surface monitoring,

127 as a technique for a cost-effective high-resolution monitoring of sediment redistribution (Eitel et al., 2011;  
128 Hänsel et al., 2016) which can be achieved by a simple installation of one device in the field. ToF-based  
129 cameras illuminate the targeted object with a light source for a known amount of time and then estimate the  
130 distance between the camera and the object by measuring the time needed for the reflected light to reach the  
131 camera sensor (Sarbolandi et al., 2018).


132 In our study we developed, tested and applied a cost-effective Time-of-Flight camera for automated  
133 monitoring of the rainfall and animal-driven sediment redistribution burrows and burrow embedding areas by  
134 burrowing animals with high temporal (four times a day) and spatial (6 mm) resolution. For this, we equipped  
135 several plots in remote study sites in the Chilean arid and mediterranean climate zone. We selected these  
136 sites in order to analyse sediment redistribution by burrowing activity of vertebrates under different rainfall  
137 regimes and as these sites have been shown to be particularly strongly affected by burrowing activity  
138 (Grigusova et al., 2021). We estimated the burrowing intensity and its dependence on rainfall. Then, we  
139 quantified the daily sediment redistribution within burrow and burrow embedding area. We analysed the  
140 impacts of animal burrowing activity and rainfall on the sediment redistribution and quantified the volume of  
141 sediment which is additionally incorporated to the hillslope sediment flux due to the presence of burrows.  
142 Finally, we estimated sediment redistribution on a burrow scale and upscaled sediment redistribution rates to  
143 the entire hillslopes.

144

## 145 2. Study sites

146 Our study sites were located in the Chilean Coastal Cordillera in two climate zones (Fig. 1): in the National  
147 Park Pan de Azúcar (further as Pan de Azúcar or PdA) and the National Park La Campana (further as La  
148 Campana or LC). The Las Lomitas site in PdA is located in the arid climate zone of the Atacama Desert with a  
149 precipitation rate of 12 mm year<sup>-1</sup>, and it has a mean annual temperature of 16.8 °C (Übernicket et al., 2021a).  
150 Here, the vegetation cover is below 5%, and it is dominated by small desert shrubs, several species of cacti  
151 (*Eulychnia breviflora*, *Copiapoa atacamensis*) and biocrusts (Lehnert et al., 2018). LC is located in the  
152 mediterranean climate zone with a precipitation rate of 367 mm year<sup>-1</sup> and a mean annual temperature of  
153 14.1 °C (Übernicket et al., 2021a). LC is dominated by an evergreen sclerophyllous forest with endemic palm  
154 trees, *Jubaea chilensis*. Both research sites have a granitic rock base, and the dominating soil texture is sandy  
155 loam (Bernhard et al., 2018). In PdA, the study setup consisted of one north-facing and one south-facing  
156 hillslope. The hillslope inclinations were ~20°, and a climate station was located ~15 km from the camera sites.  
157 In LC, the setup consisted of two north-facing and one south-facing hillslopes. The hillslope inclinations were  
158 ~25°, and a climate station was located ~250 m from the south-facing hillslope (Übernicket et al., 2021a).  
159 Among the most common vertebrate burrowing animals are in PdA carnivores (*Lycalopex culpaeus*, *Lycalopex*  
160 *griseus*); marsupials and rodents (*Phyllotis xanthopygus*, *Phyllotis limatus*, *Abrothrix andinus*) (Cerqueira,  
161 1985; Jimenez et al., 1992) and in LC rodents (*Octodon degus*, *Rattus norvegicus* and *Phyllotis darwini*) and  
162 carnivores (*Lycalopex griseus*) (Muñoz-Pedrerros et al., 2018)

163



164  
 165 **Figure 1.** Location of the cameras and climate stations on which this study was based. Black points show the  
 166 location of the research sites in Chile. The green points represent the camera plots, and the blue points the  
 167 climate stations: (a) Location of study sites in Chile: PdA stands for Pan de Azúcar, LC for La Campana; (b)  
 168 Study setup in Pan de Azúcar; (c) Study setup in LC. The background images in (b) and (c) are orthophotos  
 169 created from WorldView-2 data from 19 July 2019. For exact latitude and longitude see Table A2.

170  
 171 **3. Methodology**

172 **3.1 Time-of-Flight (ToF) principle**

173 A Time-of-Flight-based camera illuminates an object with a light source, usually in a non-visible  
 174 spectrum, such as near-infrared, for a precise length of time. ToF cameras rely on the principle of measuring  
 175 the phase shift, with different options to modulate the light source to be able to measure the phase shift. The  
 176 here employed cameras used pulse-based modulation, meaning the light pulse was first emitted by the  
 177 camera, then reflected from the surface, and finally measured by the camera using two temporary windows.  
 178 The opening of the first window is synchronized with the pulse emission i.e. the receiver opens the window  
 179 with the same  $\Delta t$  as the emitted pulse. Then, the second window is opened, for the same duration  $\Delta t$ , which is  
 180 synchronised with the closing of the first window. The first temporary window thus measures the incoming  
 181 reflected light while the light pulse is also still emitting from the camera. The second temporary window  
 182 measures the incoming reflected light when no pulse is emitting from the camera. The captured photon number  
 183 (i.e. measured by electrical charge) in both windows can be related according to equation 1 and the distance  
 184 from the camera to the object can then be calculated as follows:

185 
$$d = \frac{1}{2} * c * t * \left( \frac{g_1}{g_1 + g_2} \right) \quad . \quad (1)$$

186 In Eq. (1),  $d$  (m) is the distance from the camera to the object,  $c$  ( $\text{m s}^{-1}$ ) is the speed of light ( $299,792,458 \text{ m s}^{-1}$ ),  
 187  $t$  (s) is the overall time of the illumination and measurement,  $g_1$  is the ratio of the reflected photons to all  
 188 photons accumulated in the first window, and  $g_2$  the ratio of the reflected photons to all photons accumulated  
 189 in the second window (Li, 2014; Sarbolandi et al., 2018).

190 The sensor in our camera came from Texas Instruments and the data scan contained information on  
 191 320 x 240 points. The camera field of view (FOV) and the spatial resolution of the scans depended on the  
 192 height of the camera above the surface and camera orientation. The distance was calculated for every point,  
 193 and the object was saved in binary format as a collection of 3D points with x-, y- and z-coordinates. The point  
 194 clouds taken by the camera were transformed from the binary format to an ASCII format. Each point in the  
 195 point cloud was assigned to an x-, y- and z-coordinate. The coordinates were distributed within a three-  
 196 dimensional Euclidian space, with the point at the camera nadir (the centre of the camera sensor) being the  
 197 point of origin of the 3D Cartesian coordinate system. x- and y-coordinates describe the distance to the point  
 198 of origin (m). z-coordinate describes the distance (m) from the object to the camera. The lowest point of the  
 199 scanned surface thus has the highest z-coordinate value.

200

### 201 3.2 Data processing

202 The distortion caused by the hillslope and the camera angle was corrected for each point cloud as  
 203 follows:

$$204 z_{cor} = z_{uncor} - \tan(\alpha + \beta) * (y_1 - y_i) \quad . \quad (2)$$

205 In Eq. (2),  $z_{cor}$  is the corrected distance (m) between the camera and surface (m),  $z_{uncor}$  is the uncorrected z-  
 206 coordinate (m),  $\alpha$  is the tilt angle of the camera ( $^{\circ}$ ),  $\beta$  is the surface inclination ( $^{\circ}$ ), and  $y_i$  (m) is the distance  
 207 between each point, and the point with i) an y-coordinate = 0 and ii) the same x-coordinate as the respective  
 208 point. The most frequent errors were identified and treated as follows. Due to the ambient light reaching the  
 209 camera sensor, the z-coordinate values of some of the points were incorrect (scattering error). To remove this  
 210 error, a threshold value was calculated for each point cloud:

$$211 \Omega = mean_{z_{cor}-coordinate} \pm sd_{z_{cor}-coordinate} \quad . \quad (3)$$

212 In Eq. (3),  $\Omega$  is the threshold value,  $mean_{z_{cor}-coordinate}$  is the average value, and  $sd_{z_{cor}-coordinate}$  is the standard  
 213 deviation of the corrected z-coordinates (m). Then, all points with a z-coordinate above and below this value  
 214 were deleted. Point clouds with more than 50% of points above the threshold value  $\Omega$  were also not considered  
 215 for further processing. A drift error occurred when the z-coordinate values of around one-third of the point  
 216 clouds decreased by several centimetres from one point cloud to another. Here, the average z-coordinate of  
 217 ten point clouds before and after the drift were calculated, and the difference was added to z-coordinates of  
 218 the points affected by the drift. The corrected height values were then transformed into a digital surface model  
 219 (DSM).

220

### 221 3.3 Accuracy of the ToF cameras

222 The accuracy of the ToF camera was tested under laboratory conditions by recreating similar surface  
 223 conditions as in the field (sloping surface, covered by sediment). An artificial mound using sediment extracted  
 224 from a riverbank in central Germany was used, mimicking a mound created by a burrowing animal. During the  
 225 test, the camera was installed 100 cm above the surface. The camera FOV was 3 m<sup>2</sup> and the scan spatial  
 226 resolution was 6 mm. The surface was scanned twice by the ToF camera. Then 100 – 450 cm<sup>3</sup> of sediment  
 227 was manually extracted from the mound. The volume of the extracted sediment was measured by a measuring  
 228 cup. After extraction, the surface was again scanned twice by the camera. The experiment was repeated 45  
 229 times with varying amounts of extracted sediment. The scans were transformed to point clouds in VoxelViewer-  
 230 0.9.10, and the point clouds were corrected according to Eq. (2) and (3). The z-coordinates of the two point  
 231 clouds before and two point clouds after the extraction were averaged. The standard deviation of the z-

232 coordinate of the two scans was 0.06 cm. Figure A1 shows the spatially distributed standard deviation. The  
 233 deviation increases from the centre towards the corners of the scan. The mound was outlined and only the  
 234 points representing the mound were used in the further analysis. The point clouds were then transformed into  
 235 DSMs, and the differences between the time steps were calculated. A scan was taken of a smooth surface  
 236 (linoleum floor) and a point cloud was created from the data. Then, we fitted a plane into the point cloud and  
 237 calculated the distance between the plane and the camera sensor. The standard variation (0.17 cm) in the  
 238 distance measurements was saved. Solely, the differences between the DSMs below this variation were  
 239 considered in the calculation of the detected sediment extraction. The detected extracted sediment volume  
 240 was then calculated for each experiment as follows:

$$241 \quad Vol_{detected} = \sum_p^1 (DSM_{before} - DSM_{after}) * res^2 \quad , \quad (4)$$

242 In Eq. (4),  $Vol_{detected}$  is the volume of the extracted sediment as detected by the camera ( $cm^3$ ),  $p$  is the number  
 243 of pixels,  $DSM_{before}$  (cm) is the DSM calculated from the scan taken before the extraction,  $DSM_{after}$  (cm) is the  
 244 DSM calculated from the scan taken after the extraction,  $res$  (cm) is the resolution of the scan, which was 0.6  
 245 cm. To evaluate the camera's accuracy, the measured volume of the extracted sediment was compared to the  
 246 volume detected by the camera. The camera's accuracy was estimated between the detected volume and  
 247 measured volume as follows:

$$248 \quad MAE = \sum_1^n \frac{(Vol_{detected} - Vol_{measured})}{area} \quad . \quad (5)$$

249 In Eq. (5),  $MAE$  ( $cm^3/cm^2$ ) is the mean absolute error,  $n$  is the number of scans,  $Vol_{measured}$  ( $cm^3$ ) is the volume  
 250 of the extracted sediment measured by the measuring cup, and the  $area$  is the total surface area monitored  
 251 by the camera ( $cm^2$ ).


252

### 253 3.4 Installation of the cameras in the field

254 We installed 8 custom-tailored ToF-based cameras on 4 hillslopes in two climate zones in areas  
 255 including visible signs of bioturbation activity (burrows) and areas without visible signs of bioturbation (Fig. 2).  
 256 The cameras were installed in LC on the north-facing upper hillslope (LC-NU), north-facing lower hillslope (LC-  
 257 NL), south-facing upper hillslope (LC-SU) and the south-facing lower hillslope (LC-SL); in PdA on the north-  
 258 facing upper hillslope (PdA-NU), north-facing lower hillslope (PdA-NL), south-facing upper hillslope (PdA-SU)  
 259 and south-facing lower hillslope (PdA-SL). The custom-tailored cameras were installed during a field campaign  
 260 in March 2019, the monitoring took place for seven months, and the data were collected in October 2019. The  
 261 construction consisted of a 3D ToF-based sensor from Texas Instruments (Li, 2014), a RaspberryPi single board  
 262 computer (SBC), a timer, a 12 V 12 Ah battery and three 20 W solar panels for unattended operation (Fig. 2).  
 263 Solar panels were located at the camera pole and were recharging the battery via a charge controller. The  
 264 camera was located approximately one meter above the surface, facing the surface with a tilt angle of 10  
 265 degrees. The timer was set to close the electric circuit 4 times a day: at 1 a.m., 5 a.m., 8 a.m. and 10 p.m. At  
 266 these times, the camera and the computer were turned on for 15 minutes. The camera turned on and took five  
 267 scans delayed one second from each other and sent them to the SBC. Each camera had its own WiFi (Wireless  
 268 Fidelity) and the data could be read from the SBC via Secure Shell (SSH). The cameras collected the data for  
 269 the time period of 7 months.

270





271

272 **Figure 2.** Scheme and photo example of a Time-of-Flight-based camera installation in the field. The photo  
 273 example is from upper north-facing hillslope in La Campana. Black boxes describe single installation parts.  
 274 Purple descriptions are the variables needed for the correction of the scans. Roof, entrance and mound  
 275 describe parts of the burrow. The x-, y- and z-coordinates are 3D coordinates identifying the position of each  
 276 point in space, where the x-coordinate is the length, y-coordinate is the width and the z-coordinate is the  
 277 distance between the camera sensor and the surface.  $\alpha$  is the inclination of the camera, and  $\beta$  is the surface  
 278 inclination.

279


### 280 3.5 Delineation of burrows and burrow embedding areas

281 The surface area scanned by the cameras was divided by a delineation scheme into burrows (B) and  
 282 burrow embedding areas (EM). The burrows included three sub-areas: (i) mound (M), (ii) entrance (E) and (iii)  
 283 burrow roof (R). “Mound” describes the sediment excavated by the animal while digging the burrow. “Entrance”  
 284 describes the entry to the animal burrow up to the depth possible to obtain via the camera. “Burrow roof”  
 285 describes the part of the sediment above and uphill the burrow entrance (BANCROFT et al., 2004). During the  
 286 burrow’s creation, sediment was not only excavated but also pushed aside and uphill the entrance, which  
 287 created the burrow roof. We assume that this elevated microtopographical feature then forms an obstacle for  
 288 sediment transported from uphill, which leads to its accumulation in this area. The remaining surface within  
 289 the camera’s FOV was burrow embedding area. Please note, that this area may still be affected by the  
 290 burrowing activity of the animal and is not completely unaffected by the animal.

291 For the delineation, we used the DSM calculated from the point cloud, and a slope layer calculated  
 292 from the DSM (Horn, 1981). The DSM had a size of 4 m<sup>2</sup> a resolution of 0.6 cm. Entrance was assigned to an  
 293 area determined by a search algorithm starting at the lowest point of the DSM (pixel with the highest z-  
 294 coordinate value). We increased the circular buffer around the starting point by one pixel until the average  
 295 depth of the new buffer points was not higher than the height of the camera above the surface, or until the  
 296 slope of at least 50% of the new buffer points was not 0. Then, we masked all pixels within the buffer with a  
 297 depth lower than the average depth of the points within the buffer, which had a slope that was 0. The remaining  
 298 pixels belonged to the entrance area. Then, the surface scan was divided into an uphill and downhill part with

299 regards to the entrance position. Both the uphill and the downhill parts were subdivided into 16 squares, so  
300 that each of the four quadrants within the 2D grid (x- and y-axis) contained four squares. The squares had size  
301 of 0.5 m<sup>2</sup>.

302 To delineate the mound in the downhill part, we first identified the highest points (pixel with the lowest z-  
303 coordinate value) within all 16 squares. We then calculated the distance of these maxima to the entrance, and  
304 the pixel located nearest to the entrance was identified as the highest point of the mound (i.e., seed point).  
305 Consecutively, we increased the circular buffer around the seed point by one pixel until the average depth of  
306 the new buffer points was not lower than the height of the camera above the surface, or until the slope of at  
307 least 50% of the new buffer points was not 0. Then, we masked all pixels within the buffer with a depth higher  
308 than the average depth of the points within the buffer, which had a slope that was 0. The remaining pixels were  
309 classified as mound area. To delineate burrow roof, we used the same approach as for the delineation of  
310 mound and applied it on the uphill part of the surface scan. We used the DEM and slope layers for the  
311 delineation for several reasons. The distance from the surface to the camera was the most important parameter  
312 to derive (i) the deepest point of the entrance and (ii) the highest point of the mound or burrow roof, as this  
313 was (mostly) the closest point to the camera. After the angle correction of the z-coordinate according to chapter  
314 3.2., the surface inclination of the areas without burrow was 0°, while the angle between the border of the  
315 burrow entrance or mound and the burrow embedding surface was above 0°. Because neither the entrance  
316 nor the mound have a perfect circular form, we would largely overestimate or underestimate the entrance or  
317 mound size. Overestimate by not stopping the search algorithm until the angle between all new points of the  
318 buffer to the rest of the buffer was 0°. Underestimate by stopping the algorithm when the angle of one point of  
319 the buffer to the nearest point of the buffer was 0°. The value of 50% thus minimized the error. All pixels that  
320 were not classified during the entire delineation process were treated as burrow embedding areas.  
321 The position and the boundaries of entrance, mound and burrow roof were validated visually (Fig. 3 and A2).



322  
323 **Figure 3.** Corrected digital surface model of the camera on the upper north-facing hillslope in La Campana  
324 with delineated areas. The point of origin of the coordinate system is at the camera nadir. Distance refers to  
325 the distance between surface and camera. The red line delineates the burrow entrance, blue the mound and  
326 orange the burrow roof. The area which was outside of any delineated area was classified as burrow  
327 embedding area. The arrow indicates a downhill direction of the hillslope.

328  
329 In LC, the burrows always consisted of an entrance, mound and burrow roof. In PdA, there was no  
330 burrow roof on the upper hillslopes. Burrows without a burrow roof were located on shallower parts of the

331 hillslopes (up to an inclination of 5°), and the angle of the burrow entrance to the ground was ~90°. Burrows  
332 with a burrow roof were located on steeper parts of the hillslopes (with an inclination above 5°), and the angle  
333 of the burrow entrance to the ground was ~45°.

334

### 335 **3.6 Calculation of animal-caused and rainfall-caused sediment redistribution**

336 We pairwise compared the DSMs of each scan with the scan saved before and identified 3 types of  
337 sediment redistribution which occurred in the time period between these images. The 3 types of redistribution  
338 were: a) animal caused; b) rainfall-caused; c) both animal and rainfall caused.

339 The animal-caused sediment redistribution occurred when the animal actively reworked sediment  
340 within its burrow. Following five prerequisites had to be met when the sediment redistribution was caused  
341 solely by the animal: (i) as the animal excavates sediment from the entrance, the depth of the entrance must  
342 increase in the second scan; (ii) as the excavated sediment accumulates on the mound, the height of the  
343 mound must increase in the second scan; (iii) as the burrowing might lead to an expansion or a collapse of the  
344 burrow roof, an increase or decrease of the burrow roof must occur between the scans; (iv) as the animal only  
345 digs within his burrow, no changes must occur between the two scans within the **burrow embedding area** by  
346 the animal; (v) no rainfall occurred during this period.

347 The rainfall-caused sediment redistribution was calculated as follows: From the data from the climate  
348 stations (Übernicker et al., 2021a), we calculated the daily precipitation in mm. The sediment redistribution  
349 recorded immediately and within five scans before and after a rainfall event is defined to be the result of the  
350 rainfall event. This was necessary as the climate stations are located up to a 15 km distance from the cameras  
351 (Fig. 1). To attribute sediment redistribution to rainfall event, three preconditions had to be met: (i) A rainfall  
352 event occurred; (ii) sediment is eroded from burrow roof, mound and the **embedding area**; (iii) sediment is  
353 accumulated within the burrow entrance.

354 To attribute sediment redistribution to a combination of animal activity and rainfall, four preconditions  
355 had to be met: (i) A rainfall event occurred; (ii) sediment is eroded from embedding area; (iii) the height of  
356 burrow roof and mound decreased or increased; (iv) the depth of burrow entrance increased.

357 The animal-caused sediment redistribution was calculated as the sediment volume excavated from  
358 the entrance. Animal excavation always increased depth of the burrow entrance. The rainfall-caused sediment  
359 redistribution was calculated as the sediment volume which eroded from the burrow roof and mound. During  
360 a rainfall event, sediment eroding from burrow roof might accumulate within burrow entrances. In this case,  
361 the depth of the burrow entrance decreased. No sediment could erode from the entrance during a rainfall  
362 event. Decreased depth of a burrow entrance always points to sediment redistribution caused by rainfall,  
363 increased depth of burrow entrance always means redistribution by animals. Rainfall-caused redistribution  
364 always occurred before animal-caused redistribution, as without erosion caused by rainfall, the animals did not  
365 need to reconstruct their burrows.

366

### 367 **3.7 Calculation of daily sediment mass balance budget**

368 The volume of the redistributed sediment was calculated daily and was then cumulated from the first  
369 day of monitoring. For the calculation of the daily sediment redistribution, the change in the surface level  
370 detected by the camera was calculated first. For each day, the scans from the day before and after the  
371 respective day were averaged and subtracted. The average standard deviation of the z-coordinate of these  
372 scans was 0.06 cm. As described in Section 2.2., all values with a difference below and above the threshold

373 value of 0.2 cm were set to 0. The redistributed sediment volume was then calculated from the surface change  
 374 for each pixel as follows:

$$375 \text{ Vol}_{redistributed} = (S_b - S_a) * res^2 \quad (6)$$

376 In Eq. (6),  $\text{Vol}_{redistributed}$  ( $\text{cm}^3 \text{ pixel}^{-1}$ ) is the volume of the calculated redistributed sediment,  $S_b$  (cm) the scan  
 377 before,  $S_a$  (cm) the scan after the rainfall event and  $res$  is the spatial resolution (cm). Using the daily volume  
 378 of the redistributed sediment per pixel, we calculated the daily mass balance budget by summing the volume  
 379 of sediment eroding or accumulating within each delineated area.

380

### 381 3.8 Calculation of the overall volume of redistributed sediment after the period of 7 months

382 From the camera data, we calculated the average cumulative volume of redistributed sediment for the  
 383 period of 7 months within burrows ( $\text{Vol}_{burrows}$  ( $\text{cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ )) and burrow embedding ( $\text{Vol}_{embedding}$  ( $\text{cm}^3$   
 384  $\text{cm}^{-2} \text{ 7 months}^{-1}$ )) areas and the average sediment volume redistributed (excavated) by the animal ( $\text{Vol}_{exc}$  ( $\text{cm}^3$   
 385  $\text{cm}^{-2} \text{ 7 months}^{-1}$ )), separately for each site. We estimated the volume of sediment that was redistributed during  
 386 rainfall events due to the presence of the burrow ( $\text{Vol}_{add}$  ( $\text{cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ )).  $\text{Vol}_{add}$  was calculated as the  
 387 difference in the redistributed sediment volume between burrows and burrow embedding areas according to  
 388 Eq. (7).

$$389 \text{ Vol}_{add} = (\text{Vol}_{affected} - \text{Vol}_{unaffected}) \quad (7)$$

390 Additionally, we calculated the average volume of the redistributed sediment per burrow ( $\text{Vol}_{per burrow}$  [ $\text{cm}^3$   
 391  $\text{burrow}^{-1} \text{ 7 months}^{-1}$ ]).

$$392 \text{ Vol}_{per burrow} = (\text{Area}_{burrow} * \text{Vol}) \quad (8)$$

393 In Eq. (8),  $\text{Area}_{burrow}$  ( $\text{cm}^2$ ) is the average size of the burrows that are monitored by the cameras;  $\text{Vol}$  is  $\text{Vol}_{burrow}$   
 394 ( $\text{cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ ),  $\text{Vol}_{exc}$  ( $\text{cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ ) or  $\text{Vol}_{add}$  ( $\text{cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ ).

395 We then upscaled the  $\text{Vol}_{burrow}$  ( $\text{cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ ),  $\text{Vol}_{exc}$  ( $\text{cm}^3 \text{ cm}^{-2} \text{ 7 months}^{-1}$ ) and  $\text{Vol}_{add}$  ( $\text{cm}^3$   
 396  $\text{cm}^{-2} \text{ 7 months}^{-1}$ ) to the hillslope using the following approach. Hillslope-wide upscaling of the results  
 397 generated in this study was performed by using a previous estimation of vertebrate burrow density (Grigusova  
 398 et al., 2021). In this study, the density of burrows was measured in situ within eighty 100  $\text{m}^2$  plots and then  
 399 upscaled to the same hillslopes on which the cameras were located by applying machine-learning methods,  
 400 using the UAV-data as predictors. For upscaling, we applied a random forest model with recursive feature  
 401 elimination. The model was validated by a repeated Leave-One-Out cross validation. The density of vertebrate  
 402 burrows was between 6 and 12 100  $\text{m}^2$  in LC and between 0 and 12 100  $\text{m}^2$  in Pan de Azúcar. Using the  
 403 hillslope-wide predicted vertebrate burrow densities ( $\text{Dens}_{burrow}$  (number of burrows 100  $\text{m}^{-2}$ )) from Grigusova  
 404 et al. 2021, we estimated the volume of redistributed sediment for each pixel of the raster layers ( $\text{Vol}_{per pixel}$   
 405 ( $\text{cm}^3 \text{ m}^{-2} \text{ 7 months}^{-1}$ )) according to Eq. (9):

$$406 \text{ Vol}_{per pixel} = \text{Vol}_{per burrow} * \text{Dens}_{burrow} \quad (9)$$

407 The average hillslope-wide volume of redistributed sediment ( $\text{Vol}_{hillslope-wide}$  ( $\text{m}^3 \text{ ha}^{-1} \text{ 7 months}^{-1}$ )) was  
 408 then estimated as follows:

$$409 \text{ Vol}_{hillslope-wide} = \sum_1^m \text{Vol}_{per pixel} * 0.001 \quad (10)$$

410 In Eq (10),  $m$  is the number of pixels.

411

## 412 4. Results

### 413 4.1 Camera accuracy and data availability

414 The accuracy between the measured extracted sediment volume and sediment volume calculated  
415 from the camera scans was very high (MAE = 0.023 cm<sup>3</sup> cm<sup>-2</sup>, R<sup>2</sup> = 0.77, SD = 0.02 cm<sup>3</sup> cm<sup>-2</sup>, Fig. A3). The  
416 accuracy between the calculated and measured extracted sediment was higher when the two scans taken  
417 before as well as after the extraction of the sediment were averaged and the sediment volume was estimated  
418 using these averaged scans. When calculating the redistributed sediment from solely one scan before and  
419 after extraction, the accuracy slightly decreased (MAE = 0.081 cm<sup>3</sup> cm<sup>-2</sup>, R<sup>2</sup> = 0.64). The cameras tended to  
420 overestimate the volume of redistributed sediment. Six out of eight custom-tailored cameras collected data  
421 over the seven-month period (Table A2). One camera collected data for a period of three months and one  
422 camera stopped working a few days after installation. The quantity of usable point clouds taken at 1 a.m., 5  
423 a.m. and 10 p.m. was higher than of point clouds taken at 8 a.m. Approximately 20% of points was removed  
424 from the point clouds before final analysis due to the high scattering at the point cloud corners. After data  
425 filtering (see Section 3.2.), 1326 scans were usable and for 86% of the days, at least one usable scan was  
426 available. The usable scans were distributed continuously within the monitoring period.


427

### 428 4.3 Mass balance of redistributed sediment

429 The cameras detected (i) sediment redistribution directly following rainfall events and (ii) due to the  
430 burrowing activity in times without rainfall (Fig. 3, A4 and A5). In all cases, burrows (entrance, burrow roof and  
431 mound) exhibited higher sediment redistribution rates than burrow embedding areas. In addition, the volume  
432 of redistributed sediment by animal activity was higher after a rainfall event occurred.

433 In the following, the dynamics are exemplary explained for four cameras. Animal burrowing activity  
434 was detected seven times by the camera LC NU (Fig. 4a, A4, A5) during the monitoring period, by an increase  
435 in sediment volume in the area delineated as mound. Simultaneously, the burrow entrance showed signs of  
436 modification and sediment accumulation, but these changes were less clear. Overall, the volume of the  
437 excavated soil varied. From April until June, up to 0.5 cm<sup>3</sup> cm<sup>-2</sup> of sediment was excavated by the animal and  
438 accumulated on the mound. From June until September, animal burrowing activity was detected at four time  
439 slots (5 June 2019, 9 June 2019, 1 July 2019 and 18 August 2019) and sediment volume of up to 2 cm<sup>3</sup> cm<sup>-2</sup>  
440 accumulated each time on the mound, burrow roof and within the entrance. During the rainfall events of up to  
441 20 mm day<sup>-1</sup> on 16 June 2019, 27 mm day<sup>-1</sup> on 29 June 2019 and 7 mm day<sup>-1</sup> on 13 July 2019, sediment  
442 volume of up to 4 cm<sup>3</sup> cm<sup>-2</sup> eroded, especially from the burrow roof and the mound while a sediment volume  
443 of up to 1 cm<sup>3</sup> cm<sup>-2</sup> accumulated within the entrance during each rainfall event. Camera LC-SL (Fig. A4, A5)  
444 showed burrowing activities eight times and sediment volumes of up to 3 cm<sup>3</sup> cm<sup>-2</sup> accumulated within the  
445 entrance and burrow roof. The camera detected sediment erosion of up to 2 cm<sup>3</sup> cm<sup>-2</sup> after a rainfall event of  
446 27 mm day<sup>-1</sup> on 27 July 2019. On the south-upper hillslope, the camera detected animal burrowing activity six  
447 times, with a sediment accumulation of up to 3 cm<sup>3</sup> cm<sup>-2</sup> (Fig. A2 and A3).

448 In contrast, camera PdA-NU pointed to animal burrowing activity up to 15 times where up to 1 cm<sup>3</sup> cm<sup>-2</sup>  
449 of sediment volume was redistributed from the entrance to the mound (Fig. 4b, A4, A5). At the end of June  
450 on 27 June 2019, a rainfall event of 1.5 mm day<sup>-1</sup> occurred and up to 2 cm<sup>3</sup> cm<sup>-2</sup> of sediment eroded from the  
451 burrow roof and accumulated within the burrow entrance. We observed increased sediment redistribution by  
452 the animal after the rainfall events. Camera PdA-SL evenly revealed animal burrowing activity up to 15 times  
453 ((Fig. A4, A5)). The burrowing had a strong effect on the sediment redistribution. The rainfall event of 1.5 mm  
454 day<sup>-1</sup> on 27 June 2019 did not cause any detectable surface change.



455

456

457 **Figure 4.** Examples of the mass balance of redistributed sediment for burrows and burrow embedding areas  
 458 (a) The record of the camera on the upper north-facing hillslope in La Campana showed that larger rainfall  
 459 events cause a negative sediment balance (sediment loss), followed by a phase of positive sediment mass  
 460 balance after approximately 3 days due to sediment excavation; (b) The record of the camera on the upper  
 461 north-facing in Pan de Azúcar hillslope showed a similar pattern to the camera on the upper north-facing  
 462 hillslope, but the phase of positive mass balance was delayed in comparison. The blue line is the daily  
 463 precipitation in mm day<sup>-1</sup>, and “X” marks the days at which animal burrowing activity was detected. Mass  
 464 balances for all cameras are displayed in Fig. A2 and A3.


465

466 The analysis of cumulative volume of the redistributed sediment caused by burrowing animal activity  
 467 and rainfall over the monitored period of seven months for all eight cameras showed a heterogeneous pattern.


468 In LC, the cumulative volume of the sediment excavated by the animal within the burrow roof and  
 469 mound increased continuously (Fig. 5, A7). Especially between the rainfall events from June until August, a  
 470 cumulative volume of on average 6.5 cm<sup>3</sup> cm<sup>-2</sup> was excavated by the animal. We calculated that, on average,  
 471 8.53 cm<sup>3</sup> cm<sup>-2</sup> cumulatively eroded from the burrow roof and mound; while 2.44 cm<sup>3</sup> cm<sup>-2</sup> sediment volume  
 472 accumulated within the entrance (Fig. 5, A7). These results indicate that 28% of sediment eroding from the  
 473 burrow roof accumulated within the entrance, while over 62% of sediment eroded downhill. Averaged over all

474 camera scans, 338% more sediment was redistributed by rain within burrow compared to the burrow  
475 embedding area (Fig. 6).

476 In PdA, cameras continuously detected animal burrowing activity and excavation of the sediment (Fig.  
477 A7). The volume of the detected excavated sediment increased steadily within all cameras. The cumulative  
478 sediment accumulation surpasses the sediment eroded due to the rainfall. The volume of the sediment eroded  
479 within the burrows was 40% higher than within the burrow embedding areas. The results show that  
480 approximately 50% of the eroded sediment accumulated within the entrance (Fig. 6).  
481



482  
483 **Figure 5.** Examples of the cumulative volume of redistributed sediment within burrows and burrow embedding  
484 areas caused by animal burrowing activity or rainfall in mediterranean La Campana: (a) Upper north-facing  
485 hillslope; (b) Lower south-facing hillslope. Positive values indicate sediment accumulation. Negative values  
486 indicate sediment erosion. E is the burrow entrance; M is the mound; R is burrow roof; EM is the burrow  
487 embedding area. Cumulative volumes for all cameras are in Fig. A7.



488

489 **Figure 6.** Cumulative volume of the redistributed sediment for all cameras. Positive values indicate sediment  
 490 accumulation. Negative values indicate sediment erosion. Whiskers indicate the median of sediment  
 491 redistribution. E is the burrow entrance; M the mound; R is the burrow roof; EM is burrow embedding area; LC  
 492 stands for National Park La Campana in the mediterranean climate zone; PdA stands for National Park PdA in  
 493 the arid climate zone.

494

#### 495 **4.4 Volume of redistributed sediment**

496 The average size of the burrows was 84.36 cm<sup>2</sup> (SD = 32.54 cm<sup>2</sup>) in LC and 91.35 cm<sup>2</sup> in PdA (SD = 8.53  
 497 cm<sup>2</sup>). The animals burrowed on average 1.2 times month<sup>-1</sup> in LC and 2.33 times month<sup>-1</sup> in PdA. The volume  
 498 of the excavated sediment was 102.22 cm<sup>3</sup> month<sup>-1</sup> in LC and 124.89 cm<sup>3</sup> month<sup>-1</sup> in PdA. Each time the  
 499 animals burrowed, they excavated 42 cm<sup>3</sup> sediment volume in LC and 14.33 cm<sup>3</sup> sediment volume in PdA.  
 500 The burrowing intensity increased in winter after the rainfall occurrences in LC and stayed constant during the  
 501 whole monitoring period in PdA. The burrows deteriorate after rainfall events with a rate of 73.03 cm<sup>3</sup> month<sup>-1</sup>  
 502 or 63.90 cm<sup>3</sup> event<sup>-1</sup> in LC and 10.53 cm<sup>3</sup> month or 24.57 cm<sup>3</sup> event<sup>-1</sup>.

503

504 The overall volume of the sediment excavated by the animal and redistributed during rainfall events  
 505 varied between the sites (Table 1). The volume of the sediment redistributed by the animal was lower in LC  
 506 than in PdA. However, on the hillslope scale, a higher total area-wide volume of excavation was calculated for  
 LC compared to PdA, due to the higher burrow density in LC. The volume of the sediment redistributed within




507 burrows during rainfall events was higher in LC than in PdA. The volume of additionally redistributed sediment  
 508 due to the presence of burrows was higher in LC than in PdA (Table 1, Fig. 7).

509

510 **Table 1.** Summary of the volume of redistributed sediment, according to area and disturbance type.  $Vol_{exc}$   
 511 describes volume of the sediment excavated by the animals.  $Vol_{burrow}$  describes volume of the sediment  
 512 redistributed during rainfall events within burrows.  $Vol_{add}$  describes the difference in redistributed sediment  
 513 volume within burrows and burrow embedding areas during rainfall.

<b>Disturbance</b>	<b>Area</b>	<b>PdA</b>	<b>LC</b>
<b><math>Vol_{exc}</math></b>	Burrow	9.57 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>	8.53 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>
	Per burrow	874.22 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>	715.52 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>
	Hillslope-wide	0.11 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>	0.39 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>
<b><math>Vol_{burrow}</math></b>	Burrow	-1.15 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>	-6.09 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>
	Per burrow	-73.71 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>	-511.22 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>
	Hillslope-wide	-0.03 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>	-0.28 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>
<b><math>Vol_{add}</math></b>	Burrow	-0.69 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>	-4.30 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>
	Per burrow	-28.21 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>	-361.20 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>
	Hillslope-wide	-0.01 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>	-0.2 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>

514



515

516

517

518

519

520

521

522

523

524

525

526

**Figure 7.** Example of the hillslope-wide volume of redistributed sediment for a time period of 7 months on the south-facing hillslope in La Campana: (a) Density of burrows as estimated by Grigusova et al. (2021); (b) Volume of the sediment excavated by the animals; (c) Volume of the sediment redistributed during rainfall events within burrows; (d) Volume of additionally redistributed sediment during rainfall events due to the presence of the burrows. The values were calculated per burrow as stated in Section 3.7. by subtracting the sediment volume redistributed within burrows from the sediment volume redistributed within burrow embedding area and then upscaled. The letters in brackets indicate if the upscaling was conducted using data from burrows or burrow embedding areas. “B” stands for burrow. By “EM-B”, the redistribution calculated within burrow embedding areas was subtracted from the redistribution calculated within burrows to obtain the additional volume of redistributed sediment due to the burrows’ presence.

527

## 528 **5. Discussion**

529 Our results showed that the custom-made ToF device is a suitable tool for high-resolution, automated  
530 monitoring of surface changes, applicable also in remote areas. The ability of a continuous observation of  
531 sediment redistribution over a longer time during our study provided new insights into the importance of  
532 burrowing animals for sediment redistribution. Our research reveals that the presence of vertebrate burrows  
533 increases hillslope sediment redistribution rates much more than previously assumed (up to 208%). We  
534 showed that the quantity of animal-related sediment redistribution, however, varied with rainfall occurrence,  
535 with an increase in sediment redistribution between 40% in the arid research area and 338% percent in the  
536 mediterranean research area.

537

### 538 **5.1 Suitability of the ToF method for surface monitoring**

539 The here proposed monitoring technique enables an automatic monitoring of surface changes on a  
540 microtopographic scale, and its measurement continuity allows for the analysis of ongoing  
541 biogeomorphological processes in high temporal resolution.

542 With regard to the costs, measurement frequency and sampling autonomy, the custom-made ToF  
543 device stands in contrast to earlier studies that used laser scanning technology to monitor microtopographic  
544 changes (Table A5). Previous studies mainly applied expensive laser scanning for the estimation of sediment  
545 redistribution, and the research sites had to be personally revisited for each of the measurements (Eltner et  
546 al., 2016a; Eltner et al., 2016b; Hänsel et al., 2016; Nasermoaddeli and Pasche, 2008). The estimated costs  
547 in studies using time-lapse photogrammetry were similar to our study (up to 5000 USD) (Blanch et al., 2021;  
548 Eltner et al., 2017; Galland et al., 2016; James and Robson, 2014; Kromer et al., 2019; Mallalieu et al., 2017).  
549 However, for time-lapse monitoring, several devices needing different viewing angles increases installation  
550 efforts significantly.

551 In terms of data quality, our ToF device is more precise or comparable to those employed in other  
552 studies. The accuracy of the camera ( $R^2 = 0.77$ ) was in the range of previous studies ( $R^2 = 0.26$ – $0.83$  (Eitel et  
553 al., 2011), Table A5). The horizontal point spacing of our cameras was 0.32 cm, and the maximum number of  
554 points per  $\text{cm}^2$  was 8.5. These values are similar to previous studies in which the used devices had a horizontal  
555 point spacing in the range of 0.25–0.57 cm (Kaiser et al., 2014; Nasermoaddeli and Pasche, 2008)) (Table  
556 A5), and the maximum number of points per  $\text{cm}^2$  in a range of 1 point–25 points  $\text{cm}^{-2}$  (Eitel et al., 2011; Longoni  
557 et al., 2016) (Table A5).

558 Our cameras tended to slightly overestimate or underestimate the volume of redistributed sediment.  
559 This error occurs when the pulse reflects from several vertical objects such as walls or, in our case, branches  
560 or stones and then enters the camera sensor. This phenomenon was also observed in previous studies  
561 applying laser scanners and is inevitable if the goal is to study surface changes under natural field conditions  
562 (Ashcroft et al., 2014; Kukko and Hyypä, 2009). During operation of the cameras, we learnt that our newly  
563 developed instruments are particularly capable of delivering usable scans at night. This is likely due to the  
564 strong scattered sunlight reaching the camera sensor during the day, blurring the data (Li, 2014). Thus, in  
565 future studies, we recommend focusing on nocturnal operation to prevent light contamination from the  
566 surroundings.

567 We could thus prove that ToF cameras are a suitable and cost-effective method for a continuous  
568 monitoring of sediment redistribution at a microtopographic scale without the need of time, labour and cost  
569 intensive laser scanning/time-lapse photogrammetry campaigns.


570

## 571 5.2. Sediment Redistribution

572 Our research reveals that the presence of vertebrate burrows generally increases hillslope sediment  
573 redistribution. We show, however, that the ratio between the sediment redistribution caused by rainfall within  
574 burrow and burrow embedding areas varies between climate zones. Sediment redistribution within burrow  
575 areas was 40% higher at the arid research site, and at the mediterranean research site, it was 338% higher  
576 when compared to burrow embedding area (Table A6).

577 By monitoring microtopographical changes in a high spatio-temporal resolution, we found that the  
578 occurrence of larger rainfall events played a two-fold, accelerating role in influencing sediment redistribution  
579 (Fig. 5, A4). Firstly, rainfall-runoff eroded burrow material caused increased sediment loss. This was followed  
580 by animal burrowing activity after the rainfall. This means that rainfall triggered animal burrowing activity which  
581 was very likely related to a lower burrowing resistance of the soil due to the increased soil moisture (Herbst  
582 and Bennett, 2006; Romañach et al., 2005; Rutin, 1996). This double feedback led to frequently occurring but  
583 small redistribution rates. However, cumulatively, the mechanism increased downhill sediment fluxes. Previous  
584 studies most likely missed this low magnitude but frequent surface processes due to a lower monitoring  
585 duration and frequency, or artificial laboratory conditions, and thus, did not quantify the full volume of  
586 redistributed sediment associated with burrowing activity. To quantify all occurred sediment redistribution  
587 processes, a continuous surface monitoring, like the here presented, is needed.

588



589

590 **Figure 8.** Scheme of animal-driven and rainfall-driven sediment redistribution processes in both investigated  
591 climate zones: (a) Describes the initial surface of the burrow before the start of a sediment redistribution  
592 process, and (b) the animal excavation process in the arid climate zone. Here, due to rarely occurring rainfall  
593 events, sediment redistribution is mostly controlled by the animal burrowing activity; (c) describes the initial  
594 burrow surface in the mediterranean climate zone, (d) the process of sediment redistribution during a rainfall  
595 event and (e) the subsequent animal burrowing activity. Burrowing is triggered by decreased soil resistance  
596 due to the increased soil moisture after rainfall as well as by sediment accumulation within the burrow's  
597 entrance. Burrowing activity leads to a new supply of sediment being excavated to the surface. In the  
598 mediterranean climate zone, sediment redistribution is controlled by both animal burrowing activity and rainfall.  
599 The alternating excavation and erosion process ultimately lead to an increase in redistribution rates.

600

601 Our results indicate an up to 338% increase in the sediment volume redistributed during rainfall events  
602 measured within burrows when compared to burrow embedding areas. In contrast to our result, the maximum  
603 increase estimated in previous studies was 208% (Table A6, (Imeson and Kwaad, 1976). The two climate  
604 zones also show different patterns: In the mediterranean climate, the contribution of animals' (vertebrates')  
605 burrowing activity appear larger than previously observed by using field methods such as erosion pins or  
606 splash traps (from -3% until - 208%, Table A6, (Black and Montgomery, 1991; Hazelhoff et al., 1981; Imeson  
607 and Kwaad, 1976). In contrast, in arid PdA, our study found a much smaller increase (40%, Table A6) in the  
608 sediment volume redistributed during rainfall events measured within burrows when compared to burrow  
609 embedding areas. This is lower than previously estimated (125%, Table A6, (Black and Montgomery, 1991).  
610 However, solely one rainfall event above 0.2 mm day<sup>-1</sup> occurred during our monitoring period. Hence, we  
611 conclude that the contribution of burrowing activity of animals to hillslope sediment transport is much larger in  
612 areas with frequent rainfall events than previously thought, while it has been realistically estimated by previous  
613 studies for areas with rare rainfall events (Table A6).

614 Magnitudes of sediment volume redistributed within burrows similar to our results were previously  
615 obtained solely in studies applying rainfall simulators. These studies estimated an increase in the volume of  
616 sediment redistributed during rainfall events, measured within burrows when compared to burrow embedding  
617 areas, to be between 205% and 473% (Table A6, (Chen et al., 2021; Li et al., 2018). However, a rainfall  
618 simulator can only provide data on surface processes within a plot of a few m<sup>2</sup> in size and under ideal laboratory  
619 conditions while ignoring the uphill microtopography, vegetation cover and distribution (Iserloh et al., 2013),  
620 which were shown to reduce erosion rates. More importantly, the rainfall intensity on hillslopes decreases with  
621 (i) the angle of incidence of the rain, (ii) the inclination of the surface and (iii) the relative orientation of the  
622 sloping surface to the rain vector (Sharon, 1980). When simulating a rainfall event with the same rainfall volume  
623 as in the field, the rain is induced directly over the treated surface and has thus a higher velocity which leads  
624 to an increased splash erosion than under natural conditions (Iserloh et al., 2013). We thus propose that the  
625 rainfall experiments overestimate the erosion rate while the correct erosion rate can be measured solely under  
626 field conditions.

627 Cumulative sediment redistribution within burrow roof, mound and entrance was, on average, 28%  
628 lower than cumulative sediment redistribution only within the mound and the burrow roof (Figure A7). These  
629 results suggest that 28% of the eroded sediment from animal mounds and burrow roofs is re-accumulated  
630 within the burrow entrance during rainfall-runoff events, and the remaining 62% is incorporated into overall  
631 hillslope sediment flux. Our numbers contrast with previous studies, which quantified that about 58% of the

632 sediment excavated by animals will accumulate back in the burrow entrance and only 42% is incorporated to  
 633 downhill sediment flux (Andersen, 1987; Reichman and Seabloom, 2002). Hence, our results indicate not only  
 634 higher redistribution rates within burrows by burrowing animals but also point to much higher supply of  
 635 sediment to the downhill sediment flux than previously thought.

636 Our cost-effective ToF device provides data on surface changes in a high spatio-temporal resolution.  
 637 The high temporal resolution was able to unravel ongoing low magnitude but frequent animal excavation and  
 638 erosion processes. The high spatial resolution enabled us to estimate the exact volume of sediment fluxes  
 639 from the burrows downhill. The here presented results indicate that the contribution of burrowing animals on  
 640 the burrow as well as on the hillslope scale was much higher than previously assumed. Our results can be  
 641 integrated into long-term soil erosion models that rely on soil processes and improve their accuracy by  
 642 including animal-induced surface processes on microtopographical scales in their algorithms.

643  
 644

645 **Funding:** This study was funded by the German Research Foundation, DFG [grant numbers  
 646 BE1780/52-1, LA3521/1-1, FA 925/12-1, BR 1293-18-1], and is part of the DFG Priority Programme  
 647 SPP 1803: EarthShape: Earth Surface Shaping by Biota, sub-project “Effects of bioturbation on rates  
 648 of vertical and horizontal sediment and nutrient fluxes”.

649 **Institutional Review Board Statement:** Not applicable.

650 **Informed Consent Statement:** Not applicable.

651 **Acknowledgments:** We thank CONAF for the kind support provided during our field campaign.

652 **Competing interests:** There is no conflict of interest.

653 **Author contribution:** JB, AL and SA planned the campaign; PG and SA performed the measurements; PG  
 654 analysed the data and wrote the manuscript draft; AL, JB, NF, RB, KÜ, LP, CR, DK and PP reviewed and edited  
 655 the manuscript.

656 **Code/Data availability:** Code and all raw data can be provided by the corresponding author upon request.

657

## 658 Appendices

659 **Table A1.** List of abbreviations

$\alpha$ [°]	Tilt angle of the camera
$b$ [°]	Surface inclination
$\Omega$	Threshold value for the scan scattering error
<b>B</b>	Burrow
$\text{Area}_{\text{burrow}}$	mean in the field measured size of the burrows which are monitored
<b>Area</b>	total surface area monitored by the camera
<b>BD</b>	Bulk density
$c$ [m/s]	Speed of light
<b>D</b>	Distance from the camera to the object
$\text{Dens}_{\text{burrow}}$	Burrow density
<b>DSM</b>	Digital surface model
$\text{DSM}_{\text{after}}$	DSM calculated from the scan taken after the extraction
$\text{DSM}_{\text{before}}$	DSM calculated from the scan taken before the extraction

<b>EM</b>	Burrow embedding area
<b>Entrance</b>	entrance to the animal burrow
<b>g [-]</b>	ratio [-] of the reflected photons to all photons
<b>LC</b>	National Park LC
<b>LC-NL</b>	Camera in LC on the lower north-facing hillslope
<b>LC-NU</b>	Camera in LC on the upper north-facing hillslope
<b>LC-SL</b>	Camera in LC on the lower south-facing hillslope
<b>LC-SU</b>	Camera in LC on the upper south-facing hillslope
<b>MAE</b>	Mean absolute error
<b>MAP [°]</b>	Mean annual precipitation
<b>m.a.s.l.</b>	Meters above sea level
<b>MAT</b>	Mean annual temperature
<b>mClay [%]</b>	Mean content of clay
<b>mean<sub>z-coordinate</sub></b>	Mean value of the z-coordinates
<b>Mound</b>	the sediment excavated by the animal while digging the burrow
<b>mSand [%]</b>	Mean content of sand
<b>mSilt [%]</b>	Mean content of silt
<b>n</b>	Number of scans
<b>PdA</b>	National Park Pan de Azúcar
<b>PdA-NL</b>	Camera in PdA on the lower north-facing hillslope
<b>PdA-NU</b>	Camera in PdA on the upper north-facing hillslope
<b>PdA-SL</b>	Camera in PdA on the lower south-facing hillslope
<b>PdA-SU</b>	Camera in PdA on the upper south-facing hillslope
<b>Res</b>	Resolution
<b>Roof</b>	sediment pushed aside and uphill the entrance during burrow creation
<b>S<sub>a</sub></b>	scan after the rainfall event
<b>S<sub>b</sub></b>	scan before the rainfall event
<b>SBC</b>	Single board computer
<b>sd<sub>z-coordinate</sub></b>	standard deviation of the z-coordinates
<b>SSH</b>	Secure shell
<b>t [s]</b>	Overall time of camera illumination
<b>TOC [%]</b>	Total organic carbon
<b>ToF</b>	Time-of-Flight
<b>Vol<sub>burrow</sub></b>	volume of redistributed sediment within burrow
<b>Vol<sub>detected</sub></b>	volume of the extracted sediment as detected by the camera
<b>Vol<sub>add</sub></b>	difference in redistributed sediment volume between burrows and burrow embedding areas
<b>Vol<sub>exc</sub></b>	Volume of the sediment excavated by the animal
<b>Vol<sub>hillslope-wide</sub></b>	Hillslope-wide volume of redistributed sediment
<b>Vol<sub>measured</sub></b>	volume of the extracted sediment measured by the measuring cup

<b>Vol<sub>per burrow</sub></b>	Volume of redistributed sediment per burrow
<b>Vol<sub>per pixel</sub></b>	Volume of redistributed sediment per pixel
<b>Vol<sub>redistributed</sub></b>	volume of the calculated redistributed sediment
<b>Vol<sub>embedding</sub></b>	volume of redistributed sediment within <b>burrow embedding area</b>
<b>y<sub>i</sub></b>	distance of the point to the point of origin at the camera nadir
<b>Z<sub>cor</sub></b>	Corrected z-coordinate
<b>Z<sub>uncor</sub></b>	Uncorrected z-coordinate

660

661 **Table A2.** Number of usable scans for each camera

Camera	Latitude	Longitude	Number of scans	Percentage of usable scans taken at 1am / 5am / 8am / 10pm	Time period
<b>PdA-NU</b>	-25.98131	-70.61666	238	29 / 27 / 20 / 24	18.3.-18.9.
<b>PdA-NL</b>	-25.98277	-70.61278	52	24 / 0 / 40 / 36	27.3.-31.5
<b>PdA-SU</b>	-25.97477	-70.61641	351	30 / 26 / 32 / 11	16.3.-19.9.
<b>PdA-SL</b>	-25.97177	-70.61409	167	48 / 38 / 7 / 8	16.3.-19.9.
<b>LC-NU</b>	-32.95230	-71.06231	215	37 / 20 / 8 / 33	9.3.-9.9.
<b>LC-NL</b>	-32.93928	-71.08613	3	-	6.3.-12.9
<b>LC-SU</b>	-32.93078	-71.09066	160	22 / 28 / 26 / 25	28.3.-22.5
<b>LC-SL</b>	-32.93110	-71.08987	167	27 / 25 / 22 / 26	16.3.-19.9.

662

663 **Table A3.** Summary of the volume of redistributed sediment, according to area and disturbance type. Vol<sub>exc</sub>  
664 describes volume of the sediment excavated by the animals. Vol<sub>burrow</sub> describes volume of the sediment  
665 redistributed during rainfall events within **burrows**. Vol<sub>add</sub> describes the difference in redistributed sediment  
666 volume within **burrows and burrow embedding area** during rainfall.

Disturbance	Area	PdA	LC
<b>Vol<sub>exc</sub></b>	<b>Burrow</b>	16.41 cm <sup>3</sup> cm <sup>-2</sup> year <sup>-1</sup>	14.62 cm <sup>3</sup> cm <sup>-2</sup> year <sup>-1</sup>
	Per burrow	1498.66 cm <sup>3</sup> burrow <sup>-1</sup> year <sup>-1</sup>	1226.61 cm <sup>3</sup> burrow <sup>-1</sup> year <sup>-1</sup>
	Hillslope-wide	0.18 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>	0.67 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
<b>Vol<sub>burrow</sub></b>	<b>Burrow</b>	-1.97 cm <sup>3</sup> cm <sup>-2</sup> year <sup>-1</sup>	-10.44 cm <sup>3</sup> cm <sup>-2</sup> year <sup>-1</sup>
	Per burrow	-126.36 cm <sup>3</sup> burrow <sup>-1</sup> year <sup>-1</sup>	-876.38 cm <sup>3</sup> burrow <sup>-1</sup> year <sup>-1</sup>
	Hillslope-wide	-0.05 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>	-0.48 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
<b>Vol<sub>add</sub></b>	<b>Burrow</b>	-1.18 cm <sup>3</sup> cm <sup>-2</sup> year <sup>-1</sup>	-7.37 cm <sup>3</sup> cm <sup>-2</sup> year <sup>-1</sup>
	Per burrow	-48.36 cm <sup>3</sup> burrow <sup>-1</sup> year <sup>-1</sup>	-619.2 cm <sup>3</sup> burrow <sup>-1</sup> year <sup>-1</sup>
	Hillslope-wide	-0.02 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>	-0.34 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>

667

668

669 **Table A4.** Summary of the volume of redistributed sediment, according to area and disturbance type. Vol<sub>exc</sub>  
670 describes volume of the sediment excavated by the animals. Vol<sub>burrow</sub> describes volume of the sediment



671 redistributed during rainfall events **within burrows**. Vol<sub>add</sub> describes the difference in redistributed sediment  
 672 volume within **burrows and burrow embedding areas** during rainfall.

<b>Disturbance</b>	<b>Area</b>	<b>PdA</b>	<b>LC</b>
<b>Vol<sub>exc</sub></b>	<b>Burrow</b>	9.57 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>	8.53 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>
	Per burrow	874.22 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>	715.52 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>
	Hillslope-wide	0.11 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>	0.39 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>
<b>Vol<sub>burrow</sub></b>	<b>Burrow</b>	-1.15 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>	-6.09 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>
	Per burrow	-73.71 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>	-511.22 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>
	Hillslope-wide	-0.03 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>	-0.28 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>
<b>Vol<sub>add</sub></b>	<b>Burrow</b>	-0.69 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>	-4.30 cm <sup>3</sup> cm <sup>-2</sup> 7 months <sup>-1</sup>
	Per burrow	-28.21 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>	-361.20 cm <sup>3</sup> burrow <sup>-1</sup> 7 months <sup>-1</sup>
	Hillslope-wide	-0.01 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>	-0.2 m <sup>3</sup> ha <sup>-1</sup> 7 months <sup>-1</sup>

673  
 674 **Table A5.** Review of studies which used laser scanners for the estimation of surface processes.

<b>Reference</b>	<b>R<sup>2</sup></b>	<b>Error</b>	<b>Horizontal point spacing</b>	<b>Points per cm<sup>2</sup></b>	<b>Model</b>	<b>Price</b>
<b>Our results</b>	0.77	0.15 cm	0.32 cm	8.5	Texas Instruments OPT3101	900 USD
<b>(Eitel et al., 2011)</b>	0.23- 0.86	0.07 cm	NA	25	Leica ScanStation 2	102 375 USD
<b>(Eltner et al., 2013)</b>	NA	0.4 cm	NA	6.4	Riegl LMS- Z420i	16 795 USD
<b>(Kaiser et al., 2014)</b>	NA	NA	0.57 cm	NA	Riegl LMS- Z420i	16 795 USD
<b>(Longoni et al., 2016)</b>	NA	NA	NA	1	Riegl LMS- Z420i	16 795 USD
<b>(Morris et al., 2011)</b>	NA	0.5 cm	NA	NA	Maptek I-Site 4400LR	240 000 USD
<b>(Nasermoaddeli and Pasche, 2008)</b>	NA	0.2 cm	0.25 cm	NA	Leica Cyrax HDS 2500	4500 USD
<b>(Thomsen et al., 2015)</b>	NA	NA	0.4 cm	NA	Leica ScanStation 2	102 375 USD

675  
 676 **Table A6.** Review of studies which estimated the sediment redistribution within **burrows and burrow embedding**  
 677 **areas** and the proposed impact.

Reference	Climate	Animals	Method	Monitoring period	Frequency	Burrows	Burrow embedding area	Impact
(Imeson and Kwaad, 1976)	continental	rodents	erosion pins	15 months	monthly	20 mm		NA
(Imeson and Kwaad, 1976)	continental	rodents	splash boards	15 months	monthly	91.75g 24.49 cm <sup>-2</sup> = 3.75 cm <sup>3</sup> cm <sup>-2</sup>	94g	-3%
(Imeson and Kwaad, 1976)	continental	rodents	rainfall simulation (7.5 cm / hour intensity)	One-time measurement	NA	0.2 g – 0.73 g	0.009 g – 0.23 g	+208 %
(Imeson, 1977)	continental	vertebrates	rainfall simulation	One-time measurement	NA	0.18-0.3 100 J <sup>-1</sup> m <sup>-2</sup> rain	0.146 100 J <sup>-1</sup> m <sup>-2</sup> rain	+123 %
(Hazelhoff et al., 1981)	continental	earthworms	splash traps	12 months	monthly	NA	NA	+180 %
(Black and Montgomery, 1991)	arid	pocket gopher	erosion pins	10 months	2 months	NA	NA	+125 %
(Hakonson, 1999)	temperate	pocket gophers	rainfall simulator (60 mm / hour)	2 years	2 – 3 weeks	2.4 – 8.7 mg ha <sup>-1</sup>	4.4 – 15 mg ha <sup>-1</sup>	-43%
(Li et al., 2018)	temperate	mole crickets	rainfall simulation (36 mm / hour)	One time measurement	15 measurements	22.1 g 115 cm <sup>-2</sup> = 5.2 cm <sup>3</sup> cm <sup>-2</sup>	5 g 123 cm <sup>-2</sup> = 1.09 cm <sup>3</sup> cm <sup>-2</sup>	+473 %
(Li et al., 2018)	temperate	mole crickets	rainfall simulation (36 mm / hour)	One time measurement	15 measurements	35.3 g 220.5 cm <sup>-2</sup> = 6.24 cm <sup>3</sup> cm <sup>-2</sup>	5 g 123 cm <sup>-2</sup> = 1.09 cm <sup>3</sup> cm <sup>-2</sup>	+473 %

(Chen et al., 2021)	lab chinese zocor	rainfall simulation (80 mm / hour)	One-time measurements	3 measurements	2,69 g cm <sup>-2</sup> = 2.69 cm <sup>3</sup> cm <sup>-2</sup>	0,88 g cm <sup>-2</sup> = 0.88 cm <sup>3</sup> cm <sup>-2</sup>	+205 %
---------------------	-------------------------	--	--------------------------	-------------------	--	--	--------

678

679

680 **Table A7.** Review of studies which estimated the sediment redistribution within burrows, average burrow  
681 density as found in the literature and area-wide yearly contribution of burrowing animals to sediment  
682 redistribution.

Climate	Animals	Burrows	Average burrow density	Average burrow size	Area-wide redistribution
Continental	rodents	91.75g cm <sup>-2</sup> = 3.75 cm <sup>3</sup> cm <sup>-2</sup> (Imeson and Kwaad, 1976)	24.49 14 625 m <sup>-2</sup> = 0.02 m <sup>-2</sup> (Pang and Guo, 2017)	24.49 cm <sup>2</sup> (Imeson and Kwaad, 1976)	0.183 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
Temperate	mole crickets	22.1 g cm <sup>-2</sup> = 5.2 cm <sup>3</sup> cm <sup>-2</sup> (Li et al., 2018)	115 cm <sup>2</sup> 405 ha <sup>-1</sup> (Castner and Fowler, 1984)	115 cm <sup>2</sup> (Li et al., 2018)	0.24 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
Temperate	mole crickets	35.3 g cm <sup>-2</sup> = 6.24 cm <sup>3</sup> cm <sup>-2</sup> (Li et al., 2018)	220.5 cm <sup>2</sup> 405 ha <sup>-1</sup> (Castner and Fowler, 1984)	220.5 cm <sup>2</sup> (Li et al., 2018)	0.56 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
Lab	chinese zocor	2,69 g cm <sup>-2</sup> = 2.69 cm <sup>3</sup> cm <sup>-2</sup> (Chen et al., 2021)	94.69 2500m <sup>-2</sup> = 0.04 m <sup>-2</sup> = 400 ha <sup>-1</sup>	1256 cm <sup>2</sup>	1.35 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>


683

684 **Table A8.** Review of studies which estimated the volume of sediment excavated by burrowing animals.

Climate	Animals	Method	Monitoring period	Frequency	volume of the excavated sediment
(Black and Montgomery, 1991)	and arid porcupines	mound volume	3 years	yearly	0.2 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
(Black and Montgomery, 1991)	and arid isopods	mound volume	3 years	yearly	0.11 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>


<b>(Black and Montgomery, 1991)</b>	arid	pocket gopher	mound volume	2 years	3 model runs	0.05 – 0.11 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
<b>(Rutin, 1996)</b>	subtropical	scorpions	mound volume	6 months	2-29 days	0.42 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
<b>(Hall et al., 1999)</b>	alpine	rodents	mound volume	1 year	yearly	0.02 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
<b>(Hall et al., 1999)</b>	alpine	bears	mound volume	1 year	yearly	0.49 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>
<b>(Yoo et al., 2005)</b>	arid	pocket gopher	mound volume	1 year	One model run	0.1-0.2 m <sup>3</sup> ha <sup>-1</sup> year <sup>-1</sup>

685  
686




687  
688  
689  
690  
691  
692

**Figure A1.** Standard deviation of the z-coordinate of unprocessed five scans showed exemplary for the camera on the upper north-facing hillside. SD is standard deviation. The error increases with distance from the camera nadir point. The standard deviation was here calculated from scans before any corrections.



693


694 **Figure A2.** Delineation of the areas. The point of origin of the coordinate system is at the camera nadir. Depth  
 695 is the distance between the surface and the camera. Red is the outline of the burrow entrance. Green is the  
 696 outline of mound. Orange is the outline of burrow roof. Area which is not outlined is **burrow embedding area**.  
 697 Arrow indicates downhill direction of the hillslope. (a) LC-NU. (b) LC-NL (c) LC-SU. (d) LC-SL. (e) PdA-NU. (f)  
 698 PdA-NL. (g) PdA-SU. (h) PdA-SL.



699

700 **Figure A3.** a) Estimation of Time-of-Flight camera accuracy based on averaging two surface scans before and  
 701 after the sediment extraction under controlled conditions. The x-axis shows the exact sediment volume  
 702 measured with a cup. The y-axis represents the volume of the sediment calculated from the camera scans  
 703 (according to Equation (4)). The blue line is the linear regression calculated from the measured and detected  
 704 volume. The green shadow shows the confidence interval of 95% for the linear regression slope.  $^{***}p \leq 0.001$ .  
 705 MAE is the mean absolute error, SD is standard deviation and  $R^2$  the coefficient of determination. b) Measured  
 706 sediment volume subtracted from the detected sediment volume for all measurements.

707




708

709 **Figure A4.** Sediment mass balance for the period of 7 months separately for burrows and burrow embedding

710 areas as measured by the cameras. (a) LC-NU. (b) LC-SU. (c) LC-SL. (d) PdA-NU. (e) PdA-NL.

711 (f) PdA-SU. (g) PdA-SL. For abbreviations see Table A1.



712


713 **Figure A5.** Sediment mass balance for the period of 7 months separately for all delineated areas as measured

714 by the cameras. (a) LC-NU. (b) LC-SU. (c) LC-SL. (d) PdA-NU. (e) PdA-NL. (f) PdA-SU. (g) PdA-SL. For

715 abbreviations see Table A1.

716






717


718 **Figure A6.** Examples of surface scans showing the digital surface model (DSM) before a rainfall event (a, c)  
 719 at two camera locations in La Campana, and the calculated volume of redistributed sediment (b, d)  
 720 after the rainfall event: (a) DSM of a scan from the camera on the upper north-facing hillslope in La Campana; (b)  
 721 Detected sediment redistribution ( $\text{cm}^3 \text{cm}^{-2}$ ) on the upper north-facing hillslope in La Campana after a rainfall  
 722 event of  $17.2 \text{ mm day}^{-1}$ ; (c) DSM of a scan from the camera on the upper south-facing hillslope in La Campana;  
 723 (d) Detected sediment redistribution ( $\text{cm}^3 \text{cm}^{-2}$ ) on the upper south-facing hillslope after a rainfall event of  $17.2$   
 724  $\text{mm day}^{-1}$ . Red is the outline of the burrow entrance. Green is the outline of mound. Orange is the outline of  
 725 the burrow roof. The area which is not outlined is **burrow embedding area**. Redistribution is the volume of the  
 726 redistributed sediment, either accumulated (positive value) or eroded (negative value) per  $\text{cm}^3 \text{cm}^{-2}$ . After the  
 727 rainfall events, sediment mostly accumulated within the burrow entrance or near mounds and eroded from  
 728 burrow roofs and mounds.

729

730




731  
 732 **Figure A7.** Cumulative volume of redistributed sediment for all cameras. Positive values indicate sediment  
 733 accumulation. Negative values indicate sediment erosion. Whiskers are the median sediment redistribution. E  
 734 is the burrow entrance. M is the mound. R is burrow roof. EM is burrow embedding area. LC is mediterranean  
 735 climate zone. PdA is arid climate zone. (a) LC-NU. (b) LC-SU. (c) LC-SL. (d) PdA-NU. (e) PdA-NL. (f) PdA-  
 736 SU. (g) PdA-SL. For abbreviations see Table A1.



737  
 738 **Figure A8.** Hillslope-wide volume of redistributed sediment for a time period of one year in LC. (a-d) North-  
 739 facing hillslope. (e-h) South-facing hillslope. (a) and (e) Density of burrows as estimated by Grigusova et al.  
 740 2021. (b) and (f) Volume of the sediment excavated by the animals. (c) and (g) Volume of the sediment  
 741 redistributed during rainfall events within burrows. (d) and (h) Volume of additionally redistributed sediment  
 742 during rainfall events due to presence of the burrows. The values were calculated per burrow as stated in  
 743 section 3.7 by subtracting the sediment volume redistributed within burrows from the sediment volume

744 redistributed within burrow embedding area and then upscaled. B stays for burrow, EM stays for burrow  
 745 embedding area.  
 746



747  
 748 **Figure A9.** Hillslope-wide volume of redistributed sediment for a time period of one year in Pan de Azúcar. (a-  
 749 d) North-facing hillslope. (e-h) South-facing hillslope. (a) and (e) Density of burrows as estimated by Grigusova  
 750 et al. 2021. (b) and (f) Volume of the sediment excavated by the animals. (c) and (g) Volume of the sediment  
 751 redistributed during rainfall events within burrows. (d) and (h) Volume of additionally redistributed sediment

752 during rainfall events due to presence of the burrows. The values were calculated per burrow as stated in  
753 section 3.7 by subtracting the sediment volume redistributed within burrow from the sediment volume  
754 redistributed within burrow embedding area and then upscaled. B stays for burrow, EM stays for burrow  
755 embedding area by the burrowing animal.

756

## 757 References

- 758 Afana, A., Solé-Benet, A., Pérez, J.L., 2010. Determination of Soil Erosion Using Laser Scanners (accessed  
759 22 December 2021).
- 760 Andersen, D.C., 1987. *Geomys Bursarius* Burrowing Patterns: Influence of Season and Food Patch  
761 Structure. *Ecology* 68, 1306–1318. <https://doi.org/10.2307/1939215>.
- 762 Ashcroft, M.B., Gollan, J.R., Ramp, D., 2014. Creating vegetation density profiles for a diverse range of  
763 ecological habitats using terrestrial laser scanning. *Methods Ecol Evol* 5, 263–272.  
764 <https://doi.org/10.1111/2041-210X.12157>.
- 765 BANCROFT, W.J., HILL, D., ROBERTS, J.D., 2004. A new method for calculating volume of excavated  
766 burrows: the geomorphic impact of Wedge-Tailed Shearwater burrows on Rottneest Island. *Funct Ecology*  
767 18, 752–759. <https://doi.org/10.1111/j.0269-8463.2004.00898.x>.
- 768 Bernhard, N., Moskwa, L.-M., Schmidt, K., Oeser, R.A., Aburto, F., Bader, M.Y., Baumann, K., Blanckenburg,  
769 F. von, Boy, J., van den Brink, L., Brucker, E., Büdel, B., Canessa, R., Dippold, M.A., Ehlers, T.A.,  
770 Fuentes, J.P., Godoy, R., Jung, P., Karsten, U., Köster, M., Kuzyakov, Y., Leinweber, P., Neidhardt, H.,  
771 Matus, F., Mueller, C.W., Oelmann, Y., Oses, R., Osses, P., Paulino, L., Samolov, E., Schaller, M.,  
772 Schmid, M., Spielvogel, S., Spohn, M., Stock, S., Stroncik, N., Tielbörger, K., Übernickel, K., Scholten, T.,  
773 Seguel, O., Wagner, D., Kühn, P., 2018. Pedogenic and microbial interrelations to regional climate and  
774 local topography: New insights from a climate gradient (arid to humid) along the Coastal Cordillera of  
775 Chile. *CATENA* 170, 335–355. <https://doi.org/10.1016/j.catena.2018.06.018>.
- 776 Black, T.A., Montgomery, D.R., 1991. Sediment transport by burrowing mammals, Marin County, California.  
777 *Earth Surf. Process. Landforms* 16, 163–172. <https://doi.org/10.1002/esp.3290160207>.
- 778 Blanch, X., Eltner, A., Guinau, M., Abellan, A., 2021. Multi-Epoch and Multi-Imagery (MEMI) Photogrammetric  
779 Workflow for Enhanced Change Detection Using Time-Lapse Cameras. *Remote Sensing* 13, 1460.  
780 <https://doi.org/10.3390/rs13081460>.
- 781 Castner, J.L., Fowler, H.G., 1984. Distribution of Mole Crickets (Orthoptera: Gryllotalpidae: *Scapteriscus*) and  
782 the Mole Cricket Parasitoid *Larra bicolor* (Hymenoptera: Sphecidae) in Puerto Rico. *The Florida*  
783 *Entomologist* 67, 481. <https://doi.org/10.2307/3494730>.
- 784 Cerqueira, R., 1985. The Distribution of *Didelphis* in South America (Polyprotodontia, Didelphidae). *Journal*  
785 *of Biogeography* 12, 135. <https://doi.org/10.2307/2844837>.
- 786 Chen, M., Ma, L., Shao, M.'a., Wei, X., Jia, Y., Sun, S., Zhang, Q., Li, T., Yang, X., Gan, M., 2021. Chinese  
787 zokor (*Myospalax fontanierii*) excavating activities lessen runoff but facilitate soil erosion – A simulation  
788 experiment. *CATENA* 202, 105248. <https://doi.org/10.1016/j.catena.2021.105248>.
- 789 Coombes, M.A., 2016. Biogeomorphology: diverse, integrative and useful. *Earth Surf. Process. Landforms*  
790 41, 2296–2300. <https://doi.org/10.1002/esp.4055>.
- 791 Corenblit, D., Corbara, B., Steiger, J., 2021. Biogeomorphological eco-evolutionary feedback between life  
792 and geomorphology: a theoretical framework using fossorial mammals. *Die Naturwissenschaften* 108,  
793 55. <https://doi.org/10.1007/s00114-021-01760-y>.
- 794 Eitel, J.U.H., Williams, C.J., Vierling, L.A., Al-Hamdan, O.Z., Pierson, F.B., 2011. Suitability of terrestrial laser  
795 scanning for studying surface roughness effects on concentrated flow erosion processes in rangelands.  
796 *CATENA* 87, 398–407. <https://doi.org/10.1016/j.catena.2011.07.009>.
- 797 Eltner, A., Kaiser, A., Abellan, A., Schindewolf, M., 2017. Time lapse structure-from-motion photogrammetry  
798 for continuous geomorphic monitoring. *Earth Surf. Process. Landforms* 42, 2240–2253.  
799 <https://doi.org/10.1002/esp.4178>.
- 800 Eltner, A., Kaiser, A., Castillo, C., Rock, G., Neugirg, F., Abellán, A., 2016a. Image-based surface  
801 reconstruction in geomorphometry – merits, limits and developments. *Earth Surf. Dynam.* 4, 359–389.  
802 <https://doi.org/10.5194/esurf-4-359-2016>.
- 803 Eltner, A., Mulsow, C., Maas, H.-G., 2013. QUANTITATIVE MEASUREMENT OF SOIL EROSION FROM  
804 TLS AND UAV DATA. *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.* XL-1/W2, 119–124.  
805 <https://doi.org/10.5194/isprsarchives-XL-1-W2-119-2013>.
- 806 Eltner, A., Schneider, D., Maas, H.-G., 2016b. INTEGRATED PROCESSING OF HIGH RESOLUTION  
807 TOPOGRAPHIC DATA FOR SOIL EROSION ASSESSMENT CONSIDERING DATA ACQUISITION  
808 SCHEMES AND SURFACE PROPERTIES. *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.* XLI-  
809 B5, 813–819. <https://doi.org/10.5194/isprsarchives-XLI-B5-813-2016>.
- 810 Gabet, E.J., Reichman, O.J., Seabloom, E.W., 2003. The Effects of Bioturbation on Soil Processes and

811 Sediment Transport. *Annu. Rev. Earth Planet. Sci.* 31, 249–273.  
812 <https://doi.org/10.1146/annurev.earth.31.100901.141314>.

813 Galland, O., Bertelsen, H.S., Guldstrand, F., Girod, L., Johannessen, R.F., Bjugger, F., Burchardt, S., Mair,  
814 K., 2016. Application of open-source photogrammetric software MicMac for monitoring surface  
815 deformation in laboratory models. *J. Geophys. Res. Solid Earth* 121, 2852–2872.  
816 <https://doi.org/10.1002/2015JB012564>.

817 Grigusova, P., Larsen, A., Achilles, S., Klug, A., Fischer, R., Kraus, D., Übernickel, K., Paulino, L., Pliscoff, P.,  
818 Brandl, R., Farwig, N., Bendix, J., 2021. Area-Wide Prediction of Vertebrate and Invertebrate Hole  
819 Density and Depth across a Climate Gradient in Chile Based on UAV and Machine Learning. *Drones* 5,  
820 86. <https://doi.org/10.3390/drones5030086>.

821 Hakonson, T.E., 1999. The Effects of Pocket Gopher Burrowing on Water Balance and Erosion from Landfill  
822 Covers. *J. environ. qual.* 28, 659–665. <https://doi.org/10.2134/jeq1999.00472425002800020033x>.

823 Hall, K., Boelhouwers, J., Driscoll, K., 1999. Animals as Erosion Agents in the Alpine Zone: Some Data and  
824 Observations from Canada, Lesotho, and Tibet. *Arctic, Antarctic, and Alpine Research* 31, 436–446.  
825 <https://doi.org/10.1080/15230430.1999.12003328>.

826 Hancock, G., Lowry, J., 2021. Quantifying the influence of rainfall, vegetation and animals on soil erosion and  
827 hillslope connectivity in the monsoonal tropics of northern Australia. *Earth Surf. Process. Landforms* 46,  
828 2110–2123. <https://doi.org/10.1002/esp.5147>.

829 Hänsel, P., Schindewolf, M., Eltner, A., Kaiser, A., Schmidt, J., 2016. Feasibility of High-Resolution Soil  
830 Erosion Measurements by Means of Rainfall Simulations and SfM Photogrammetry. *Hydrology* 3, 38.  
831 <https://doi.org/10.3390/hydrology3040038>.

832 Hazelhoff, L., van Hoof, P., Imeson, A.C., Kwaad, F.J.P.M., 1981. The exposure of forest soil to erosion by  
833 earthworms. *Earth Surf. Process. Landforms* 6, 235–250. <https://doi.org/10.1002/esp.3290060305>.

834 Herbst, M., Bennett, N.C., 2006. Burrow architecture and burrowing dynamics of the endangered Namaqua  
835 dune mole rat (*Bathyergus janetta*) (Rodentia: Bathyergidae). *Journal of Zoology* 270, 420–428.  
836 <https://doi.org/10.1111/j.1469-7998.2006.00151.x>.

837 Horn, B.K.P., 1981. Hill shading and the reflectance map. *Proc. IEEE* 69, 14–47.  
838 <https://doi.org/10.1109/PROC.1981.11918>.

839 Imeson, A.C., 1977. Splash erosion, animal activity and sediment supply in a small forested Luxembourg  
840 catchment. *Earth Surf. Process. Landforms* 2, 153–160. <https://doi.org/10.1002/esp.3290020207>.

841 Imeson, A.C., Kwaad, F.J.P.M., 1976. Some Effects of Burrowing Animals on Slope Processes in the  
842 Luxembourg Ardennes. *Geografiska Annaler: Series A, Physical Geography* 58, 317–328.  
843 <https://doi.org/10.1080/04353676.1976.11879941>.

844 Iserloh, T., Ries, J.B., Arnáez, J., Boix-Fayos, C., Butzen, V., Cerdà, A., Echeverría, M.T., Fernández-Gálvez,  
845 J., Fister, W., Geißler, C., Gómez, J.A., Gómez-Macpherson, H., Kuhn, N.J., Lázaro, R., León, F.J.,  
846 Martínez-Mena, M., Martínez-Murillo, J.F., Marzen, M., Mingorance, M.D., Ortigosa, L., Peters, P.,  
847 Regüés, D., Ruiz-Sinoga, J.D., Scholten, T., Seeger, M., Solé-Benet, A., Wengel, R., Wirtz, S., 2013.  
848 European small portable rainfall simulators: A comparison of rainfall characteristics. *CATENA* 110, 100–  
849 112. <https://doi.org/10.1016/j.catena.2013.05.013>.

850 James, M.R., Robson, S., 2014. Sequential digital elevation models of active lava flows from ground-based  
851 stereo time-lapse imagery. *ISPRS Journal of Photogrammetry and Remote Sensing* 97, 160–170.  
852 <https://doi.org/10.1016/j.isprsjrs.2014.08.011>.

853 Jimenez, J.E., Feinsinger, P., Jaksi, F.M., 1992. Spatiotemporal Patterns of an Irruption and Decline of Small  
854 Mammals in Northcentral Chile. *Journal of Mammalogy* 73, 356–364. <https://doi.org/10.2307/1382070>.

855 Jones, C.G., Gutiérrez, J.L., Byers, J.E., Crooks, J.A., Lambrinos, J.G., Talley, T.S., 2010. A framework for  
856 understanding physical ecosystem engineering by organisms. *Oikos* 119, 1862–1869.  
857 <https://doi.org/10.1111/j.1600-0706.2010.18782.x>.

858 Kaiser, A., Neugirg, F., Rock, G., Müller, C., Haas, F., Ries, J., Schmidt, J., 2014. Small-Scale Surface  
859 Reconstruction and Volume Calculation of Soil Erosion in Complex Moroccan Gully Morphology Using  
860 Structure from Motion. *Remote Sensing* 6, 7050–7080. <https://doi.org/10.3390/rs6087050>.

861 Kinlaw, A., Grasmueck, M., 2012. Evidence for and geomorphologic consequences of a reptilian ecosystem  
862 engineer: The burrowing cascade initiated by the Gopher Tortoise. *Geomorphology* 157–158, 108–121.  
863 <https://doi.org/10.1016/j.geomorph.2011.06.030>.

864 Kromer, R., Walton, G., Gray, B., Lato, M., Group, R., 2019. Development and Optimization of an Automated  
865 Fixed-Location Time Lapse Photogrammetric Rock Slope Monitoring System. *Remote Sensing* 11, 1890.  
866 <https://doi.org/10.3390/rs11161890>.

867 Kukko, A., Hyypä, J., 2009. Small-footprint Laser Scanning Simulator for System Validation, Error  
868 Assessment, and Algorithm Development. *photogramm eng remote sensing* 75, 1177–1189.  
869 <https://doi.org/10.14358/PERS.75.10.1177>.

870 Larsen, A., Nardin, W., Lageweg, W.I., Bätz, N., 2021. Biogeomorphology, quo vadis? On processes, time,  
871 and space in biogeomorphology. *Earth Surf. Process. Landforms* 46, 12–23.  
872 <https://doi.org/10.1002/esp.5016>.

- 873 Le Hir, P., Monbet, Y., Orvain, F., 2007. Sediment erodability in sediment transport modelling: Can we  
874 account for biota effects? *Continental Shelf Research* 27, 1116–1142.  
875 <https://doi.org/10.1016/j.csr.2005.11.016>.
- 876 Lehnert, L.W., Thies, B., Trachte, K., Achilles, S., Osses, P., Baumann, K., Schmidt, J., Samolov, E., Jung, P.,  
877 Leinweber, P., Karsten, U., Büdel, B., Bendix, J. 2018. A Case Study on Fog/Low Stratus Occurrence at  
878 Las Lomitas, Atacama Desert (Chile) as a Water Source for Biological Soil Crusts. *Aerosol Air Qual. Res.*  
879 18, 254–269. <https://doi.org/10.4209/aaqr.2017.01.0021>.
- 880 Li, G., Li, X., Li, J., Chen, W., Zhu, H., Zhao, J., Hu, X., 2019a. Influences of Plateau Zokor Burrowing on Soil  
881 Erosion and Nutrient Loss in Alpine Meadows in the Yellow River Source Zone of West China. *Water* 11,  
882 2258. <https://doi.org/10.3390/w11112258>.
- 883 Li, L., 2014. Time-of-Flight Camera – An Introduction. Technical White Paper.  
884 <https://www.ti.com/lit/wp/sloa190b/sloa190b.pdf> (accessed 22 December 2021).
- 885 Li, T., Jia, Y., Shao, M.'a., Shen, N., 2019b. *Camponotus japonicus* burrowing activities exacerbate soil  
886 erosion on bare slopes. *Geoderma* 348, 158–167. <https://doi.org/10.1016/j.geoderma.2019.04.035>.
- 887 Li, T., Shao, M.'a., Jia, Y., Jia, X., Huang, L., 2018. Small-scale observation on the effects of the burrowing  
888 activities of mole crickets on soil erosion and hydrologic processes. *Agriculture, Ecosystems &*  
889 *Environment* 261, 136–143. <https://doi.org/10.1016/j.agee.2018.04.010>.
- 890 Li, T.C., Shao, M.A., Jia, Y.H., Jia, X.X., Huang, L.M., Gan, M., 2019c. Small-scale observation on the effects  
891 of burrowing activities of ants on soil hydraulic processes. *Eur J Soil Sci* 70, 236–244.  
892 <https://doi.org/10.1111/ejss.12748>.
- 893 Longoni, L., Papini, M., Brambilla, D., Barazzetti, L., Roncoroni, F., Scaioni, M., Ivanov, V., 2016. Monitoring  
894 Riverbank Erosion in Mountain Catchments Using Terrestrial Laser Scanning. *Remote Sensing* 8, 241.  
895 <https://doi.org/10.3390/rs8030241>.
- 896 MALLALIEU, J., CARRIVICK, J.L., QUINCEY, D.J., SMITH, M.W., JAMES, W.H.M., 2017. An integrated  
897 Structure-from-Motion and time-lapse technique for quantifying ice-margin dynamics. *J. Glaciol.* 63, 937–  
898 949. <https://doi.org/10.1017/jog.2017.48>.
- 899 Meysman, F.J.R., Boudreau, B.P., Middelburg, J.J., 2003. Relations between local, nonlocal, discrete and  
900 continuous models of bioturbation. *Journal of Marine Research* 61, 391–410.  
901 <https://doi.org/10.1357/002224003322201241>.
- 902 Morris, R.H., Buckman, S., Connelly, P., Dragovich, D., Ostendorf, B., and Bradstock, R.A., 2011. The dirt on  
903 assessing post-fire erosion in the Mount Lofty Ranges: comparing methods.
- 904 Muñoz-Pedrerros, A., Yáñez, J., Norambuena, H.V., Zúñiga, A., 2018. Diet, dietary selectivity and density of  
905 South American grey fox, *Lycalopex griseus*, in Central Chile. *Integrative zoology* 13, 46–57.  
906 <https://doi.org/10.1111/1749-4877.12260>.
- 907 Nasermoaddeli, M.B., Pasche, E., 2008. Application of terrestrial 3D scanner in quantification of the  
908 riverbank erosion and deposition. [https://www.tuhh.de/t3resources/wb/Publicationen/MA-](https://www.tuhh.de/t3resources/wb/Publicationen/MA-Veroeffentlichungen/nasermoaddelli/riverflow2008.pdf)  
909 [Veroeffentlichungen/nasermoaddelli/riverflow2008.pdf](https://www.tuhh.de/t3resources/wb/Publicationen/MA-Veroeffentlichungen/nasermoaddelli/riverflow2008.pdf) (accessed 22 December 2021).
- 910 Pang, X.P., Guo, Z.G., 2017. Plateau pika disturbances alter plant productivity and soil nutrients in alpine  
911 meadows of the Qinghai-Tibetan Plateau, China. *Rangel. J.* 39, 133. <https://doi.org/10.1071/RJ16093>.
- 912 Reichman, O.J., Seabloom, E.W., 2002. The role of pocket gophers as subterranean ecosystem engineers.  
913 *Trends in Ecology & Evolution* 17, 44–49. [https://doi.org/10.1016/S0169-5347\(01\)02329-1](https://doi.org/10.1016/S0169-5347(01)02329-1).
- 914 Richards, P.J., Humphreys, G.S., 2010. Burial and turbulent transport by bioturbation: a 27-year experiment  
915 in southeast Australia. *Earth Surf. Process. Landforms* 21, n/a-n/a. <https://doi.org/10.1002/esp.2007>.
- 916 Ridd, P.V., 1996. Flow Through Animal Burrows in Mangrove Creeks. *Estuarine, Coastal and Shelf Science*  
917 43, 617–625. <https://doi.org/10.1006/ecss.1996.0091>.
- 918 Romañach, S.S., Reichman, O.J., Seabloom, E.W., 2005. Seasonal influences on burrowing activity of a  
919 subterranean rodent, *Thomomys bottae*. *Journal of Zoology* 266, 319–325.  
920 <https://doi.org/10.1017/S0952836905006941>.
- 921 Rutin, J., 1996. The burrowing activity of scorpions (*Scorpio maurus palmatus*) and their potential  
922 contribution to the erosion of Hamra soils in Karkur, central Israel. *Geomorphology* 15, 159–168.  
923 [https://doi.org/10.1016/0169-555X\(95\)00120-T](https://doi.org/10.1016/0169-555X(95)00120-T).
- 924 Sarbolandi, H., Plack, M., Kolb, A., 2018. Pulse Based Time-of-Flight Range Sensing. *Sensors (Basel,*  
925 *Switzerland)* 18. <https://doi.org/10.3390/s18061679>.
- 926 Schiffers, K., Teal, L.R., Travis, J.M.J., Solan, M., 2011. An open source simulation model for soil and  
927 sediment bioturbation. *PloS one* 6, e28028. <https://doi.org/10.1371/journal.pone.0028028>.
- 928 Sharon, D., 1980. The distribution of hydrologically effective rainfall incident on sloping ground. *Journal of*  
929 *Hydrology* 46, 165–188. [https://doi.org/10.1016/0022-1694\(80\)90041-4](https://doi.org/10.1016/0022-1694(80)90041-4).
- 930 Thomsen, L.M., Baartman, J.E.M., Barneveld, R.J., Starkloff, T., Stolte, J., 2015. Soil surface roughness:  
931 comparing old and new measuring methods and application in a soil erosion model. *SOIL* 1, 399–410.  
932 <https://doi.org/10.5194/soil-1-399-2015>.
- 933 Übernickel, K., Ehlers, T.A., Paulino, L., Fuentes Espoz, J.-P., 2021a. Time series of meteorological stations  
934 on an elevational gradient in National Park La Campana, Chile.

935 Übernicketl, K., Pizarro-Araya, J., Bhagavathula, S., Paulino, L., Ehlers, T.A., 2021b. Reviews and syntheses:  
936 Composition and characteristics of burrowing animals along a climate and ecological gradient, Chile.  
937 *Biogeosciences* 18, 5573–5594. <https://doi.org/10.5194/bg-18-5573-2021>.  
938 Voiculescu, M., Ianăș, A.-N., Germain, D., 2019. Exploring the impact of snow vole (*Chionomys nivalis*)  
939 burrowing activity in the Făgăraș Mountains, Southern Carpathians (Romania): Geomorphic  
940 characteristics and sediment budget. *CATENA* 181, 104070.  
941 <https://doi.org/10.1016/j.catena.2019.05.016>.  
942 Wei, X., Li, S., Yang, P., Cheng, H., 2007. Soil erosion and vegetation succession in alpine Kobresia steppe  
943 meadow caused by plateau pika—A case study of Nagqu County, Tibet. *Chin. Geograph.Sc.* 17, 75–81.  
944 <https://doi.org/10.1007/s11769-007-0075-0>.  
945 Wilkinson, M.T., Richards, P.J., Humphreys, G.S., 2009. Breaking ground: Pedological, geological, and  
946 ecological implications of soil bioturbation. *Earth-Science Reviews* 97, 257–272.  
947 <https://doi.org/10.1016/j.earscirev.2009.09.005>.  
948 Yair, A., 1995. Short and long term effects of bioturbation on soil erosion, water resources and soil  
949 development in an arid environment. *Geomorphology* 13, 87–99. [https://doi.org/10.1016/0169-](https://doi.org/10.1016/0169-555X(95)00025-Z)  
950 [555X\(95\)00025-Z](https://doi.org/10.1016/0169-555X(95)00025-Z).  
951 Yoo, K., Amundson, R., Heimsath, A.M., Dietrich, W.E., 2005. Process-based model linking pocket gopher  
952 (*Thomomys bottae*) activity to sediment transport and soil thickness. *Earth Surf. Process. Landforms* 33,  
953 917. <https://doi.org/10.1130/G21831.1>.  
954  
955