## Supplemental Information for "Intensified paraglacial slope failures due to accelerating downwasting of a temperate glacier in Mt. Gongga, Southeastern Tibet Plateau"

Yan Zhong<sup>1,2</sup>, Qiao Liu<sup>1</sup>, Matthew Westoby<sup>3</sup>, Yong Nie<sup>1</sup>, Francesca Pellicciotti<sup>4</sup>, Bo Zhang<sup>5</sup>, Jialun Cai<sup>5</sup>, Guoxiang Liu<sup>5</sup>, Haijun Liao<sup>1,2</sup>, Xuyang Lu<sup>1</sup>

<sup>1</sup> Institute of Mountain Hazards and Environment, Chinese Academy of Sciences, Chengdu 610041, China

<sup>2</sup> College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup> Department of Geography and Environmental Sciences, Engineering and Environment, Northumbria University, Newcastle upon Tyne NE1 8ST, UK

<sup>4</sup> Swiss Federal Institute for Forest, Snow and Landscape Research WSL, 8903 Birmensdorf, Switzerland <sup>5</sup> Department of Surveying and Geo-Informatics, Faculty of Geosciences and Environmental Engineering, Southwest Jiaotong University, Chengdu 611756, China

Correspondence to: Qiao Liu (liuqiao@imde.ac.cn)

## **Supplemental Figures**

5



15

Fig. S1: Spatial extents of the four repeated UAV surveys by Qiao Liu et al. and the distribution of tie points selected for imagery co-registration. Note the 'tie points' outside the UAV mapping areas were only used for extrapolation but not for co-registration.



20 Fig. S2: Tie points for manually tracked the displacement of the four slopes (B1-B4 in Fig. 6) in UAV imagery by Qiao Liu et al.



Fig. S3: Detailed of small magnitude rockfalls in UAV- and field-photographs by Qiao Liu et al.





