# Effect of rock uplift and Milankovitch timescale variations in precipitation and vegetation cover on catchment erosion rates

<sup>3</sup> Hemanti Sharma<sup>1</sup>, Todd A Ehlers<sup>1\*</sup>, Christoph Glotzbach<sup>1</sup>, Manuel Schmid<sup>1</sup> Katja Tielbörger<sup>2</sup>

4 <sup>1</sup>Department of Geosciences, University of Tübingen, 72076 Tübingen, Germany

5 <sup>2</sup>Department of Biology, Plant Ecology Group, University of Tübingen, Auf der Morgenstelle 5, 72076 Tübingen, Germany

6 Correspondence to: Todd A. Ehlers (todd.ehlers@uni-tuebingen.de)

7 Abstract. Catchment erosion and sedimentation are influenced by variations in the rates of rock uplift (tectonics), and periodic 8 fluctuations in climate and vegetation cover. This study focuses on quantifying the effects of changing climate and vegetation 9 on erosion and sedimentation over distinct climate-vegetation settings applying the Landlab-SPACE landscape evolution model. As catchment evolution is subjected to tectonic and climate forcings at millennial to million-year time-scales, the 10 11 simulations are performed for different tectonic scenarios and periodicities in climate-vegetation change. We present a series of generalized experiments that explore the sensitivity of catchment hillslope and fluvial erosion and sedimentation for 12 13 different rock uplift rates (0.05 mm a<sup>-1</sup>, 0.1 mm a<sup>-1</sup>, 0.2 mm a<sup>-1</sup>) and Milankovitch climate periodicities (23 kyr, 41 kyr and 100 14 kyr). Model inputs were parameterized for two different climate and vegetation conditions at two sites in the Chilean Coastal 15 Cordillera at  $\sim 26^{\circ}$ S (arid and sparsely vegetated) and  $\sim 33^{\circ}$ S (Mediterranean). For each setting, steady state topographies were produced for each uplift rate before introducing periodic variations in precipitation and vegetation cover. Following this, the 16 17 sensitivity of these landscapes was analysed for 3 Myr in a transient state. Results suggest that regardless of the uplift rate, 18 transients in precipitation and vegetation cover resulted in transients in erosion rates in the direction of change in precipitation 19 and vegetation. While the transients in sedimentation were observed to be in the opposite direction of change in the 20 precipitation and vegetation cover, with phase lags of  $\sim 1.5 - 2.5$  kyr. These phase lags can be attributed to the changes in plant functional type (PFT) distribution induced by the changes in climatic as well as the regolith production rate. These effects are 21 22 most pronounced over longer period changes (100 kyr) and higher rock uplift rates (0.2 mm yr<sup>-1</sup>). This holds true for both 23 vegetation and climate settings considered. Furthermore, transient changes in catchment erosion due to varying vegetation and 24 precipitation were between ~35%-110% of the background (rock uplift) rate and would be measureable with commonly used techniques (e.g. sediment flux histories, cosmogenic nuclides). Taken together, we find that vegetation-dependent erosion and 25 sedimentation are influenced by Milankovitch timescale changes in climate, but that these transient changes are superimposed 26 27 upon tectonically driven rates of rock uplift.

28 Keywords: vegetation dynamics, climate change, tectonics, landscape evolution modelling, SPACE, Landlab

# 29 1 Introduction

30 The pioneering work of G.K. Gilbert (Gilbert, 1877) highlighted that surface uplift, climate, and biota (amongst other things) 31 jointly influence catchment-scale rates of weathering and erosion. In recent decades a wide range of studies have built upon 32 these concepts and quantified different ways in which climate, tectonics, or vegetation cover influence rates of erosion and 33 sedimentation. For example, recent work highlights that denser vegetation or lower precipitation both decrease erosion (Alonso 34 et al., 2006; Bonnet and Crave, 2003; Huntley et al., 2013; McPhillips et al., 2013; Miller et al., 2013; Perron, 2017; Schaller 35 et al., 2018; Starke et al., 2020; Tucker, 2004). In addition, periodic changes in climate (such as changes driven by Milankovitch timescale orbital variations) have also been recognized as influencing rates of catchment erosion and sedimentation (Braun et 36 al., 2015; Hancock and Anderson, 2002; Hyun et al., 2005; Schaller et al., 2004) although our ability to measure orbital 37 38 timescale induced erosional changes can be challenging (e.g. Schaller and Ehlers, 2006; Whipple, 2009). Several studies have also documented how the combined effects of either climate and vegetation change, or variable rates of rock uplift and climate change (including glaciation) impact catchment scale processes (Herman et al., 2010; Mishra et al., 2019; Schmid et al., 2018; Tucker, 2004; Yanites and Ehlers, 2012). Taken together, the previous studies have found that the long-term development of topography (such as over million-year time scales) is in many situations sensitive to the tectonic, climate, and vegetation history of the region, and that competing effects of different coeval processes (e.g. climate change and tectonics) exist, but are not well understood.

45 Quantification of climate, vegetation, and tectonic effects on catchment erosion is challenging because these processes are 46 confounded and can, if coupled, have opposing effects on erosion and/or sedimentation. For example, precipitation has both 47 direct (positive) as well as indirect effects on erosion that operate via vegetation cover. Namely, plants require water to grow and survive, vegetation cover is usually positively affected by precipitation both on a global scale (i.e. when comparing biomes 48 49 across latitudinal gradients) as well as on a regional or local scale (e.g. Huxman et al., 2004; Sala et al., 1988; Zhang et al., 50 2016). Though vegetation cover is also influenced by temperature, seasonality and many other abiotic factors such as soil type 51 and thickness, the positive relationship of biomass and cover with water availability is rather general. For example, in dry 52 ecosystems such as hot deserts or Mediterranean systems, vegetation cover is primarily limited by water availability and is 53 therefore very low. As precipitation increases, vegetation cover increases rapidly, although water availability can still be the 54 limiting factor as well as other factors (Breckle, 2002). In temperate systems, where water is abundant and soils are well 55 developed, plant growth is primarily limited by low winter temperatures. Overall, the relationship between precipitation and 56 vegetation cover follows a saturation curve with large sensitivity (e.g. measured as rain use efficiency- RUE) to precipitation 57 in arid to Mediterranean systems and low sensitivity in temperate or tropical systems (Gerten et al., 2008; Huxman et al., 2004;

58 Yang et al., 2008; Knapp et al., 2017).

59 Previous modelling and observational studies have made significant progress in understanding the interactions between surface 60 processes and either climate (Dixon et al., 2009; Routschek et al., 2014; Seybold et al., 2017; Slater and Singer, 2013), 61 vegetation (Acosta et al., 2015; Amundson et al., 2015; Istanbulluoglu and Bras, 2005) or coupled climate-vegetation dynamics 62 (Dosseto et al., 2010; Jeffery et al., 2014; Mishra et al., 2019; Schmid et al., 2018). Over geologic (millennial to million-year) 63 timescales, observational studies of these interactions are impossible (or require proxy data) and numerical modeling approaches provide a means to explore interactions between climate, vegetation, tectonics, and topography. The first 64 65 observational study of this kind suggested that high MAP (mean annual precipitation) is associated with denser vegetation and hence resulting in lower erosion rates (Langbein and Schumm, 1958). One of the first numerical modeling studies 66 implementing a vegetation-erosion coupling was conducted by Collins et al. (2004). This study was followed by work from 67 Istanbulluoglu and Bras (2006), which quantified the effect of vegetation on landscape relief and drainage formation. More 68 69 recently, work by Schmid et al. (2018) included the effects of transient climate and vegetation coupled with a landscape 70 evolution model to predict topographic and erosional variations over millennial- to million-year timescales. However, Schmid et al., (2018) presented a simplified approach to consider hillslope and detachment-limited fluvial erosion and only considered 71 72 a homogeneous substrate. Other studies have documented that sediment or bedrock erosion by rivers is not dominated purely 73 by detachment-limited (Howard, 1994) or transport-limited fluvial erosion (Willgoose et al., 1991). Rather, it often involves a 74 combination of, or transition between, both conditions (e.g., Pelletier, 2012). Given this, treatment of bedrock erosion and 75 sediment transport for mixed bedrock-alluvial streambeds provides a more realistic framework for understanding the influence 76 of climate, vegetation, and tectonic processes on topographic development. Recent work (Shobe et al., 2017) presented an 77 additional component (SPACE) to the Landlab surface process model. SPACE allows for the simulation of mixed detachment-78 transport limited fluvial processes, including separate layers for bedrock and loose sediment. Finally, the sensitivity of 79 topography to different rock uplift rates in variable climate-vegetation settings has not yet been investigated. The combined 80 interactions of tectonics (rock uplift) and variable climate and vegetation warrant investigation given the significant influence

- 81 of rock uplift on mean elevation, erosion rates and river channel profiles (Kirby and Whipple, 2012; Turowski et al., 2006)
- 82 and hillslopes.



Figure 1: The representative study areas in the Chilean Coastal Cordillera used for the model setup. The model parameters were loosely tuned to the climate and vegetation conditions in these areas (Schmid et al., 2018). The Pan de Azucar area in the north neighbours the Atacama Desert and has sparse vegetation cover (10%) and an arid climate (30 mm yr<sup>-1</sup>). The La Campana area in south has a Mediterranean climate and ecosystem with more abundant vegetation (70%) and precipitation (350 mm yr<sup>-1</sup>). These two study areas are part of the German EarthShape priority research program (<u>www.earthshape.net</u>).

89 In this study, we complement the previous work and investigate the transient landscape response for mixed bedrock-alluvial systems. We do this for different rates of rock uplift and periodic changes (Milankovitch cycles) in precipitation and vegetation. 90 91 Our focus is on erosion and sedimentation changes occurring over millennial to million-year timescales. Sub-annual to decadal 92 scale changes are beyond the scope of this study. More specifically, this study evaluates the following two hypotheses: First, 93 if vegetation cover and climate vary on Milankovitch timescales, then any increases or decreases in catchment erosion will be 94 more pronounced over longer (e.g. 100 kyr) rather than shorter (e.g. 21 kyr) periodicities due to the longer duration of change 95 imposed. Second, if increasing rates of tectonic uplift cause increases in catchment erosion rates, then any periodic variations in climate and vegetation cover will be muted (lower amplitude) at higher uplift rates because the effect of rock uplift on 96 erosion will outweigh climate and vegetation change effects. Given the complexity of this problem, we investigate these 97 98 hypotheses through numerical landscape evolution modelling using a step-wise increase in model complexity where the 99 contributions of individual processes (i.e. climate, or vegetation, or tectonics) are identified separately before looking into the fully coupled system and resulting interactions. We apply a two-dimensional coupled detachment-transport limited landscape 100 evolution model for fluvial processes. In addition, hillslope diffusion (Johnstone and Hilley, 2014) and weathering and soil 101 102 production (Ahnert, 1977) processes are considered. Although this study is primarily focused on documenting the predicted sensitivity of catchments to variations in tectonics, climate, and vegetation change - we have tuned our model setup to the 103 104 conditions along the Chilean Coastal Cordillera (Fig. 1) which features a similar tectonic setting, but an extreme climate and 105 ecological gradient. This was done to provide realistic parameterizations for vegetation cover and precipitation in different 106 ecological settings. This area is also part of the German-Chilean priority research program, EarthShape: Earth Surface Shaping

107 by Biota (www.earthshape.net) where extensive ongoing research is occurring.

### 108 2 Methods

- We apply the landscape evolution model, Landlab (Hobley et al., 2017) using the SPACE 1.0 module of Shobe et al. (2017) for detachment *vs.* transport limited fluvial processes. The Landlab/SPACE programs were modified for vegetation dependent hillslope and fluvial erosion using the approach of Schmid et al. (2018). In general, the geomorphic processes considered involve weathering and regolith production calibrated to the Chilean Coastal Cordillera observations of Schaller et al. (2018), vegetation dependent coupled detachment-transport limited fluvial erosion, and depth dependent hillslope diffusion. The model parameters (i.e., bedrock and sediment erodibility and diffusion coefficient) in the simulations are based on those of Schmid et al. (2018). A detailed explanation of the weathering, erosion, sediment transport and deposition processes is provided in
- 116 Appendix A, and a summary of model parameters used is given in Table 1.

# 117 2.1 Model setup and scenarios considered

- 118 The model consists of a 10 km by 10 km rectangular grid with 100 m node spacing (Fig. 2a), with a total domain area of 100
- 119 km<sup>2</sup>. We conducted generalized simulations that are loosely tuned to the climate and vegetation conditions in two areas in
- 120 the Chilean Coastal Cordillera (Fig. 1) which have predominantly granitoid lithology (van Dongen et al., 2018; Kojima et al.,
- 121 2017; Oeser et al., 2018; Rossel et al., 2018). These areas exhibit a large climate and vegetation gradient ranging from and
- 122 arid climate (MAP: 30 mm) and sparse vegetation (10%) in Pan de Azucar National Park to a wetter Mediterranean climate
- 123 (MAP: 35 cm) with more abundant vegetation (70%) in La Campana National Park.
- 124 Bedrock elevation and sediment cover thickness are considered as separate layers to quantify simultaneous bedrock erosion 125 and sediment entrainment across the model domain. Simulations were conducted for 15 Myr to generate a steady-state 126 topography with the mean values of precipitation and vegetation cover for the two study areas. The rates of rock uplift are kept 127 constant during the steady-state simulations, and subsequently in the transient stage with oscillating vegetation cover and 128 precipitation. After the development of a steady-state topography, transient forcings in vegetation cover and mean annual 129 precipitation (MAP) (Fig. 2b) were introduced for 3 Myr. Vegetation cover varied by  $\pm 10\%$  around the mean value used to 130 develop the steady-state topography. The 10% vegetation cover variation is based on the dynamic vegetation modelling results 131 of Werner et al. (2018) for the Chilean Coastal Cordillera. They found that from the Last Glacial Maximum to present that 132 vegetation cover in the region varied by ~10%. The periodicity of vegetation change varied between simulations (Table 1).
- 133 Changes in vegetation cover are driven by climatic variations, where MAP has been shown to be much more influential than 134 temperature changes, especially in relatively dry regions (e.g. Mowll et al., 2015) and in grasslands (e.g. Sala et al., 1988).
- 135 Many previous studies have shown that annual primary production (ANPP) and associated vegetation cover increases linearly
- 136 (Mowll et al., 2015; Xia et al., 2014) or in an asymptotic manner with MAP (Huxman et al., 2004; Smith et al., 2017; Yang et
- 137 al., 2008; Zhang et al., 2016; Knapp et al., 2017). These findings are also highly consistent among different approaches such
- 138 as global (Gerten et al., 2008) or regional (Zhang et al., 2016) models, field and remotely-sensed observations across biomes
- 139 and among years (Huxman et al., 2004; Xia et al., 2014; Yang et al., 2008), as well as rapid vegetation responses to rainfall
- 140 manipulation experiments (Smith et al., 2017). An asymptotic relationship appears the more common case, especially when
- 141 looking at warm and dry ecosystems, i.e. regions up to approx. 600mm MAP (Huxman et al., 2004; Mowll et al., 2015). Here,
- 142 it has been demonstrated that the sensitivity of ANPP to MAP decreases from more water-limited systems such as deserts to
- 143 Mediterranean and temperate regions (Huxman et al., 2004; Yang et al., 2008). Namely, the same increase in MAP will yield
- 144 much larger increase in vegetation cover in dry regions than in wetter ones. To implement these effects, we use an empirical
- 145 approach based on vegetation-precipitation relationships observed in the Chilean Coastal Cordillera (see Schmid et al. 2018
- 146 for details) to estimate what mean annual precipitation rates are associated with different vegetation cover amounts (Fig. 2b,
- 147 2c).



149Figure 2: Model geometry and climate and vegetation forcings used in this study. (a) A simple representation of the model setup150with a square grid, and catchment outlet in the lower left corner. (b) Graphical representation of the magnitude and pattern of151fluctuations imposed on vegetation (top) and precipitation (bottom) during the transient state of the model. Red rectangles represent152one cycle, whose effects are discussed in detail. (c) Graphical representation of precipitation and vegetation cover correlation from153the Chilean study areas and used as the empirical bases for how precipitation rates vary for +/-10% changes in vegetation cover154(Schmid et al., 2018).

The effects of vegetation cover on hillslope and fluvial processes are modified from the approach of Schmid et al, (2018), see also Appendix, and Table 1. Briefly, we applied a slope and depth-dependent linear diffusion rule following the approach of

- 157 Johnstone and Hilley (2014). The diffusion coefficient  $(K_d)$  is defined as a function of the bare soil diffusivity  $(K_b)$  and
- 158 exponentially varies with vegetation cover following the approach of Istanbulluoglu (2005) and Dunne et al. (2010). Fluvial
- 159 erosion is estimated for a two-layer topography (i.e., bedrock and sediment are treated explicitly) in the coupled detachment –
- 160 transport limited model. Bedrock erosion and sediment entrainment are calculated simultaneously in the model following the

- 161 approach of Shobe et al. (2017). The effects of vegetation cover on fluvial erosion were implemented using the approach of
- 162 Istanbulluoglu (2005) and Schmid et al. (2018) and by introducing the effect of a vegetation dependent Manning's roughness.
- 163 The sediment and bedrock erodibility ( $K_{vs}$  and  $K_{vr}$ , respectively) are influenced by the fraction of vegetation cover V (see
- 164 appendix for governing equations). Figure 3 shows the range of resulting diffusion coefficients (Kd) and sediment and bedrock
- 165 erodibility (K<sub>vs</sub>, K<sub>vb</sub>, respectively) values considered in this study. The exponential and power-law relationships producing
- 166 these values, respectively, are a source of non-linearity that are manifested in the results discussed in subsequent sections.



167

168Figure 3: Graphical representation of the range of vegetation dependent diffusion coefficient (Kd, left y-axis), sediment erodibility169(Kvs), and bedrock erodibility (Kvb) values considered in this study (see Appendix for governing equations). The combined

170 erodibility is referred to as Kv (right y-axis).







As the study areas exhibit similar granitoid lithology, the erosional parameters (Table. 1) are kept uniform for both the study areas. However, parameters based on climate conditions namely soil production rate (Schaller et al., 2018), MAP and vegetation cover (Schmid et al., 2018), are different for these areas. The vegetation cover and precipitation rate are kept uniform

- 179 across the model domain due to low to moderate relief in target catchments (~750 m for Pan de Azucar and ~1500 m in La
- 180 Campana).
- 181 The model scenarios considered were designed to provide a stepwise increase in model complexity to identify how variations
- 182 in precipitation, vegetation cover, or rock uplift rate influence erosion and sedimentation. The model scenarios include:
- Influence of oscillating precipitation and constant vegetation cover, on erosion and sedimentation (Fig. 4a, Fig. 5, Section
   3.1).
- 185 2. Influence of constant precipitation and oscillating vegetation cover, on erosion and sedimentation (Fig. 4b, Fig. 6, Section3.2).
- 187 3. Influence coupled oscillations in precipitation and vegetation cover, on erosion and sedimentation (Fig. 4c, Fig. 7, Section
  188 3.3).
- 189 4. Influence of different periodicities of precipitation/vegetation change on erosion and sedimentation (Fig. 8, Section 3.4).
- 190 5. Influence of rock uplift rate and oscillating precipitation/vegetation on erosion sedimentation (Fig. 9, Section 3.5).
- 191 Table 1. Landscape evolution model input parameters used and corresponding units.

Model Parameters	Values
Grid size	10 [km] x 10 [km], dx: 100 [m]
Model runtime (totalTime)	Steady-state: 15 [Ma], Transient state: 3 [Ma]
Rock uplift rates (U)	0.05 [mm a <sup>-1</sup> ], 0.1 [mm a <sup>-1</sup> ], 0.2 [mm a <sup>-1</sup> ]
Periodicities (sinePeriod)	23 [kyr], 41 [kyr], 100 [kyr] (Milankovitch cycles)
Initial sediment thickness (H_initial)	0 [m]
Bedrock erodibility (Kr)	2 x 10 <sup>-9</sup> [m <sup>-1</sup> ]
Sediment erodibility (Ks)	2 x 10 <sup>-8</sup> [m <sup>-1</sup> ]
Soil production/transport decay depth (h*)	0.5 [m]
Reach scale bedrock roughness (H*)	1 [m]
Porosity ( $\phi$ )	0.2 [-]
Fraction of fine sediments (Ff)	0.2 [-]
Effective terminal settling velocity (Vs)	10 [m a <sup>-1</sup> ]
m, n	0.6, 1 [-]
Bedrock erosion threshold stream power ( $\omega_cr$ )	5 x 10 <sup>-4</sup> [m a <sup>-1</sup> ]
Sed. entr. threshold stream power ( $\omega_cs$ )	5 x 10 <sup>-5</sup> [m a <sup>-1</sup> ]
Maximum sediment production rate (W <sub>o</sub> )	9.7 x 10 <sup>-6</sup> [m yr <sup>-1</sup> ] (10% Veg. cover, 1.3 x 10 <sup>-4</sup> [m yr <sup>-1</sup> ] (70% Veg. cover)
Mean annual precipitation (P)	0.03 [m yr <sup>-1</sup> ] (10% Veg. cover), 0.35 [m yr <sup>-1</sup> ] (70% Veg.
Bare soil diffusivity (K <sub>b</sub> )	$0.01 \ [m^2 \ yr^{-1}]$
Exponential decay coefficient (a)	0.3 [-]
Critical channel formation area (Acrit)	1 x 10 <sup>6</sup> [m <sup>2</sup> ]
Reference vegetation cover (V <sub>r</sub> )	1 (100%)
Manning's number for bare soil (n <sub>s</sub> )	0.01 [-]
Manning's number for ref. vegetation $(n_v)$	0.6 [-]
Scaling factor for vegetation influence (w)	1 [-]

192 The porosity (0.2) used in this study is lower than usual range for soil (0.3 - 0.4), as sediment produced as a result of weathering 193 in the study areas is a mixture of fine and coarse grained regolith (Schaller et al., 2020). Manning's numbers for bare soil and reference vegetation cover are same as used by Schmid et al., (2018). The rate of rock uplift is kept temporally and spatially 194 195 constant (0.05 mm  $a^{-1}$ ) for both study areas, for the simulations in scenarios 1 - 4. This is done in order to minimize the effect 196 of tectonics on topography to isolate the sensitivity of geomorphic processes to changing precipitation and vegetation cover. In scenario 5, the effect of different rock uplift rates (i.e., 0.05 mm a<sup>-1</sup>, 0.1 mm a<sup>-1</sup> and, 0.2 mm a<sup>-1</sup>) is studied in combination 197 198 with the coupled oscillations in precipitation and vegetation cover. The rock uplift rate used in the scenarios 1-4 is estimated 199 from the findings of Melnick (2016) and Avdievitch et al. (2018), which suggests the modern and paleo uplift and exhumation rates of < 0.1 mm a<sup>-1</sup> for the study areas and northern Coastal Cordillera in general. Similarly, the periodicity of oscillations 200 for precipitation and vegetation cover are kept constant (23 kyr) for model scenarios 1, 2, 3, and 5. In scenario 4, the effect of 201 202 different periodicities (i.e., 23 kyr, 41kyr, and 100 kyr) is studied in combination with coupled oscillations in precipitation and 203 vegetation cover. The periodicities of oscillations are based on Milankovitch cycles (Ashkenazy et al., 2010; Hyun et al., 204 2005). In the simulations with variations in either vegetation cover or climate, a perfect sinusoidal function is used to demonstrate the oscillations in precipitation rates for both catchments (Fig. 4a,4b). However, in case of coupled oscillations in 205 vegetation cover and climate, an asymmetric sinusoidal function is used for precipitation rates (Fig. 4c). This is done due to 206 207 the observed non-linear relationships between changing vegetation cover and precipitation in Figure 2. The non-linearity stems from the fact that in high vegetation cover settings (e.g., 70%, Fig. 2) a large increase in precipitation is needed to increase 208 209 vegetation cover by 10% compared to a smaller decrease in precipitation required to reduce vegetation cover by 10%.

# 210 2.2 Boundary and Initial conditions

211 An initial low relief (< 1 m) random noise topography was applied to the model grid at the start of the simulations. The initial

- 212 topographies had a slight initial topographic slope of  $\approx 1.4 \times 10^{-5}$  (Fig. 2a). The boundaries on all sides of the domain were
- 213 closed (no flow), except the south-west corner node which was an outlet node. From these conditions, the steady-state
- 214 topography was calculated over 15 Myr model time, and the resulting bedrock elevation and sediment thickness were used as
- 215 input for the transient scenarios described in section 2.1.

# 216 3 Results

In the following sections, we focus our analysis on the mean catchment sediment thickness (i.e. the combined thickness of soil and regolith) over the entire domain, mean bedrock erosion rates (excluding sediment erosion), mean sediment entrainment rates and the mean catchment erosion rates. The mean catchment erosion rates are the sum of bedrock erosion and sediment entrainment rates. To simplify the presentation of result, results are shown only for the first cycle of transient climate and vegetation change. Results from the first cycle were representative of subsequent cycles (not shown), and no longer-term variations or trends in erosion/sedimentation were identified or warrant discussion.

# 223 3.1 Influence of oscillating precipitation and constant vegetation cover on erosion and sedimentation (Scenario 1)

In this scenario, with a rock uplift rate of 0.05 mm a<sup>-1</sup> and 23kyr periodicity in precipitation, the mean catchment sediment entrainment rates follow the pattern of change in precipitation (Fig. 5a, b), but with an offset (phase lag) between the maxima and minima of entrainment and precipitation. A higher variation in the range of sediment entrainment rates (i.e., - 0.036 mm yr<sup>-1</sup> – 0.043 mm yr<sup>-1</sup>, Fig. 5b) is observed for simulations with 10% vegetation cover. Negative values in sediment entrainment rates correspond to sediment deposition rates during drier periods. The peak in sediment entrainment rates (e.g. 0.043 mm yr<sup>-1</sup> for 10% veg., and ~0.038 mm yr<sup>-1</sup> for 70% veg., Fig. 5b), is observed with a time lag of (~ -2 kyr) before the peak in maximum

- 230 precipitation in both the 10% and 70% vegetation cover simulations. This result suggests that as precipitation increases 231 sediment is readily entrained where available in the catchment until bedrock is locally exposed. The changes in mean catchment 232 sediment thickness (Fig. 5c) are influenced by changes in the sediment entrainment and precipitation rates, but with a lag time 233 between the maximum in precipitation and the minimum in sediment thickness. The lowest mean catchment sediment thickness 234 (e.g. ~0.97 m for 10% veg., and ~1.9 m for 70% veg., Fig. 5c) also occurs with a time lag of (~3 kyr) after the peak in precipitation rates, for both the 10% and 70% vegetation cover simulations. The same time lag ( $\sim$ 3 kyr) is observed in the peak 235 in mean catchment bedrock erosion (e.g. ~0.087 mm yr<sup>-1</sup> for 10% veg. and ~0.1 mm yr<sup>-1</sup> for 70% veg., Fig. 5d) and coincides 236 with when the minimum sediment cover is present and more bedrock is exposed for erosion. As we use the total change in 237 238 bedrock elevation to estimate bedrock erosion rates, the loss in bedrock due to weathering (exponential) is also accounted for. The phase lag in bedrock erosion and sediment thickness can be attributed to exponential weathering, which is discussed in 239 240 detail in section 4.2. Finally, the mean-catchment erosion rates follow the pattern of change in precipitation rates (Fig. 5a, 5e) 241 without a phase lag. The maximum erosion rates are similar in range for both the 10% and 70% vegetation cover simulations (e.g. ~  $0.12 \text{ mm yr}^{-1}$ , Fig. 5e). However, in the 10% vegetation cover simulation, the minimum in the mean catchment erosion 242 rate decreases more (e.g. to ~ 0.01 mm yr<sup>-1</sup>, Fig. 5e) relative to the higher vegetation cover scenario. The different decreases 243 in the minimum erosion rate between the two vegetation cover amounts corresponds to the differences in precipitation rates 244 (Figs 5a, 4a). 245
- 246



Figure 5: Temporal evolution of catchment averaged predictions for scenario 1 described in the text (section 3.1). Graphical representation of mean catchment sedimentation and erosion to (a) oscillating precipitation [mm yr<sup>-1</sup>] and constant vegetation cover [-] in terms of (b) sediment entrainment [mm yr-1], (c) sediment thickness [m], (d) bedrock erosion [mm yr-1], (e) mean erosion rates [mm yr<sup>-1</sup>] for entire catchment. The periodicity of climate and vegetation oscillations is 23 kyr with rate of rock uplift as 0.5 mm yr<sup>-</sup> 1.

The absence of a phase lag between the mean-catchment erosion and precipitation rates reflects that the combined sediment entrainment and bedrock erosion rates when added together track the overall trend in precipitation rate changes, but the individual components (sediment vs. bedrock) respond differently.

# 256 **3.2 Influence of constant precipitation, oscillating vegetation cover, on erosion and sedimentation (Scenario 2)**

257 Results from this scenario with constant mean annual precipitation (at the mean value of the previous scenario) and oscillating

258 vegetation cover (Fig. 4b. 6a) show a starkly different catchment response from scenario 1 (section 3.1). The sediment

- 259 entrainment rates show for both simulations (Fig. 6b) a small decrease in entrainment as vegetation cover increases (e.g. ~ -
- 260 0.05 mm yr<sup>-1</sup> for 10% veg., and ~ -0.01 mm yr<sup>-1</sup> for 70% veg., Fig. 6b. As vegetation cover decreases later in the cycle,
- entrainment rates increase (e.g. to  $\sim 0.13$  mm yr<sup>-1</sup> for 10% veg., and to 0.01 mm yr<sup>-1</sup> for 70% veg., Fig. 6b). The larger magnitude
- 262 of increase in entrainment for the 10% vegetation cover case corresponds to the minimum (0%) in vegetation cover where the

263 potential for erosion is the highest. In the 10% vegetation cover simulation, the lowest mean catchment sediment thickness

264 was observed ~1.5 kyr after the minimum in vegetation cover (Fig. 6c).

265



Figure 6: Temporal evolution of catchment averaged predictions for scenario 2 described in the text (section 3.2). Graphical representation of mean catchment sedimentation and erosion to (a) constant precipitation [mm yr<sup>-1</sup>] and oscillating vegetation cover [-] in terms of (b) sediment entrainment [mm yr<sup>-1</sup>], (c) sediment thickness [m], (d) bedrock erosion [mm yr<sup>-1</sup>], (e) mean erosion rates [mm yr<sup>-1</sup>] for entire catchment. The periodicity of climate and vegetation oscillations is 23 kyr with rate of rock uplift as 0.5 mm yr<sup>-</sup> 1.

272 The range of mean catchment sediment thickness varies significantly in the simulations (e.g. ~0.72 m - ~1.38 m for 10% veg., and  $\sim 2.2 \text{ m} - \sim 2.3 \text{ m}$  for 70% veg., Fig. 6c). The same time lag ( $\sim 1.5 \text{ kyr}$ ) is observed between the peak in mean catchment 273 274 bedrock erosion rates (Fig. 6d) and the minimum in vegetation cover. This is most likely due to the maximum exposure of 275 bedrock for erosion when catchment average sediment thicknesses are at their minimum. Also, the first phase of the cycle is 276 mainly depositional while bedrock erosion (including weathering) is observed, which happens partly in places where there is no deposition. Finally, mean catchment erosion rates (Fig. 6e) are significantly affected ( $\sim +0.25$  mm yr<sup>-1</sup>) by oscillating 277 278 vegetation cover in simulations with a mean 10% vegetation. For the 70% vegetation cover simulation, a similar maximum in 279 erosion occurs also during the minimum in vegetation, but is far less dramatic, presumably due to the still somewhat large

- 280 (60%) amount of vegetation-cover present. Although the relief and slopes are lower in sparsely vegetated catchment (10% V),
- 281 significantly higher erosion rates are observed as precipitation is kept constant at 30 mm yr<sup>-1</sup>, while the vegetation cover was
- 282 reduced to 0%. This can be attributed to low (bedrock/sediment) stream power thresholds.

# 283 3.3 Influence of coupled oscillations of precipitation and vegetation cover, on erosion and sedimentation (Scenario 3)

284 The catchment response to coupled oscillations in precipitation rate and vegetation cover (Fig. 4c) on erosion and 285 sedimentation represents a composite of the effects discussed in the previous 2 sections (Fig. 7). For example, the mean catchment sediment entrainment rates have a peak in entrainment rates (~1.5 kyr) prior to the peak in climate/vegetation values. 286 287 A similar effect was noted for scenarios 1 (Fig. 5, section 3.1). As the precipitation rates and vegetation cover decrease later in the cycle (Fig. 7a), the sediment entrainment rates increase. In more detail, the 70% vegetation cover simulations show a 288 modest increase similar to that observed in scenario 1 (Fig. 5b), whereas the 10% vegetation cover shows are sharp peak in 289 290 the sediment entrainment rates when 0% vegetation cover is present. This later observation is similar what is observed for 291 scenario 2 (Fig. 6b, section 3.2). Thus, in the case of co-varying precipitation rates and vegetation cover, the response observed 292 in terms of sediment entrainment is not predicted to be the same for all degrees of vegetation cover, and depends heavily on 293 the initial vegetation cover of the system around which variations occur.

294



Figure 7: Temporal evolution of catchment averaged predictions for scenario 3 described in the text (section 3.3). Graphical representation of mean catchment sedimentation and erosion to (a) coupled oscillations in precipitation [mm yr<sup>-1</sup>] and vegetation cover [-] in terms of (b) sediment entrainment [mm yr<sup>-1</sup>], (c) sediment thickness [m], (d) bedrock erosion [mm yr<sup>-1</sup>], (e) mean erosion rates [mm yr<sup>-1</sup>] for entire catchment. The periodicity of climate and vegetation oscillations is 23 kyr with rate of rock uplift as 0.5 mm yr<sup>-1</sup>.

Mean catchment sediment thicknesses in the 10% vegetation cover simulation show a modest response and vary between 1.16 m - 1.24 m (Fig. 7c), and with a time lag of ~2.5 kyr between the peak in precipitation/vegetation and minimum sediment thickness. This lag is also observed in the case of the 70% vegetation cover simulation, but with a higher amplitude of change in sediment thickness (e.g. 2 m - 2.22 m, Fig. 7c). A similar trend in time lags between the peaks in climate/vegetation and bedrock erosion (Fig. 6d) are also present. These observations for variations in sediment thickness again represent the combined effects of the results discussed in section 3.1 and 3.2 (Figs. 5c, 6c).

- 307 The amplitude of change in bedrock erosion is between 0.05 mm yr<sup>-1</sup> 0.06 mm yr<sup>-1</sup> for 10% veg., and 0.05 mm yr<sup>-1</sup> 0.08
- 308 mm yr<sup>-1</sup> for 70% veg. (Fig. 7d). The bedrock erosion response for both simulations represents a composite of the effects shown
- 309 in the previous two scenarios (sections 3.1, 3.2). Here the increase in time lag in the maximum in erosion rates (most notable
- 310 for the 70% vegetation cover simulation) resembles the effect of a large increase in precipitation rates (compared Fig. 5d) for
- 311 the first part of the cycle. Whereas, the second peak in bedrock erosion visible in the 10% vegetation cover scenario more

- 312 closely resembles the effects shown in Fig 6d when the vegetation cover goes to 0%, and the landscape is increasingly sensitive
- 313 to erosion with whatever runoff (albeit little) is available.
- 314 Finally, the mean catchment erosion rates (Fig. 7e) again show the combined effects of the sediment entrainment rate and bedrock erosion histories previously discussed (Figs. 7b, d). In the simulation with 70% initial vegetation cover, the mean 315 316 catchment erosion rates follow the pattern of changes in precipitation rates (e.g. ranging from 0.04 mm yr<sup>-1</sup> to 0.1 mm yr<sup>-1</sup>, Fig. 7e, see also Fig. 5e). A similar trend is present in the first half of the cycle in the simulation with 10% vegetation cover, 317 but with much lower magnitudes (i.e., 0.05 mm yr<sup>-1</sup> to 0.06 mm yr<sup>-1</sup>, Fig. 7e). However, during the second half of the cycle, 318 319 the erosion rates increase up to  $\sim 0.06$  mm yr<sup>-1</sup> and have a second peak at  $\sim 17-18$  kyr for the 10% vegetation simulation when 320 the vegetation cover is at 0%. The previous result is however in contradiction to the detachment-limited results shown in Fig 17 of Schmid et al. (2018), who found that erosion rates decreased to 0 mm yr<sup>-1</sup> for the period of no vegetation cover and 321 minimum precipitation rate of (~10 mm yr<sup>-1</sup>). This contradiction is related to the increase in sediment entrainment at this time 322 323 (Fig. 7b) which heavily influences the mean erosion. The detachment-limited approach of Schmid et al. (2018) could not 324 account for this, and will be discussed in detail in section 4.2. To summarize, as discussed previously the locations of the 325 maximums and minimums in the mean erosion rate and the shape of the curves (Fig. 7e) can be linked to different times in the 326 climate and vegetation history when either the effects of variable precipitation rate or vegetation cover dominate the mean 327 catchment erosional response.

### 328 3.4 Influence of the periodicity of precipitation/vegetation variations on erosion and sedimentation (Scenario 4)

329 Here we show the influence of different periodicities (23, 41, and 100 kyr) in precipitation and vegetation change on catchment erosion and sedimentation for the cases of a 10% mean vegetation cover (Fig. 8) and 70% vegetation cover (Fig. 9). We find 330 331 higher variations in mean sediment entrainment rates (Fig. 8b, 9b) for both the 10% and 70% vegetation cover simulations for 332 the shorter periodicities (23 and 41 kyr). However, the phase lag in the peaks of sediment entrainment and precipitation rates was higher for longer periodicities (e.g. ~9%, ~16.2%, ~19% in 23 kyr, 43 kyr, and 100 kyr, respectively) for the 10% 333 334 vegetation cover case (Fig. 8b). These phase lags are however, dampened in the highly vegetated landscapes (70%) at longer 335 periods (i.e., ~9%, ~9.5%, ~14% in 23 kyr, 43 kyr, and 100 kyr respectively, Fig. 9b). In a landscape with 10% vegetation 336 cover, the simulation with longer periodicity (100 kyr) shows higher variations in mean catchment sediment thickness (e.g. 337 1.14 cm - 1.25 cm, Fig. 8c). This is mimicked in the landscape with 70% vegetation cover, with the range of sediment thickness between 1.95 cm - 2.27 cm (Fig. 9c). A similar trend with higher amplitude of change is also observed for bedrock erosion 338 rates in the sparsely vegetated landscape (10%) with values ranging from 0.05 mm a<sup>-1</sup> to 0.062 mm yr<sup>-1</sup> (Fig. 8d) for longer 339 periodicity (100 kyr). The same pattern is observed in highly vegetated landscapes (70%), with the values of bedrock erosion 340 341 rates ranging from 0.045 mm yr<sup>-1</sup> to 0.094 mm yr<sup>-1</sup> (Fig. 9d) for the longer periodicity (100 kyr).



Figure 8: Temporal evolution of catchment averaged predictions for scenario 4 described in the text (section 3.4). Graphical representation of mean catchment sedimentation and erosion to (a) different periodicities of coupled oscillations in precipitation [mm yr<sup>-1</sup>] and vegetation cover [-] in terms of (b) sediment entrainment [mm yr<sup>-1</sup>], (c) sediment thickness [m], (d) bedrock erosion [mm yr<sup>-1</sup>], (e) mean erosion rates [mm yr<sup>-1</sup>] for entire catchment. The rate of rock uplift is kept constant as 0.5 mm yr<sup>-1</sup>. The

347 simulations represent 10% initial vegetation cover.



348

Figure 9: Temporal evolution of catchment averaged predictions for scenario 4 described in the text (section 3.4). Graphical representation of mean catchment sedimentation and erosion to (a) different periodicities of coupled oscillations in precipitation [mm yr<sup>-1</sup>] and vegetation cover [-] in terms of (b) sediment entrainment [mm yr<sup>-1</sup>], (c) sediment thickness [m], (d) bedrock erosion [mm yr<sup>-1</sup>], (e) mean erosion rates [mm yr<sup>-1</sup>] for entire catchment. The rate of rock uplift is kept constant as 0.5 mm yr<sup>-1</sup>. The simulations represent 70% initial vegetation cover.



# 356 **3.5 Influence of rock uplift rate and oscillating precipitation/vegetation on erosion sedimentation (Scenario 5)**

Here we investigate the response of mean catchment erosion and sedimentation for different rates of rock uplift (i.e., 0.05 mm a<sup>-1</sup>, 0.1 mm a<sup>-1</sup>, 0.2 mmyr<sup>-1</sup>) for the 10% vegetation cover (Fig. 10) and 70% vegetation cover (Fig. 11) scenarios. To simplify the presentation and comparison of results, the periodicity of precipitation and vegetation change is kept the same as section 3.3 (i.e., 23 kyr). In general, the results discussed below demonstrate that the transient catchment response to coupled oscillations in precipitation rate and vegetation cover are similar in shape regardless of the rock uplift rate. The magnitude of change in mean catchment erosion associated with precipitation and vegetation changes increases with increasing uplift rate,

despite an identical amount of vegetation and precipitation change imposed (Figs. 10a, 11a) on each rock uplift rate simulation.



Figure 10: Temporal evolution of catchment averaged predictions for scenario 5 described in the text (section 3.5). Graphical representation of mean catchment sedimentation and erosion with different rates of rock uplift [mm a<sup>-1</sup>] to (a) coupled oscillations in precipitation [mm yr<sup>-1</sup>] and vegetation cover [-] in terms of (b) sediment entrainment [mm yr<sup>-1</sup>], (c) sediment thickness [m], (d) bedrock erosion [mm yr<sup>-1</sup>], (e) mean erosion rates [mm yr<sup>-1</sup>] for entire catchment. The periodicity of climate and vegetation oscillations is 23 kyr. The simulations represent 10% initial vegetation cover.



Figure 11: Temporal evolution of catchment averaged predictions for scenario 5 described in the text (section 3.5). Graphical representation of mean catchment sedimentation and erosion with different rates of rock uplift [mm a<sup>-1</sup>] to (a) coupled oscillations in precipitation [mm yr<sup>-1</sup>] and vegetation cover [-] in terms of (b) sediment entrainment [mm yr<sup>-1</sup>], (c) sediment thickness [m], (d) bedrock erosion [mm yr<sup>-1</sup>], (e) mean erosion rates [mm yr<sup>-1</sup>] for entire catchment. The periodicity of climate and vegetation oscillations is 23 kyr. The simulations represent 70% initial vegetation cover.

In more detail, the temporal pattern of changes in sediment entrainment rates (Fig. 10b, 11b) is similar for all uplift rates considered, but the amplitude of change increases as the uplift rate increases. In addition, the phase lag between the peaks in sediment entrainment rates and maximum precipitation rates in the 10% vegetation simulation (Fig. 10b) varies with the rock uplift rate. For example, the peaks in sediment entrainment rates have phase lag of ~ -4 kyr, -2.5 kyr, and -2 kyr for rock uplift rates of 0.2 mm a<sup>-1</sup>, 0.1 mm a<sup>-1</sup>, and 0.05 mm a<sup>-1</sup> respectively (Fig. 10b) in first half of the vegetation/precipitation oscillation. However, the phase lags are overall shorter in highly vegetated landscapes (70%) (e.g. ~ -3 kyr, -2 kyr, -1 kyr) before the maximum in precipitation for rock uplift rates of 0.2 mm a<sup>-1</sup>, 0.1 mm a<sup>-1</sup>, and 0.05 mm a<sup>-1</sup> respectively (Fig. 11b).

383 For the landscape with 10% vegetation cover, the simulation with the highest rates of rock uplift  $(0.02 \text{ mm a}^{-1})$  showed lower

384 mean catchment sediment thickness (e.g.  $\sim 0.5 \text{ m} - \sim 0.6 \text{ m}$ , Fig. 10c). In contrast, the slowest rock uplift simulation (0.05 mm

 $a^{-1}$  had thicker sediment thickness of ~1.16 m - ~1.24 m, (Fig. 10c). The same pattern was observed in the catchment with

11c). These results for sediment thickness variations reflect that higher rock uplift rates result in steeper slopes (not shown) and higher mean catchment erosion rates (Figs. 10e, 11e) such that regolith production rates are outpaced by erosion and therefore result in thinner sediment thicknesses. Also, the thicker sediment thickness for lower uplift rates could be an integrated result of slightly lower erosion rates compared to sediment production rates over the whole 15 Myr model runtime (steady state). This result is akin to the observational results from Heimsath et al. (1997).

Temporal variations in bedrock and mean catchment erosion rates are similar to those described in section 3.3 (Fig. 7) for the sparsely and more heavily vegetated conditions. The primary difference is that at high rock uplift rates the amplitude of bedrock or mean catchment erosion increases (Figs. 10d,e; 11d,e). To summarize, these results highlight that regardless of the rock uplift rate, similar temporal changes are observed in sediment entrainment or thickness, and in bedrock and catchment

- 396 erosion for oscillating precipitation rates and vegetation cover. However, the amplitude of change (or absolute change) in
- 397 entrainment and erosion rates increases with increases in rock uplift rate. This will be discussed in detail in section 4.4.

#### 398 4 Discussion

In this section, we synthesize the results from previous section (scenarios 1-5) in detail. We further investigate the effects of coupled climate and vegetation oscillations (Scenario 3) on the occurrence of erosion and sedimentation on spatial scale.

### 401 **4.1 Differences in effects between oscillating vegetation or precipitation**

402 Here the sensitivity of erosion and sedimentation to variable precipitation and/or vegetation cover is analysed. In the scenario 403 with oscillating precipitation and constant vegetation cover, sparsely vegetated landscapes (10%) are eroding slowly during periods of lower precipitation. This might be attributed to the dependency of the bedrock erosion and sediment entrainment on 404 405 the amount of water available through precipitation, which in turn affects the erosion rates. The mean erosion in this scenario 406 is dominated by bedrock erosion with a significant contribution from sediment entrainment. Also, the mean erosion rates over 407 one climate oscillation cycle are observed to be slightly higher ( $\sim 20\%$ ) than mean erosion rates at steady state for sparsely vegetated landscape (10% V). For densely vegetated landscape (70% V), this difference is significant (i.e., 50% higher mean 408 409 erosion rates during a transient cycle, in comparison to steady state). This implies the non-linearity of the erosion response to 410 changes in MAP, which is significantly higher in densely vegetated landscape where amplitude of change in MAP (e.g., 260 411 mm - 720 mm) is much higher than drier landscapes (e.g., 10 mm - 60 mm).

412 Similarly, in a scenario with constant precipitation and variable vegetation cover, sparsely vegetated landscapes (10%) are observed to be much more sensitive in terms of erosion rates during periods of no vegetation cover. The amplitude of erosional 413 414 change was ten times higher than that of densely vegetated landscapes. The mean erosion in sparsely vegetated landscapes is 415 dominated equally by bedrock erosion (Fig. 6d) and sediment entrainment, due to the higher availability of bare soil. This 416 justifies the argument of a higher sensitivity of sparsely vegetated landscapes to erosion and sedimentation. This result confirms 417 the findings of Yetemen et al. (2015) (see Fig. 2g), which suggests that shear stress (erosion) decreases significantly (1 to 0.1) 418 as the total grass cover (vegetation) is increased from 0% (bare soil) to 20% grass cover. Also, a small change in vegetation 419 cover in densely vegetated landscapes would not result in significant differences in erosional processes. Unlike previous 420 scenario (oscillating precipitation and constant vegetation cover), we do not observe non-linearity in erosion response to the changes in vegetation cover (i.e., mean erosion rates over one transient cycle are equal to steady state mean erosion rates). 421

In general, mean catchment sediment thickness is observed to be inversely proportional to precipitation, owing to higher stream
 power. This in turn translates to a higher sediment flux during wetter periods. The influence of oscillating precipitation and

424 constant vegetation cover on sediment thickness is slightly higher in simulations with sparse vegetation cover. In simulations

425 with constant precipitation and oscillating vegetation cover, the sensitivity of sediment thickness is much higher in landscapes

- 426 with sparse vegetation. This can be attributed to an absence of vegetation cover. A decreased impact of oscillating vegetation
- 427 cover on sediment thickness occurs in landscapes with denser vegetation cover and demonstrates that surface processes in
- 428 these settings are not highly dependent on changes in vegetation density. This has been explained by Huxman et al. (2004),
- 429 who found that vegetation cover responds to MAP variations in wet and dry systems during dry years.

# 430 **4.2** Synthesis of coupled oscillations of precipitation and vegetation cover simulations

The sensitivity of erosion and sedimentation to coupled oscillations in precipitation and vegetation cover (scenario 3, section 431 3.3) indicates that mean catchment erosion rates (Fig. 7e) are correlated with precipitation for densely vegetated landscapes 432 433 (70%). This is owed to the dominating effect of mean annual precipitation changes (from 26 cm yr<sup>-1</sup> to 72 cm yr<sup>-1</sup>) on erosion over vegetation cover change (from 60% to 80%, Fig. 7a) in these landscapes. This can be attributed to the higher amplitude 434 435 of precipitation oscillations in these simulations required to change vegetation cover by +/-10% (Fig. 2b). In the case of a sparsely vegetated landscape (10%), mean erosion rates (Fig. 7e) are also correlated to precipitation, but only for the first half 436 of the cycle when vegetation cover is present. However, mean erosion rates increase rapidly in the second half of the cycle 437 when MAP decreases (from 60 mm yr<sup>-1</sup> to 10 mm yr<sup>-1</sup>, Fig. 7a), and vegetation cover magnitudes decrease (from 20% to 0%, 438 Fig. 7a). This inverse correlation between precipitation and erosion can be attributed to increasing susceptibility of the surface 439 440 to sediment entrainment as vegetation cover decreases to bare soil, even with very low precipitation rates. The non-linearity 441 of erosion response to changes in MAP is reduced by half (in comparison to changing climate and constant vegetation) in 442 coupled simulations.

- Thus, the temporal evolution of mean erosion rates between the heavily (70%) and sparsely (10%) vegetated landscapes varies depending on the initial vegetation state of the catchment. As a result, correlated and anti-correlated relationships between precipitation, vegetation cover, and erosion are predicted and are the result of precipitation or vegetation exerting a dominant or subsidiary influence on catchment erosion at different times in the catchment history and for different catchment precipitation and vegetation cover conditions. This prediction is consistent with observed correlations of vegetation cover and catchment average erosion rates recently documented along the western Andean margin by Starke et al. (2020).
- 449 The lag behavior observed in sediment entrainment, thickness and bedrock erosion is explained in additional simulations we 450 conducted (results not shown for brevity) where the weathering (regolith production) function was turned off in the model 451 simulations (see Fig. A1 in appendix). In these simulations, we did not observe any significant phase lags in maximum and minimum of erosion rates, sediment thickness and vegetation cover/precipitation. Also, the erosion rates for sparsely vegetated 452 453 catchment (10% V), drops to a minimum during the phase of bare soil and minimum precipitation (10 mm yr<sup>-1</sup>). Hence, 454 sediment supply through weathering can be attributed to double peaks observed in mean catchment sediment entrainment rates 455 (Fig. 7b) and erosion rates (7e). When there is no explicit weathering / regolith production involved in the model simulations, 456 sediment supply for entrainment is significantly reduced. As a result, entrainment rates are observed to be two orders of 457 magnitude lower than bedrock erosion, hence entrainment rates are not shown in Fig. A1. This implies that weathering plays 458 a major role in leading to the phase lags observed in above results.

# 459 **4.3 Differences between the periodicities of climate and vegetation cover oscillations**

The periodicity of change in climate will mainly affect vegetation via the lag-time it takes for the vegetation to respond i.e., if the vegetation structure does not change (e.g., grasslands or forests), then grasslands are very flexible (Bellard et al., 2012; Kelly and Goulden, 2008; Smith et al., 2017). Grasslands can plastically respond from year to year while forests may die off and be replaced by grasslands when it becomes drier and vice-versa. This change in vegetation type might lead to the fluctuations in sedimentation and erosion rates due to periodicity of change in climate and vegetation cover.

# 465 4.4 The effect of rock uplift rate on signals of varying precipitation and vegetation cover

466 No difference in erosion rates was identified between the two different vegetation/precipitation simulations for a given uplift 467 rate when the erosion rate is averaged over the full period of vegetation/precipitation change. In a steady state landscape, erosion rates are equal to the rock uplift rates according to the law of continuity of mass (e.g., Tucker et al., 2001). This means 468 469 that steady state landscapes experience higher erosion rates with higher uplift rates. However, the mean catchment erosion 470 rates shown in Fig. 10e, 11e show temporal variations in the erosion rate driven by oscillations in the precipitation rate and 471 vegetation. When average erosion rates are calculated over a complete cycle of the oscillation, the mean erosion rates are 472 slightly higher than rock uplift rates, owing to the non-linearity of erosion response to changes in MAP. This result indicates that any climate or vegetation driven changes in erosion will not be evident when observed over too long a period time, but 473 474 might introduce shorter-term transients (high or low) depending on the climate/vegetation cycle of change. This finding is significant for observational studies seeking to measure the predictions shown in this study. More specifically, 475 476 thermochronometer dating approaches used to quantify denudation rates over million-year timescales will be hard pressed to 477 measure any signal of how climate or vegetation change on Milankovitch timescales influence denudation. Rather, the rate of 478 tectonic rock uplift or exhumation (in the case of erosion rates equalling the rock uplift rate) will be measured. In contrast, 479 observational techniques sensitive to decadal (e.g. sediment fluxes) or millennial (e.g. cosmogenic radio nuclides measured from river terraces) can be sensitive to timescales less than the period of oscillation and are more like to record transient 480 481 catchment erosion rates influenced by variations in precipitation or vegetation cover.

482 The vegetation and precipitation driven transients in mean catchment erosion rates predicted by this study were large enough 483 to be measured by some observational techniques. For example, in sparsely vegetated landscapes the half amplitude of change in erosion rates (from steady-state values) slightly decreases as the uplift rate increases. A higher magnitude of change in 484 transient erosion rates (from stead-state values) is found in densely vegetated landscapes and is again slightly decreased as the 485 486 uplift rate increases. Previous work by Schaller and Ehlers (2006) investigated the ability of denudation rates calculated from 487 cosmogenic radionuclides measured in a sequence of fluvial terraces to record periodic (Milankovitch timescale) variations in 488 denudation rates. The magnitude of change in predicted transient erosion rates described above is above the detection limit 489 reported by Schaller and Ehlers (2006), particularly when the mean catchment denudation rate is  $\sim 0.1$  mm yr<sup>-1</sup> or higher. Thus, 490 the predictions suggested in this study are testable in field-based studies, and other methods such as basin sedimentation rate 491 histories (e.g. determined from magneto-stratigraphy, optically stimulated luminescence, or other methods) also hold potential.

### 492 4.5 Spatial changes in where erosion and sedimentation changes occur

In the previous sections, our analysis focused on the spatially averaged response of the catchment in terms of changes in sedimentation and erosion. Here, we discuss the same model results as previously presented for but show two examples (for two different vegetation covers) of the spatial variations of erosion and sediment thickness within the catchments. This provides a basis for understanding where in the catchment changes are occurring.



497

Figure 12: Two-dimensional map-view representation of changes in topographic elevations [m] (1<sup>st</sup> row), sediment thickness [m] (2<sup>nd</sup> row), and erosion rates [mm yr<sup>-1</sup>] (3<sup>rd</sup> row). These changes are represented with respect to steady state conditions (1<sup>st</sup> column), for maximum (2<sup>nd</sup> column) and minimum (3<sup>rd</sup> column) values of precipitation and vegetation in an oscillation cycle. The simulations represent 10% initial vegetation cover.

502 Spatial variations in the pattern of erosion and sedimentation in the simulations with 23 kyr coupled precipitation and 503 vegetation oscillations, and a rock uplift rate of 0.05 mm a<sup>-1</sup>, are shown in the topographic elevation, sediment thickness, and 504 erosion rate changes for both the maximum and minimum in precipitation and vegetation cover. In the simulations with sparse 505 vegetation cover (10%) (Fig. 12) at the maximum in precipitation and vegetation cover, erosion rate changes from steady state are  $\sim 0.03$  mm yr<sup>-1</sup> in valleys and  $\sim 0.01$  mm yr<sup>-1</sup> on hillslopes. At the minimum in precipitation and vegetation cover, erosion 506 rate changes from steady state are higher in valleys than hillslopes. This may be attributed to an absence of vegetation during 507 508 this period, where the surface (bedrock or sediment) is readily available for erosion even with lower precipitation rates. The 509 sediment thickness is observed to be slightly higher in the streambeds and valleys for streams with larger accumulation area. 510 However, the smaller streams have lower sediment thickness compared to connected hillslopes. For example, higher sediment 511 thickness (~1.24 m) is observed near the catchment outlet in the lower-left corner of the domain. At the maximum in precipitation and vegetation cover cycle, the landscape experiences a slightly higher contrast in sediment thickness compared 512 513 to the steady-state condition, whereby a net lowering of the sediment layer is observed of approximately 2 cm to 5 cm on the 514 hillslopes and ~6 cm near the catchment outlet. This can be attributed to higher sediment fluxes during this period. At the minimum in the precipitation and vegetation cover cycle, the landscape experiences a slight difference from the steady state 515 516 sediment thickness ( $\sim 2$  cm lowering) except for deposition in higher order streams (up to  $\sim 2$  cm) near the catchment outlet.



517

518 Figure 13. Two-dimensional map-view representation of changes in topographic elevations [m] (1<sup>st</sup> row), sediment thickness [m] (2<sup>nd</sup> 519 row), and erosion rates [mm yr<sup>-1</sup>] (3<sup>rd</sup> row). These changes are represented with respect to steady state conditions (1<sup>st</sup> column), for 520 maximum (2<sup>nd</sup> column) and minimum (3<sup>rd</sup> column) values of precipitation and vegetation in an oscillation cycle. The simulations 521 represent 70% initial vegetation cover.

522 In the simulations with dense vegetation cover (70%) (Fig. 13), erosion rate changes from steady state conditions are higher 523 during the maximum in the precipitation and vegetation cover cycle with higher magnitudes ( $\sim 0.08$  mm yr<sup>-1</sup> in valleys and up to ~0.02 mm yr<sup>-1</sup> on hillslopes and ridges) due to the higher precipitation rates. At minimum precipitation and vegetation cover 524 525 magnitudes (P = 26 cm; V = 60%), erosion rate changes are reduced (up to -0.03 mm yr<sup>-1</sup>) in valleys and (up to -0.01 mm yr<sup>-1</sup>) 526 <sup>1</sup>) on hillslopes in comparison to the erosion rates at steady state. Sediment thickness is observed to be relatively higher in the streambeds and valleys (~2.25 m) than the hillslopes. It is contrastingly higher in the lowlands than the areas at higher 527 528 elevations. At maximum precipitation and vegetation cover (maximum in the cycle) sediment thickness is ~10 cm lower on hillslopes and up to ~30 cm lower in valleys. The same trend with lower amplitude is evident for the minimum in the 529 530 precipitation and vegetation cover cycle. This implies that at higher vegetation cover, sediment thickness is significantly reduced as a result of higher sediment flux during the peak in precipitation rates. This in turn signifies the dominance of 531 532 precipitation changes over vegetation cover change in highly vegetated landscapes.

# 533 4.6 Comparison to previous studies

Results presented in this study document a higher sensitivity of catchment erosion and sedimentation of sparsely vegetation landscapes (10%) to changes in vegetation cover, whereas densely vegetated (70%) landscapes are more responsive to changes in precipitation than vegetation changes. This confirms the broad findings of Schmid et al. (2018) and Yetemen et al. (2019),

which suggest vulnerability of erosion rates in sparsely vegetated landscapes to changes in vegetation cover and, that for 537 538 densely vegetated landscapes, sensitivity to the changes in MAP. However, there are differences between the results of Schmid 539 et al. (2018) and this study, particularly for the temporal changes in erosion rates we observe for the sparse vegetation cover 540 (10%) scenario with coupled precipitation/vegetation cover oscillations. More specifically, previous results from the 541 detachment limited model shown in Fig. 17 of Schmid et al. (2018) show that catchment erosion rates in sparsely vegetated 542 landscapes decrease as the precipitation and vegetation cover increases in the first part of a cycle. In the second part of the 543 cycle when precipitation and vegetation decrease to their minimum Schmid et al. (2018) predict erosion rates are ~0 mm yr<sup>-1</sup>. 544 However, in the coupled detachment-transport fluvial erosion model presented here (SPACE), we observe a different behavior and erosion rates slightly increase as precipitation and vegetation cover increase (from 0.05 mm yr<sup>-1</sup> to 0.065 mm yr<sup>-1</sup>, Fig. 545 7(e)), rather than decrease. This difference is due to higher sediment entrainment rates we predict during the period of no 546 547 vegetation and low precipitation (10 mm  $yr^{-1}$ ), which is a result of higher vulnerability of bare soil to erosion, even with very 548 low precipitation rates. Therefore, the application of a detachment limited, vs. coupled detachment-transport limited modelling 549 approach has bearing on the predicted response, and when comparing results to natural systems care should be taken in which

550 approach is used.

Previous geochemistry related observational studies from the Chilean Coastal Cordillera (EarthShape study areas, 551 552 www.earthshape.net) are also available for comparison to this study. For example, the steady-state sediment thickness in our simulations for 10% and 70% initial vegetation cover are predicted to be higher than the field observations reported by Schaller 553 554 et al. (2018) and Oeser et al. (2018), who reported a ~20 cm and ~60 cm depth of mobile sediment layers on hillslopes in the 555 Pan de Azucar and La Campana study areas, respectively. Also, the natural topography is steeper, with higher relief and rock 556 uplift rates might be different. Spatial variations in vegetation also occur (e.g., in La Campana), with higher vegetation density 557 along valleys, which might lead to the discrepancies between the observed and predicted sediment thickness. However, the trend in our results (higher sediment thickness for densely vegetated (70%) landscapes) follows the findings of Oeser et al. 558 559 (2018) who document that sediment increase with increasing mean annual precipitation and vegetation in Chilean Coastal 560 Cordillera.

In addition, previous field studies (Oeser et al., 2018; Owen et al., 2011; Schaller et al., 2018) applied cosmogenic nuclides to estimate the denudation and soil production rates in the Chilean Coastal Cordillera. They suggest an increase in soil production rates from arid zones in the north to wet tropical zones in the south of the Chilean Coastal Cordillera. These findings are consistent with the predicted increase in sediment depths (e.g. 1.24 m for V = 10% and 2.22 m for V = 70%, Fig. 7(b)) in our study. Finally, the effects of rock uplift and precipitation rates on topography and erosion rates, as documented by Bonnet and Crave (2003) and Lague et al. (2003) show a linear relationship between mean topographic elevation and rock uplift rate for steady-state conditions.

### 568 4.7 Model limitations

The model setup used in this study was intended to quantify the sensitivity of hillslope and fluvial erosion, and sediment transport and depositional processes for different climates with variations in precipitation rates and vegetation cover over Milankovitch time scales. This study was designed as an incremental step forward from previous modelling studies (Collins et al., 2004; Istanbulluoglu and Bras, 2005; Istanbulluoglu and Bras, 2006; Schmid et al., 2018).

There are several simplifying assumptions made in our modelling approach that warrant discussion and potential investigation in future studies. For example, this study assumed uniform vegetation cover and lithology for the entire catchment. The assumption of uniform vegetation cover in the catchment is likely reasonable given that relatively small  $(10x10 \text{ km}^2)$  size of catchments investigated and the modest topographic relief produced (between ~75-600m, Fig. 10a). Although temperature and precipitation (and therefore vegetation cover) can vary with elevation, the generally low relief of the catchments in this study do not make this a major concern. Due to the long (geologic) timescales considered in this study and computational considerations, mean annual precipitation rates were applied and stochastic distributions of precipitation could not be considered. While our approach is common for landscape evolution modelling studies conducted on geologic timescales, we recognize that in some settings (such as the arid region of this study, Fig. 1) precipitation events are rare, stochastic in nature, and might have an influence in the results presented here. This is a caveat that warrants future investigation.

The vegetation-erosion parameterization considered in this study follows from that of Istanbulluoglu and Bras (2006) and 583 Schmid et al. (2018). In this parameterization the total vegetation cover of the catchment is considered only, rather than the 584 585 distribution of vegetation cover by individual plant functional types (e.g. grass, shrubs, trees) that would have different Manning's coefficients associated with them. The 'total vegetation cover' approach used in our (and previous) work is a 586 reasonable starting point for understanding landscape evolution over large spatial and temporal scales because: a) more detailed 587 588 observations about the changes in the distribution of plant functional types over Milankovitch timescales is not available and 589 would be poorly constrained, and b) empirical relationships between total vegetation cover and precipitation are available and 590 easily implemented (e.g. Fig. 2b). However, future work should focus on exploring how the temporal and spatial distribution 591 of different plant functional types during changing climate impacts catchment erosion given that recent work (Mishra et al., 592 2019; Starke et al., 2020) has identified this as important. This limitation can be handled in future studies with the full coupling of a dynamic vegetation models, such as LPJ-GUESS (Smith et al., 2014; Werner et al., 2018) to a landscape evolution model 593 594 for the explicit treatment of how different vegetation types change temporally and spatially within a catchment and influence 595 catchment erosion. Also, the 'total vegetation cover' in the model is not disturbed by flow and entrainment, which were 596 observed to have a large impact on the results of Collins et al. (2004) and Istanbulluoglu and Bras (2005). If the vegetation 597 cover was spatio-temporally influenced by above processes in our simulations, the resulting erosion and sedimentation would have been hybrid between sparse (10% V) and densely vegetated (70% V) catchments, with vegetation losses in channels. The 598 599 time-scale for the current study was based on Milankovitch cycles, to address the effects of periodicity on erosion and 600 sedimentation. However, the effects of seasonal (sub annual) variations in precipitation (Istanbulluoglu and Bras, 2006; 601 Yetemen et al., 2015) and satellite derived vegetation cover (with catchment variable plant function type distributions) also 602 warrant future investigation to identify if coupled seasonal variations in vegetation cover and precipitation influence catchment 603 erosion.

Finally, the results of this study rely upon the vegetation-erosion parameterizations described in section 2 and the appendix (see also Fig. 3). While there is an observational basis for these relationships (see section 1.1, 1.2 in Appendix). There are, frankly, a sparse number of field studies available robustly constraining how different vegetation types and amounts influence hillslope and surface water erosional processes. Thus, we consider the erosional parameterizations used here as hypotheses (rather than robust geomorphic transport laws) that warrant investigation in future field or flume studies.

#### 609 5 Summary and Conclusions

610 In this study, we investigate the effects of variable vegetation cover and climate over Milankovitch timescales on catchment scale erosion and sedimentation. Simulations were presented to document if these transients are muted (lower amplitude) at 611 612 higher rock uplift rates. The approach used here complements previous studies by using a coupled fluvial detachment-transport limited and hillslope diffusion landscape evolution model, and also investigates the degree to which transient effects of 613 vegetation cover and precipitation are measurable in observational studies. The main conclusions deduced from this study are: 614 615 i. The step-wise increase in complexity of the model simulations was essential for identifying temporal changes in 616 catchment erosion and sediment thickness. A non-linear response in erosion and sediment thickness to varying 617 precipitation and vegetation cover was observed and results were dependent on the initial vegetation and precipitation

- state of the catchment. The sources of non-linearity stem from: a) a non-linear relationship between precipitation
  changes required to cause +/-10% change in vegetation cover (Fig. 2); and b) exponential and power-law relationships
  in the prescribed vegetation dependent hillslope and fluvial, respectively, geomorphic transport laws (Fig. 3, see also
  Appendix).
- 622ii.Analysis of results for covarying precipitation and vegetation cover indicate that erosion and sedimentation in densely623vegetated landscapes (V = 70%) are more heavily influenced by changes in precipitation than changes in vegetation624cover. This is due to the higher amplitude of precipitation change needed to cause variations in vegetation cover in625densely vegetated settings (Fig. 5a, 7e).
- Analysis of results for covarying precipitation and vegetation cover indicate that erosion and sedimentation in sparsely
   vegetated landscapes (V = 10%) are more sensitive to variable vegetation cover with constant precipitation rates (Fig.
   6, 7e), particular when precipitation rates decrease and vegetation cover approaches 0%.
- iv. Concerning the first hypotheses stated in the introduction: We found the effect of Milankovitch periodicity variations
   on the amplitude of change in sediment thickness and bedrock erosion is more pronounced for longer climate and
   vegetation oscillations (100 kyr) in both climate and vegetation settings. This finding confirms the hypothesis.
   Furthermore, periodicity effects on erosion and sediment thickness are larger in densely (70%) vegetated landscapes
   than sparsely (10%) vegetated landscapes, thereby indicating a sensitivity of the response to the biogeographic zone
   the changes are imposed on.
- 635 v. With respect to our second hypothesis: all transient forcings in precipitation and vegetation cover explored in this 636 study resulted in variations in erosion and sediment thickness around the mean erosion rate, which is determined by 637 the rock uplift rate. As rock uplift rates increased from 0.05 mm a<sup>-1</sup> to 0.2 mm a<sup>-1</sup>, the effects of periodic changes in precipitation and vegetation cover on erosion rates became more pronounced, and were between about 35% to 110%, 638 639 respectively, of the background rock uplift rate. This finding negates the hypothesis, and suggests that regardless of 640 the tectonic setting considered (within the range of rock uplift rates explored here) erosional transients from varying 641 precipitation and vegetation cover occur, but the detection of these changes requires measurement of erosion rates 642 integrating over short time scales such that the average (tectonically driven) mean erosion rate is not recovered.
- vi. Finally, in comparison to previous studies, the 35% to 110% transient changes in erosion rate documented here are
  at, or above, the detection limit for measurement cosmogenic radionuclides in river sediments preserved in fluvial
  terraces, but would be undetectable with bedrock thermochronometer dating techniques that average erosion rates
  over longer timescales. The potential to measure vegetation related transient changes in erosion rates with cosmogenic
  nuclides is highest in settings with higher rock uplift rates (e.g. 0.1 mm a<sup>-1</sup>, 0.2 mm a<sup>-1</sup>) and at longer (41 to 100 kyr)
  periodicities.

# 649 Appendix

#### 650 1 Effect of vegetation and precipitation on hillslope and fluvial erosion

The approach followed in our study follows the law of continuity of mass (e.g., Tucker et al., 2001). It states that the rate of change in topographic elevation (z) is defined as follows:

653 
$$\frac{\partial z}{\partial t} = U - \frac{\partial z}{\partial t} (fluvial) + \frac{\partial z}{\partial t} (hillslope),$$
 (1)

where, U is uplift rate [m yr<sup>-1</sup>], t is time [yr]. The second and third terms on right-hand side refer to the rate change in topographic elevation due to fluvial and hillslope processes respectively.

# 656 1.1 Vegetation dependent hillslope processes

657 The rate of change in topography due to hillslope diffusion (Fernandes and Dietrich, 1997; Martin, 2000) is defined as follows:

658 
$$\frac{\partial z}{\partial t}(hillslope) = \nabla q_s$$
, (2)

where  $q_s$  is sediment flux along the slope *S*. We applied slope and depth-dependent linear diffusion rule following the approach of Johnstone and Hilley (2014) such that:

661 
$$q_s = K_d S d_* \left( 1 - e^{-\frac{H}{d_*}} \right),$$
 (3)

where  $K_d$  is diffusion coefficient [m<sup>2</sup> yr<sup>-1</sup>],  $d_*$  is sediment transport decay depth [m], and *H* denotes sediment thickness. The diffusion coefficient is defined as a function of vegetation cover present on hillslopes, which is estimated following the approach of Istanbulluoglu (2005), Dunne et al. (2010) and (Schmid et al., 2018) as follows:

$$665 \quad K_d = K_b e^{-(\alpha V)} \,, \tag{4}$$

where  $K_d$  is defined as a function of vegetation cover *V*, an exponential decay coefficient  $\alpha$ , and linear diffusivity  $K_b$  for bare soil.

# 668 1.2 Vegetation dependent fluvial processes

The fluvial erosion is estimated for a two-layer topography (i.e., bedrock and sediment are treated explicitly) in the coupled detachment – transport limited model, SPACE 1.0 (Shobe et al., 2017). Bedrock erosion and sediment entrainment are calculated simultaneously in the model. Total fluvial erosion is defined as:

$$672 \quad \frac{\partial z}{\partial t} \left( fluvial \right) = \frac{\partial R}{\partial t} + \frac{\partial H}{\partial t}, \tag{5}$$

where, left-hand side denotes the total fluvial erosion rate. The first and second terms on right-hand side denote the bedrock erosion rate and sediment entrainment rate, respectively.

675 The rate of change of height of bedrock *R* per unit time  $[m yr^{-1}]$  is defined as:

$$676 \quad \frac{\partial R}{\partial t} = U - E_r \,, \tag{6}$$

677 where  $E_r$  [m yr<sup>-1</sup>], is the volumetric erosion flux of bedrock per unit bed area.

The change in sediment thickness H [m] per unit time [yr] was calculated following Davy and Lague (2009) and Shobe et al. (2017). It is defined as a fraction net deposition rate and solid fraction sediments, as follows:

$$680 \quad \frac{\partial H}{\partial t} = \frac{D_s - E_s}{1 - \emptyset},\tag{7}$$

where,  $D_s$  [m yr<sup>-1</sup>] is the deposition flux of sediment,  $E_s$  [m yr<sup>-1</sup>] is volumetric sediment entrainment flux per unit bed area, and  $\phi$  is the sediment porosity.

683 Following the approach of Shobe et al. (2017),  $E_s$  and  $E_r$  given by:

684 
$$E_s = (K_s q^m S^n - \omega_{cs}) \left( 1 - e^{-\frac{H}{H_*}} \right),$$
 (8)

685 
$$E_r = (K_r q^m S^n - \omega_{cr}) e^{-H/H_*}$$
, (9)

where,  $K_s$  [m<sup>-1</sup>] and  $K_r$  [m<sup>-1</sup>] are the sediment erodibility and bedrock erodibility parameters respectively. The threshold stream power for sediment entrainment and bedrock erosion are denoted as  $\omega_{cs}$  [m yr<sup>-1</sup>] and  $\omega_{cr}$  [m yr<sup>-1</sup>] in above equations. Bedrock roughness is denoted as  $H_*$  [m] and the term  $e^{-H/H_*}$  corresponds to the soil production from bedrock. With higher bedrock roughness magnitudes, more sediment would be produced.

 $K_s$  and  $K_r$  were modified in the model using the approach of Istanbulluoglu (2005) and Schmid et al. (2018) by introducing the effect of Manning's roughness to quantify the effect of vegetation cover on bed shear stress:

692 
$$\tau_v = \rho_w g(n_s + n_v)^{6/10} q^m S^n F_t$$
, (10)

693 where,  $\rho_w$  [kg m<sup>-3</sup>] and g [m s<sup>-2</sup>] are the density of water and acceleration due to gravity respectively. Manning's numbers for 694 bare soil and vegetated surface are denoted as  $n_s$  and  $n_v$ .  $F_t$  represents shear stress partitioning ratio. Manning's number for 695 vegetation cover and  $F_t$  are calculated as follows:

696 
$$n_v = n_{vr} \left(\frac{v}{v_r}\right)^w,\tag{11}$$

$$697 \quad F_t = \left(\frac{n_s}{n_s + n_p}\right)^{\frac{3}{2}},\tag{12}$$

698 where,  $n_{vr}$  is Manning's number for the reference vegetation. Here,  $V_r$  is reference vegetation cover (V = 100%) and V is local 699 vegetation cover in a model cell, w is empirical scaling factor.

Through combining stream power equation (Tucker et al., 1999; Howard, 1994; Whipple and Tucker, 1999) and the above concept of the effect of vegetation on shear stress, we follow the approach of Schmid et al. (2018) to define new sediment and bedrock erodibility parameters influenced by the surface vegetation cover on fluvial erosion, as follows:

703 
$$K_{vs} = K_s \rho_w g (n_s + n_v)^{6/10} F_t$$
, (13)

704 
$$K_{vr} = K_r \rho_w g(n_s + n_v)^{6/10} F_t$$
, (14)

where,  $K_{vs}$  [m<sup>-1</sup>] and  $K_{vr}$  [m<sup>-1</sup>] are modified sediment erodibility and bedrock erodibility respectively. These are influenced by fractional vegetation cover *V*. Hence,  $K_s$  and  $K_r$  in Eq. (8) and Eq. (9) are replaced by  $K_{vs}$  and  $K_{vr}$  to include an effect of vegetation cover on fluvial processes in the model. The trends of K<sub>d</sub>, K<sub>vs</sub> and K<sub>vr</sub> are illustrated in Fig. 3.

# Influence of coupled oscillations of precipitation and vegetation cover, on erosion and sedimentation (Scenario 3) without weathering function



710

Figure A1: Temporal evolution of catchment averaged predictions for scenario 3 (with no weathering) described in the text (section 3.3). Graphical representation of normalized mean catchment sedimentation and erosion to (a) coupled oscillations in precipitation [mm yr<sup>-1</sup>] and vegetation cover [-] in terms of (b) sediment thickness [-], (c) bedrock erosion [-], (d) mean erosion rate [-] for entire catchment. The periodicity of climate and vegetation oscillations is 23 kyr with rate of rock uplift as 0.5 mm yr<sup>-1</sup>.

# 715 Code availability

716 The code and data used in this study are freely available upon request.

# 717 Author contributions

718 HS and TAE designed the initial model setup and simulation programs. HS and TAE conducted model modifications,

719 simulation runs and analysis. MS provided assistance in program modification. KT provided insight into plant ecology in

720 Chilean Study areas and vegetation-climate change relationships. HS and TAE prepared the manuscript with contributions

721 from C.G., KT, and MS.

# 722 Competing interests

723 The authors declare that they have no competing interests.

### 724 Acknowledgements

725 H.S and T.A.E. acknowledge support by Open Access Publishing Fund of University of Tübingen. We also acknowledge

727 Foundation (DFG). In addition, T.A.E, K.T. and M.S. acknowledge support from the German priority research program (SPP-

support from the Research Training Group 1829 Integrated Hydrosystem Modelling, funded by the German Research

- 728 1803; grants EH329/14-1 and EH329/14-2 to T.A.E, and TI 338/14-1 and TI338/14-2 to K.T.). We thank Erkan Istanbulluoglu
- 729 and Omer Yetemen for their constructive reviews.

#### 730 References

726

- Acosta, V. T., Schildgen, T. F., Clarke, B. A., Scherler, D., Bookhagen, B., Wittmann, H., von Blanckenburg, F., and Strecker,
- 732 M. R.: Effect of vegetation cover on millennial-scale landscape denudation rates in East Africa, 7, 408-420,
- 733 https://doi.org/10.1130/1402.1, 2015.
- Ahnert, F.: Some comments on the quantitative formulation of geomorphological processes in a theoretical model, 2, 191– 201, https://doi.org/10.1002/esp.3290020211, 1977.
- 736 Alonso, R. N., Bookhagen, B., Carrapa, B., Coutand, I., Haschke, M., Hilley, G. E., Schoenbohm, L., Sobel, E. R., Strecker,
- 737 M. R., Trauth, M. H., and Villanueva, A.: Tectonics, Climate, and Landscape Evolution of the Southern Central Andes: the
- Argentine Puna Plateau and Adjacent Regions between 22 and 30°S, in: The Andes: Active Subduction Orogeny, edited by:
- 739 Oncken, O., Chong, G., Franz, G., Giese, P., Götze, H.-J., Ramos, V. A., Strecker, M. R., and Wigger, P., Springer Berlin
- 740 Heidelberg, Berlin, Heidelberg, 265–283, 2006.
- Amundson, R., Heimsath, A., Owen, J., Yoo, K., and Dietrich, W. E.: Hillslope soils and vegetation, 234, 122–132,
  https://doi.org/10.1016/j.geomorph.2014.12.031, 2015.
- Ashkenazy, Y., Eisenman, I., Gildor, H., and Tziperman, E.: The Effect of Milankovitch Variations in Insolation on Equatorial
  Seasonality, Journal of Climate, 23, 6133–6142, https://doi.org/10.1175/2010JCLI3700.1, 2010.
- Avdievitch, N. N., Ehlers, T. A., and Glotzbach, C.: Slow Long-Term Exhumation of the West Central Andean Plate Boundary,
  Chile, 37, 2243–2267, https://doi.org/10.1029/2017TC004944, 2018.
- Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., and Courchamp, F.: Impacts of climate change on the future of
  biodiversity, Ecology Letters, 15, 365–377, https://doi.org/10.1111/j.1461-0248.2011.01736.x, 2012.
- Bonnet, S. and Crave, A.: Landscape response to climate change: Insights from experimental modeling and implications for
  tectonic versus climatic uplift of topography, 31, 123–126, https://doi.org/10.1130/00917613(2003)031<0123:Lrtcci>2.0.Co;2, 2003.
- Braun, J., Voisin, C., Gourlan, A. T., and Chauvel, C.: Erosional response of an actively uplifting mountain belt to cyclic
  rainfall variations, Earth Surf. Dynam., 3, 1–14, https://doi.org/10.5194/esurf-3-1-2015, 2015.

- 754 Breckle, S.-W.: Walter's Vegetation of the Earth, 4th ed., Springer-Verlag Berlin Heidelberg, XX, 527 pp., 2002.
- Collins, D. B. G., Bras R. L., and Tucker G. E.: Modeling the effects of vegetation-erosion coupling on landscape evolution,
  109, https://doi.org/10.1029/2003jf000028, 2004.
- Dixon, J. L., Heimsath, A. M., and Amundson, R.: The critical role of climate and saprolite weathering in landscape evolution,
  34, 1507–1521, https://doi.org/10.1002/esp.1836, 2009.
- van Dongen, R., Scherler, D., Wittmann, H., and von Blanckenburg, F.: Cosmogenic <sup&gt;10&lt;/sup&gt;Be in river
   sediment: where grain size matters and why, 1–31, https://doi.org/10.5194/esurf-2018-83, 2018.
- Dosseto, A., Hesse, P. P., Maher, K., Fryirs, K., and Turner, S.: Climatic and vegetation control on sediment dynamics during
  the last glacial cycle, 38, 395–398, https://doi.org/10.1130/g30708.1, 2010.
- 763 Gerten, D., Luo, Y., Le Maire, G., Parton, W. J., KEOUGH, C., WENG, E., BEIER, C., CIAIS, P., CRAMER, W., DUKES,
- J. S., HANSON, P. J., KNAPP, A. A. K., LINDER, S., NEPSTAD, D., RUSTAD, L., and SOWERBY, A.: Modelled effects
  of precipitation on ecosystem carbon and water dynamics in different climatic zones, Global Change Biology, 14, 2365–2379,
  https://doi.org/10.1111/j.1365-2486.2008.01651.x, 2008.
- 767 Gilbert, G. K.: Report on the geology of the Henry Mountains, Washington, D.C., https://doi.org/10.3133/70039916, 1877.
- Hancock, G. S. and Anderson, R. S.: Numerical modeling of fluvial strath-terrace formation in response to oscillating climate,
  114, 1131–1142, https://doi.org/10.1130/0016-7606(2002)114<1131:NMOFST>2.0.CO;2, 2002.
- Heimsath, A. M., Dietrich, W. E., Nishiizumi, K., and Finkel, R. C.: The soil production function and landscape equilibrium,
  Nature, 388, 358–361, https://doi.org/10.1038/41056, 1997.
- 772 Herman, F., Rhodes, E. J., Braun, J., and Heiniger, L.: Uniform erosion rates and relief amplitude during glacial cycles in the
- 773 Southern Alps of New Zealand, as revealed from OSL-thermochronology, Earth and Planetary Science Letters, 297, 183–189,
- 774 https://doi.org/10.1016/j.epsl.2010.06.019, 2010.
- Hobley, D. E. J., Adams, J. M., Nudurupati, S. S., Hutton, E. W. H., Gasparini, N. M., Istanbulluoglu, E., and Tucker, G. E.:
- Creative computing with Landlab: an open-source toolkit for building, coupling, and exploring two-dimensional numerical
  models of Earth-surface dynamics, 5, 21–46, https://doi.org/10.5194/esurf-5-21-2017, 2017.
- 778 Howard, A. D.: A detachment-limited model of drainage basin evolution, 2261-2285, 1994.
- Huntley, B., Allen, J. R., Collingham, Y. C., Hickler, T., Lister, A. M., Singarayer, J., Stuart, A. J., Sykes, M. T., and Valdes,
  P. J.: Millennial climatic fluctuations are key to the structure of last glacial ecosystems, 8, e61963,
  https://doi.org/10.1371/journal.pone.0061963, 2013.
- 782 Huxman, T. E., Smith, M. D., Fay, P. A., Knapp, A. K., Shaw, M. R., Loik, M. E., Smith, S. D., Tissue, D. T., Zak, J. C., 783 Weltzin, J. F., Pockman, W. T., Sala, O. E., Haddad, B. M., Harte, J., Koch, G. W., Schwinning, S., Small, E. E., and Williams, 784 D. Convergence efficiency, Nature, 429, 651-654, G.: across biomes to а common rain-use 785 https://doi.org/10.1038/nature02561, 2004.

- Hyun, S., Ahagon, N., and Yoon, H.-I.: Milankovitch cycles and paleoceanographic evolution within sediments from ODP
  Sites 980 and 983 of the North Atlantic Ocean, 9, 235, https://doi.org/10.1007/BF02910583, 2005.
- Istanbulluoglu, E. and Bras, R. L.: Vegetation-modulated landscape evolution: Effects of vegetation on landscape processes,
  drainage density, and topography, 110, https://doi.org/10.1029/2004jf000249, 2005.
- Istanbulluoglu, E. and Bras, R. L.: On the dynamics of soil moisture, vegetation, and erosion: Implications of climate variability
  and change, 42, https://doi.org/10.1029/2005wr004113, 2006.
- Jeffery, M. L., Yanites, B. J., Poulsen, C. J., and Ehlers, T. A.: Vegetation-precipitation controls on Central Andean topography, 119, 1354–1375, https://doi.org/10.1002/2013jf002919, 2014.
- Johnstone, S. A. and Hilley, G. E.: Lithologic control on the form of soil-mantled hillslopes, 43, 83–86, https://doi.org/10.1130/g36052.1, 2014.
- Kelly, A. E. and Goulden, M. L.: Rapid shifts in plant distribution with recent climate change, Proc Natl Acad Sci USA, 105,
  11823, https://doi.org/10.1073/pnas.0802891105, 2008.
- Kirby, E. and Whipple, K. X.: Expression of active tectonics in erosional landscapes, 44, 54–75, https://doi.org/10.1016/j.jsg.2012.07.009, 2012.
- Knapp, A. K., Ciais, P., and Smith, M. D.: Reconciling inconsistencies in precipitation–productivity relationships: implications
  for climate change, New Phytologist, 214, 41–47, https://doi.org/10.1111/nph.14381, 2017.
- Kojima, S., Soto, I., Quiroz, M., Valencia, P., and Fernandez, I.: Geological and Geochemical Characteristics of the IntrusionRelated Vein-Type Gold Deposits in the El Morado District, Coastal Cordillera, Northern Chile, 67, 197–206,
  https://doi.org/10.1111/rge.12129, 2017.
- Lague, D., Crave, A., and Davy, P.: Laboratory experiments simulating the geomorphic response to tectonic uplift, 108, ETG
  3-1-ETG 3-20, https://doi.org/10.1029/2002jb001785, 2003.
- McPhillips, D., Bierman, P. R., Crocker, T., and Rood, D. H.: Landscape response to Pleistocene-Holocene precipitation
  change in the Western Cordillera, Peru:10Be concentrations in modern sediments and terrace fills, 118, 2488–2499,
  https://doi.org/10.1002/2013jf002837, 2013.
- 810 Melnick, D.: Rise of the central Andean coast by earthquakes straddling the Moho, 9, 401–407, 811 https://doi.org/10.1038/ngeo2683, 2016.
- Miller, S. R., Sak, P. B., Kirby, E., and Bierman, P. R.: Neogene rejuvenation of central Appalachian topography: Evidence
  for differential rock uplift from stream profiles and erosion rates, 369–370, 1–12, https://doi.org/10.1016/j.epsl.2013.04.007,
  2013.
- Mishra, A. K., Placzek, C., and Jones, R.: Coupled influence of precipitation and vegetation on millennial-scale erosion rates
  derived from 10Be, 14, e0211325, https://doi.org/10.1371/journal.pone.0211325, 2019.

- 817 Mowll, W., Blumenthal, D. M., Cherwin, K., Smith, A., Symstad, A. J., Vermeire, L. T., Collins, S. L., Smith, M. D., and
- 818 Knapp, A. K.: Climatic controls of aboveground net primary production in semi-arid grasslands along a latitudinal gradient
- 819 portend low sensitivity to warming, Oecologia, 177, 959–969, https://doi.org/10.1007/s00442-015-3232-7, 2015.
- 820 Oeser, R. A., Stroncik, N., Moskwa, L.-M., Bernhard, N., Schaller, M., Canessa, R., van den Brink, L., Köster, M., Brucker,

821 E., Stock, S., Fuentes, J. P., Godoy, R., Matus, F. J., Oses Pedraza, R., Osses McIntyre, P., Paulino, L., Seguel, O., Bader, M.

822 Y., Boy, J., Dippold, M. A., Ehlers, T. A., Kühn, P., Kuzyakov, Y., Leinweber, P., Scholten, T., Spielvogel, S., Spohn, M.,

823 Übernickel, K., Tielbörger, K., Wagner, D., and von Blanckenburg, F.: Chemistry and microbiology of the Critical Zone along

- 824 a steep climate and vegetation gradient in the Chilean Coastal Cordillera, CATENA, 170, 183-203,
- 825 https://doi.org/10.1016/j.catena.2018.06.002, 2018.
- Owen, J. J., Amundson, R., Dietrich, W. E., Nishiizumi, K., Sutter, B., and Chong, G.: The sensitivity of hillslope bedrock erosion to precipitation, 36, 117–135, https://doi.org/10.1002/esp.2083, 2011.
- Pelletier, J. D.: Fluvial and slope-wash erosion of soil-mantled landscapes: detachment- or transport-limited?, 37, 37–51,
  https://doi.org/10.1002/esp.2187, 2012.
- Perron, J. T.: Climate and the Pace of Erosional Landscape Evolution, 45, 561–591, https://doi.org/10.1146/annurev-earth060614-105405, 2017.
- Rossel, K., Aguilar, G., Salazar, E., Martinod, J., Carretier, S., Pinto, L., and Cabré, A.: Chronology of Chilean Frontal
  Cordillera building from geochronological, stratigraphic and geomorphological data insights from Miocene intramontanebasin deposits, 30, 289–310, https://doi.org/10.1111/bre.12221, 2018.
- Routschek, A., Schmidt, J., and Kreienkamp, F.: Impact of climate change on soil erosion A high-resolution projection on
  catchment scale until 2100 in Saxony/Germany, 121, 99–109, https://doi.org/10.1016/j.catena.2014.04.019, 2014.
- Sala, O. E., Parton, W. J., Joyce, L. A., and Lauenroth, W. K.: Primary Production of the Central Grassland Region of the
  United States, Ecology, 69, 40–45, https://doi.org/10.2307/1943158, 1988.
- Schaller, M. and Ehlers, T. A.: Limits to quantifying climate driven changes in denudation rates with cosmogenic
  radionuclides, 248, 153–167, https://doi.org/10.1016/j.epsl.2006.05.027, 2006.
- 841 Schaller, M., Blanckenburg, F. von, Hovius, N., Veldkamp, A., van den Berg, M. W., and Kubik, P. W.: Paleoerosion Rates
- from Cosmogenic 10Be in a 1.3 Ma Terrace Sequence: Response of the River Meuse to Changes in Climate and Rock Uplift,
- 843 The Journal of Geology, 112, 127–144, https://doi.org/10.1086/381654, 2004.
- 844 Schaller, M., Ehlers, T. A., Lang, K. A. H., Schmid, M., and Fuentes-Espoz, J. P.: Addressing the contribution of climate and 845 Cordillera 489, vegetation cover on hillslope denudation, Chilean Coastal  $(26^{\circ}-38^{\circ}S),$ 111 - 122,https://doi.org/10.1016/j.epsl.2018.02.026, 2018. 846
- Schaller, M., Dal Bo, I., Ehlers, T. A., Klotzsche, A., Drews, R., Fuentes Espoz, J. P., and van der Kruk, J.: Comparison of
  regolith physical and chemical characteristics with geophysical data along a climate and ecological gradient, Chilean Coastal
  Cordillera (26 to 38\degree\,S), 6, 629–647, https://doi.org/10.5194/soil-6-629-2020, 2020.

850 Schmid, M., Ehlers, T. A., Werner, C., Hickler, T., and Fuentes-Espoz, J.-P.: Effect of changing vegetation and precipitation

851 on denudation - Part 2: Predicted landscape response to transient climate and vegetation cover over millennial to million-year

- 852 timescales, 6, 859–881, https://doi.org/10.5194/esurf-6-859-2018, 2018.
- Seybold, H., Rothman, D. H., and Kirchner, J. W.: Climate's watermark in the geometry of stream networks, 44, 2272–2280,
  https://doi.org/10.1002/2016gl072089, 2017.
- Shobe, C. M., Tucker, G. E., and Barnhart, K. R.: The SPACE 1.0 model: A Landlab component for 2-D calculation of
  sediment transport, bedrock erosion, and landscape evolution, 1–38, https://doi.org/10.5194/gmd-2017-175, 2017.
- Slater, L. J. and Singer, M. B.: Imprint of climate and climate change in alluvial riverbeds: Continental United States, 19502011, 41, 595–598, https://doi.org/10.1130/g34070.1, 2013.
- Smith, B., W\aarlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., and Zaehle, S.: Implications of incorporating N
  cycling and N limitations on primary production in an individual-based dynamic vegetation model, 11, 2027–2054,
  https://doi.org/10.5194/bg-11-2027-2014, 2014.
- Smith, M. D., Wilcox, K. R., Power, S. A., Tissue, D. T., and Knapp, A. K.: Assessing community and ecosystem sensitivity
  to climate change toward a more comparative approach, Journal of Vegetation Science, 28, 235–237,
  https://doi.org/10.1111/jvs.12524, 2017.
- Starke, J., Ehlers, T. A., and Schaller, M.: Latitudinal effect of vegetation on erosion rates identified along western South
  America, Science, 367, 1358, https://doi.org/10.1126/science.aaz0840, 2020.
- Tucker, G. E.: Drainage basin sensitivity to tectonic and climatic forcing: implications of a stochastic model for the role of entrainment and erosion thresholds, 29, 185–205, https://doi.org/10.1002/esp.1020, 2004.
- Tucker, G. E., Gasparini, N. M., Lancaster, S. T., and Bras, R. L.: Modeling Floodplain Dynamics and Stratigraphy:
  Implications for Geoarchaeology, 1999.
- Turowski, J. M., Lague, D., Crave, A., and Hovius, N.: Experimental channel response to tectonic uplift, 111, n/a-n/a, https://doi.org/10.1029/2005jf000306, 2006.
- Werner, C., Schmid, M., Ehlers, T. A., Fuentes-Espoz, J. P., Steinkamp, J., Forrest, M., Liakka, J., Maldonado, A., and Hickler,
  T.: Effect of changing vegetation and precipitation on denudation Part 1: Predicted vegetation composition and cover over
- the last 21 thousand years along the Coastal Cordillera of Chile, 6, 829–858, https://doi.org/10.5194/esurf-6-829-2018, 2018.
- Whipple, K. X.: The influence of climate on the tectonic evolution of mountain belts, Nature Geoscience, 2, 730–730,
  https://doi.org/10.1038/ngeo638, 2009.
- Whipple, K. X. and Tucker, G. E.: Dynamics of the stream-power river incision model: Implications for height limits of
  mountain ranges, landscape response timescales, and research needs, 104, 17661–17674,
  https://doi.org/10.1029/1999jb900120, 1999.
- Willgoose, G., Bras, R. L., and Rodriguez-Iturbe, I.: A coupled channel network growth and hillslope evolution model: 1.
  Theory, 27, 1671–1684, https://doi.org/10.1029/91WR00935, 1991.

Xia, J., Chen, J., Piao, S., Ciais, P., Luo, Y., and Wan, S.: Terrestrial carbon cycle affected by non-uniform climate warming,
Nature Geoscience, 7, 173–180, https://doi.org/10.1038/ngeo2093, 2014.

Yang, Y., Fang, J., ma, W., and Wang, W.: Relationship between variability in aboveground net primary production and
precipitation in global grasslands. Geophysical Research Letters, Geophysical Research Letters - GEOPHYS RES LETT, 35,
https://doi.org/10.1029/2008GL035408, 2008.

Yanites, B. J. and Ehlers, T. A.: Global climate and tectonic controls on the denudation of glaciated mountains, Earth and
Planetary Science Letters, 325–326, 63–75, https://doi.org/10.1016/j.epsl.2012.01.030, 2012.

Yetemen, O., Istanbulluoglu, E., and Duvall, A. R.: Solar radiation as a global driver of hillslope asymmetry: Insights from an
ecogeomorphic landscape evolution model, Water Resources Research, 51, 9843–9861,
https://doi.org/10.1002/2015WR017103, 2015.

Yetemen, O., Saco, P. M., and Istanbulluoglu, E.: Ecohydrology Controls the Geomorphic Response to Climate Change,
Geophysical Research Letters, 46, 8852–8861, https://doi.org/10.1029/2019GL083874, 2019.

895 Zhang, Y., Xiao, X., Guanter, L., Zhou, S., Ciais, P., Joiner, J., Sitch, S., Wu, X., Nabel, J., Dong, J., Kato, E., Jain, A. K.,

896 Wiltshire, A., and Stocker, B. D.: Precipitation and carbon-water coupling jointly control the interannual variability of global

land gross primary production, Scientific Reports, 6, 39748, https://doi.org/10.1038/srep39748, 2016.

898