

Temporal changes in the debris flow threshold under the effects of ground freezing and sediment storage on Mt. Fuji

Fumitoshi Imaizumi¹, Atsushi Ikeda², Kazuki Yamamoto³, Okihiro Ohsaka⁴

¹Faculty of Agriculture, Shizuoka University, Shizuoka, 422-8429, Japan

5 ²Faculty of Life and Environmental Sciences, University of Tsukuba, 305-8572, Japan

³Bureau of Economic Affairs, Shizuoka city office, Shizuoka, 422-8429, Japan

⁴Shizuoka Professional University of Agriculture, Japan, 424-8701

Correspondence to: Fumitoshi Imaizumi (imaizumi@shizuoka.ac.jp)

Abstract. Debris flows are one of the most destructive sediment transport processes in mountainous areas because of their
10 large volume, high velocity, and kinematic energy. Debris flow activity varies over time and is affected by changes in
hydrogeomorphic processes in the initiation zone. To clarify temporal changes of debris flow activities in cold regions, the
rainfall threshold for the debris flow occurrence was evaluated in Osawa failure at a high elevation on Mt. Fuji, Japan. We
conducted field monitoring of the ground temperature near a debris flow initiation zone to estimate the presence or absence of
seasonally frozen ground during historical rainfall events. The effects of ground freezing and the accumulation of channel
15 deposits on the rainfall threshold for debris flow occurrence were analyzed using rainfall records and annual changes in the
volume of channel deposits since 1969. Statistical analyses showed that the intensity-duration threshold during frozen periods
was clearly lower than that during unfrozen periods. A comparison of maximum hourly rainfall intensity and total rainfall also
showed that debris flows during frozen periods were triggered by a smaller magnitude of rainfall than during unfrozen periods.
Decreases in the infiltration rate due to the formation of frozen ground likely facilitated the generation of overland flow,
20 triggering debris flows. During unfrozen periods, the ~~rainfall threshold was~~ *value of maximum hourly rainfall intensity
triggering sediment transfer was* higher when the volume of channel deposits was larger. ~~Increases in the water content in
channel deposits caused by the infiltration of rainfall is likely important for the debris flow occurrence during unfrozen periods.~~
The results suggest that the occurrence of frozen ground and the sediment storage volume need to be monitored and estimated
for better debris flow disaster mitigation in cold regions.

25 1 Introduction

Debris flows are one of the most destructive sediment transport processes in mountain areas because of their large volume and
high velocity and kinematic energy (Scott et al., 2004; Theule et al., 2018; Song et al., 2019). Debris flow risks are increasing
in cold mountainous regions under climate change due to changes in hydrological processes on hillslopes, increases in the
magnitude of rainfall events, and those in sediment supply activity in their initiation zones (Stoffel and Huggel, 2012; Stoffel
30 et al., [2014a](#), [2014b](#)). Because debris flows are usually triggered by heavy rainfall events (Scott et al., 2004; Sidle and Chigira,

2004; Dowling and Santi, 2014; Zhang et al., 2019), estimation of the rainfall thresholds triggering debris flows is essential for debris flow hazard mitigation measures, such as the installation of early warning systems (Pan et al., 2018; Hürlimann et al., 2019).

Many studies have attempted to determine rainfall thresholds by analyzing rainfall records in periods with and without a debris flow (e.g., Cannon et al., 2011; Hürlimann et al., 2019). The intensity-duration (ID) threshold is one of the most commonly used rainfall thresholds (Caine, 1980; Guzzetti et al., 2008; Staley et al., 2012; Zhou and Tang, 2014). Other factors, such as cumulative rainfall depth and the return time of rainfall, have also been used to estimate rainfall thresholds (Fiorillo and Wilson, 2004; Imaizumi et al., 2006). Although many studies have calculated a single rainfall threshold for each torrent and region, the threshold varies with time and is affected by changes in hydrological processes and the volume of debris flow material (Jakob et al., 2005; Schlunegger et al., 2009; Chen et al., 2012; Theule et al., 2012). Therefore, temporal changes in the rainfall threshold need to be understood to evaluate future risks of debris flows under climate change.

Debris flow activity is controlled by hydrological characteristics on hillslopes and debris flow material (i.e., channel deposits and talus slopes) in debris flow initiation zones (Bovis and Dagg, 1988; Staley et al., 2014; Loye et al., 2016). ~~Wildfires~~ ~~Decreases in~~ the infiltration rate of rainfall into the ground, ~~resulting results in~~ the generation of overland flow that leads to debris flows (Cannon et al., 2001; Shakesby and Doerr, 2006; Ebel et al., 2012; Staley et al., 2014). The hydraulic conductivity of debris flow material also controls the initiation of debris flows (Bovis and Dagg, 1988).

~~In cold regions,~~ Ground freezing changes hydrological processes in watersheds (Blackburn et al., 1990; Iwata et al., 2010; Coles and McDonnell, 2018-). ~~with the~~ infiltration rate and hydraulic conductivity are generally low in periods with a frozen soil layer (Blackburn et al., 1990; Iwata et al., 2010). Thus, the discharge of shallow runoff during the frozen period is higher than during the unfrozen period (Shanley and Chalmers, 1999; Coles and McDonnell, 2018). ~~Space for liquid water in soil matrix decreases by formation of ice (Zhao and Gray, 1990; Watanabe and Flury, 2008), possibly affecting initiation condition of debris flow.~~ However, the effect of ground freezing on the rainfall threshold of debris flow remains poorly understood.

The volume of debris flow material in the initiation zones of debris flow is controlled by the sediment supply from hillslopes and the evacuation of sediment by debris flows and fluvial processes (Imaizumi et al., 2006; Berger et al., 2011a; Theule et al., 2012; Imaizumi et al., 2019; Rengers et al., 2020a). Debris flow frequency is generally high in torrents with high sediment supply activity (May, 2002; Jakob et al., 2005). Bovis and Jakob (1999) classified debris flow basins into weathering-limited basins, which require a long sediment recharge period for the occurrence of a debris flow, and transport-limited basins, in which the debris flow occurrence is primarily controlled by hydroclimatic events. These studies imply that the volume of debris flow material in the watershed needs to be interpreted to improve the accuracy of debris flow prediction.

Mount Fuji, which is the most famous mountain of Japan, has frequently experienced severe debris flow disasters, as well as slush avalanches, which are mixtures of water and snow that erode large amounts of sediment as they move downstream (Hanaoka et al., 2007). Sediment accumulated in gully bottoms, which is eroded by overland flow during rainfall events, is the main source of debris flow material. Many disasters around Mt. Fuji have occurred in spring and late autumn, despite precipitation in those seasons being lower than in summer and early autumn (Hanaoka et al., 2007). Although previous

コメントの追加 [A1]: Rev.1-4

コメントの追加 [A2]: Rev.1-4,1-5

コメントの追加 [A3]: Rev.2-6

65 interpretations have implied that seasonal ground freezing affects temporal changes in the rainfall threshold of debris flow
(Anma, 2007), this has not been supported by quantitative analyses due to the lack of field monitoring data. On Mt. Fuji, the
sediment storage volume in debris flow torrents has also changed over time because of the sediment supply from hillslopes
and the evacuation of sediment by debris flows (Hanaoka et al., 2007).

70 The overall aim of this study was to clarify temporal changes in the rainfall threshold for the debris flow occurrence in a
cold region under the effects of seasonal ground freezing and changes in the sediment storage volume. We conducted field
monitoring of ground temperatures on Mt. Fuji to estimate the ground freezing conditions during historical rainfall events that
triggered or did not trigger debris flows and slush avalanches. The specific objectives of the study were to estimate periods of
ground freezing over 50 years using an empirical model based on field monitoring, calculate the rainfall thresholds in periods
with and without frozen ground given changes in the sediment storage volume, and understand the hydrological processes
75 triggering debris flows in relation to the formation of frozen ground and the volume of debris flow material.

2 Study site

The Osawa failure, which is a huge gully (length 2.1 km, width 500 m, and depth 150 m), is the most active debris flow torrent
on Mt. Fuji. The Osawa failure is located on the western slope of Mt. Fuji (Figs. 1a, 1b), where a westerly wind dominates.
Results of age determination using ^{14}C indicated that this gully started to form circa 3,000 years ago (Iwatsuka and Machida,
80 1962). The Osawa failure increased in ~~size-length and width~~ following a catastrophic failure circa 1,000 years ago (Hanaoka
et al., 2007). The total volume of sediment derived from the gully is estimated to be 75,000,000 m³. The west side of Mt. Fuji
is underlain by an alternation of scoria and basaltic lava, whereas thick tephra layers occur on the east side (Miyaji, 1998).
Unconsolidated scoria layers have been actively eroded by freeze-thaw activity, overland flows, and strong winds. As the
erosion of basal scoria layers proceeds, the lava outcrop becomes unstable due to overhanging, resulting in episodic failure of
85 the outcrop of up to 30 m in thickness. Between 1970 to 2004, the estimated total volume of sediment evacuated from the
Osawa failure was 5,020,000 m³ (Hanaoka et al., 2007).

The Osawa failure has a cold climate due to its high elevation. The monthly average air temperature around the debris flow
initiation zone (3,200 m a.s.l.) is lower than 0°C from November to April (Fig. 2). The maximum daily air temperature is also
below 0°C from November to April, ~~facilitating the development of seasonally frozen ground~~. The average annual precipitation
90 from 2010 to 2019 was 2,737 mm yr⁻¹. Heavy rainfall (total rainfall >100 mm) occurs during the “Baiu” rainy season (i.e.,
monsoon season from June to July) and during typhoon crossings (August to October). Precipitation in winter (December to
February) accounts for only 8% of the annual precipitation. This relatively dry climate in winter promotes seasonal ground
freezing under negative air temperatures, because the thermal insulation by thick snow is inhibited except for the leeward
sites depressions accumulating, where drifted snow accumulates (Ikeda et al., 2012). Spring (March to April) is the main
95 snowfall season on Mt. Fuji, and the study on the summit revealed that the drifted snow prevented both further ground cooling
and the following warming in the following early summer. In contrast to the leeward sites on the summit, snow insulation in

コメントの追加 [A4]: Rev.2-7

コメントの追加 [A5]: Rev.1-7

コメントの追加 [A6]: Rev.2-8


書式変更: 上付き

the Osawa failure must be limited in the main snowfall season, because snow disappearance in mid-April, 2018, and late April, 2019, on the debris flow initiation zone (except for the narrow valley bottom) was observed by a distant-view camera installed at the western foot of the mountain by Shizuoka Prefecture.

100 In the Osawa failure, the slope gradients of the scoria and lava layers are 35–50° and 40–90°, respectively. Talus slopes, with a slope gradient of 35–40°, are distributed at the foot of the hillslopes. Most parts of the channel are generally covered by channel deposits, which are the main sources of debris flow material (Figs. 1c, 1d). Sediment supply processes are different between scoria and basalt. Scoria is mainly transported into channels by soil creep and surface wash, while basalt is transported by rockfall and dry ravel in relation to the failure of steep outcrops and secondary movement of deposits on hillslopes. The channel gradient in the gully ranges from 20° (the lower part of the Osawa failure) to 35° (channel head). Debris flows and slush avalanches in the Osawa failure generally run down Osawa Creek to the Osawa alluvial fan, which is located approximately 6 km west of the gully. Some small-scale debris flows terminate in the Osawa failure and the channel sections between the gully and alluvial fan.

コメントの追加 [A7]: Rev.1-2

コメントの追加 [A8]: Rev.1-6



110 Figure 1: Topographic map and photographs of the Osawa failure. (a) Topographic map of the area around the Osawa failure and Osawa Creek. The area shown in (d) is located within the black-outlined box. (b) Full view of Mt. Fuji. The location of the Osawa failure is indicated by the white dashed line. (c) Channel deposits in the Osawa failure. The location where this photograph was taken is indicated by the red arrow in (d). (d) Topographic map of the area around the Osawa failure. RG and TM indicate the locations of a rain gauge and ground temperature sensors, respectively. The summit meteorological station is located on the triangle of the mountain peak.

コメントの追加 [A9]: Rev.1-9




Figure 2: Average monthly precipitation at 2,330 m a.s.l. (RG in Fig. 1d) and air temperature around the debris flow initiation zone (3,200 m a.s.l.) in the Osawa failure from 2010 to 2019. Air temperature around the debris flow initiation zone was estimated from that data recorded at the summit station (3,775 m a.s.l.) and the lapse rate (0.55). The solid air temperature curve indicates monthly average temperature, while the upper and lower bounds of the shaded area indicate the average daily maximum and minimum temperatures, respectively, in each month.

120

125

130

135

The Ministry of Land, Infrastructure, Transport, and Tourism (MLIT) has been monitoring debris flows and slush avalanches in the Osawa failure and Osawa Creek since 1969. The runout timing of previous debris flows was observed with video cameras and water level gauges at Osawagawa Bridge Observatory (490 m a.s.l.) and Iwadoi Observatory (900 m a.s.l.), as well as with wire sensors and video cameras at Otaki Observatory (1,500 m a.s.l., Fig. 1a). Some debris flows were just monitored at one or two observatories because of the absence of monitoring devices, problems encountered during monitoring, or the short travel distance of the debris flows. The MLIT has also distinguished slush avalanches from debris flows based on field surveys just after each event and the presence or absence of snow in the flow as captured by video cameras. A total of 35 debris flows and eight slush avalanches were monitored from 1969 to 2019 (Fuji Sabo Office, 2017; Fuji Sabo Office, 2020). Four of the 35 debris flows were reclassified into slush avalanches, because snowpack over debris flow initiation zone was identified by a distant-view camera. The maximum total discharge of a single debris flow, which was estimated with a video camera, ultrasonic water level gauge, and Doppler flow-velocity sensor at Iwadoi observatory, was $3 \times 10^5 \text{ m}^3$ (Fuji Sabo Office, 2017).

コメントの追加 [A10]: Rev.2-9

3 Methodology

3.1 Ground temperature monitoring

140

A depth profile of the ground temperature was obtained at an elevation of 3,220 m a.s.l. along the northern boundary of the Osawa failure (TM in Fig. 1d). The elevation of the monitoring point was selected based on the known initiation points of many debris flows, which were determined in an analysis of aerial photographs and airborne Light Detection and Ranging (LiDAR) digital elevation models (DEMs). A 2-m-deep hole was drilled in welded scoria. Then, temperature sensors (TMC-HD, Onset) were installed at 10 different depths (ground surface, 0.1, 0.25, 0.5, 0.75, 1.0, 1.25, 1.5, 1.75, and 2 m vertically)

and connected to data loggers (U12-008, Onset). Errors in temperature measurement after calibration of the sensors were within $\pm 0.1^\circ\text{C}$. Monitoring was conducted from September 30, 2017 to September 17, 2019, with temperature recorded at intervals of 1 hr. The sensor at the ground surface was covered by pebbles to match its albedo to that of the natural ground surface.

The occurrence of diurnal freeze-thaw activity and the formation of seasonally frozen ground were determined based on observed ground temperature data. The period with diurnal freeze-thaw activity was determined to be the period during which the ground surface temperature rose above and fell below 0°C within 1 calendar day, whereas the period with seasonal ground freezing was determined to be the period during which the ground temperature was equal to or below 0°C over multiple days.

3.2 Estimation of frozen periods

To estimate the presence or absence of seasonally frozen ground during rainfall events before the monitoring of ground temperature began, the downward progress of the freezing and thawing fronts at the debris flow initiation zone since 1969 was estimated using air temperature records at the summit of Mt. Fuji (3,775 m a.s.l.), which were provided by the Japan Meteorological Agency (JMA). Because the ground-freezing period is spatially variable and affected by slope orientation, elevation, and snow distribution (e.g., Bayard et al., 2005; Kneisel et al., 2015), it is difficult to precisely estimate the frozen period at an exact debris flow initiation point, which differs among debris flow events inside the Osawa failure. Therefore, we estimated the frozen period at TM (Fig. 1d) using the simple degree-day method to conduct an initial evaluation of the effect of ground freezing on the debris flow occurrence.

First, the ground surface temperature at the debris flow initiation zone TM was estimated using a linear ordinary least squares regression equation for the relationship between daily average air temperature at the summit station and the monitored daily average ground surface temperature at TM from September 30, 2017 to September 17, 2019. Although the amplitude of diurnal changes in the ground surface temperature was generally higher than that of air temperature, we used the empirical model of the relationship between air temperature and ground surface temperature due to their high correlation ($R^2 = 0.94$). Then, the downward progress of the freezing and thawing fronts was obtained using the degree-day method, which was obtained from an approximate solution of the unsteady heat conduction equation from freezing degree days and thawing degree days, respectively, which were calculated from the estimated daily average ground surface temperature (Yamazaki, 1998; Miao et al., 2019; Walvoord et al., 2019) as follows (Yamazaki, 1998; Miao et al., 2019; Walvoord et al., 2019):

$$D_f = c_f(-\sum T(t))^{1/2}, T(t) < 0 \quad (1)$$

$$D_m = c_m(\sum T(t))^{1/2}, T(t) > 0 \quad (2)$$

where D_f is the depth of the freezing front, $T(t)$ is the estimated daily average ground surface temperature at TM, D_m is the depth of the thawing front, and c_f and c_m are coefficients derived from thermal conduction, moisture content ratio, heat of fusion and other parameters, are constant. The coefficients c_f and c_m , which reflect temporal changes in the freezing and thawing

コメントの追加 [A11]: Rev.1-8

コメントの追加 [A12]: Rev.1-9

コメントの追加 [A13]: Rev.1-10, 12

コメントの追加 [A14]: Rev.1-12

コメントの追加 [A15]: Rev.1-11

書式変更: 上付き/下付き(なし)

コメントの追加 [A16]: Rev.1-13

コメントの追加 [A17]: Rev.1-12

コメントの追加 [A18]: Rev.2-10

書式変更: フォント: 斜体

コメントの追加 [A19]: Rev.1-13

depths as calculated from monitoring data, were used for the estimation of the freezing and thawing fronts ~~before field monitoring began.~~ This degree-day method assumes that sediment characteristics and the moisture content ratio are spatially and temporally constant.

コメントの追加 [A20]: Rev.1-13

Air temperature at the Fuji City meteorological station (20 km south of the Osawa failure), which is also maintained by the JMA, was used to estimate the ground temperature in the period for which data from the summit station were unavailable (0.9% of 1969–2019), although the elevation of this station (66 m a.s.l.) is much lower than that of the debris flow initiation zone. The regression equation for the relationship between the air temperature at the Fuji City station and ground surface temperature at the debris flow initiation zone (TM) was used to estimate the freezing and thawing lines in a depth-time graph ($R^2=0.93$). In this study, the periods between freezing and thawing lines are referred to as frozen periods, whereas the periods between thawing and subsequent freezing lines are referred to as unfrozen periods.

The site TM, which lacks snow cover almost all year-round, was chosen as a safe location instead of sites along the inaccessible paths of the debris flows in the Osawa failure. Except for the gully bottom, snow cover in the Osawa failure is generally shallow or absent, ~~even in~~ until mid-winter, due to its slope orientation (windward side of Mt. Fuji), steep topography, and low precipitation (Fig. 2). However, snow can accumulate in spring, although no quantitative data regarding the snow distribution were available. Therefore, we ~~preliminarily did not consider snow cover to estimate~~ the freezing and thawing lines based on the TM data. Thermal differences caused by snow cover differences between TM and the initiation points of debris flow are then discussed.

コメントの追加 [A21]: Rev.1-2

3.3 Rainfall monitoring

From May 1, 1969, the MLIT has observed precipitation at 2,330 m a.s.l. using a tipping-bucket rain gauge, with a precision of 1 mm, close to the Osawa failure. The rain gauge, which has an internal heating system, can also monitor winter precipitation. The observation of precipitation was occasionally interrupted by problems with the monitoring system in winter (mainly from December to April) until 2012, because of difficulties with maintenance. Hence, we could not obtain precipitation data associated with seven debris flows and one slush avalanche.

The time period used to separate different rainfall events (inter event time definition, IETD) affects the magnitude and accuracy of the rainfall threshold triggering landslides and debris flows (Bezak et al., 2016; Hong et al., 2018). The IETD in previous studies generally ranged from 6 to 24 h; it is affected by catchment size and the available data sets (Guzzetti et al., 2007, 2008; Bezak et al., 2016; Imaizumi et al., 2017). In this study, we compared rainfall thresholds obtained by three different inter event times (6, 12, and 24 h) to obtain the appropriate IETD in the Osawa failure.

コメントの追加 [A22]: Rev.1-16

書式変更: フォント: (日) MS 明朝, (言語 1) 日本語

Rainfall thresholds appropriately separating rainfall events with and without debris flows were determined by calculating the critical successful index (CSI, threat score) using the following equation (e.g., Liao et al., 2011; Formetta et al., 2016; Staley et al., 2017):

$$CSI = \frac{tp}{tp + fp + fn} \quad (3)$$

210 where tp is the number of true positive events, fp is the number of false positive events, and fn is the number of false negative events. In this study, the ID threshold, a widely used concept (e.g., Caine 1980; Guzzetti et al., 2008; Hürlimann et al., 2019), was employed to determine the importance of rainfall intensity and duration on the debris flow occurrence. The ID threshold is expressed by the following equations:

$$I = \alpha D^\beta \quad (4)$$

215 where I is rainfall intensity, D is rainfall duration, and α and β are the constant and slope of the power law curve, respectively. In addition to the ID threshold producing the highest CSI value, the ID threshold for the lower boundary of debris flow occurrence was obtained by quantile regressions in the 2nd percentile (Guzzetti, 2007; 2008; Saito et al., 2010). The rainfall threshold was also obtained by comparing the maximum hourly intensity and total rainfall to further improve our understanding of the debris flow initiation mechanism. Appropriate rainfall thresholds were determined under various conditions in the debris

220 flow initiation zone, such as the presence or absence of a frozen layer and large or small volumes of sediment storage. The uncertainty of the rainfall thresholds was accessed by 1,000 bootstrap resamplings. Debris flows and slush avalanches were likely to occur in the period with and without a snowpack in their initiation zones, respectively (Decaulne, 2007; Eckerstorfer and Christiansen, 2012). We did not distinguish between debris flows and slush avalanches to obtain rainfall thresholds, because the presence or absence of a snowpack during rainfall events was unclear, especially during events without significant sediment transfer. Rainfall events with a total rainfall <10 mm were not analyzed in this study, because they appeared to be unrelated to the occurrence of debris flows and slush avalanches. The time period used to separate different rainfall events was set to 24 h, which was longer than in many other studies (e.g., Bezak et al., 2016; Imaizumi et al., 2017). This was because the time required for flood concentration in the Osawa failure is likely longer than for many other debris flow torrents due to the large catchment area. Antecedent rainfall expressed by the following equation was also calculated for each rainfall events:

$$230 \quad AR = \sum_{i=1}^n K^i R_i \quad (5)$$

where AR is antecedent rainfall, K is the decay coefficient, R_i is daily precipitation in the i -th day preceding to the debris flow event, and n is the number of days used for calculation of the antecedent rainfall. The decay coefficient K and number of days n ranged from 0.7 to 1 and from 2 days to 4 weeks, respectively, in previous studies (e.g., Chen et al., 2005; Jakob et al., 2012; Guo et al., 2013; Peng et al., 2015; Bel et al., 2017). In this study, the K and n were set to 0.8 and 7 days, respectively, which were commonly used values in previous studies.

3.4 Changes in the volume of channel deposits

The MLIT measured the topography of the entire Osawa failure annually using aerial photogrammetry from 1970-1971 to 2006 and airborne LiDAR from 2007 to 2019 (Fuji Sabo Office, 2017, 2020; Table 1; Figure 3). Two measurements were

コメントの追加 [A23]: Rev.1-14

書式変更: フォント: 斜体

書式変更: フォント: 斜体

コメントの追加 [A24]: Rev.1-15

コメントの追加 [A25]: We noticed that events with total rainfall of 10 mm were also excluded from analyses in the previous version of the manuscript. Therefore, we have recomputed threshold values.

書式変更: フォント: 斜体

書式変更: フォント: 斜体, 下付き

書式変更: フォント: 斜体

書式変更: フォント: 斜体

書式変更: フォント: 斜体

書式変更: フォント: (日) MS 明朝, (言語 1) 日本語

コメントの追加 [A26]: Rev.1-21

240 conducted in each of 2000, 2004, and 2011, when large debris flows occurred. The measurements were mainly conducted from
 245 September to October, when weather conditions were stable and there was no snowpack. In the period prior to 1979, channel
 bed topography along cross-sectional lines with intervals of 50–to 100 m was assessed annually via photogrammetry using
 monochromatic photographs. In the period 1980–2006, DEMs with a grid size of 10×10 m or 5×10 m were constructed
 annually via photogrammetry using monochromatic and color aerial photographs, with scales of 1/3,000 to 1/10,000. The
 horizontal and vertical errors of the control points obtained at feature intersections in the period 1980–2006 were in the range
 of 0.15 to 0.30 m.

Since 2007, DEMs with a grid size of 0.5 to 1.0 m have been constructed from point clouds obtained via airborne LiDAR
 scanning, with a point density of 2.25 to 4.0 points m^{-2} . The accuracy of the measurements as determined by a comparison
 with ground control points on the alluvial fan was <0.1 m. The root mean square error of the elevation data between adjacent
 flight courses was also <0.1 m. Errors in the measurements were likely larger on steeper terrain, especially at the basalt outcrops
 in the Osawa failure.

250 Fiji Sabo Office (2020) estimated the volume of channel deposits inside the Osawa failure using a DEM in November 2000,
 when the channel bed elevation was lowest deposits were absent in a large parts of the channels, due to the evacuation of channel
 deposits by debris flows, thus determining the base topography of the channel bed (Fig. 3). In the 1974–1975, 1979–1980, and
 1998–1999 periods, when topography was measured using aerial photographs, the loss of a large volume of channel deposits
 (>100,000 m^3) was estimated despite there being no debris flow signals at the Otaki, Iwadoi, and Osawagawa Bridge
 observatories. Errors in the estimation of sediment volume as well as the occurrence of debris flows terminating in the section
 between the Osawa failure and the observatories possibly affected the evaluated loss of channel deposits. Hence, our analysis
 of the volume of channel deposits contains some inherent errors for the period in which data were obtained using aerial
 photography surveys.

260 In this study, we obtained rainfall thresholds for two volume classes of channel deposits: $\geq 350,000$ m^3 and $< 350,000$ m^3 .
 The boundary of the two classes was set to 350,000 m^3 to ensure that a statistically sufficient number of debris flow events
 could be allocated to both volume classes. Hence, this value itself does not have physical meaning relevant to the occurrence
 of debris flow. The inherent errors in the estimation of the volume of channel deposits may affect the misclassification of
 channel deposit class, especially in the period 1993–1997 when the volume of the channel deposits was close to the threshold
 of volume classes.

Table 1: General description of ~~Aerial~~ the aerial photogrammetry and airborne LiDAR measurements

Period	Number of measurements	Measurement method	Data type
December 1971 to October 1979	9	Aerial photogrammetry	Topography along cross-sectional lines with intervals of 50–to 100 m
October 1980 to November 2000	22	Aerial photogrammetry	DEM with a grid size of 10×10 m

コメントの追加 [A27]: Rev.1-21

コメントの追加 [A28]: Rev.1-17

コメントの追加 [A29]: Rev.1-18

コメントの追加 [A30]: Rev.1-20

コメントの追加 [A31]: Rev.1-19

書式変更: 左揃え

書式変更: 左揃え

September 2001 to October 2006	7	Aerial photogrammetry	DEM with a grid size of 10 × 5 m
October 2007 to October 2017	12	Airborne LiDAR	DEM with a grid size of 0.5 m
August 2018 to August 2019	2	Airborne LiDAR	DEM with a grid size of 1.0 m

書式変更: 左揃え

書式変更: 左揃え

書式変更: 左揃え

270




Figure 3: Temporal changes in the volume of channel deposits in the Osawa failure (original data from Fuji Sabo Office, 2020).

275 4 Results

4.1 Changes in ground temperature

In the shallower part of the ground (i.e., ≤ 0.25 m deep), diurnal freeze-thaw activity was frequently observed in autumn and spring, when the diurnal amplitude of the ground surface temperature was evident, whereas seasonally frozen ground was observed in winter (Figs. 4 and 5). At depths between 0.25 to 1.25 m, seasonally frozen ground occurred over long periods up to four months (Fig. 5). The minimum ground temperatures recorded each winter were lowest at the surface and increased toward the deeper part of the ground (Fig. 4), and the ground below 1.75 m was not frozen in both the 2017–2018 and 2018–2019 throughout the monitoring frozen periods (Fig. 5). Temporal patterns in the ground temperature were different between the 2017–2018 and 2018–2019 frozen periods (Fig. 4). In 2017–2018, ground temperature above a depth of 1.5 m rapidly increased during a heavy rainfall event on March 5 (total rainfall and maximum hourly rainfall of 89 mm and 18 mm h⁻¹, respectively), which triggered a slush avalanche. Although the seasonally frozen layer was retained for more than 1 month after the rainfall event, the ground temperature at a depth >0.5 m did not fall much below 0°C. In spring 2019, when heavy

280

285

コメントの追加 [A32]: Rev.2-11

rainfall events were not observed, the ground temperature below 0.5 m was lower than that in spring 2018. After ground warming on March 20–21 and April 20–24, 2019, ground cooling was observed, because the spring was colder than in the previous year. In both winters, seasonally frozen ground to a depth of 1.25 m persisted until mid-June in 2018 and early July in 2019 (Fig. 4). The clear diurnal changes in ground surface temperature in both the 2017–2018 and 2018–2019 frozen periods indicated that the snowpack was never sufficiently deep to prevent ground cooling in the observed winters.

290

コメントの追加 [A33]: Rev.1-2




Figure 4: Temporal changes in the ground temperature at 3,220 m a.s.l. from September 1, 2017 to September 1, 2019. Daily precipitation is also shown in the figure.




Figure 5: Comparison of observed and estimated periods of diurnal freeze-thaw activity and seasonal ground freezing at 3,221 a.s.l. The timing of thawing in 2019 could not be estimated using data from the summit station due to monitoring equipment failure. Periods of ground freezing were estimated using summit station data, whereas the end of the frozen period in 2019 was estimated using data from the Fuji City station.

The most appropriate coefficient c_f that could adequately replicate the gives the minimum standard error between the monitored and estimated downward progress of the freezing line timing of freezing in the depths between 0.25 to 1.25 m (= 3.6 days) was $5.5 \text{ m } ^\circ\text{C}^{-1/2} \text{ d}^{-1/2}$ (Fig. 5). Except in the near-surface ($\leq 0.25 \text{ m}$) and deep layers ($> 1.5 \text{ m}$), the positions of the freezing front were adequately estimated based on air temperatures measured at the summit and Fuji City stations (Fig. 5). The disagreement between the estimated freezing and observed non-freezing states at the depth of 1.75 m implied the presence of a thermal buffer, such as a layer with a high water content.

The rapid and simultaneous increase in ground temperature from negative values to the melting point (Fig. 4) indicated that thawing of the porous volcanic ground was controlled by not only heat conduction but also heat advection. In particular, infiltrated rainwater on March 5, 2018 passed through the seasonal frozen layer and released latent heat by partly freezing in the voids (Fig. 4). Ikeda et al. (2012) observed partial but rapid thawing of the frozen layer on the summit due to heavy rainfalls, mostly from tropical cyclones, in late summer, although thawing due to rains in early summer was unclear at TM in 2018 and 2019 (Fig. 4). Taking this into account, we calculated the coefficient c_m as an empirical value rather than a physical parameter. The empirical c_m was $6.8 \text{ m } ^\circ\text{C}^{-1/2} \text{ d}^{-1/2}$ when Eq. (2) best reproduced the time of thawing timing of the depths between 0.75 and at a depth of 1.25 m deep (Fig. 5).

The 1.5 m-deep point thawed much earlier than the 1.25 m-deep point (Fig. 5), which indicates that geothermal heat thawed ground upward around the deepest position of the seasonal frozen layer. However, most of the frozen layer thawed downward because the heat input from the surface was much larger than the geothermal heat flow. In this study, periods of ground freezing

コメントの追加 [A34]: Rev.1-23

コメントの追加 [A35]: Rev.1-22

コメントの追加 [A36]: Rev.2-12

コメントの追加 [A37]: Rev.1-23

コメントの追加 [A38]: Rev.1-22

コメントの追加 [A39]: Rev.2-11

could be roughly estimated from the time when the freezing front started to progress downward from the ground surface, and the time when the estimated thawing front reached ~~not a depth of~~ but a depth of 1.25 m (Fig. 5).

4.2 Changes in channel bed topography

Channel bed topography clearly changed with time and was affected by the sediment supply from hillslopes and the evacuation of channel deposits by debris flows (Fig. 6). From September 2013 to October 2014 (Fig. 6a), no debris flow was monitored at the observatories. During this period, aggradation of the channel bed of up to 5 m in association with sediment supply from the surrounding hillslopes was evident. From September to November of 2011, the channel deposit zone was significantly eroded by a debris flow that occurred on November 9, 2011 (Fig. 6b). During this period, debris flows occurred in some tributaries without any failure of the surrounding hillslopes (Fig. 6b). Sediment produced by the failure of basaltic lava outcrops was deposited just below the outcrops without evolving into a debris flow. Hence, past debris flows in the Osawa failure were mainly caused by the evacuation of channel deposits rather than the direct mobilization of landslide sediment.

Figure 6: Changes in channel bed topography. (a) From September 2013 to October 2014, no debris flow was observed. (b) From September to November of 2011, a debris flow occurred on November 9, 2011. The dashed line indicates the area with channel deposits.

コメントの追加 [A40]: Rev.1-24

4.3.2 Effects of ground freezing on the conditions required to initiate a debris flow Seasonal pattern of debris flow occurrence of debris flow

In the Osawa failure, the rainfall pattern is different between frozen and unfrozen periods (Fig. 6). In the frozen periods, hourly rainfall intensity exceeds 30 mm h^{-1} several times in each year, while the hourly rainfall intensity seldom exceeds 30 mm h^{-1} in frozen periods (Fig. 6). Antecedent rainfall in unfrozen periods is also higher than that in frozen periods. Antecedent rainfall sometimes exceeds 150 mm in unfrozen periods, while the antecedent rainfall is generally lower than 100 mm in frozen periods (Fig. 6).

The numbers of debris flows and slush avalanches were highest in November and March, respectively (Fig. 7). Although average monthly precipitation in August (278 mm) was higher than that in November (115 mm; Fig. 2), no debris flow was observed in August. Just one debris flow and one slush avalanche occurred in mid-winter (i.e., January and February), when precipitation was mostly snow. Debris flows occurred during rainfall events with high hourly rainfall intensity (Fig. 6). However, it is difficult to clearly separate debris flows and no flow events based on a single rainfall index. For example, on September 4, 2011, debris flow was not triggered by a rainfall event with high rainfall intensity (38 mm h^{-1}) and high total rainfall (595 mm) as well as 62 mm of antecedent rainfall prior to the event. In contrast, on November 19, 2011, a debris flow was triggered by low rainfall intensity (19 mm h^{-1}), moderate total rainfall (164 mm), and no antecedent rainfall.


コメントの追加 [A41]: Rev.2-3

書式変更: 上付き

コメントの追加 [A42]: Rev.1-1


コメントの追加 [A43]: Rev.2-4

書式変更: 上付き



350

Figure 6: Seasonal changes in the rainfall pattern in the period from 2011 to 2012, when debris flows occurred successively. The timing of the debris flow events is also illustrated in the figure. Frozen periods were estimated by the degree-day method.



355

Figure 7: Monthly frequency of debris flows and slush avalanches.

4.3 Effects of ground freezing on the conditions required to initiate a debris flow

Figures 7a-8a to 78c shows the influence of rainfall intensity and duration on sediment transfer (i.e., debris flows and slush avalanches), which tended to occur during the conditions represented in the upper-right quadrant of all plots regardless of the inter event time IETD. This indicated that the magnitude of rainfall events that triggered sediment transfer was larger than that of an ordinary rainfall event. However, the distributions of rainfall events that did or did not trigger sediment transfer could not be completely separated. The length of the IETD affected the rainfall duration rather than the rainfall intensity, with rainfall duration becoming longer with an increasing IETD. The CSI for the ID threshold was significantly low ($0.21 < 0.25$). The slope of the power law curve was gentler with increasing in IETD. Among the three IETDs, 24 h resulted in the highest

360

365 CSI value. ~~Many~~ Much of the sediment transfers occurred during the event with low antecedent rainfall (i.e., <50 mm). On the other hand, sediment transfer did not occur during some rainfall events with high antecedent rainfall (i.e., >100 mm).

コメントの追加 [A44]: Rev.1-16

コメントの追加 [A45]: Rev.1-2

Rainfall thresholds for the occurrence of sediment transfer were also obtained by comparing total rainfall and maximum hourly rainfall intensity (Figs. 78d to 78f). Regardless of the IETD, the total rainfall and maximum rainfall intensity that produced the highest CSI value were 4206 mm and 42 mm h⁻¹, respectively. However, the distributions of rainfall events that did or did not trigger sediment transfer could not be completely separated well (CSI = < 0.157). The distribution of no flow events with high antecedent rainfall (i.e., >100 mm) overlapped with that of sediment transfer events with low antecedent rainfall (i.e., <50 mm). Although the value amount of antecedent rainfall is varies affected by the setting of K and n in Eq. (4), Fig. 8 implies that the antecedent rainfall was not a critical factor affecting the occurrence of sediment transfer. The CSI did not differ among the IETDs (Figs. 8d-8f). Hereafter, the IETD was set to 24 h, which resulted in the highest CSI in the intensity-duration analysis.

コメントの追加 [A46]: Rev.2-13

コメントの追加 [A47]: Rev.1-2

コメントの追加 [A48]: Rev.1-16




Figure 8: Rainfall thresholds for triggering debris flows and slush avalanches. Comparison of the duration and intensity of rainfall events with or without debris flows and slush avalanches that were calculated with an inter event time definition (IETD) of (a) 6 h, (b) 12 h, and (c) 24 h. Dashed lines indicate the rainfall thresholds with the highest critical successful index (CSI). Comparison of the total rainfall and maximum hourly rainfall intensity of rainfall events with or without debris flows and slush avalanches that were calculated with an IETD of (d) 6 h, (e) 12 h, and (f) 24 h.

When separate ID diagrams were plotted for the frozen and unfrozen periods, respectively separately, the distribution of rainfall events that triggered sediment transfer during frozen periods was positioned further to the lower-left side than in the corresponding diagram for unfrozen periods (Figs. 7b9a-7e). Thus, the ID threshold in frozen periods was lower than that in unfrozen periods. The CSI values of the ID thresholds in the frozen and unfrozen periods were 0.42 and 0.1516, respectively. Therefore, the CSI in frozen periods was substantially improved compared to the CSI for all periods (0.25, Fig. 8c). The

frequency distribution of constant α in Eq. 4 was obtained by 1,000 bootstrap resamplings and clearly differed between the frozen and unfrozen periods, while slope β widely overlapped between the two periods (Figs. 9b, 9c). The lower limit of rainfall triggering sediment transfer, which was obtained by the 2nd percentile regression lines, in the frozen periods was lower than that in the unfrozen periods over a wide range of rainfall durations. This also indicated that the rainfall threshold for sediment transfer in the frozen periods was lower than that in the unfrozen periods. Some no flow events were distributed in the upper-right side of the sediment transfer plots in the ID diagram, especially in the unfrozen periods. Therefore, the upper limit of rainfall events did not produce sediment transfer could not be determined in the Osawa failure. Slush avalanches were observed only during frozen periods. Although the distribution of slush avalanches, which were observed only during frozen periods, overlapped with that of debris flows, all slush avalanche events were represented on the lower left side of the threshold line.

書式変更: フォント: 10 pt, 太字 (なし), 斜体 (なし)

書式変更: フォント: 10 pt

書式変更: フォント: 10 pt

書式変更: フォント: 10 pt

コメントの追加 [A49]: Rev.1-3

書式変更: 上付き

コメントの追加 [A50]: Rev.2-1




Figure 9: Intensity-duration thresholds for triggering debris flows and slush avalanches. (a) Comparison of the duration and intensity of rainfall events with or without debris flows and slush avalanches for rainfall events in frozen and unfrozen periods. Solid lines in the plots indicate the rainfall threshold, which optimally separated rainfall events that induced debris flows and slush avalanches from those that did not. Dashed lines in the plots indicate 2nd percentile regression lines that represent the lower limit condition for the occurrence of sediment transfer. Frequencies of (b) constant α and (c) the slope of the power law curve β , which produced the highest critical successful index (CSI), obtained by 1,000 bootstrap resamplings are also shown in the figure. Arrows indicate values that produced the highest CSI in the analysis using the original data set (solid line in (a)).


書式変更: 上付き

Rainfall thresholds for the occurrence of sediment transfer were also obtained by comparing total rainfall and maximum hourly rainfall intensity (Fig. 7d). The total rainfall and maximum rainfall intensity that produced the highest CSI value were 314 mm and 42 mm h⁻¹, respectively. However, the distributions of rainfall events that did or did not trigger sediment transfer could not be completely separated (CSI = 0.15). As with the ID diagrams, the maximum hourly intensity and total rainfall of rainfall events in the frozen and unfrozen periods were plotted separately (Figs. 7e, 7f10a). In the frozen periods, the threshold values of total rainfall and maximum hourly rainfall intensity were 420-99 mm and 22-19 mm h⁻¹, respectively. The slush avalanches were distributed in the lower-left side of the diagram compared to the debris flows. In the unfrozen periods, the

書式変更: インデント: 最初の行: 0 mm

threshold values of total rainfall and maximum hourly rainfall were higher than those in the frozen periods (314 mm and 42 64 mm h⁻¹, respectively). Both of the debris flows triggered by a maximum hourly rainfall <30 mm in unfrozen periods 415 occurred within the 10-day period prior to the estimated subsequent frozen period. Thus, it is possible that the ground was frozen when these debris flows occurred. The CSI values of the rainfall thresholds in the frozen and unfrozen periods (0.38 and 0.2736, respectively) were better than those obtained using all rainfall events (0.4516; Fig. 8f). The frequency 420 distributions of maximum hourly intensity and total rainfall obtained by bootstrapping did not overlap between the frozen and unfrozen periods (Figs. 910b, 910c), indicating that the difference in threshold was not affected by the presence or absence of specific rainfall events.

コメントの追加 [A51]: Rev.1-3



425 **Figure 10: Thresholds of total rainfall and maximum hourly rainfall intensity triggering debris flows and slush avalanches. (a) Comparison of the total rainfall and the maximum hourly rainfall intensity of rainfall events with or without debris flows and slush avalanches for rainfall events in the frozen and unfrozen periods. The solid lines in the plots indicate the rainfall thresholds that optimally separated rainfall events that induced debris flows and slush avalanches and those that did not. Frequencies of (b) maximum rainfall intensity and (c) total rainfall that produced the highest critical successful index (CSI), obtained by 1,000 bootstrap resamplings. Arrows indicate the values that produced the highest CSI in the analysis using the original data set (same as values in (a)).**

4.4 Effects of the volume of channel deposits on the conditions required to initiate a debris flow

During frozen periods, the ID threshold in the periods with large volume of channel deposits was not very different between 435 periods with large and higher than in periods with small volumes of channel deposits (Fig. 811a). However, as a result of bootstrapping, the frequency distributions of both α and β overlapped between periods with large and small volumes of channel deposits (Figs. 11b, 11c). Hence, a high uncertainty exists in the effect of the volume of channel deposits on the ID threshold. The CSI values of the thresholds for small and large volumes of channel deposits were 0.4542 and 0.3846, respectively. Therefore, the CSI was not improved from when the ID threshold was determined without considering the volume of channel

440 deposits (0.42, Fig. 7b9a). During unfrozen periods, the ID threshold in periods with a large volume of channel deposits was clearly slightly higher than that in periods with a small volume of channel deposits (Fig. 8b11d). As with the situation during frozen periods, the frequency distributions of both α and β were overlapped between periods with large and small volumes of channel deposits (Figs. 11e, 11f). The CSI values for both volume groups (0.20 and 0.14 for small and large volumes of channel deposits, respectively) were not significantly higher than the values obtained without considering the volume of channel deposits (0.165, Fig. 7e9a).

445 As with the ID threshold, the maximum hourly intensity and total rainfall that triggered debris flows were not clearly affected by the volume of channel deposits during the frozen periods (Figs. 9a12a, 9b, Table 2). Frequency distributions of maximum hourly rainfall and total rainfall in bootstrapping had a clearer peak than those of α and β of the ID threshold (Figs. 12b, 12c). By contrast, the thresholds of maximum hourly rainfall intensity in the unfrozen periods were very different and were dependent on the volume of channel deposits (Figs. 9e12d, 9d12e, Table 2). A larger hourly rainfall intensity and total rainfall were needed for the debris-flow occurrence of debris flows in periods with a large volume of channel deposits. Many rainfall events that exceeded the rainfall-hourly intensity threshold in periods with a small volume of deposits did not trigger debris flows in periods with a large volume of deposits. The threshold of total rainfall during unfrozen periods was less clear than that of hourly rainfall intensity (Figs. 12d, 12f). The CSI of the rainfall thresholds obtained separately for each sediment volume class (0.30 and 0.60 for large and small volumes of channel deposits, respectively) was improved from that calculated for all rainfall levels, regardless of the sediment volume (0.27, Fig. 7f).

コメントの追加 [A52]: Rev.1-3

コメントの追加 [A53]: Rev.1-3




Figure 8101: Comparison of the duration and intensity of rainfall events with or without debris flows and slush avalanches, analyzed according to the volume of channel deposits. The boundary between the two volume classes was set to 350,000 m³. (a) Comparison during Frozen periods. The frequencies of (b) constant α and (c) slope of the power law curve β , which produced the highest critical successful index (CSI), obtained by 1,000 bootstrap resamplings during frozen periods. (d) Unfrozen-Comparison during unfrozen periods. (a) Comparison during frozen periods. Frequencies of the (e) constant α and (f) slope of the power law curve β obtained by bootstrap resampling during unfrozen periods.

460




Figure 9112: Comparison of the total rainfall and maximum hourly rainfall in rainfall events with or without debris flows and slush avalanches, plotted according to the volume class of channel deposits. The boundary between the two volume classes was set to 350,000 m³. (a) Comparison during Frozen periods with a small volume of deposits (<350,000 m³). (b) Frozen periods with large volume of deposits (≥350,000 m³). (c) Unfrozen periods with a small volume of deposits. (d) Unfrozen periods with a large volume of deposits. Dashed lines in the plots indicate the rainfall thresholds, which that optimally separated rainfall events that induced debris flows and slush avalanches and those that did not. Frequencies of (b) maximum hourly rainfall intensity and (c) total rainfall, which produced the highest critical successful index (CSI), obtained by 1,000 bootstrap resamplings during frozen periods. (e) Comparison during unfrozen periods. The values of the rainfall thresholds are listed in Table 1. Frequencies of (be) maximum rainfall intensity and (ef) total rainfall obtained by bootstrap resampling.

Table 2: Rainfall thresholds for triggering debris flows and slush avalanches, obtained via comparisons of maximum hourly rainfall intensity and total rainfall.

Conditions	Number of rainfall events	Number of debris flows and slush avalanches	Rainfall threshold		CSI
			Maximum hourly rainfall (mm h ⁻¹)	Total rainfall (mm)	
Frozen / small volume of deposits	342	16	17	99	0.43
Frozen / large volume of deposits	460	10	23	121	0.37
Unfrozen / small volume of deposits	377	4	38	248	0.30

コメントの追加 [A54]: We have removed this table because we described rainfall threshold and CSI in Figures.

Unfrozen / large volume of deposits	457	4	67	314	0.60
-------------------------------------	-----	---	----	-----	------

5. Discussion

Our simple model using ground surface temperature could roughly reproduce the period of ground freezing at site TM (Fig. 5), although subsurface heterogeneity, infiltration of rainfall, and upward movement of sensible heat were not considered in the model. The snowpack, which affects the period of ground freezing in many cold regions (e.g., Zhang, 2005; Luo, 2018), was not considered important to the formation of frozen ground at the monitoring point, because diurnal changes in the ground surface temperature (Fig. 4) indicated that the snowpack was never sufficiently deep to prevent ground cooling in the observed were clear even in mid-winters (Fig. 4). Similarly, the shallow snow depth in winter due to the strong winds and steep topography under the dry climate in winter probably prevented snow accumulation on around the windward steep slopes such as the whole Osawa failure (Ikeda et al., 2012), which likely resulted in the limited influence of the snowpack on the period of ground freezing observed in this study. However, snowfalls from March to April may affect the thermal situation differently among topographic conditions. Air temperatures from March to April fluctuated around 0 °C, which had a direct impact on the diurnal freeze-thaw of the near-surface at site TM (Fig. 4). This situation hindered further downward freezing at the bottom of the frozen layer and thawing below the diurnal freeze-thaw layer (Fig. 5). In the same period, where there was thick snow insulating the ground from the atmosphere, the snow must have prevented both further cooling and thawing of the existing frozen ground. Thus, the thickness of the frozen layer formed in winter might be independent of the snow thickness in spring. In addition, the ground at TM thawed after snow disappearance in the debris flow initiation zone (mid-April, 2018 and late April, 2019). These conditions suited our expansion of the simplistic thermal model to the distinction between frozen and unfrozen periods over in the failure area the initiation, although further studies on various spatial and temporal heterogeneities will be required zone. Although the ground freezing and snow conditions of channel deposits at very bottom of the valley may differ from those at TM, the effect of ground freezing could be demonstrated by the estimation of freezing periods at TM (Figs. 9, 10). This implies that the water supply from hillslopes, which have a greater area than the valley bottom, is important to the occurrence of sediment transfer.

Because hourly rainfall intensity and total rainfall in the unfrozen period was higher than in frozen periods (Figs. 6, 9), sediment transfers can occur more easily in the frozen periods in terms of the rainfall characteristics. However, the amount of sediment transfer in frozen periods was larger than that in unfrozen periods (33 and 9, respectively) because of the lower rainfall thresholds was significantly different between the frozen and unfrozen periods in the frozen periods (Figs. 8, 7, 9, 10). Therefore, the formation of frozen ground is a critical factor affecting the initiation seasonal changes of in the susceptibility a to debris flow sediment transfer in the Osawa failure. In unfrozen periods, many rainfall events exceeding rainfall thresholds in frozen periods did not trigger sediment transfer, also implying the importance of frozen ground freezing in the occurrence of sediment transfer (Figs. 8, 9, 10). Bovis and Dagg (1988) implied that the hydraulic conductivity of channel deposits affects

書式変更: インデント: 最初の行: 0 字

コメントの追加 [A55]: Rev.1-25

コメントの追加 [A56]: Rev.1-2, Rev.1-26

コメントの追加 [A57]: Rev.1-26

コメントの追加 [A58]: Rev.1-12

コメントの追加 [A59]: Rev.1-27

コメントの追加 [A60]: Rev.1-1

the temporal changes in debris flow activity. Based on constant head permeability tests in the laboratory, Nanzaka and Iwata (1989) reported that the hydraulic conductivity of frozen deposits in the Osawa failure (3 to 22 mm h⁻¹) was lower than that of unfrozen deposits (720 mm h⁻¹). Decreases in the infiltration rate due to ground freezing, which results in the occurrence of overland flow, have been reported in other regions (Blackburn et al., 1990; Iwata et al., 2010; Coles and McDonell, 2018).

Consequently, the generation of overland flow that was affected by the low infiltration rate of deposits in frozen periods likely resulted in the debris flow occurrence during smaller rainfall events (Fig. 710). Ice melt, which increases the water supply to debris flow material, caused by rainfall is another potential factor lowering the rainfall threshold. The effect of the volume of channel deposits on the rainfall threshold was unclear in the frozen periods (Figs. 811a, 912a), suggesting that debris flows in these periods were triggered by hydrological processes near the ground surface rather than the saturation of entire channel deposits (Fig. 10a13a). Increasing in the debris flow frequency due to decreasing decreases in the infiltration rate has also been reported in the regions affected by wild fire (Staley et al., 2017; McGuire et al., 2018; Rengers et al., 2020b). Because a rainfall index representing the total water supply to the basin (i.e., total rainfall) was also an important factor in the triggering of debris flows (Fig. 7e120a), a sufficient amount of stream discharge may also be needed for the occurrence of debris flow.

As with the Osawa failure, slush avalanches and debris flows have been recorded in other cold regions other than the Osawa failure (Decaulne and Saemundsson, 2006; Decaulne, 2007). In contrast to a debris flow, the presence of a snowpack is necessary for the occurrence of a slush avalanche (Decaulne, 2007; Eckerstorfer and Christiansen, 2012). The magnitude of rainfall events triggering slush avalanches were smaller than those that triggered debris flows (Fig. 710a), because snowmelt during rainfall events increases the water supply to debris flow material (Decaulne and Saemundsson, 2006). Because snow in a slush avalanche sometimes melts as it moves to lower elevations due to the increasing temperature (Anma, 2007), some sediment transfer events classified as debris flows would originally be have been slush avalanches in the initiation zone. That could be one of the reasons why the distributions of slush avalanches and debris flows according to rainfall factors overlapped in Fig. 79a.

コメントの追加 [A61]: Rev.2-3

コメントの追加 [A62]: Rev.2-3




Figure 4013: Schematic diagram of the hydrological characteristics of the Osawa failure. (a) Frozen periods. The layer with the ice in soil matrix is colored by light blue. (b) Unfrozen periods.

コメントの追加 [A63]: Rev.1-28

In contrast to the frozen periods, overland flow rarely occurred in the unfrozen periods because of the high infiltration rate of the deposits. Therefore, an increase in the groundwater level in channel deposits would likely influence the occurrence of debris flows (Fig. 10b13b). The slope of the ID threshold in environments in which the rainfall duration was important was steeper than in environments in which rainfall intensity was important (Guzzetti et al., 2008). A-The steeper threshold line in the unfrozen periods relative to the frozen periods (Fig. 811) indicated that rainfall duration was the important factor in the unfrozen periods. A-large rainfall magnitudeA high hourly rainfall intensity was needed for the occurrence of debris flows when a large volume of sediment was stored in the initiation zone (Figs. 8, 912b). The relationship between the rainfall threshold and the volume of sediment stored was different from that reported in weathering-limited debris flow basins (Bovis and Jakob, 1999; Larsen et al., 2006). Because debris flows occurred even in periods with a small volume of channel deposits stored in the channel (Fig. 3), the threshold of sediment volume needed for the occurrence of a debris flow, which has been reported in weathering-limited basins (Bovis and Jakob, 1999), was not clear in the Osawa failure. The relationship between the rainfall threshold and the storage volume in the Osawa failure was also different from that in transport-limited basins, in which the rainfall threshold can be given as a constant value regardless of the storage volume (Bovis and Jakob, 1999; Schlunegger et al., 2009). Channel deposits in the Osawa failure had a thickness of up to 10 m in some periods and were also characterized by a high infiltration rate. The ratio of the thickness of the saturated layer to the entire thickness of the deposits is a factor controlling the movement of cohesionless deposits in steep channels, because the ratio determines the balance between shear stress and shear strength (Imaizumi et al., 2016, 2017). A large water supply is needed for the movement of thick deposits. Hence, the volume of channel deposits likely has a negative effect on the debris flow occurrence in unfrozen periods.

Plots of rainfall events with and without debris flow events overlapped to a certain extent, even after consideration of the frozen periods and volume of channel deposits (Figs. 811, 912). The magnitude of some rainfall events that triggered debris flows in unfrozen periods was as low as those in the frozen periods (Fig. 710), indicating that it is was difficult to accurately estimate when frozen periods occur. Because periods of ground freezing differ with slope aspect and elevation, it is difficult to determine if the ground was frozen at the initiation point for all events. Another potential factor reducing the CSI was the spatial variability of precipitation in the Osawa failure, which has a difference of elevation of up to 1,500 m. Although most of the channel deposits in the Osawa failure consistently face west (Fig. 1d), slope aspect controls debris flow activity in other regions (Tillery and Rengers, 2020; Ding et al., 2020) due to its impact on the duration of seasonal ground freezing (Dashtseren et al., 2014), weathering rate of the rock wall (Rengers et al., 2020a), distribution of water (Xu et al., 2013), and geometric characteristics (Ding et al., 2020). The ID threshold for the Osawa failure was higher than those measured for many other torrents globally (e.g., Guzzetti et al., 2008; Bezak et al., 2016; Hürlimann et al., 2019). Although the lower limit for debris flow in the Osawa failure was not far from the ID thresholds for other regions, rainfall levels above the lower limit without the debris flow occurrence pushed the threshold upward. In general, the lower limit lines for debris flow are lower than the lines separating rainfall events with and without debris flows (Cannon et al., 2008; Bezak et al., 2016). Because the debris flow

コメントの追加 [A64]: Rev.2-14

コメントの追加 [A65]: Rev.1-29

コメントの追加 [A66]: Rev.2-2

~~threshold is often the lower limit line (Cannon et al., 2008; Guzzetti et al., 2008), the threshold in the Osawa failure gully is higher.~~ The observatories are located far from the debris flow initiation zone, resulting in debris flows just a short distance away being missed. This might have also affected the higher threshold line.

~~Under climate change, increases in the amount of sediment delivered to channels by the melting of permafrost and increases in extreme rainfall events are considered to be potential factors increasing the threat of debris flows on cold mountains (Staffler et al., 2008; Stoffel et al., 2014a, 2014b). The results of our study imply that climate change also impacts seasonal debris flow activity via shifts in the periods of seasonal ground freezing.~~

コメントの追加 [A67]: Rev.2-5

6. Summary and Conclusion

To determine the effects of frozen ground on temporal changes in the rainfall threshold for the occurrence of debris flows and slush avalanches, we estimated periods of seasonal ground freezing based on field monitoring conducted close to the initiation zone in the Osawa failure on Mt. Fuji. We also evaluated the influence of sediment storage volume in the debris flow initiation zone on the rainfall threshold. The ID threshold in periods of ground freezing was clearly lower than in periods without ground freezing. A comparison of the maximum hourly rainfall intensity and total rainfall during rainfall events also showed that debris flows occurred during periods of seasonal ground freezing with rainfall events of a smaller magnitude. Decreases in the infiltration rate due to the formation of frozen ground facilitated the generation of overland flow, which triggered debris flows.

The volume of channel deposits also affected the rainfall threshold, especially during unfrozen periods. The rainfall threshold of hourly rainfall intensity in periods with large volumes of channel deposits was higher than that in periods with small volumes of channel deposits, indicating that a large water supply is needed for the debris flow occurrence when channel deposits are thick. Increases in the water content in channel deposits caused by the infiltration of rainfall are important for the debris flow occurrence during unfrozen periods, whereas hydrological processes near the ground surface are more important in frozen periods.

Our study determined that the rainfall threshold is variable even in the same torrent due to the influence of the formation of frozen ground and changes in sediment storage volume. Therefore, the occurrence of frozen ground and the sediment storage volume need to be monitored and estimated to improve debris flow disaster mitigation in cold regions. Because periods of ground freezing and the sediment supply ratio are strongly affected by climatic conditions, future changes in the debris flow threshold should be carefully monitored for continuous sediment flow-associated disaster mitigation under climate change.

Data availability

~~Because this study was commissioned by Ministry of Land, Infrastructure, Transport and Tourism (MLIT), Japan, The the ground temperature data that support the findings of this study are available from the corresponding author, Fumitoshi~~

Imaizumi, upon reasonable request and agreement by [Ministry of Land, Infrastructure, Transport and Tourism, Japan](#) MLIT. Rainfall and topographic data are available upon agreement by MLIT, which holds the copyright of the data.

コメントの追加 **[A68]**: Rev.1-30

Author contributions

600 FI and OO conceived and designed the study. AI designed and conducted field monitoring of ground freezing. KY contributed analyses of historical rainfall data. All of the authors conducted field motoring. FI contributed to the writing of the paper.

Competing interests

The authors declare that they have no conflict of interest

Acknowledgements

605 This study was funded by Ministry of Land, Infrastructure, Transport and Tourism, Japan. This study was also supported by JSPS Grant Numbers 18H02235 and 19K01156. Airborne DEMs and rainfall data was provided by Fuji Sabo Office, Chubu Regional Bureau, Ministry of Land, Infrastructure, Transport and Tourism, Japan. [We are grateful to Tomohiro Egusa for his advice on statistical analysis. The author acknowledges Francis Rengers and one anonymous reviewer for their constructive reviewers that improved this paper.](#)

References

- 610 Anma, S.: Lahars and Slush Lahars on the Slopes of Fuji Volcano. Aramaki et al. Ed., Fuji Volcano, 285–301 (in Japanese with English abstract), 2007.
- Bayard, D., Stähli, M., Parriaux, A., and Flühler, H: The influence of seasonally frozen soil on the snowmelt runoff at two Alpine sites in southern Switzerland, *J. Hydrol.*, 309, 66–84. <https://doi.org/10.1016/j.jhydrol.2004.11.012>, 2005
- 615 [Bel, C., Liébault, F., Navratil, O., Eckert, N., Bellot, H., Fontaine, F., and Laigle, D.: Rainfall control of debris-flow triggering in the Réal Torrent, Southern French Prealps, *Geomorphology*, 291, 17-32, <https://doi.org/10.1016/j.geomorph.2016.04.004>, 2017.](#)
- Berger, C., McArdell, B. W., and Schlunegger, F.: Sediment transfer patterns at the Illgraben catchment, Switzerland: Implications for the time scales of debris flow activities, *Geomorphology*, 125, 421–432. <https://doi.org/10.1016/j.geomorph.2010.10.019>, 2011
- 620 [Bezák, N., Šraj, M., and Mikoš, M.: Copula-based IDF curves and empirical rainfall thresholds for flash floods and rainfall-induced landslides. *Journal of Hydrology*, 541, 272-284 doi:10.1016/j.jhydrol.2016.02.058, 2016.](#)

- Blackburn, W. H., Pierson, F. B., and Seyfried, M. S.: Spatial and temporal influence of soil frost on infiltration and erosion of sagebrush rangeland, *J. Am. Water Res. As.*, 26(6), 991–997. <https://doi.org/10.1111/j.1752-1688.1990.tb01434.x>, 1990.
- 625 [Bovis, M.J. and Dagg, B. R.: A model for debris accumulation and mobilization in steep mountain streams, *Hydrol. Sci. J.*, 33, 589–604, 1988.](https://doi.org/10.1002/(SICI)1096-9837(199910)24:11<1039::AID-ESP29>3.0.CO;2-U)
- Bovis, M. J., and Jakob, M.: The roll of debris supply conditions in predicting debris flow activity, *Earth Surf. Process. Landf.*, 24, 1039–1054. [https://doi.org/10.1002/\(SICI\)1096-9837\(199910\)24:11<1039::AID-ESP29>3.0.CO;2-U](https://doi.org/10.1002/(SICI)1096-9837(199910)24:11<1039::AID-ESP29>3.0.CO;2-U), 1999.
- ~~Bovis, M.J. and Dagg, B. R.: A model for debris accumulation and mobilization in steep mountain streams, *Hydrol. Sci. J.*, 33, 589–604, 1988.~~
- 630 Caine, N.: The rainfall intensity–duration control of shallow landslides and debris flows, *Geogr. Ann. A* 62, 23–27, 1980.
- Cannon, S. H., Kirkham, R. K., and Parise, M.: Wildfire-related debris-flow initiation processes, Storm King Mountain, Colorado, *Geomorphology*, 39, 171–188, [https://doi.org/10.1016/S0169-555X\(00\)00108-2](https://doi.org/10.1016/S0169-555X(00)00108-2), 2001.
- Cannon S. H., Boldt, E. M., Laber, J.L., Kean, J.W., and Staley, D. M.: Rainfall intensity-duration thresholds for postfire debris-flow emergency-response planning, *Nat. Hazards*, 59, 209–236, <https://doi.org/10.1007/s11069-011-9747-2>, 2011.
- 635 Chen, H. X., Zhang, L. M., Chang, D. S., and Zhang, S.: Mechanisms and runout characteristics of the rainfall triggered debris flow in Xiaojiaogou in Sichuan Province, China, *Nat. Hazards*, 62, 1037–1057. <https://doi.org/10.1007/s11069-012-0133-5>, 2012.
- [Cheng, C. Y., Chen, T. C., Yu, F. C., Yu, W. H., and Tseng, C. C.: Rainfall duration and debris-flow initiated studies for real-time monitoring, *Environ. Geol.* 47, 715–724, <https://doi.org/10.1007/s00254-004-1203-0>, 2005.](https://doi.org/10.1007/s00254-004-1203-0)
- 640 Coles, A. E., and McDonnell, J. J.: Fill and spill drives runoff connectivity over frozen ground, *J. Hydrol.*, 558, 115–128. <https://doi.org/10.1016/j.jhydrol.2018.01.016>, 2018.
- [Dashtseren, A., Ishikawa, M., Iijima, Y., and Jambaljav, Y.: Temperature Regimes of the Active Layer and Seasonally Frozen Ground under a Forest-Steppe Mosaic, Mongolia, *Permafrost. Periglac. Process.*, 25, 295–306, <https://doi.org/10.1002/ppp.1824>, 2014.](https://doi.org/10.1002/ppp.1824)
- 645 Decaulne, A.: Snow-avalanche and debris-flow hazards in the fjords of north-western Iceland, mitigation and prevention, *Nat. Hazards*, 41, 81–98. <https://doi.org/10.1007/s11069-006-9025-x>, 2007.
- ~~Ebel, B. A., Moody, J. A., and Martin, D. M.: Hydrologic conditions controlling runoff generation immediately after wildfire, *Water Resour. Res.*, 48, W03529, <https://doi.org/10.1029/2011WR011470>, 2012.~~
- Decaulne, A., and Saemundsson, T.: Geomorphic evidence for present-day snow-avalanche and debris-flow impact in the Icelandic Westfjords. *Geomorphology*, 80, 80–93, <https://doi.org/10.1016/j.geomorph.2005.09.007>, 2006.
- 650 [Ding, M., Tang, C., and Miao, C.: Response analysis of valley settlements to the evolution of debris flow fans under different topographic conditions: a case study of the upper reaches of Min River, China, *Bull. Eng. Geol. Environ.*, 79, 1639–1650, <https://doi.org/10.1007/s10064-019-01641-9>](https://doi.org/10.1007/s10064-019-01641-9)
- [Dowling, C. A., and Santi, P.: Debris flows and their toll on human life: a global analysis of debris-flow fatalities from 1950 to 2011, *Nat. Hazards*, 71, 203–227, <https://doi.org/10.1007/s11069-013-0907-4>, 2014.](https://doi.org/10.1007/s11069-013-0907-4)
- 655

Ebel, B. A., Moody, J. A., and Martin, D. M.: Hydrologic conditions controlling runoff generation immediately after wildfire, *Water Resour. Res.*, 48, W03529, <https://doi.org/10.1029/2011WR011470>, 2012

660 Eckerstorfer, M., and Christiansen, H. H.: Meteorology, topography and snowpack conditions causing two extreme mid-winter slush and wet slab avalanche periods in high Arctic maritime Svalbard. *Permafrost. Periglac. Process.*, 23, 15–25, <https://doi.org/10.1002/ppp.734>, 2012.

Fiollio, F., and Wilson, R. C.: Rainfall induced debris flows in pyroclastic deposits, Campania (southern Italy), *Eng. Geol.* 75, 263–289, <https://doi.org/10.1016/j.enggeo.2004.06.014>, 2004.

Fuji Sabo Office: Monitoring reports of sediment transport monitoring at Mt. Fuji in 2016 financial year, Fuji Sabo Office, Ministry of Land, Infrastructure, Transport and Tourism, Japan (in Japanese), 2017.

665 Fuji Sabo Office: Monitoring reports of sediment transport monitoring at Mt. Fuji in 2019 financial year, Fuji Sabo Office, Ministry of Land, Infrastructure, Transport and Tourism, Japan (in Japanese), 2020.

Formetta, G., Capparelli, G., and Versace, P.: Evaluating performance of simplified physically based models for shallow landslide susceptibility, *Hydrol. Earth Syst. Sci.*, 20, 4585–4603, <https://doi.org/10.5194/hess-20-4585-2016>, 2016.

670 Guo X. J., Cui, P., and Li, Y.: Debris flow warning threshold based on antecedent rainfall: a case study in Jiangjia Ravine, Yunnan, China, *J. Mount. Sci.*, 10, 305–314, <https://doi.org/10.1007/s11629-013-2521-z>, 2013.

Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: Rainfall thresholds for the initiation of landslides in central and southern Europe, *Meteorol. Atmos. Phys.*, 98, 239–267, <https://doi.org/10.1007/s00703-007-0262-7>, 2007.

Guzzetti, F., Peruccacci, S., Rossi, M., and Stark, C. P.: The rainfall intensity–duration control of shallow landslides and debris flows: an update, *Landslides*, 5, 3–17, <https://doi.org/10.1007/s10346-007-0112-1>, 2008.

675 Dowling, C. A., and Santi, P.: Debris flows and their toll on human life: a global analysis of debris flow fatalities from 1950 to 2011, *Nat. Hazards*, 71, 203–227, <https://doi.org/10.1007/s11069-013-0907-4>, 2014.

Hanaoka, M., Tomita, Y., and Ito, M.: Osawa Kuzure (failure) and debris flows in Mt. Fuji, Aramaki et al. Ed., *Fuji Volcano*, 407–425 (in Japanese with English abstract), 2007.

Hürlimann, M., Coviello, V., Bel, C., Guo, X., Berti, M., Graf, C., Hübl, J., Miyata S., Smith, J. B., and Yin, H. Y.: Debris-flow monitoring and warning: Review and examples, *Earth-Sci. Rev.*, 199, 102981, <https://doi.org/10.1016/j.earscirev.2019.102981>, 2019.

Hong, M., Kim, J. and Jeong, S.: Rainfall intensity-duration thresholds for landslide prediction in South Korea by considering the effects of antecedent rainfall, *Landslides* 15, 523–534, <https://doi.org/10.1007/s10346-017-0892-x>, 2018.

685 Ikeda, A., Iwahaba, T., and Sueyoshi, T.: Year-round monitoring of shallow ground temperatures at high altitudes of Mt. Fuji with a critical discussion on the popular belief of rapid permafrost degradation. *J. Geogr. (Chigaku Zasshi)*, 121, 306–331, <https://doi.org/10.5026/jgeography.121.306> (in Japanese with English abstract), 2012.

Imaizumi, F., Hayakawa, Y.S., Hotta, N., Tsunetaka, H., Ohsaka, O., and Tsuchiya, S.: Relationship between the accumulation of sediment storage and debris-flow characteristics in a debris-flow initiation zone, Ohya landslide body, Japan, *Nat. Hazards Earth Syst. Sci.*, 17, 1923–1938. <https://doi.org/10.5194/nhess-17-1923-2017>, 2017.

書式変更: 英語 (米国)

書式変更: 英語 (米国)

書式変更: 英語 (米国)

書式変更: 英語 (米国)

- 690 Imaizumi, F., Sidle, R. C., Tsuchiya, S., and Ohsaka, O.: Hydrogeomorphic processes in a steep debris flow initiation zone—, *Geophys. Res. Lett.*, 33, L10404, <https://doi.org/10.1029/2006GL026250>, 2006.
- Imaizumi F., Tsuchiya, S., and Ohsaka, O.: Field observations of debris-flow initiation processes on sediment deposits in a previous deep-seated landslide site, *J. Mount. Sci.*, 13, 213–222, <https://doi.org/10.1007/s11629-015-3345-9>, 2016.
- 695 Imaizumi, F., Masui, T., Yokota, Y., Tsunetaka, H., Hayakawa, Y.S., and Hotta, N.: Initiation and runout characteristics of debris flow surges in Ohya landslide scar, Japan, *Geomorphology*, 339, 58–69, <https://doi.org/10.1016/j.geomorph.2019.04.026>, 2019.
- Iwata, Y., Hayashi, M., Suzuki, S., Hirota, T., and Hasegawa, S.: Effects of snow cover on soil freezing, water movement, and snowmelt infiltration: A paired plot experiment, *Water Resour. Res.*, 46, W09504, <https://doi.org/10.1029/2009WR008070>, 2010.
- 700 Iwatsuka, S., and Machida, H.: The development of Osawa Valley, Mt. Fuji —The fundamental study on the Development of radial valleys on Volcano—, *J. Geogr. (Chigaku Zasshi)*, 729, 3–18 (in Japanese with English abstract). <https://doi.org/10.5026/jgeography.71.143>, 1962.
- Jakob, M., Bovis, M., and Oden, M.: The significance of channel recharge rates for estimating debris-flow magnitude and frequency—, *Earth Surf. Process. Landf.*, 30, 755–766, <https://doi.org/10.1002/esp.1188>, 2005.
- 705 [Jakob, M., Owen, T., Simpson, T.: A regional real-time debris-flow warning system for the District of North Vancouver, Canada, *Landslides*, 9, 165–178. <https://doi.org/10.1007/s10346-011-0282-8>, 2012.](https://doi.org/10.1007/s10346-011-0282-8)
- Kneisel, C., Emmert, A., Polich, P., Zollinger, B., and Egli M.: Soil geomorphology and frozen ground conditions at a subalpine talus slope having permafrost in the eastern Swiss Alps, *Catena*, 133, 107–118, <https://doi.org/10.1016/j.catena.2015.05.005>, 2015.
- 710 Larsen, I. J., Pederson, J. L., and Schmidt, J. C.: Geologic versus wildfire controls on hillslope processes and debris flow initiation in the Green River canyons of Dinosaur National Monument, *Geomorphology*, 81, 114–127, <https://doi.org/10.1016/j.geomorph.2006.04.002>, 2006.
- Liao, Z. H., Hong, Y., Kirschbaum, D., Adler, R. F., Gourley, J. J., and Wooten, R.: Evaluation of TRIGRS (transient rainfall infiltration and grid-based regional slope-stability analysis)'s predictive skill for hurricane-triggered landslides: a case study in Macon County, North Carolina—, *Nat. Hazards*, 58, 325–339. <https://doi.org/10.1007/s11069-010-9670-y>, 2011.
- 715 Loye, A., Jaboyedoff, M., Theule, J. I., and Liébault, F.: Headwater sediment dynamics in a debris flow catchment constrained by high-resolution topographic surveys, *Earth Surf. Dynam.*, 4, 489–513, <https://doi.org/10.5194/esurf-4-489-2016>, 2016.
- Luo, D., Jin, H., Marcheko, S. S., and Romanovsky, V E.: Difference between near-surface air, land surface and ground surface temperatures and their influences on the frozen ground on the Qinghai-Tibet Plateau—, *Geoderma*, 312, 74–85, <https://doi.org/10.1016/j.geoderma.2017.09.037>, 2018.
- 720 May, C. L.: Debris flows throughout different forest age classes in the central Oregon coast range, *J. A. Water. Resour. As.*, 38, 1097–1113. <https://doi.org/10.1111/j.1752-1688.2002.tb05549.x>, 2002.

- McGuire, L. A., Rengers, F. K., Kean, J. W., Staley, D. M., and Mirus, B. B.: Incorporating spatially heterogeneous infiltration capacity into hydrologic models with applications for simulating post-wildfire debris flow initiation, *Hydrol. Process.*, 32, 1173–1187, <https://doi.org/10.1002/hyp.11458>, 2018.
- Miao, Y., Shi, Y., Zhuang, H. Y., Wang, S. Y., and Liu, H. B.: Influence of Seasonal Frozen Soil on Near-Surface Shear Wave Velocity in Eastern Hokkaido, Japan, *Geophys. Res. Lett.*, 46, 9497–9508, <https://doi.org/10.1029/2019GL082282>, 2019.
- Miyaji, N.: Histories of younger Fuji volcano, *J. Geol. Soc. of Japan*, 94, 433–452 (in Japanese with English abstract), 1988.
- Nanzaka, J., and Iwata, Y.: Debris flow and frozen soil layer at the Osawa collapse in Mt. Fuji, *J. Japan Soc. Eros. Control Eng.*, 42(4), 23–27 (in Japanese), https://doi.org/10.11475/sabo1973.42.4_23, 1989.
- Pan H. L., Jiang, Y. J., Wang, J., and Ou G. Q.: Rainfall threshold calculation for debris flow early warning in areas with scarcity of data, *Nat. Hazards Earth Syst. Sci.*, 18, 1395–1409, <https://doi.org/10.5194/nhess-18-1395-2018>, 2018.
- ~~Peng, J. B., Fan, Z. J., Wu, D., Zhuang, J. Q., Dai, F. C., Chen, W. W., and Zhao, C. et al.: 2015: Heavy rainfall triggered loess-mudstone landslide and subsequent debris flow in Tianshui, China, *Eng. Geol.*, 186, 79–90, <https://doi.org/10.1016/j.enggeo.2014.08.015>, 2015.~~
- ~~Rengers, F.K., Kean, J.W., Reitman, N.G., Smith, J.B., Coe, J. A., and McGuire, L. A.: The Influence of Frost Weathering on Debris Flow Sediment Supply in an Alpine Basin, *J. Geophysical Res. Earth Surf.*, <https://doi.org/10.1029/2019JF005369>, 2020a.~~
- Rengers, F. K., McGuire, L. A., Oakley, N. S., Kean, J. W., Staley, D. M., and Tang, H.: Landslides after wildfire: initiation, magnitude, and mobility, *Landslides*, 17, 2631–2641, <https://doi.org/10.1007/s10346-020-01506-3>, 2020b.
- ~~Sidle, R. C., and Chigira, M.: Landslides and debris flows strike Kyushu, Japan, *EOS*, 85, 145–151, <https://doi.org/10.1029/2004EO150001>, 2011.~~
- ~~Saito, H., Nakayama, D., and Matsuyama, H.: Relationship between the initiation of a shallow landslide and rainfall intensity-duration thresholds in Japan, *Geomorphology*, 118, 167–175, <https://doi.org/10.1016/j.geomorph.2009.12.016>, 2010.~~
- Schlunegger, F., Badoux, A., McArdeell, B. W., Gwerder, C., Schnydrig, D., Rieke-Zapp, D., and Molnar, P.: Limits of sediment transfer in an alpine debris-flow catchment, Illgraben, Switzerland, *Quat. Sci. Rev.*, 28, 1097–1105, <https://doi.org/10.1016/j.quascirev.2008.10.025>, 2009.
- Scott, K. M., Vallance, J. W., Kerle, N., Luis Macías, J., Strauch, W., and Devoli, G.: Catastrophic precipitation-triggered lahar at Casita volcano, Nicaragua: occurrence, bulking and transformation, *Earth Surf. Process. Landf.*, 30, 59–79, <https://doi.org/10.1002/esp.1127>, 2014.
- Shakesby R. A., and Doerr S. H.: Wildfire as a hydrological and geomorphological agent, *Earth-Sci. Rev.*, 74, 269–307, <https://doi.org/10.1016/j.earscirev.2005.10.006>, 2006.
- Shanley, J. B., and Chalmers, A.: The effect of frozen soil on snowmelt runoff at Sleepers River, Vermont, *Hydrol. Process.*, 13, 1843–1857. [https://doi.org/10.1002/\(SICI\)1099-1085\(199909\)13:12<1843::AID-HYP879>3.0.CO;2-G](https://doi.org/10.1002/(SICI)1099-1085(199909)13:12<1843::AID-HYP879>3.0.CO;2-G), 1999.
- ~~Sidle, R. C., and Chigira, M.: Landslides and debris flows strike Kyushu, Japan, *EOS*, 85, 145–151, <https://doi.org/10.1029/2004EO150001>, 2011.~~

書式変更: フォント: (日) MS 明朝, (言語 1) 日本語

書式変更: フォント: (日) MS 明朝, (言語 1) 日本語

Song, D., Zhou, G. G. D., Xu, M., Choi, C. E., Li, S., and Zheng, Y.: Quantitative analysis of debris-flow flexible barrier capacity from momentum and energy perspectives, *Eng. Geol.*, 251, 81–92, <https://doi.org/10.1016/j.enggeo.2019.02.010>, 2019.

760 [Staffler, H., Pollinger, R., Zischg, A., and Mani, P.: Spatial variability and potential impacts of climate change on flood and debris flow hazard zone mapping and implications for risk management, *Nat. Hazards Earth Syst. Sci.*, 8, 539–558, <https://doi.org/10.5194/nhess-8-539-2008>, 2008.](#)

Staley, D. M., Kean, J. W., Cannon, S. H., Schmidt, K. M., and Laber, J. L.: Objective definition of rainfall intensity–duration thresholds for the initiation of post-fire debris flows in southern California, *Landslides*, 10, 547–562, <https://doi.org/10.1007/s10346-012-0341-9>, 2012.

765 Staley, D. M., Wasklewicz, T. A., [and Kean, J. W.](#): Characterizing the primary material sources and dominant erosional processes for post-fire debris-flow initiation in a headwater basin using multi-temporal terrestrial laser scanning data–, *Geomorphology*, 214, 324–338, <https://doi.org/10.1016/j.geomorph.2014.02.015>, 2014.

770 Staley, D. M., Negri, J. A., Kean, J. W., Laber, J. L., Tillery, A. C., and Youberg, A. M.: Prediction of spatially explicit rainfall intensity–duration thresholds for post-fire debris-flow generation in the western United States. *Geomorphology*, 278, 149–162, <https://doi.org/10.1016/j.geomorph.2016.10.019>, 2017.

Stoffel, M., [and Huggel, C.](#): Effects of climate change on mass movements in mountain environments. *Prog. in Phys. Geogr.*, 36, 421–439, <https://doi.org/10.1016/j.scitotenv.2014.02.102>, 2012.

775 [Stoffel, M., Tiranti, D., and Huggel, C.: Climate change impacts on mass movements—Case studies from the European Alps. *Sci. Total Environ.*, 493, 1255–1266, <https://doi.org/10.1016/j.scitotenv.2014.02.102>, 2014a.](#)

Stoffel, M., Mendlik, T., Schneuwly-Bollschweiler, M., and Gobiet A.: Possible impacts of climate change on debris-flow activity in the Swiss Alps, *Clim. Change*, 122, 141–155, <https://doi.org/10.1007/s10584-013-0993-z>, 2014b.

780 Theule, J. I., Li’ebault, F., Loye, A., Laigle, D., and Jaboyedoff, M.: Sediment budget monitoring of debris-flow and bedload transport in the Manival Torrent, SE France, *Nat. Hazards Earth Syst. Sci.*, 12, 731–749, <https://doi.org/10.5194/nhess-12-731-2012>, 2012.

Theule, J. I., Crema, S., Marchi, L., Cavalli, M., and Comiti, F.: Exploiting LSPIV to assess debris-flow velocities in the field, *Nat. Hazards Earth Syst. Sci.*, 18, 1–13, <https://doi.org/10.5194/nhess-18-1-2018>, 2018.

[Tillery, A., and Rengers, F.: Controls on debris-flow initiation on burned and unburned hillslopes during an exceptional rainstorm southern New Mexico, *Earth Surf. Process. Landf.*, 45, 1051–1066, <https://doi.org/10.1002/esp.4761>, 2019.](#)

785 [Yamazaki, T., Nishida, A., & Kondo, J.: Seasonal frost depth of ground with the bare surface, snow cover and vegetation, *J. Japan. Soc. Snow Ice*, 60\(3\), 213–224, <https://doi.org/10.5331/seppyo.60.213> \(in Japanese\), 1998.](#)

Walvoord M. A., Voss, C. I., Ebel, E. A., and Minsley, B. J.: Development of perennial thaw zones in boreal hillslopes enhances potential mobilization of permafrost carbon, *Environ. Res. Lett.*, 14, 015003, <https://doi.org/10.1088/1748-9326/aaf0cc>, 2019.

書式変更: フォント: (日) MS 明朝, (言語 1) 日本語

790 [Watanabe, K., and Flury, M.: Capillary bundle model of hydraulic conductivity for frozen soil, *Water Resour. Res.*, 44, W12402, <https://doi.org/10.1029/2008WR007012>, 2008.](#)

[Xu, W., Yu, W., Jing, S., Zhang, G., and Huang, J.: Debris flow susceptibility assessment by GIS and information value model in a large-scale region, Sichuan Province \(China\), *Nat. Hazards*, 65, 1379–1392, <https://doi.org/10.1007/s11069-012-0414-z>, 2013.](#)

795 [Yamazaki, T., Nishida, A., and Kondo, J.: Seasonal frost depth of ground with the bare surface, snow cover and vegetation, *J. Japan. Soc. Snow Ice*, 60\(3\), 213–224, <https://doi.org/10.5331/seppyo.60.213> \(in Japanese\), 1998.](#)

[Zhang, K., Wang, S., Bao, H., and Zhao, X: Characteristics and influencing factors of rainfall-induced landslide and debris flow hazards in Shaanxi Province, China, *Nat. Hazards Earth Syst. Sci.*, 19, 93–105, 2019.](#)

Zhang, T.: Influence of the seasonal snow cover on the ground thermal regime: An overview, *Rev. Geophys.*, 43, RG4002, <https://doi.org/10.1029/2004RG000157>, 2005.

800 [Zhao, L. and Glay, D. M.: A parametric expression for estimating infiltration into frozen soils, *Hydrol. Process.*, 11, 1761–1775, \[https://doi.org/10.1002/\\(SICI\\)1099-1085\\(19971030\\)11:13<1761::AID-HYP604>3.0.CO;2-O\]\(https://doi.org/10.1002/\(SICI\)1099-1085\(19971030\)11:13<1761::AID-HYP604>3.0.CO;2-O\), 1998.](#)

Zhou, W., and Tang, C.: Rainfall thresholds for debris flow initiation in the Wenchuan earthquake-stricken area, southwestern China, *Landslides*, 11, 877–887, <https://doi.org/10.1007/s10346-013-0421-5>, 2014.

805

書式変更: 英語 (米国)

書式変更: 英語 (米国)

書式変更: フォント: (日) MS 明朝, (言語 1) 日本語