

Rapid Holocene bedrock canyon incision of Beida River, North Qilian Shan, China

Yiran Wang¹, Michael E. Oskin¹, Youli Li², Huiping Zhang³

¹Department of Earth and Planetary Sciences, University of California, Davis, California, USA

5 ²College of Urban and Environmental Sciences, Peking University, Beijing, China

³Institute of Geology, China Earthquake Administration, Beijing, China

Correspondence to: Yiran Wang (yrwwang@ucdavis.edu)

Abstract. Located at the transition between monsoon and westerly dominated climate systems, major rivers draining the western North Qilian Shan incise deep, narrow canyons into latest Quaternary foreland basin sediments of the Hexi Corridor.

10 Field surveys ~~and previously published geochronology~~ show that the Beida River incised 125 m at the mountain front over the Late Pleistocene and Holocene at an average rate of 6 m/kyr. We hypothesize that a steep knickzone, with 3% slope, initiated at the mountain front and has since retreated to its present position, 10 km upstream. ~~Terrace-Additional terrace-dating results suggestsuggests that~~ this knickzone formed around the mid-Holocene, over a duration of less than 1.5 kyr, during which incision accelerated ~~from 6 m/kyr~~ to at least 25 m/kyr. These incision rates are much ~~largerfaster~~ than the uplift rate across the
15 North Qilian fault, which suggests a climate-related increase in discharge drove rapid incision over the Holocene and formation of the knickzone. Using the relationship between incision rates and the amount of base level drop, we show the maximum duration of knickzone formation to be \sim 700 yr and the minimum incision rate to be 50 m/kyr. ~~This~~~~We interpret that this~~ period of increased river incision ~~is the result of increasing excess discharge, which likely~~ corresponds to a pluvial lake-filling event at the terminus of the Beida River and correlates with a wet period driven by strengthening of the Southeast Asian Monsoon.


20 1 Introduction


An incising river responds to tectonic or climatic perturbation by adjusting its slope, expressed by formation of knickpoints or knickzones (Crosby and Whipple, 2006; Tucker and Whipple, 2002; Whittaker, 2012), and through changes of its channel width (Finnegan et al., 2005). Understanding the evolution and migration of knickzones, channel width, and the coupling between these adjustments, is important in unravelling the type, duration, and amplitude of a perturbation (Attal et al., 2011;

25 Berlin and Anderson, 2007; Bishop et al., 2005). Previous studies on headward migrating knickpoints focus on the role of tectonic uplift or a base level fall, and usually regard climate conditions and channel width as constant (e.g. Tucker and Whipple, 2002; Crosby and Whipple, 2006; Haviv et al., 2006; Wobus et al., 2006). Here we present a case of steep, quickly retreating knickzones within the western North Qilian Shan, formed under the combined influence of climatic change and lithologic control. Through modelling of incision of the Beida River, as recorded by its profile and stream terraces preserved along its
30 course, we suggest this knickzone was formed during a short period, ~~4-5~~~~four to five~~ thousand years before present, under an

exceedingly fast incision rate. This is most likely to be the result of an increase in river discharge, and perhaps a commensurate decrease of sediment supply.

In western China, the North Qilian Shan is the source of several northeast flowing rivers with deep canyons incised across the mountain-basin boundary (Figure 1a). In the western North Qilian Shan, three major tributaries of the Hei He drainage: Maying, Hongshuiba, and Beida River, carve ~~deeply~~deep canyons into the foreland sediments and across the fault-controlled boundary with the bedrock hinterland, forming prominent knickzones within the hinterland that are tens of meters high (Figure 2b-d). As one of these deeply incised rivers, the Beida River is characterized by a prominent knickzone which separates its profile into three patches (upper patch 1, knickzone patch 2, and lower patch 3; Figure 3a). Each patch can be distinguished by different channel slopes and channel widths: gentle and wide upper patch, steep and narrow knickzone, and a lower patch with a gentle slope similar to the upper patch, but a narrower channel. The successive generation and retreat of these patches corresponds to different boundary conditions (cf. Royden and Perron, 2013), and together record the incision history at the mountain front.





45 **Figure 1** a. Digital elevation map of the **research area and the Hei He drainage system**. **Glacial/Active glacial coverage mapped** based on Raup et al., (2007). Inset figure: Location of **our research area** with respect to monsoon and westerly moisture sources.






Figure 2 b-d. River and maximum elevation profiles of three major tributaries of the Hei He drainage, and swath profiles of maximum topography the North Qilian Shan surrounding the rivers. Swaths are, Maximum elevation sampled from swaths 16 km in width, centred on each river. Locations of active strands of the North Qilian fault denote foreland-hinterland junction at mountain front. All three rivers exhibit deep incision below the top of foreland-basin sediments. Red highlights knickzone reach within bedrock. Dashed line in **CD is reconstructed profile of the Beida River from the ca. ~24 ka **T1 terrace tread**, prior to onset of rapid incision phase.**

50








Figure 2 a) ~~2~~ Map of the Beida River within the north Qilian Shan, with survey points and geochronology sample sites indicated. b) ~~2~~ Photo of Beida River canyon and sample site a and b within the foreland, looking upstream. Thick layers of gravel are exposed on the canyon walls. c) ~~2~~ Photo of Beida River canyon and T1 terraces near the mountain front, looking east (location in figure 2a). In this photo, the 24 kyr T1 terrace tread is ~40 m above the 9.5 kyr ~~inset T1~~ terrace tread. d) ~~2~~ Photo of Beida River canyon and sample site ~~eB-03~~ inside the mountains, looking upstream. This photo shows that the Beida River deeply incises into the bedrock below the terrace fill. e-h. cross sections of the Beida River terraces at each sample site.

2 Geological background

The Qilian Shan form the ~~northeastern~~northeaster margin and the youngest growing portion of the Tibetan plateau (Tapponnier et al., 2001). Thermochronology studies suggest that uplift of the North Qilian Shan commenced around the mid-Miocene, with an exhumation rate of ~1 m/yr (Jolivet et al., 2001; Zheng et al., 2010; He et al., 2017, 2021). The Hexi Corridor, north of the Qilian Shan, consists of a chain of foreland basins. Bordering arid central Asia, the Qilian Shan and Hexi Corridor occupy the transition zone between Southeast Asian Monsoon and westerlies (An et al., 2001; Wei and Gasse, 1999). The monsoon brings summer rain inland while the mid-latitude westerlies bring dry air and a small amount of water vapor in winter. ~~The monsoon influence diminishes westward as the annual precipitation within the Hexi Corridor decreases from 300-400 mm in the east to <100 mm in the west (Meng et al., 2012). At high altitude within the Qilian Shan (>4000 m), the precipitation is significantly greater, with an overall trend that also decreases presently from east (>700 mm) to west (~300 mm)~~Monsoon influence diminishes and annual precipitation declines from east to west, both within the Hexi Corridor and at high altitude within the Qilian Shan (Meng et al., 2012; Shi et al., 2006). At the mountain front bounding the western Hexi Corridor, the annual precipitation declines from 200-300 mm/yr at the Maying River to 100-200 mm/yr at the Beida River. At high altitude (>4000 m) within the western North Qilian Shan, there is a similar annual precipitation gradient from east to west, from 300-400 mm/yr to 200-300 mm/yr (Qiang et al., 2016; Geng et al., 2017). The maxima of precipitation occur at approximately 4200 m elevation (Chen et al., 2018). The modern glacial equilibrium line altitude of Qilian Shan increases from 4400 to 5000 m from northeast to southwest (Shi, 2011), reflecting the westward decrease in precipitation. Between the year 2005 to 2010, within the Qilian Shan there were 2684 glaciers with a total area of 1600 km² and an ice volume of 84 km³ (Guo et al., 2014; Sun et al., 2015). These glaciers covered approximately 4% of the landscape above 4000 m elevation. The extent of these glaciers has fluctuated repeatedly throughout the Quaternary. Dating of moraines suggests that glacial advances have occurred during the little ice age (~1300-1850 A.D.), MIS (Marine Isotope Stage) 2, MIS 4, MIS 6, and MIS 12; some glacial expansion may have occurred during MIS 3 as well (Shi et al., 2006). The glacial equilibrium line altitude of Qilian Shan during the Last Glacial Maximum (LGM) is estimated to be as much as 400 m lower than present (Hu et al., 2014; Owen et al., 2003; Shi et al., 2006; Zhao et al., 2001; Zhou et al., 2002).

The Hei He ~~(river)~~ forms the largest drainage basin in the North Qilian Shan, and terminates within the Juyanze paleolake basin, north of the Hexi Corridor. Sediment and core records from the Juyanze paleolake basin indicate frequent dry-wet oscillations over the past 11,000 yr (Hartmann and Wünnemann, 2009; Herzschuh et al., 2004; Mischke et al., 2002, 2005). The highest lake level occurred during the early-Holocene ~~(~10700-8900 yr B.P., ~20 m deep)~~, and the highest mid-Holocene

90 lake level (~15–17 m deep) occurred during 5400-3900 yr B.P. and peaked at about 4200 yr BP (East Juyanze lake, Hartmann and Wünnemann, 2009).

Three major tributaries, Beida, Hongshuibai, and Maying, join the Hei He from the south and west (Figure 1, table 1). In the hinterland, the three rivers flow through Pre-Cambrian and Paleozoic meta-sedimentary and meta-igneous rocks; (Figure S1); in the foreland, these rivers ~~incise into Quaternary deposit~~ sediments of into the Hexi Corridor basins. The North Qilian range overthrusts the southern margin of the Hexi Corridor, placing metasedimentary rocks against the Quaternary basin deposits. Presently, the channels of the Maying, Hongshiba, and Beida River ~~each incise over 100m~~ have incised deeply into Late Pleistocene alluvial fans of the proximal foreland basin, and ~~by a similar depth~~ into correlative fill terraces and bedrock within the range (Hetzl et al., 2019; Yang et al., 2020; Wang et al., 2020). ~~A~~, forming a prominent knickzone ~~occurs~~ along each river, ~~presently~~ located ~10-15 km upstream of the mountain front (Figure 2). ~~1b-d~~. The canyons of the three rivers are deepest at the mountain front, ~125 m, ~190 m, and ~240 m for Beida, Hongshuibai, and Maying River, respectively (Figure S2). The depths of these river canyons gradually decrease basinward until 25-30 km downstream, where the rivers form active alluvial fans.

105 ~~As the largest tributary of the Hei He, the Beida River is ~360 km long, occupying 6880 km² drainage area within the Qilian Shan (Ding et al. 1999; Jiuquan History Compilation Committee, 1998). Table 1 hydrological information of the three major rivers draining the western North Qilian Shan (Gansu Province Local Chronicles Compilation Committee, 1998)~~

	<u>Length (km)</u>	<u>Drainage area (km²)</u>	<u>Annual discharge (x 10⁸ m³)</u>	<u>Glacial coverage (km²); contribution to annual discharge</u>
<u>Beida</u>	<u>243</u>	<u>6910</u>	<u>6.53</u>	<u>137; 16%</u>
<u>Hongshuibai</u>	<u>87</u>	<u>1580</u>	<u>2.87</u>	<u>131; 35%</u>
<u>Maying</u>	<u>34</u>	<u>619</u>	<u>1.16</u>	<u>20; 12%</u>

Along the Beida River, at least two generations of fill terraces (T1 and T2) are preserved well and continuously in the hinterland and extend to the foreland basin. Our previous research (Wang et al., 2020) suggests that T1 was abandoned ~~at~~ after 24±3 kyr B.P., ~~and T2 during the last glacial period; T2~~ was abandoned ~~at~~ after 144±30 kyr B.P., during the penultimate glacial period.

110 Flights of terraces inset below both ~~main~~ T1 and T2 terrace treads, ~~marking mark~~ progressive degradation of the terrace fill and incision into underlying bedrock. Along the mountain front, strands of the North Qilian fault cut the terraces of Beida River, offsetting T1 and T2 vertically by 14.8 m and >59.7 m, respectively. In the hinterland, the T2 terrace profile reveals a long wavelength fold (~30km) with maximum uplift of ~120 m relative to T1. Combined, the fault offset and folding indicate a maximum uplift rate of ~1 m/kyr at the fold crest, and a horizontal shortening rate of 1.4±0.4 m/kyr across the North Qilian

115 Shan (Wang et al., 2020).

3 Methods

3.1 Field survey

The late Pleistocene T1 fill terrace, up to 60 m thick, is preserved continuously along the narrow, ~~lower~~ bedrock gorge of the Beida River. This terrace grades to an extensive alluvial fan deposit emanating from the mountain front, with minor disruption from reverse fault offsets (Figure 42). The Beida river gorge cuts across the fault-controlled basin boundary, forming a narrow slot canyon up to 125 m deep within the foreland-basin fan gravels. We mapped and surveyed the terrace sequence and the course of Beida River inside the mountain range using a laser rangefinder (~0.3 m distance accuracy, 0.25° inclination accuracy) and differential GPS. Wherever possible, the terrace tread (top of the fluvial gravel), terrace strath (base of fluvial gravel), and present riverbed were measured together (Figure 3, Table S1). In the foreland, we extract terrace elevations and the river profile from an 8 m resolution digital elevation model, produced by the Polar Geospatial Center (Shean, 2017). For the hinterland tributaries, we extract river profiles from the same 8 m resolution DEM. Present bedrock channel widths were measured from Google Earth imagery at 100 m intervals along the river course. We measured the width of the water surface from imagery acquired during the wet season, mostly between July to September, 2010 to 2016. Due to limited data availability, a few measurements were obtained from imagery acquired in May and October (Table S2).

3.2 Geochronology

The abandonment age of T1 was dated to be 24 ± 3 kyr by combining optically stimulated luminescence (OSL) and Terrestrial Cosmogenic Nuclide (^{10}Be) exposure ages (Figure 32, sample site aB-01; Wang et al., 2020). To document the post-24 kyr incision history of Beida River within the hinterland, we collected charcoal samples from the fine sand and silty overbank deposits on three inset terraces (Figure 32, sample site e, d, eB-03, B-04, B05, Figure S2S3). These overbank deposits were deposited after terrace formation, but before incision was sufficient to isolate the terrace surface from flood events. ~~An OSL sample (Figure 3 sample site b) was collected on an even lower inset terrace in~~ At one site within the foreland basin-, ~~we collected an OSL sample~~ from the bottom of the loess covering ~~the a low inset terrace deposits.~~ (Figure 2, sample site B-02). Ten charcoal samples were measured at the Keck Carbon Cycle AMS Facility at UC Irvine. The results were calibrated with IntCal14 calibration curve (Reimer et al., 2013) (Table 42). The OSL sample (BD-O-12) was processed and measured at the State Key Laboratory of Earthquake Dynamics, China Earthquake Administration. The equivalent doses (D_e) for the pure fine-grained quartz were determined by the simplified multiple aliquot regenerative-dose (SMAR) protocol (Table 23, Table S3, Figure S4S4).

Table 2 ^{14}C age of Beida River terraces

Sample site <u>site</u> ¹	Coordinates	Sample name	fraction \pm Modern	$D^{14}\text{C}$ \pm (‰)	^{14}C age \pm (BP)	Calibrated age \pm	
						1σ (BP)	2σ (BP)

eB-03 (T1'; 115 m)	39.60126°, 98.01365°	BDC-3	0.3462	0.0028	-653.8	2.8	8520	70	9473-9544	9332-9340 9404-9632 9645-9657
		BDC-4	0.3393	0.0008	-660.7	0.8	8680	20	9557-9630 9647-9653	9552-9679
	BDC-5	0.3378	0.0008	-662.2	0.8	8720	20	9611-9699	9561-9737	
	BDC-6	0.3320	0.0009	-668.0	0.9	8855	25	9895-9949 9990-10012 10025-10038 10061-10134	9784-9848 9861-9878 9883-9966 9982-10155	
eB-04 (T1'b; 90 m)	39.57600°, 97.99493°	BDC-8	0.5973	0.0011	-402.7	1.1	4140	15	4617-4652 4669-4703 4757-4765 4784-4809	4580-4726 4752-4770 4780-4815
		BDC-9	0.5715	0.0011	-428.5	1.1	4495	20	5054-5077 5105-5136 5163-5189 5213-5228 5231-5251 5257-5281	5047-5147 5153-5202 5210-5288
	BDC-10	0.5862	0.0011	-413.8	1.1	4290	15	4844-4856	4839-4862	
eB-05 (T1'a; 42 m)	39.498771°, 97.971940°	BDC-11	0.4702	0.0012	-529.8	1.2	6060	20	6893-6944	6807-6811 6856-6979
		BDC-12	0.4497	0.0029	-550.3	2.9	6420	60	7309-7419	7185-7186 7246-7439
		BDC-14	0.4725	0.0010	-527.5	1.0	6025	20	6800-6815 6845-6901	6797-6934

145 1. Inside the bracket shows the terrace that the sample collected from, and the height of the terrace above the riverbed. Note that labels a, b, and indicate local terrace ordering, and cannot be correlated to terraces upstream and downstream.

Table 3 OSL age of loess covering terrace tread

Sample Site	Sample no-name ¹	Coordinates	U /ppm	Th /ppm	K (%)	Water Content (%)	Dose Rate (Gy/ka)	Equivalent Dose ¹ (Gy)	Age ⁺² (ka)
bB-02	BD-O-12	98.02299, 39.64376	2.34±0.10	9.32±0.28	1.67±0.06	0	3.5±0.3	11.4±0.7	3.2±0.2

1. Grain size <100 μm

±2. Uncertainties in equivalent dose, dose rate and age determinations are expressed at the 1σ confidence level.

150

3.3 Bedrock incision model

We apply the concept of slope patches (Royden & Perron, 2013) to model the evolution of the Beida River stream profile. The formation of a slope patch is based on stream power, which has the form

$$\frac{dz}{dt} = K \left(\frac{QS}{W} \right)^n, \quad (1)$$

155

where z is the channel elevation, t is time, ~~A is drainage area~~, Q is river discharge, S is channel slope, W is channel width, and K and n are an empirical erosional efficiency and exponent, respectively (Tucker & Whipple, 2002; Whipple & Tucker, 1999). A slope patch forms in the bedrock channel immediately upstream of the channel outlet, with channel slope that develops in balance with the rate of base-level fall. Setting $\frac{dz}{dt}$ to the incision rate, I , at the mountain front, we rearrange equation 1 to solve for this channel slope:

160

$$S = \left| \frac{dz}{dx} \right| = \left(\frac{I}{K} \right)^{\frac{1}{n}} \left(\frac{W}{Q} \right) \quad (2)$$

165

~~During formation of a slope patch, river profile elevation is found by integrating equation 2 over its finite span x_b to x .~~ For the case of the Beida River, no major tributary enters along its lower 30 km long course; the drainage area of the Beida River, measured from 30 m SRTM DEM, at the river outlet (mountain front) is $\sim 6.91 \times 10^9$ km², while the drainage area above the knickzone is $\sim 6.84 \times 10^9$ km², ~~leading to a ignorable~~ difference of $\sim 1\%$. Thus, we assume that Q does not vary spatially along the channel course and may be treated as constant over an incision phase, though it may vary from one phase to the next. We also assume that channel width (W) remains constant within a given slope patch. During formation of a slope patch, river profile elevation is thus found by integrating equation 2 over its finite span x_b to x :

$$z(x) = \left(\frac{I}{K} \right)^{\frac{1}{n}} \left(\frac{W}{Q} \right) (x - x_b) + z_b = S(x - x_b) + z_b \quad (3)$$

170

where x_b and z_b are the horizontal position and elevation ~~of the channel outlet, respectively~~ at base level, respectively. We define base level as the bedrock-basin transition at the mountain front.

We model the bedrock incision history of the Beida river as a consequence of varying ~~discharge~~incision rate over time. ~~Each slope patch along its course corresponds to a past discharge condition at base level.~~ Once formed, a slope patch behaves as a kinematic wave, retaining its gradient as it retreats upstream (Perron and Royden, 2013). The elevation of the (n-1)th slope patch (the patch formed one stage before present) may thus be cast as a function of its slope during formation, $S_{(n-1)}$ and an effective base-level elevation $z_{b(n-1)}$ of the slope patch projected to the outlet position. This base level may be predicted by correcting the present base level elevation, z_b , by the difference in the amount of incision across neighboring patches n and n-1,

$$z_{b(n-1)} = z_b + (I_{n,j} - I_{n-1,j})t_j. \quad (4)$$

$I_{n,j}$ is the incision rate of patch n, currently being formed during time interval t_j , directly upstream of the outlet. $I_{n-1,j}$ is the incision rate of patch n-1 during that time interval t_j . Note that ~~because discharge has changed,~~ the ~~latter~~ incision rate ~~is~~may be different than the incision rate during formation of patch n-1 (i.e. faster for an increase in discharge). However, varying discharge over time is not required for our model, only changes of channel slope driven by varying of incision rate at the mountain front. For the (n-2)th patch, the effective base level contains two correction terms (see z_{b1} , z_{b2} of figure [4e3c](#)),

$$z_{b(n-2)} = z_b + (I_{n,j} - I_{n-2,j})t_j + (I_{n-1,j-1} - I_{n-2,j-1})t_{j-1}. \quad (5)$$

This may be generalized to additional slope patches, each corrected by the incision rate differences between patches. We apply equations 4 and 5, combined with the incision recorded in stream terraces adjacent to the Beida river, to constrain its incision-rate history. Note that eqs. 4 and 5 are derived on the premise that discharge (Q) is constant during each incision phase, and that channel width (W) may vary between different patches but remains constant within a patch – i.e., that eq. 3 is valid.

4 Results

190 4.1 Beida River ~~patches and terraces~~profiles


Presently, the 30 km reach of the Beida River upstream of the mountain front is entirely contained within a bedrock channel. Channel slopes, measured directly from fitting the long profile, show a knickzone between 10 to 12 km upstream of the mountain front. (Figure 3). The knickzone divides the river profile into three patches: patch 1, upstream of the knickzone, with slope of 0.013; patch 2, the knickzone itself, with slope of 0.029; patch 3, below the knickzone with slope of 0.012 (Figure 3c). Channel width also varies along the course of the Beida River ~~(Figure 4d).~~ In patch 1, the channel width ranges between 14 to 140 m, with an average of 43 m; in patch 2, the channel width ranges between 8 to 29 m, with an average of 17 m; in patch 3, the channel width ranges between 14 to 70 m, with an average of 30 m (Figure 4d)–3d). In the foreland, the river incises into the 24 kyr T1 terrace, which is part of an extensive alluvial fan surface. The river canyon is deepest at the mountain front, and its depth gradually decreases downstream, indicating a decrease of river gradient within the foreland from 24 kyr B.P. to present. Near the mountain front, the slope of the riverbed is presently ~0.010, while the slope of the adjacent alluvial fan surface is ~0.013.

North of the mountain front, the T1 fill terrace merges into the extensive T1 alluvial fan. Within the mountain range, the T1 fill ranges from 60 to 80 m of thickness, and consists of unconsolidated medium to poorly sorted, well rounded boulder cobble conglomerate and sandy conglomerate. The lithology of the sediments mainly consists of quartzite, granite, slate, and limestone.

205 T1 treads are very well preserved, only covered by 1–2 m loess cap except at tributary junctions, where alluvial fans are deposited upon the tread. The T1 tread presently lies 125 m above present riverbed of patch 3 at the mountain front, and 60 m above riverbed of patch 1. Bedrock below the T1 strath is exposed continuously for 30 km upstream of the mountain front. The most prominent inset terrace, T1', is preserved continuously at an elevation 10 to 15 m lower than T1 tread, both inside the mountain range and in the foreland basin. ¹⁴C dating of charcoal samples at site c (Figure 3, 4a) indicates abandonment of

210 T1' prior to 9.5 ± 0.16 kyr BP. Inside the mountain range, several levels of inset terraces are preserved along the river, generally between a few meters to 40 m below the T1' tread, recording progressive incision of the Beida River. An inset terrace at site e, situated 42 m above the present riverbed of patch 1, yielded an age of 6.9 ± 0.07 kyr. At site d, an inset terrace with tread elevated 90 m above the patch 3 riverbed yielded an age of 4.7 ± 0.12 kyr (Figure 3, 4a, Table 1). In the foreland, the river incised into the basin deposits. T1' and other minor inset terraces of T1 occur 10–30 m below the T1 tread and 80–100 m above present river level. Terraces are absent between this elevation and 30–40 m above the riverbed (Figure 4a, 4b). An OSL

215 sample was collected from an inset terrace 2.5 km downstream of the mountain front, with its tread 37 m above the present riverbed, yielding an age of 3.2 ± 0.20 kyr from the bottom of the loess cover (site b; Table 2). Notably, the difference in elevation and age of site b and d indicates a pulse of very rapid incision of 43 m in 1.5 ± 0.3 kyr, or an average of 28 m/kyr, uncorrected for differential uplift between the two sites.







220

Major tributaries of the Beida River all have knickpoints at their junction with the main stem (Figure 4). Below the main-stem knickzone, these tributary knickpoints are mostly over 100 m high and form waterfalls. Above the main stem knickzone, the tributary knickzones are steep but graded to the main stem, and only half as high as those downstream of the main-stem

225

knickzone. By projecting the tributary profiles to the main stem, we find that the tributary junctions define a single, graded main-stem profile without evidence for a knickzone (Figure 4).




230 Figure 3 a) Longitudinal profile of the Beida River channel, with terrace treads (T2, T1 and all T1 inset terrace treads) and strath elevations, ~~with~~ sample sites and ages also indicated. Dash lines are inferred prior river profiles corresponding to the sample site elevations, with the relative height above present base level annotated (not corrected for tectonic uplift). b) terrace elevations (red) and channel elevations (open circles) along the Beida River within the foreland basin. Note sparse terrace record below T1/T1'. c) Three patches of Beida river profile, projected to effective base level at the mountain front. d) Present bedrock channel width measured at 100m intervals along the course of the Beida River. The width measurements that are within ~500 m upstream & downstream of the dam are excluded.

235 ~~4.2 Exposure of T1 bedrock strath~~


240 Though the terrace fill underlying the T1 tread is up to ~60-80 m thick, we find that the Beida river has not re-excavated its former channel. Instead, the river incised primarily along the western edge of its former canyon, and the paleocanyon fill and former walls of its bedrock gorge are preserved beneath the T1 fill deposit along much of the river course we surveyed (Figure 5, S3). Thus, bedrock incision likely commenced very soon after the onset of incision of T1. This is most clear for the ~3-4 km long reach immediately upstream of the North Qilian fault, where the wide T1 terrace treads (24 kyr) and T1' terrace tread (9.5 kyr) are preserved on the east bank of the Beida River, while only bedrock is exposed on the west bank. Farther upstream, exposures at several locations show that entire paleo-channel has been preserved beneath T1 and T1'. Our surveys further show that all observed terrace straths are located immediately beneath the prominent 9.5 kyr T1' terrace (Figure 4a). Therefore, though it is difficult to pinpoint the exact starting time of the bedrock incision, based on the terrace distribution and bedrock exposure along the channel, we suggest that the onset of bedrock incision along the Beida River may have started as early as ~9.5 kyr B.P., the age of the inset T1' terrace.

245

a



b



c







Figure 4 a. Map of the tributaries of the Beida River. Red delineates the main stem, blue delineates tributaries. b. Longitudinal profiles of the tributaries of the Beida River, as numbered in a. The main-stem knickzone is highlighted in red. Dashed red line connects the projection of tributary streams to the junction with the main stem based on stream gradient above each tributary knickzone.

4.2 Beida River terrace development

North of the mountain front, the 24 kyr T1 fill terrace merges into the extensive T1 alluvial fan. Though the river has incised an over 100 m deep canyon, no bedrock is exposed in the foreland. Within the mountain range, the T1 fill ranges from 60 to

260 80 m of thickness, and consists of unconsolidated medium to poorly sorted, well-rounded boulder-cobble conglomerate and sandy conglomerate. The lithology of the sediments mainly consists of quartzite, granite, slate, and limestone. T1 treads are very well preserved, only covered by 1-2 m loess cap except at tributary junctions, where alluvial fans are deposited upon the tread. The T1 tread presently lies ~115 m above the riverbed immediately north of the northernmost strand of the North Qilian fault, ~125-130 m above the present riverbed of patch 3 at the mountain front, up to ~130-135 m above the patch 3 upstream of the mountain front, and ~60 m above the riverbed of patch 1. Bedrock below the T1 strath is exposed continuously in the hinterland for at least 30 km along the Beida River. The most prominent inset (cut) terrace, T1', is preserved continuously at an elevation 7 to 20 m lower than T1 tread, both inside the mountain range and in the foreland basin. ¹⁴C dating of charcoal samples at site B-03 (Figure 2a, 2f, 3a) indicates abandonment of T1' occurred after 9.5 ± 0.16 kyr BP. Several levels of local inset (cut) terraces are preserved along the river, generally between a few meters to ~40 m below the T1' tread. Different from T1 and T1' terraces, these inset terraces are local features formed as the river incised and meandered (Merritts et al., 1994) and therefore it is difficult or impossible to correlate these to other terraces upstream and downstream. All inset terraces are cut into the T1 fill. No inset terraces appear to have been formed as the river cut below the level of the T1 fill. On top of all the inset terrace treads is half to 1 m thick fine sand layers with small gravel (<10 mm) layers in between, which we interpret as overbank sediments deposited during floods soon after the terrace surface was abandoned. Loess, 0.5 to 1.5 m thick, is deposited atop the overbank deposits.

275 Selected inset terraces were dated to constrain the incision history of the Beida River. An inset terrace at site B-05, situated 42 m above the present riverbed of patch 1, yields an age of 6.9 ± 0.07 kyr (Figure 2a, 2h, 3a, Table 2). At site B-04, an inset terrace with tread elevated 90 m above the patch 3 riverbed and ~24 m below the T1' tread yields an age of 4.7 ± 0.12 kyr (Figure 2a, 2g, 3a, Table 2). In the foreland, a suite of terraces was formed as the river first incised into the basin deposits. Here T1' and other younger inset terraces occur between 7 and 30 m below the T1 tread and 80 to 100 m above present river level. Terraces are absent between 80 m and ~30-40 m above the riverbed (Figure 3a, 3b). An OSL sample collected from basal loess capping an inset terrace tread, 37 m above the present riverbed, yielded an age of 3.2 ± 0.40 kyr (site B-02 of figure 2a, 2e, 3a; Table 3).

285 We find that the Beida River incised its present bedrock gorge adjacent to its former canyon filled by T1. This appears to have started with formation of the T1' terrace at 9.5 kyr. Upstream of the mountain front, the T1' terrace tread is preserved continuously on the east bank for 3.6 km, while only bedrock is exposed on the west bank. Though the T1 terrace fill is up to ~60-80 m thick in this area, bedrock is exposed in the canyon walls 10 to 15 m below T1' and 20 to 30 m below the top of the T1 fill. Farther upstream, mapped exposures show that the entire paleo-channel fill has been preserved beneath T1 and T1' for several channel reaches (Figure 5, S5).




Figure 5 Bedrock incision and terrace preservation along the Beida River. a. Terrace map of the lower reach of Beida River. T1 terraces are all preserved on the east side of the canyon. b. Terrace profiles of Beida River. Letters correspond to reaches where terrace mapping indicates that the original paleocanyon is preserved and the river incised into the adjacent bedrock soon after the abandonment of T1' (See Figure S3S5). c. Photo of the lower reach of Beida River, looking downstream, with paleochannel axis and paleo-canyon wall annotated. Here the river incised mostly into bedrock along the west edge of the T1 terrace fill deposit.

4.3 Incision rate estimation with terrace records

The incision rate at the mountain front ~~for each stage~~ may be calculated from the ages and relative heights of the T1 inset terraces. Because the sample sites are scattered along the river, the heights of these terraces cannot be compared directly. Instead, we reconstruct the elevation of the river channel at 9.5, 6.9, and 4.7 kyr B.P., by projecting the paleo-channel profile through the three sample sites to the mountain front based on the slope of patch 1 (Figure 4a3a). The heights of these paleo channels above the present riverbed elevation at the mountain front are 110 m, 96 m, 80 m, respectively (Table 34). After correction for faulting at the mountain front and folding of the range interior (Wang et al. 2020), the elevations without tectonic uplift at the mountain front are 102 m, 89 m, and 75 m above the present base-level riverbed, respectively (Table 4). For the foreland terraces, we project the 3). The oldest T1-2 kyr terrace (24.2 kyr) and the youngest terrace (3.2 kyr), 115 m and to the mountain front based on present alluvial channel slope, retaining an elevation of 37 m above the riverbed, a river. Because this terrace is located on the footwall of the North Qilian Fault, therefore now we do not apply an adjustment for tectonic uplift. We also do not correct this site for subsidence, because unlike for the hinterland, where we have good constraints on uplift is

305 ~~needed from deformed terraces, we do not have a marker of subsidence of the foreland. Wang et al. 2020 estimated that tectonic~~
~~subsidence is only a small fraction (<10%) of the hangingwall uplift and thus such a correction would be less than 0.5 m for~~
~~this terrace. For the 24 kyr T1 tread, we use the T1 height at the mountain front, 131 m. After adjusting for tectonic uplift (15~~
~~m), its corrected elevation is 116 m.~~ With these adjusted terrace heights, we calculate the incision rate at mountain front
between 24.2 to 9.5 kyr B.P. was ~~~0.91~~ m/kyr. ~~The incision~~ The incision rate accelerated over much of the Holocene, from ~5
310 m/kyr between 9.5 to 6.9 kyr B.P., ~~to ~6 m/kyr between 6.9 to 4.7 kyr B.P., and~~ to ~25 m/kyr, between 4.7 to 3.2 kyr B. P.
The incision rate remains high, at ~~~2512~~ m/kyr, from 3.2 kyr B. P. to present— (Figure 6). It is worth noting the 3.2 kyr terrace
date is a minimum age, which may lead to overestimation of the post-3.2 kyr incision rate and underestimation of incision rate
between 4.7 and 3.2 kyr. However, we consider the 3.2 kyr date to be a good estimate of the terrace age because loess has been
deposited continuously in this area since at least the mid-early Holocene (Küster et al., 2006). We also note that samples from
315 the loess cover atop of T1 terrace tread at site B-01 yield ages of 5.7 kyr and 6.5 kyr (Wang et al., 2020), which further supports
that loess deposition began at this time.

5 Discussion

5.1 Beida River knickzone formation and incision stages.


~~We rule out the possibility of tectonic uplift as a driver of the formation of the prominent knickzone (patch 2), for two reasons.~~
320 ~~First, we document that the knickzone was formed after the abandonment of T1 and T1', and that this knickzone was cut~~
~~largely within a newly formed bedrock channel, and not within re-excavated paleo-channel fill (Figure 4a). Therefore, it is~~
~~unlikely that the knickzone formed by re-excavation of an older bedrock channel with a pre-existing knickzone that may have~~
~~formed over a protracted period of uplift. Second, the rate of incision of the Beida River greatly exceeds the tectonic uplift~~
~~rate. Based on our previous research (Wang et al., 2020), the average vertical uplift rate at the mountain front is only ~0.6~~
325 ~~m/kyr. Additional uplift occurs via folding within the hinterland, reaching a maximum of ~15 m for the T1 terrace (vertical~~
~~displacement by the frontal fault excluded) at 5 to 10 km upstream of the range front. This folding contributes an additional~~
~~~0.6 m/kyr to the uplift rate. Compared to the ~12 m/kyr post 9.5 kyr B.P. average river incision rate, the tectonic uplift rate~~  
~~is an order of magnitude less. This suggests that the Beida River profile is and has been in a transient state since 9.5 kyr B.P.~~  
~~An alternative hypothesis is that the Beida River knickzone formed in place, related either to a change in erodibility of bedrock~~  
330 ~~or a major tributary confluence. Based on a drainage area analysis, no major tributary confluence occurs at the knickzone~~  
~~location. In the neighbouring Maying and Hongshuibai River, similar to the Beida River, present river channels also incised~~  
~~into Late Pleistocene fill terraces (22.5–26 kyr for Hongshuibai River, Hetzel et al., 2019, Yang et al., 2020), forming prominent~~  
~~knickzones 10–15 km upstream of the mountain front (Figure 2). This suggests these knickzones share similar origins,~~  
~~reflecting regional forcing. Local variations of lithology would thus be an unlikely cause for knickzone formation. In addition,~~  
335 ~~if the knickzone of the Beida River were a persistent feature of the channel, evidence should be preserved in the strath (paleo~~  
~~river bed) elevations beneath the T1 terrace. We find that T1 strath elevations shows no such sign of a knickzone (Figure 4a).~~

Therefore, neither the regional context nor local incision history suggest the presence of fixed knickzone, and this alternative hypothesis can be ruled out.

We interpret the knickzone of the Beida River as a migrating kinematic wave, related to an accelerated incision rate earlier in time and downstream of its present position. We suggest Beida River knickzone formed in response to an abrupt change in the rate of base-level lowering (Whipple and Tucker, 1999). It is likely that knickzone of the Beida River first formed at the mountain front, where bedrock is juxtaposed against Quaternary foreland basin sediments by the North Qilian fault. The knickzone is hypothesized to have formed here as a response to increased incision rate into the foreland basin sediments, thus lowering base level relative to the bedrock. The most likely cause for this increase in incision rate is a change in river hydrology during this time period, with perhaps a contribution from decreased sediment flux. Both mechanisms would increase transport capacity and promote incision of the unconsolidated foreland basin sediments. By happenstance, it appears, the Beida River incised the western edge of its channel that was filled by T1. This brought the river channel in contact with bedrock well above its former channel thalweg position below T1. This resulted in formation of a knickzone as incision of the foreland basin outpaced bedrock incision upstream of the mountain front.

Based on the slope patch theory for bedrock incision, we associate the three slope patches along the bedrock channel of the Beida River as formed during three incision stages since 9.5 kyr B.P.. Each patch is tied to the mountain front outlet where the bedrock channel transitions to an alluvial channel. During the 1<sup>st</sup> stage, relatively slow incision rate in the foreland formed the gentle slope and wide channel of patch 1. During the 2<sup>nd</sup> stage, incision rate increased in the foreland, which leads to the formation of a steeper, narrower patch 2. During the 3<sup>rd</sup> stage, incision rate in the foreland decreased, therefore the youngest patch 3 with gentle slope and wider channel formed, while the steeper patch 2 retreated upstream, replacing patch 1 (Figure 6). Combined with the terrace records, we therefore define the 1st incision stage to occur from 9.5 kyr to sometime around or after 4.7 kyr B. P., because the incision rates between 9.5 to 6.9 kyr B.P. and 6.9 to 4.7 kyr B.P. are identical within error, between 5–6 m/kyr (Figure 7). This is followed by the 2<sup>nd</sup> stage, with an incision rate of at least 25 m/kyr. The 3<sup>rd</sup>, present stage, started after 3.2 kyr B.P. with an incision rate of ~12 m/kyr. In addition to our terrace chronology, this incision history is also supported by the relative preservation of terraces along the Beida River, with a notable absence of terraces preserved between 40 and 80 m above river level (75 and 30 m below T1) within the foreland, during the period of most rapid incision inferred from the bracketing terrace record. Because the starting time and the duration of stage 2 is not well constrained due to the sparse terrace record, it is possible that stage 2 was shorter duration than 1.5 kyr and its ~25 m/kyr incision rate a minimum.





365

**Figure 6** Simplified model for Beida River incision since 9.5 kyr. The surveyed terrace treads and the paleo riverbeds we projected are corrected for tectonic uplift, therefore slightly deviated from the sample locations. The time period between 24.2 to 9.5 kyr B.P. is not included in this model because the river was mostly incising into the T1 terrace fill, and therefore the incision of the foreland and hinterland should be synchronous.


370

**Table 4** Terrace ages and relative heights.


| Terrace Age<br>(kyr)              | Original terrace height<br>(projected to mountain<br>front, m) | Tectonic<br>uplift rate <sup>1</sup>            | Terrace Height (Adjusted based on tectonic uplift)    |                         |
|-----------------------------------|----------------------------------------------------------------|-------------------------------------------------|-------------------------------------------------------|-------------------------|
| Terrace Age <sup>1</sup><br>(kyr) | Original terrace height<br>(projected to mountain<br>front, m) | Tectonic<br>uplift rate <sup>2</sup><br>(m/kyr) | Relative to present riverbed at<br>mountain front (m) | Relative to patch 1 (m) |
| 24.2 ± 3.3                        | 115 ± 3<br>131 ± 3 <sup>3</sup>                                | 0.62 ± 0.08                                     | 115<br>116 ± 3                                        | 63<br>64 ± 3            |
| 9.5 ± 0.16                        | 110 ± 2.7                                                      | 0.86 ± 0.24                                     | 102 ± 5                                               | 50 ± 5                  |
| 6.9 ± 0.07                        | 96 ± 3.2                                                       | 0.92 ± 0.25                                     | 89 ± 4.9                                              | 37 ± 5                  |
| 4.7 ± 0.12                        | 80 ± 2.4                                                       | 1.08 ± 0.29                                     | 75 ± 3.8                                              | 23 ± 3.8                |
| 3.2 ± 0.24                        | 37 ± 3                                                         | 0                                               | 37 ± 3                                                | -                       |

<sup>1</sup> terrace ages are expressed at the 2σ confidence level

<sup>2</sup>Tectonic uplift rates at the sample site are calculated based on the folding and faulting data in Wang et al., (2020).



<sup>3</sup> Here we report the terrace height at mountain front instead of the height at sample site.



375

**Figure 6 a) Terrace height above base level and the incision rate of each stage at the mountain front. b) Terrace height and incision rate of each stage above patch 1, upstream of the knickpoint. Details of elevations with correction for tectonic uplift are in Table 34.**

## 5 Discussion

### 380 **5.2 Coupled incision model for knickzone formation**

Geometric and timing relations for patches 1, 2, and 3 may be coupled to further constrain the duration and incision rate of the knickzone formation stage (stage 2). Here we formulate these relationships into two sets of constraints. Constraint 1 comes from the knickzone retreat process and the geometric relationship between patch 2 and 3. The effective base level of patch 2 ( $z_{b,2}$ , Figure 4c) is determined by the relative incision rate between patch 2 and patch 3, and the duration of the 3<sup>rd</sup> stage. Based on equation 4, we have

$$z_{b,2} = (I_3 - I_{2,3}) \times t_3 \quad (6a)$$

Because we may estimate  $I_3$  and  $t_3$  from the position of the youngest, 3.2 kyr B.P. terrace, and  $z_{b,2}$  from projection of the patch 2 river profile to the mountain front, we may calculate the incision rate along patch 2 during stage 3,  $I_{2,3}$  as

$$I_{2,3} = I_3 - z_{b,2}/t_3 \quad (6b)$$

390 It is reasonable to assume that during the knickzone formation, the incision rate along patch 2 should be greater than present incision rate,  $I_2 > I_{2,3}$ . With known  $I_3$ ,  $t_3$ , and  $z_{b,2}$  (see table S4 for details), the minimum rate of incision during knickpoint formation,  $I_2$ , is  $50 \pm 17$  m/kyr.

To find the maximum duration for the time of knickpoint formation,  $t_2$ , we introduce a second set of constraints, derived from the total incision at the mountain front since 9.5 kyr B.P. and the total incision along patch 1 since 6.9 kyr should both match the observed terrace record:

$$H_{9.5kyr} = t_1 \times I_1 + t_2 \times I_2 + t_3 \times I_3 = I_1(9.5 kyr - t_2 - t_3) + I_2 t_2 + I_3 t_3 \quad (7)$$

and

$$H_{6.9kyr} = I_1(6.9 kyr - t_2 - t_3) + I_2 t_2 + I_3 t_3 \quad (8)$$

400 In addition, the geometric relationship between patch 1 and 2 should also be satisfied. The effective base level of patch 1 ( $z_{b,1}$ ) is determined by the relative incision rate between patch 1 and patch 2, and patch 1 and patch 3, and the durations of the 2<sup>nd</sup> and 3<sup>rd</sup> stage. Based on equation 5, we have

$$z_{b,1} = (I_2 - I_{1,2})t_2 + (I_3 - I_{1,3})t_3 \quad (9)$$

Based on constraint 2 and  $I_{2min} = 50 \pm 17$  m/kyr, we calculate the maximum duration of the 2<sup>nd</sup> stage as  $700 \pm 340$  yr, which is at approximately half of the duration suggested by isolated terrace record.

### 405 **5.3.5.1 Channel width and bedrock incision rate**

Variations in channel width may exert as much influence as discharge on incision rate (Lave and Avuoa, 2000). Drainage area, and thus discharge, does not appreciably change across our research area, however the bedrock channel width varies

widely, from over 100 m (patch 1) to as low as ~10 m (patch 2). Based on bedrock channel erosional mechanisms, the channel width should scale with discharge and channel slope (Finnegan et al., 2005, 2007; Lamb et al., 2015; Turowski et al., 2007; Wobus et al., 2006; Wohl and David, 2008). Though the narrowing of patch 2, and a clear, almost factor of two increase in the average width from patch 2 to patch 3 ~~is~~, are both consistent with coupling of channel width and slope, it is obvious that the upper reach of the patch 1 is considerably wider than expected (Figure 4d3d). In addition, the cause for narrowing of channel width on patch 1, ~~a~~ over the 5 km reach upstream of the knickzone, is also not clear. Increased flow velocity and shear stress immediately above the knickzone (Haviv et al., 2006) could contribute to this narrowing, though it is unlikely that such a hydraulic effect would extend for kilometers upstream. Another possibility is that the wider reach of the Beida River is a result of lateral incision and removal of a more extensive T1 terrace fill. Though bedrock is exposed in the channel throughout patch 1, T1 terraces more extensively preserved along its widest reach.

~~Variations in channel width could help to explain the two-fold difference in incision rate between patch 3 and patch 1 (Figure 7) at almost identical channel slopes (Figure 4). The~~ Variations in channel width may exert as much influence as discharge on incision rate (Lave and Avuoc, 2000). For the Beida River, we observe that patch 1 and patch 3 share similar channel slopes (Figure 3c), but show a two-fold difference in their incision rates (Figure 6). We attribute this difference to the difference in average channel width between different patches (Figure 3d). We suggest that the narrower average width of patch 3 focuses stream power and enhances incision rate relative to patch 1 (Eq. 1). This explanation is at odds, however, ~~with the presence of the narrow, downstream narrower~~ reach of patch 1, which by this reasoning should be incising as fast as patch 3. We speculate that this reach could reflect recent removal of the T1 terrace and focusing of the channel as it inset into bedrock. We note that the channel slope here is slightly steeper than upstream, which could indicate acceleration of incision rate. At present the terrace record is insufficient to test this hypothesis.

## **5.2 Beida River knickzone formation and incision stages.**

We interpret the knickzone of the Beida River as a migrating kinematic wave, related to an accelerated incision rate earlier in time and downstream of its present position. We suggest the Beida River knickzone formed in response to an abrupt change in the rate of base-level lowering (Whipple and Tucker, 1999). It is likely that the knickzone of the Beida River first formed at the mountain front, where bedrock is juxtaposed against Quaternary foreland-basin sediments by the North Qilian fault. The knickzone is hypothesized to have formed here as a response to increased incision rate into the foreland-basin sediments, thus lowering base level relative to the bedrock. The most likely cause for this increase in incision rate is a change in discharge during this time period, with perhaps a contribution from decreased sediment flux. Both mechanisms would increase transport capacity and promote incision of the unconsolidated foreland-basin sediments (Tucker and Whipple, 2002). We suggest a change of discharge (and/or decrease of sediment flux) and the change in erodibility from foreland sediments to hinterland bedrock are the two key factors contributing to knickzone formation here. Without a change of erodibility, an increase of discharge would lead to uniform incision (if discharge is spatially invariant) of the bedrock channel, and relaxation (lowering

440 of the river gradient) for the alluvial channel. We hypothesize instead that exposure of bedrock within the channel at the fault contact with the foreland led incision of the unconsolidated foreland-basin sediments to outpace bedrock incision upstream, resulting in formation of the knickzone.

We can rule out the alternative hypothesis of tectonic uplift as a driver of the formation of the prominent knickzone (patch 2), for two reasons. First, judging by the continuity of the T1 terrace tread (Figure 3a) and our field work (Wang et al., 2020; figure S6), we can rule out the presence of an active fault under the knickzone. Second, the rate of incision of the Beida River greatly exceeds the tectonic uplift rate. Based on our previous research (Wang et al., 2020), the average vertical uplift rate at the mountain front since the abandonment of T1 is only ~0.6 m/kyr. Additional uplift occurs via folding within the hinterland, reaching a maximum of ~15 m for the T1 terrace (vertical displacement by the frontal fault excluded) at 5 to 10 km upstream of the range front. This folding contributes an additional ~0.6 m/kyr to the uplift rate. In addition, the long-term exhumation rate based on thermochronology is ~1 m/kyr (Zheng et al., 2010). In comparison, the average incision rate since the abandonment of T1 terrace (24.2 kyr) is ~5.4 m/kyr, and the average incision rate since the abandonment of T1' (9.5 kyr) is ~12 m/kyr, both much larger than the tectonic uplift or exhumation rate.

A second alternative hypothesis is that the Beida River knickzone formed at its present location, related either to a change in erodibility of bedrock or a major tributary confluence. Based on a drainage-area analysis, no major tributary confluence occurs at the knickzone location. The geologic map (Figure S1) shows no significant lithology change under the present knickzone location, which suggests local variations of lithology are unlikely to have caused knickzone formation at its present position. In addition, if the knickzone of the Beida River were a persistent feature of the channel, evidence for a buried knickzone should be preserved in the strath (paleo river bed) elevations beneath the T1 terrace. We find that T1 strath elevations show no such sign of a knickzone (Figure 3a). Therefore, neither the regional context nor local incision history suggest the presence of fixed knickzone, and this alternative hypothesis can also be ruled out.

Based on the calculated incision rates, we can divide the Beida River incision history since the abandonment of T1 into two major phases. The first phase is between 24 to 9.5 kyr B.P. with an average incision rate ~1 m/yr after correcting for tectonic uplift, representing a close to equilibrium state relative to the long-term uplift and exhumation rate of the North Qilian Shan. The second phase is between 9.5 kyr B.P. to present, with an average incision rate an order of magnitude larger than the previous phase, representing an accelerated incision state. During the first phase, the relatively slow incision rate allowed the Beida river to meander widely, not only in the foreland but also within its hinterland canyon. This meandering resulted in widening of the canyon as the river laterally eroded into the bedrock. This more or less stable state ended with the abandonment of the prominent T1' terrace. During the second phase, incision rate increased in both the foreland and the hinterland, and has been dominated by down-cutting instead of lateral erosion. Due to this change in incision rate, only local inset terraces were formed during this phase. It is also during this phase that tributary knickzones started to form in response to the increasing incision rate of the main stem. We also suggest that channel incision into bedrock began during this second phase, based on the following evidence. First, bedrock outcrops occur as high as 10 meters below the mapped remnants of the 9.5 kyr T1' tread.


Second, mapping of the T1 terrace fill shows that prior to formation of T1', the Beida River had cut laterally into bedrock along the western side of its canyon over the 3.6 km reach immediately upstream of the mountain front. Post-T1' incision occurred in this area, preserving the T1 paleo channel axis and much of its buried canyon wall below the T1 and T1' terraces. The post-9.5 kyr B.P. increase of incision rate thus led to isolation of the river course within the bedrock below T1'.

**Based on the slope patch theory for bedrock incision, we associate the three slope patches along the bedrock channel of the Beida River as formed during three incision stages over the second phase, since 9.5 kyr B.P.. In our model, each patch is projected to the mountain front outlet where the bedrock channel transitions to an alluvial channel. During the 1<sup>st</sup> stage, relatively slow incision rate in the foreland formed the gentle slope and wide channel of patch 1. During the 2<sup>nd</sup> stage, incision rate increased drastically in the foreland, which led to the formation of a steeper, narrower patch 2. During the 3<sup>rd</sup> stage, incision rate in the foreland decreased, forming the youngest patch 3 with gentle slope and wider channel, while the steeper patch 2 retreated upstream, replacing patch 1 (Figure 7). Combined with the terrace records, we therefore define the 1st incision stage to occur from 9.5 kyr to sometime around 4.7 kyr B. P., because the incision rates between 9.5 to 6.9 kyr B.P. and 6.9 to 4.7 kyr B.P. are identical within error, between ~5-6 m/kyr (Figure 6). This is followed by the 2<sup>nd</sup> stage, with an incision rate of at least ~25m/kyr.**

~~5.4 Possible causes of rapid incision of Beida River~~

~~Moderately~~The 3<sup>rd</sup>, present stage, started at or before 3.2 kyr B.P. with an incision rate of ~12 m/kyr. Because the starting time and the duration of stage 2 is not well constrained due to the sparse terrace record, it is possible that stage 2 has a shorter duration than 1.5 kyr and its ~25 m/kyr incision rate should be considered a minimum.

In addition to our terrace chronology, a three-stage incision history is also supported by the relative preservation of terraces along the Beida River, with a notable absence of terraces preserved between 40 and 80 m above river level (75 and 30 m below T1) within the foreland, during the period of most rapid incision inferred from the bracketing terrace record. These three incision stages have also left their imprints into the tributary knickzones (Figure 4). Steep knickzones, often ending at waterfalls, occur on tributaries below the main-stem knickzone, suggesting that when these tributary knickzones formed the main stem was incising very fast so that the incision rate of the tributaries could not keep pace. Gentler knickzones formed on tributaries that join upstream of the main-stem knickzone suggest slower incision rate of the main stem here, as the main-stem knickzone has not yet retreated through this area. These tributary knickzones continue to grow in height and upstream extent, as the incision rate in the main stem is still fast at present.



**Figure 7** Simplified model for Beida River incision since 9.5 kyr. The surveyed terrace treads and the paleo riverbeds we projected are corrected for tectonic uplift, therefore slightly deviated from the sample locations. The time period between 24.2 to 9.5 kyr B.P. is not included in this model because the river was mostly incising into the T1 terrace fill, and therefore the incision of the foreland and hinterland should be synchronous.

### 5.3 Coupled incision model for knickzone formation

Geometric and timing relations for patches 1, 2, and 3 may be coupled to further constrain the duration and incision rate of the knickzone formation stage (stage 2). Here we formulate these relationships into two sets of constraints. Constraint 1 comes from the knickzone retreat process and the geometric relationship between patch 2 and 3. The effective base level of patch 2 ( $z_{b2}$ , Figure 3c) is determined by the relative incision rate between patch 2 and patch 3, and the duration of the 3<sup>rd</sup> stage. Based

on equation 4, we have

$$z_{b2} = (I_3 - I_{2,3}) \times t_3 \quad (6a)$$

Because we may estimate  $I_3$  and  $t_3$  from the position of the youngest, 3.2 kyr B.P. terrace, and  $z_{b2}$  from projection of the patch 2 river profile to the mountain front, we may calculate the incision rate along patch 2 during stage 3,  $I_{2,3}$  as

$$I_{2,3} = I_3 - z_{b2}/t_3 \quad (6b)$$

It is reasonable to assume that during the knickzone formation, the incision rate along patch 2 should be greater than present incision rate,  $I_2 > I_{2,3}$ . With known  $I_3$ ,  $t_3$ , and  $z_{b2}$  (see table S4 for details), the minimum rate of incision during knickpoint formation,  $I_2$ , is  $50 \pm 17$  m/kyr.

To find the maximum duration for the time of knickpoint formation,  $t_2$ , we introduce a second set of constraints, derived from the total incision at the mountain front since 9.5 kyr B.P. and the total incision along patch 1 since 6.9 kyr should both match the observed terrace record:

$$H_{9.5kyr} = t_1 \times I_1 + t_2 \times I_2 + t_3 \times I_3 = I_1(9.5 kyr - t_2 - t_3) + I_2 t_2 + I_3 t_3 \quad (7)$$

and


$$H_{6.9kyr} = I_1(6.9kyr - t_2 - t_3) + I_{1,2}t_2 + I_{1,3}t_3 \quad (8)$$

525 In addition, the geometric relationship between patch 1 and 2 should also be satisfied. The effective base level of patch 1 ( $z_{b1}$ ) is determined by the relative incision rate between patch 1 and patch 2, and patch 1 and patch 3, and the durations of the 2nd and 3rd stage. Based on equation 5, we have

$$z_{b1} = (I_2 - I_{1,2})t_2 + (I_3 - I_{1,3})t_3 \quad (9)$$

530 Based on constraint 2 and  $I_{2min}=50 \pm 17$  m/kyr, we calculate the maximum duration of the 2<sup>nd</sup> stage as  $680 \pm 460$  yr (2  $\sigma$  confidence), which is approximately half of the duration suggested by the terrace record alone. It is worth noting that because the  $t_3$  we used in the calculation is based on a minimum age for the lowest terrace, the maximum duration of stage 2 may be even shorter and the minimum stage 2 incision rate at mountain front may be even faster than calculated here. In addition, though we are able to constrain the maximum duration of the 2<sup>nd</sup> stage to  $\sim 700$  yr, which suggests this  $<700$  yr event happened sometime between 4.7 to 3.2 kyr B.P., we do not have enough data to pinpoint when this event occurred within this 1500 yr time span.

535 To verify these model results, we ran simulations of the channel evolution using our parameter estimations and compare this with our surveyed channel profile (Figure 8). The simulation results fit well with the observed channel profile, while the modelled channel positions at the end of stage 2 have a relatively wide range of possible geometries because we have less control on the starting and ending time of stage 2. The simulation results therefore verify that the 700 yr maximum duration and 50 m/kyr minimum 2<sup>nd</sup> stage incision rate are good constraints for the Beida River incision history.



540 **Figure 8 Monte-Carlo simulation of channel position at the end of stage 2 and at present, end of stage 3, compared to the observed river profile. 500 simulations results shown. Parameters for simulations were selected from valid model results for the timing, duration, and incision rate stages 2 and 3.**



#### 545 5.4 Climate implications of the Beida River incision history

545 Based on our prior study (Wang et al., 2020), the major T1 and T2 fill terraces of the Beida River correspond to the last glacial and penultimate glacial periods, respectively. A similar pattern occurs along the Hongshuiba River, the next major river to the east of the Beida River. The T1 terrace here is similar in age as T1 at the Beida River (26.5±5.0 kyr in Yang et al., 2020; 22.5±2.2 kyr in Hetzel et al., 2019), and the T4 terrace here has similar age as the T2 terrace at the Beida River (153±12 kyr in Hetzel et al., 2019). In the eastern North Qilian Shan, terrace records of the Shagou River also show a fill-cut pattern  
550 synchronized to glacial-interglacial cycles (Pan et al., 2003). incision of the Beida River since 9.5 kyr B.P. was ~~Therefore, we suggest the formation of fill terraces along the major rivers draining the North Qilian Shan is controlled by the same mechanisms, linked to glacial-interglacial cycles. A full fill-cut cycle begins with the river backfilling its canyon with sediment during the glacial period. This is a period with elevated, glacially derived sediment flux, and perhaps a drier climate and lower~~  
555 discharge than at present. At the glacial-interglacial transition, the river transitions from deposition to erosion as sediment supply gradually declines. This corresponds to the first incision phase of the Beida River (24-9.5 kyr B.P.). The fast, second incision phase of the Bieda River starting after 9.5 kyr B.P. corresponds in time with highstands of several endorheic lakes located in the Northwest China including the Juyanze paleolake, which have been interpreted as a signal of intensification of monsoon influence (i.e., Herzschuh et al., 2006; Hartmann and Wünnemann, 2009; Jiang et al, 2008; Li et al, 2009; Rhodes et al, 1996; Wang et al, 2013). Therefore, the post-9.5 ka fast incision phase may relate to a regional increase in discharge, beginning near the end of the last glacial termination, as a result of the introduction of monsoon-derived summer moisture.

560 The mid-Holocene <680±460 yr event which punctuated by a <700±340-yr event, during which the gorge incised ~30-40 m. This the moderately fast incision of the Beida River since 9.5 kyr B.P. seems to be a unique event triggered by a result of an unusual short-term hydrologic perturbation. Similar knickzones are found in other western North Qilian rivers, i.e., Hongshuiba River and Maying River (Figure 1-2), suggesting), suggest that this was a regional event that affected the entire western North Qilian Shan.

570 Formation of knickzones at the bedrock basin boundary via an increase in incision rate into the foreland basin sediments requires an increase in water discharge or a decrease in sediment flux from upstream, either of which promotes an increase of available sediment transport capacity (excess discharge). Decrease in sediment flux could be either due to glacial retreat and reduction of glacially produced sediment supply, or due to stabilization of hillslopes due to increased vegetation coverage inside the North Qilian Shan. Both explanations could help to explain the transition from infill to incision of the T1 terrace. But neither explanation seems likely to explain the We suggest this short-term pulse of incision during stage 2.

575 One source of increasing water discharge during knickzone formation could have been glacial melt. Such a cause would increase discharge simultaneously for other rivers draining the glaciated highlands of the North Qilian Shan that display similar deep canyons and migrating knickpoints (Figure 1). However, present glacier coverage and discharge from the Beida River show that this explanation to be unlikely. The present ice cover and volume of Beida River drainage are 215.27 km<sup>2</sup> and 8.75 km<sup>3</sup>, respectively (Sun et al., 2015). With the present average annual discharge of Beida River, 0.64 km<sup>3</sup> (Ding et al., 1999), a

50% increase of discharge due to excess glacial melt would deplete stored glacial ice in merely 27 yrs. In fact, due to rising temperature, the glacial coverage of Beida River drainage has shrunk at a pace of approximately 9% per decade over the past 50 yrs (Sun et al., 2015), contributing 15% of the average discharge of Beida River. Another study on a glacier west of the Beida River drainage indicates that during Last Glacial Maximum, the paleo-glacier cover was only 34% larger than the contemporary glacier cover (Hu et al., 2014). Combining all the evidence, we suggest increasing glacial melt may have played a role but alone it is not enough to trigger a dramatic knickzone formation event in the Beida River.

The other likely source of increasing water discharge is increasing hydrologic anomaly was driven by an increase in precipitation, which could also affect neighbouring rivers. Compared to the glacial melt, precipitation has the potential of increasing water discharge to a higher level, especially in the arid-semi arid region like the North Qilian Shan. Lake records as a result of increased monsoon influence, and possibly enhanced by glacial melt due to warmer temperatures. Records of the terminal Juyanze paleo-lake suggest the highest mid-Holocene lake level appeared at 4.2 kyr B.P. (Hartmann and Wünnemann, 2009), and pollen records of the same area suggest a wet, or pluvial period between 5.4 and 3.9 kyr B.P. (Herzschuh et al., 2004). This timing correlates well with knickzone formation during our 2<sup>nd</sup> stage. Regionally, evidence for similar humid periods can also be found from Zhuyeze, a lake fed by Shiyang River of the eastern North Qilian Shan (Chen et al., 2006), Qinghai Lake (Chen et al., 2016) located within the southeast Qilian Shan, Tianchi Lake of Liupan Shan (Zhou et al., 2010), and Yanhaizi Lake of Inner Mongolia (Chen et al., 2003). Cave records from upper Hanjiang region and Qinling Mountains (Tan et al., 2018) and stratigraphic sections from the Loess Plateau also support the existence of a mid-Holocene humid period (Fang et al., 1999; Fang et al., 2003; Chen et al., 1997; Xiao et al., 2002). All of this evidence suggests there was a humid period during the mid-Holocene that correlates well with our stage 2 rapid incision period. Considering the western North Qilian Shan is located at the transition zone between the Southeast Asian Monsoon and mid-latitude westerlies, and that wet periods regionally correspond to increased monsoon influence (Chen et al., 2018; Tan et al., 2018), we hypothesize that the <700 yr incision event on the Beida River occurred at the peak of monsoon influence within the region.

In summary, combining all evidence from river incision rates, glacial coverage, and local and regional climate records, we suggest that the brief, rapid incision event recorded by the Beida River knickzone is most likely linked to a pluvial event that resulted from an increase of precipitation and possibly increased glacier melt during the mid-Holocene. The Qilian Shan and Hexi Corridor occupy the transitional zone between the Southeast Asian monsoon and the westerlies, with wet periods corresponding to increased monsoon influence (Tan et al., 2018). The combination of circumstantial evidence documented here suggests that a pluvial event occurred at the peak of monsoon influence within the westernmost North Qilian Shan. Between, we suggest the Beida River experienced the following incision and climate history since 24 kyr and 9.5 kyr B.P. After 24 kyr B.P., the Beida River drainage was under the dominant influence of the arid westerly moisture source, the river started to incise as it gradually depleted its sediment supply during the deglaciation and early interglacial period. Its incision rate (corrected for uplift) of  $1.0-9 \pm 0.8$  m/kyr was more or less in balance with the rock uplift rate, during this time. During the Early Holocene, the humid Southeast Asian monsoon expanded to the central North Qilian Shan, where it affected

the Hei He ~~main stem drainage~~ and filled Juyanze lake to ~~its highest a high~~ lake level. The expanded monsoon ~~may have mildly influenced the western North Qilian Shan, as it~~ correlates to the 1<sup>st</sup> incision stage of the Beida River when the canyon incised at a rate of  $5.6 \pm 2.2$  m/kyr. During the mid-Holocene, the ~~Asian~~ monsoon influence grew stronger, ~~starting in the Hei He drainage around 5.4 leading to 5.1 kyr B.P., and then expanding further to the western North Qilian Shan a few hundred years later regional pluvial period.~~ This peak of monsoon influence lasted less than ~~700±340~~680±460 yr in the Beida River drainage, which led to an increase of precipitation, and therefore an increase of water discharge and incision rate. This climate event led to deepening of canyons into the foreland basin, ~~prompting and prompted~~ the formation of bedrock knickzones in the Maying, Hongshuiba, and Beida River sub-basins of the Hei He. During this pluvial period, the incision rate of the Beida River increased to at least  $50 \pm 17$  m/kyr. Afterward, the discharge decreased to its present condition and river incision slowed to  $11.6 \pm 1.4$  m/kyr (Figure 89).




Figure 9 a) Schematic cross sections of Beida River channel evolution at the mountain front; b) diagram of incision rate vs. time since 24 kyr B.P.; c) climate reconstruction based on the Eastern ~~Juyan palaeo-lake~~~~Juvanze palaeolake~~ record (Herzschuh et al., 2004).

## 5.5 Glacial cycles and terrace formation in the North Qilian Shan

Though this study mainly focuses on the Holocene incision history of the Beida River, older fill terraces preserved within the North Qilian Shan suggest there are at least two other fill-cut cycles that predate the latest Late Pleistocene-Holocene cycle. Based on our prior fieldwork (Wang et al., 2020), there are two older generations correspond to fill terraces T2 and T3 (Figure 5a). T3 is undated, and only preserved in a few locations.  $^{10}\text{Be}$  and OSL dating of T2 terrace tread suggests an age of  $\sim 144$  kyr (Wang et al., 2020), which indicates the abandonment of T2 occurred during the penultimate glacial period, similar to that T1 was abandoned during the last glacial period. This suggests that the incision and deposition of fill terraces of Beida River correlates with the glacial-interglacial cycles. Though we hypothesize that knickzones present on the Hongshuibai River and Maying River also formed at the mountain front during a mid-Holocene pluvial period, the consistent locations of these knickzones 10-15 km upstream of the mountain front is somewhat surprising, considering the dissimilar sizes of these drainage basins (Table 1). One might expect that the knickzones on the Hongshuibai River and Maying River would have retreated less, as these drainage basins are smaller than that of the Beida River (Table 1). Such a comparison however does not account for the strong west to east precipitation gradient present within western North Qilian Shan, as well as the important orographic increase in precipitation. Drainage area thus becomes a poor proxy for discharge, especially when comparing across drainage basins. It is further worth noting that the Hongshuibai and Maying river canyons are deeper than that of the Beida River, suggesting an even fast rate of Holocene incision. Because knickzone retreat rate may be faster for steeper, narrower channels (Finnegan et al., 2005), the rate of abrupt base level drop may also affect the amount of retreat of these knickzones. Conversely, lesser streams draining across the mountain front and the Beida River tributaries do not exhibit evidence for deep, rapid incision. We attribute this difference to the aridity at lower elevation and that only those drainages that tap high elevations (over 4000 m) are supplied continuously during the summer by glacier melt. In summary, variations in discharge, due both to monsoon influence and orographic effects, strongly impacts the Holocene incision rate of major rivers draining the western North Qilian Shan.

~~A similar pattern occurs along the Hongshuibai River, the next major river to the east of the Beida River. The T1 terrace here is similar in age as T1 at the Beida River (26 kyr in Yang et al., 2020; 22.5 kyr in Hetzel et al., 2019), and the T4 terrace here has similar age as T2 terrace at the Beida River (153 kyr in Hetzel et al., 2019). In the eastern North Qilian Shan, terrace records of Shagou River also show a fill-cut pattern synchronized to glacial-interglacial cycles (Pan et al., 2003).~~

Therefore, we suggest the formation of fill terraces of different rivers along the North Qilian Shan is controlled by the same mechanisms, linked to glacial-interglacial cycles. In the Beida River, a full fill-cut cycle begins with the river backfilling its canyon with sediment during the glacial period. This is a period with elevated, glacially derived sediment flux, and perhaps lower discharge than at present. At the glacial-interglacial transition, the river transitions from deposition to erosion as sediment supply is gradually depleted. Fast incision commences during the early interglacial period as a result an increase of discharge from the introduction of monsoon-derived summer moisture.

## 6 Conclusions

660 The Beida River in the North Qilian Shan has incised deeply into both the bedrock and the adjacent foreland basin sediments. The incision rates indicated from terrace records and our models greatly exceed rates of tectonic uplift here. Our work demonstrates the capability of bedrock rivers in arid regions to incise deep channels and form fast retreating knickpoints within a short period. Field investigation and geomorphic mapping identify a 24 kyr fill terrace, T1, and several sets of inset terraces below. The longitudinal profile of the present river channel preserves a steep knickzone, located 10 to 12 km upstream of the  
665 mountain front. Terrace ages, and relationships between terrace treads and the riverbed, indicate that the knickzone was formed quickly after 54.7 kyr BP, likely driven by an increase of river discharge. By applying the concept of slope patches along with channel geometry and terrace records, we constrain that during the knickzone formation, the incision rate was at least  $50 \pm 17$  m/kyr and the duration of this period of increased discharge was less than  $700 \pm 340$   $680 \pm 460$  yr, which is about half of that estimated from the sparse terrace age record alone. The period of increased incision rate identified from the Beida River  
670 correlates to a pluvial period recorded at the terminal Juyanze lake. The likely cause of rapid incision of the Beida River, and adjacent rivers with similar deeply incised canyons, is the increased influence of the southeast Southeast Asian Monsoon over the Holocene, with the most rapid incision period corresponding to a peak of monsoon influence ca. 4 to 5 kyr B.P.

## Acknowledgments, Samples, and Data

675 This work was supported by the US National Science Foundation [grant number EAR-1524734] to Michael Oskin, the National Natural Science Foundation of China [grant number 41571001] to Youli Li, the Second Tibetan Plateau Scientific Expedition and Research (STEP) [grant number 2019QZKK0704] and the National Natural Science Foundation of China ([grant number 41622204; 41761144071]) to Huiping Zhang, and through Cordell Durrell Geology Field Fund to Yiran Wang. We placed our data in a permanent repository at Open Science Framework  
680 ([https://osf.io/bpvw9/?view\\_only=da77dd9c7ace4e76bf96dedc9fbaa61f](https://osf.io/bpvw9/?view_only=da77dd9c7ace4e76bf96dedc9fbaa61f)), and also upload as a zip file separately.

## References

- An, Z., Kutzbach, J. E., Prell, W. L. and Porter, S. C.: Evolution of Asian monsoons and phased uplift of the Himalaya - Tibetan plateau since Late Miocene times, *Nature*, 411(6833), 62–66, doi:10.1038/35075035, 2001.
- 685 Attal, M., Cowie, P. A., Whittaker, A. C., Hobbey, D., Tucker, G. E. and Roberts, G. P.: Testing fluvial erosion models using the transient response of bedrock rivers to tectonic forcing in the Apennines, Italy, *J. Geophys. Res. Earth Surf.*, 116(2), 1–17, doi:10.1029/2010JF001875, 2011.
- Berlin, M. M. and Anderson, R. S.: Modeling of knickpoint retreat on the Roan Plateau, western Colorado, *J. Geophys. Res. Earth Surf.*, 112(3), 1–16, doi:10.1029/2006JF000553, 2007.

- Bishop, P., Hoey, T. B., Jansen, J. D. and Lexartza Artza, I.: Knickpoint recession rate and catchment area: The case of uplifted  
690 rivers in Eastern Scotland, *Earth Surf. Process. Landforms*, 30(6), 767–778, doi:10.1002/esp.1191, 2005.
- Chen, C. T. A., Lan, H. C., Lou, J. Y. and Chen, Y. C.: The dry Holocene Megathermal in Inner Mongolia, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 193(2), 181–200, doi:10.1016/S0031-0182(03)00225-6, 2003.
- Chen, F., Bloemendal, J., Wang, J. M., Li, J. J. and Oldfield, F.: High-resolution multi-proxy climate records from Chinese  
695 Loess: Evidence for rapid climatic changes over the last 75 kyr, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 130(1–4), 323–  
335, doi:10.1016/S0031-0182(96)00149-6, 1997.
- Chen, F., Cheng, B., Zhao, Y., Zhu, Y. and Madsen, D. B.: Holocene environmental change inferred from a high-resolution  
pollen record, Lake Zhuyeze, arid China, *Holocene*, 16(5), 675–684, doi:10.1191/0959683606h1951rp, 2006.
- Chen, F., Wu, D., Chen, J., Zhou, A., Yu, J., Shen, J., Wang, S. and Huang, X.: Holocene moisture and East Asian summer  
700 monsoon evolution in the northeastern Tibetan Plateau recorded by Lake Qinghai and its environs: A review of conflicting  
proxies, *Quat. Sci. Rev.*, 154, 111–129, doi:10.1016/j.quascirev.2016.10.021, 2016.
- ~~Jiuquan History Compilation Committee: Jiuquan City Chorography, Lanzhou University Press, Lanzhou., 1998.~~
- ~~Chen, R., Han, C., Liu, J., Yang, Y., Liu, Z., Wang, L. and Kang, E.: Maximum precipitation altitude on the northern flank of  
the Qilian Mountains, northwest China, *Hydrol. Res.*, 49(5), 1696–1710, doi:10.2166/nh.2018.121, 2018.~~
- Crosby, B. T. and Whipple, K. X.: Knickpoint initiation and distribution within fluvial networks: 236 waterfalls in the Waipaoa  
705 River, North Island, New Zealand, *Geomorphology*, 82(1–2), 16–38, doi:10.1016/j.geomorph.2005.08.023, 2006.
- Ding, Y., Ye, B. and Liu, S.: Effect of Climatic Factors on Streamflow in the Alpine Catchment of the Qilian Mountains,  
*Acta Geol. Sin.*, 54(5), 431–437, 1999.
- Fang, X., Lü, L., Mason, J. A., Yang, S., An, Z., Li, J. and Zhilong, G.: Pedogenic response to millennial summer monsoon  
enhancements on the Tibetan Plateau, *Quat. Int.*, 106–107, 79–88, doi:10.1016/S1040-6182(02)00163-5, 2003.
- 710 Fang, X. M., Ono, Y., Fukusawa, H., Bao-Tian, P., Li, J. J., Dong-Hong, G., Oi, K., Tsukamoto, S., Torii, M. and Mishima,  
T.: Asian summer monsoon instability during the past 60,000 years: Magnetic susceptibility and pedogenic evidence from  
the western Chinese Loess Plateau, *Earth Planet. Sci. Lett.*, 168(3–4), 219–232, doi:10.1016/S0012-821X(99)00053-9,  
1999.
- Finnegan, N. J., Roe, G., Montgomery, D. R. and Hallet, B.: Controls on the channel width of rivers: Implications for modeling  
715 fluvial incision of bedrock, *Geology*, 33(3), 229–232, doi:10.1130/G21171.1, 2005.
- Finnegan, N. J., Sklar, L. S. and Fuller, T. K.: Interplay of sediment supply, river incision, and channel morphology revealed  
by the transient evolution of an experimental bedrock channel, *J. Geophys. Res. Earth Surf.*, 112(3),  
doi:10.1029/2006JF000569, 2007.
- ~~Gansu Geological Bureau: Regional Geology of Gansu Province, Geological Publishing House, Beijing. (in Chinese), 1989.~~
- 720 ~~Geng, H., Pan, B., Huang, B., Cao, B. and Gao, H.: The spatial distribution of precipitation and topography in the Qilian Shan  
Mountains, northeastern Tibetan Plateau, *Geomorphology*, 297, 43–54, doi:10.1016/j.geomorph.2017.08.050, 2017.~~

- Guo, W., Xu, J., Liu, S., Shangguan, D., Wu, L., Yao, X., [ZHAOZhao, J.](#), [LIULiu, Q.](#), [JANGJiang, Z.](#), [LHLi, P.](#), [WEIWei, J.](#), [BAOBao, W.](#), [YUYu, P.](#), [DINGDing, L.](#), [LHLi, G.](#), [GEGe, C.](#) and [WANGWang, Y.](#): The Second Glacier Inventory Dataset of China (Version 1.0), Cold Arid Reg. Sci. Data Cent. Lanzhou, doi:10.3972/glacier.001.2013.db, 2014.
- 725 Hartmann, K. and Wünnemann, B.: Hydrological changes and Holocene climate variations in NW China, inferred from lake sediments of Juyanze palaeolake by factor analyses, *Quat. Int.*, 194(1–2), 28–44, doi:10.1016/j.quaint.2007.06.037, 2009.
- Haviv, I., Enzel, Y., Whipple, K. X., Zilberman, E., Stone, J., Matmon, A. and Fifield, L. K.: Amplified erosion above waterfalls and oversteepened bedrock reaches, *J. Geophys. Res. Earth Surf.*, 111(4), 1–11, doi:10.1029/2006JF000461, 2006.
- 730 [He, P., Song, C., Wang, Y., Chen, L., Chang, P., Wang, Q. and Ren, B.: Cenozoic exhumation in the Qilian Shan, northeastern Tibetan Plateau: Evidence from detrital fission track thermochronology in the Jiuquan Basin. \*J. Geophys. Res. Solid Earth\*, 122\(8\), 6910–6927, doi:10.1002/2017JB014216, 2017.](#)
- [He, P., Song, C., Wang, Y., Wang, D., Chen, L., Meng, Q. and Fang, X.: Early Cenozoic activated deformation in the Qilian Shan, northeastern Tibetan Plateau: Insights from detrital apatite fission-track analysis. \*Basin Res.\*, 33\(3\), 1731–1748, doi:10.1111/bre.12533, 2021.](#)
- 735 Herzschuh, U., Tarasov, P., Wünnemann, B. and Hartmann, K.: Holocene vegetation and climate of the Alashan Plateau, NW China, reconstructed from pollen data, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 211(1–2), 1–17, doi:10.1016/j.palaeo.2004.04.001, 2004.
- [Herzschuh, U., Kürschner, H. and Mischke, S.: Temperature variability and vertical vegetation belt shifts during the last ~50,000 yr in the Qilian Mountains \(NE margin of the Tibetan Plateau, China\). \*Quat. Res.\*, 66\(1\), 133–146, doi:10.1016/j.yqres.2006.03.001, 2006.](#)
- 740 Hetzel, R., Hampel, A., Gebbeken, P., Xu, Q. and Gold, R. D.: A constant slip rate for the western Qilian Shan frontal thrust during the last 200 ka consistent with GPS-derived and geological shortening rates, *Earth Planet. Sci. Lett.*, 509, 100–113, doi:10.1016/j.epsl.2018.12.032, 2019.
- 745 Hu, G., Yi, C., Zhang, J., Liu, J., Jiang, T. and Qin, X.: Optically stimulated luminescence dating of a moraine and a terrace in Laohugou valley, western Qilian Shan, northeastern Tibet, *Quat. Int.*, 321, 37–49, doi:10.1016/j.quaint.2013.12.019, 2014.
- [Jiang, Q., Shen, J., Liu, X. and Zhang, E.: Holocene climate reconstructions of Ulungur Lake \(Xinjiang, China\) inferred from ostracod species assemblages and stable isotopes. \*Front. Earth Sci. China\*, 2\(1\), 31–40, doi:10.1007/s11707-008-0007-z, 2008.](#)
- 750 [Jolivet, M., Brunel, M., Seward, D., Xu, Z., Yang, J., Roger, F., Tapponnier, P., Malavieille, J., Arnaud, N. and Wu, C.: Mesozoic and Cenozoic tectonics of the northern edge of the Tibetan plateau: Fission-track constraints. \*Tectonophysics\*, 343\(1–2\), 111–134, doi:10.1016/S0040-1951\(01\)00196-2, 2001.](#)
- [Küster, Y., Hetzel, R., Krbetschek, M. and Tao, M.: Holocene loess sedimentation along the Qilian Shan \(China\): Significance for understanding the processes and timing of loess deposition. \*Quat. Sci. Rev.\*, 25\(1–2\), 114–125, doi:10.1016/j.quascirev.2005.03.003, 2006.](#)
- 755



- Lamb, M. P., Finnegan, N. J., Scheingross, J. S. and Sklar, L. S.: New insights into the mechanics of fluvial bedrock erosion through flume experiments and theory, *Geomorphology*, 244, 33–55, doi:10.1016/j.geomorph.2015.03.003, 2015.
- Lavé, J. and Avouac, J. P.: Active folding of fluvial terraces across the Siwaliks Hills, Himalayas of central Nepal, *J. Geophys. Res.*, 105(B3), 5735, doi:10.1029/1999JB900292, 2000.
- 760 Li, Y., Wang, N., Cheng, H., Long, H. and Zhao, Q.: Holocene environmental change in the marginal area of the Asian monsoon: A record from Zhuye Lake, NW China, *Boreas*, 38(2), 349–361, doi:10.1111/j.1502-3885.2008.00063.x, 2009.
- Meng, X., Zhang, S. and Zhang, Y.: The Temporal and Spatial Change of Temperature and Precipitation in Hexi Corridor in Recent 57 Years, *Acta Geol. Sin.*, 67(11), 1482–1492, 2012.
- Merritts, D. J., Vincent, K. R. and Wohl, E. E.: Long river profiles, tectonism, and eustasy: a guide to interpreting fluvial terraces, *J. Geophys. Res.*, 99(B7), 14031–14050, doi:10.1029/94jb00857, 1994.
- 765 Mischke, S., Fuchs, D., Riedel, F. and Schudack, M. E.: Mid to Late Holocene palaeoenvironment of Lake Eastern Juyanze (north-western China ) based on ostracods and stable isotopes, *Geobios*, 35, 99–110, 2002.
- Mischke, S., Demske, D., Wünnemann, B. and Schudack, M. E.: Groundwater discharge to a Gobi desert lake during Mid and Late Holocene dry periods, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 225(1–4), 157–172, doi:10.1016/j.palaeo.2004.10.022, 2005.
- 770 Owen, L. A., Spencer, J. Q., Haizhou, M., Barnard, P. L., Derbyshire, E., Finkel, R. C., Caffee, M. W. and Nian, Z. Y.: Timing of Late Quaternary glaciation along the southwestern slopes of the Qilian Shan, Tibet, *Boreas*, 32(2), 281–291, doi:10.1080/03009480310001632, 2003.
- Pan, B., Burbank, D. W., Wang, Y., Wu, G., Li, J. and Guan, Q.: A 900 k.y. record of strath terrace formation during glacial-interglacial transitions in northwest China, *Geology*, 31(11), 957–960, doi:10.1130/G19685.1, 2003.
- 775 Perron, J. T. and Royden, L.: An integral approach to bedrock river profile analysis, *Earth Surf. Process. Landforms*, 38(6), 570–576, doi:10.1002/esp.3302, 2013.
- Qiang, F., Zhang, M., Wang, S., Liu, Y., Ren, Z. and Zhu, X.: Estimation of areal precipitation in the Qilian Mountains based on a gridded dataset since 1961, *J. Geogr. Sci.*, 26(1), 59–69, doi:10.1007/s11442-016-1254-7, 2016.
- 780 Raup, B., Racoviteanu, A., Khalsa, S. J. S., Helm, C., Armstrong, R. and Arnaud, Y.: The GLIMS geospatial glacier database: A new tool for studying glacier change, *Glob. Planet. Change*, 56(1–2), 101–110, doi:10.1016/j.gloplacha.2006.07.018, 2007.
- Reimer, P., Bard, E., Bayliss, A., Beck, J., Blackwell, P., Ramsey, C. B., Grootes, P., Guilderson, T., Hafliðason, H., Hajdas, I., Hatté, C., Heaton, T., Hoffmann, D., Hogg, A., Hughen, K., Kaiser, K., Kromer, B., Manning, S., Niu, M., Reimer, R., Richards, D., Scott, E., Southon, J., Staff, R., Turney, C. and van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP, *Radiocarbon*, 55(4), 1869–1887, 2013.
- 785 Rhodes, T. E., Gasse, F., Lin, R., Fontes, J. C., Wei, K., Bertrand, P., Gibert, E., Mélières, F., Piotr, T., Wang, Z. and Cheng, Z. Y.: A Late Pleistocene-Holocene lacustrine record from Lake Manas, Zunggar (northern Xinjiang, western China), *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 120(1–2), 105–121, doi:10.1016/0031-0182(95)00037-2, 1996.

- 790 Royden, L. and Perron, J. T.: Solutions of the stream power equation and application to the evolution of river longitudinal profiles, *J. Geophys. Res. Earth Surf.*, 118(2), 497–518, doi:10.1002/jgrf.20031, 2013.
- Shean, D.: High Mountain Asia 8-meter DEM Mosaics Derived from Optical Imagery, Version 1, tile-224, tile-257, Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center, doi:https://doi.org/10.5067/KXOVQ9L172S2, 2017.
- 795 Shi, Y.: *New Understanding of Quaternary Glaciations in China*, Shanghai Popular Science Press, Shanghai (In Chinese)., 2011.
- Shi, Y., Cui, Z. and Su, Z.: *The Quaternary Glaciations and Environmental Variations in China*, Hebei Science and Technology Press, Shijiazhuang (In Chinese)., 2006.
- Sun, M., Liu, S., Yao, X., Guo, W. and Xu, J.: Glacier changes in the Qilian Mountains in the past half century: Based on the revised First and Second Chinese Glacier Inventory, *Acta Geogr. Sin.* (In Chinese with English Abstr., 70(9), 1402–1414, doi:10.11821/dlxb201509004, 2015.
- 800 Tan, L., Cai, Y., Cheng, H., Edwards, L. R., Gao, Y., Xu, H., Zhang, H. and An, Z.: Centennial- to decadal-scale monsoon precipitation variations in the upper Hanjiang River region, China over the past 6650 years, *Earth Planet. Sci. Lett.*, 482, 580–590, doi:10.1016/j.epsl.2017.11.044, 2018.
- 805 Tapponnier, P., Xu, Z., Roger, F., Meyer, B., Arnaud, N., Wittlinger, G. and Yang, J.: Oblique stepwise rise and growth of the tibet plateau, *Science* (80-. ), 294(5547), 1671–1677, doi:10.1126/science.105978, 2001.
- Tucker, G. E. and Whipple, K. X.: Topographic outcomes predicted by stream erosion models: Sensitivity analysis and intermodel comparison, *J. Geophys. Res. Solid Earth*, 107(B9), ETG 1-1-ETG 1-16, doi:10.1029/2001JB000162, 2002.
- Turowski, J. M., Lague, D. and Hovius, N.: Cover effect in bedrock abrasion: A new derivation and its implications for the modeling of bedrock channel morphology, *J. Geophys. Res. Earth Surf.*, 112(4), doi:10.1029/2006JF000697, 2007.
- 810 Wang, [W., Feng, Z., Ran, M. and Zhang, C.: Holocene climate and vegetation changes inferred from pollen records of Lake Aibi, northern Xinjiang, China: A potential contribution to understanding of Holocene climate pattern in East-central Asia, \*Quat. Int.\*, 311, 54–62, doi:10.1016/j.quaint.2013.07.034, 2013.](#)
- [Wang, Y., Oskin, M. E., Zhang, H., Li, Y., Hu, X. and Lei, J.: Deducing ~~Crustal Scale Reverse Fault Geometry and Slip Distribution From Folded River Terraces~~ crustal-scale reverse-fault geometry and slip distribution from folded river terraces, \*Qilian Shan, China, Tectonics\*, 39\(1\), 1–18, doi:10.1029/2019TC005901, 2020.](#)
- 815 Wei, K. and Gasse, F.: Oxygen isotopes in lacustrine carbonates of West China revisited: Implications for post glacial changes in summer monsoon circulation, *Quat. Sci. Rev.*, 18(12), 1315–1334, doi:10.1016/S0277-3791(98)00115-2, 1999.
- Whipple, K. X. and Tucker, G. E.: Dynamics of the stream-power river incision model: Implications for height limits of mountain ranges, landscape response timescales, and research needs, *J. Geophys. Res.*, 3(2), 221–222 [online] Available from: <http://dx.doi.org/10.1029/1999JB900120>; doi:10.1029/1999JB900120, 1999.
- 820 Whittaker, A. C.: How do landscapes record tectonics and climate?, *Lithosphere*, 4(2), 160–164, doi:10.1130/RF.L003.1, 2012.

- 825 Wobus, C., Whipple, K. X., Kirby, E., Snyder, N., Johnson, J., Spyropolou, K., Crosby, B. and Sheehan, D.: Tectonics from topography: Procedures, promise, and pitfalls, *Spec. Pap. 398 Tectonics, Clim. Landsc. Evol.*, 2398(October 2015), 55–74, doi:10.1130/2006.2398(04), 2006.
- Wohl, E. and David, G. C. L.: Consistency of scaling relations among bedrock and alluvial channels, *J. Geophys. Res. Earth Surf.*, 113(4), 1–16, doi:10.1029/2008JF000989, 2008.
- [Xiao, J., Nakamura, T., Lu, H. and Zhang, G.: Holocene climate changes over the desert/loess transition of north-central China, \*Earth Planet. Sci. Lett.\*, 197\(1–2\), 11–18, doi:10.1016/S0012-821X\(02\)00463-6, 2002.](#)
- 830 Yang, H., Yang, X., Huang, W., Li, A., Hu, Z., Huang, X. and Yang, H.: <sup>10</sup>Be and OSL dating of Pleistocene fluvial terraces along the Hongshuiba River: Constraints on tectonic and climatic drivers for fluvial downcutting across the NE Tibetan Plateau margin, China, *Geomorphology*, 348, doi:10.1016/j.geomorph.2019.106884, 2020.
- [Zhao, J.D., Zhou, S.Z., Cui, J.X., Pan, X.D., Xu, L.B., Zhang, X.W.: ESR chronology of Bailanghe Valley and new understanding of Qilian Shan Mountain's Quaternary glaciation. \*J. Mt. Sci.\* 19 \(6\), 481–488 \(In Chinese\), 2001.](#)
- 835 [Zheng, D., Clark, M. K., Zhang, P., Zheng, W. and Farley, K. A.: Erosion, fault initiation and topographic growth of the North Qilian Shan \(northern Tibetan Plateau\), \*Geosphere\*, 6\(6\), 937–941, doi:10.1130/GES00523.1, 2010.](#)
- Zhou, A., Sun, H., Chen, F., Zhao, Y., An, C., Dong, G., Wang, Z. and Chen, J.: High-resolution climate change in mid-late Holocene on Tianchi Lake, Liupan Mountain in the Loess Plateau in central China and its significance, *Chinese Sci. Bull.*, 55(20), 2118–2121, doi:10.1007/s11434-010-3226-0, 2010.
- 840 [Zhou, S., Li, J. and Zhang, S.: Quaternary glaciation of the Bailang River Valley, Qilian Shan, \*Quat. Int.\*, 97–98\(July 2015\), 103–110, doi:10.1016/S1040-6182\(02\)00055-1, 2002.](#)