

CLIMATE CHANGES AND THE FORMATION OF FLUVIAL TERRACES IN CENTRAL AMAZONIA INFERRED FROM LANDSCAPE EVOLUTION MODELLING

Ariel Henrique do Prado ^{1,2}, Renato Paes de Almeida ², Cristiano Padalino Galeazzi ^{2,4}, Victor Sacek ³ & Fritz Schlunegger ¹.

¹Institute of Geological Sciences, University of Bern, Bern, 3012, Switzerland.

²Instituto de Geociências, Universidade de São Paulo, São Paulo, Brazil.

³Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, São Paulo, Brazil.

⁴School of Earth and Environmental Sciences, Seoul National University, Seoul 21990, Republic of Korea.

Correspondence: Ariel Henrique do Prado (ariel.doprado@geo.unibe.ch)

Abstract. Climate changes have been considered an essential factor controlling the shaping of the recent alluvial landscapes in central Amazonia, with implications for explaining the biogeographic patterns in the region. This landscape is characterized by wide floodplains and various terrace levels at different elevations. A set of older terraces with ages between 50 and >200 kys occupy the higher portions of central Amazonia, whereas multiple terraces next to floodplains occur at lower elevations and display ages of a few thousands of years. These lower terraces, referred to as mid-lower terraces, reveal what can be perceived as a stochastic pattern both in space and time. Despite the widespread occurrence of these geomorphic features, no process-oriented analysis has been conducted to explain their formation. Here, we develop a landscape evolution model referred to as SPASE to explicitly account for fluvial erosion and deposition in combination with lateral channel migration to explore the controls on terrace development. The model results show that the higher terraces were deposited under the condition of a higher base-level for the basins upstream of the confluence between the Solimões and Negro Rivers. The subsequent decrease in the base level initiated a phase of gradual incision, thereby resulting in the current fluvial configuration. The model also predicts that high-frequency climate changes resulted in the construction of mid-lower terraces at various elevations which, however, are all situated at lower elevation than the higher terrace levels. Our model shows that dry-to-wet shifts in climate, in relation to the modern situation, yield a landscape architecture where mid-lower terrace levels are better preserved than wet-to-dry changes in climate, again if the current situation is considered as reference. Finally, our results show that fast and widespread landscape changes possibly occurred in response to high-frequency climate changes in central Amazonia, at least since the Late Pleistocene, with great implications for the distribution and connectivity of different biotic environments in the region. Because of this short time scale of response to external

perturbations, we suggest that the streams in Central Amazonia possibly also respond in rapid and sensitive ways to human perturbations.

1 Introduction

40 The lowlands of the central Amazonia host one of the largest systems of alluvial deposits on Earth, composed of extensive Quaternary fluvial terraces with wide incised valleys where modern rivers flow (Fig. 1) (Rossetti et al. 2005; Pupim et al. 2019; Sioli, 1984). These deposits are of great relevance for the reconstruction of both the evolution of the physical landscape and the patterns of biotic diversity in the region (Ribas et al. 2012; 45 2019; Pupim et al. 2019, Bicudo et al., 2019). The areas that currently display terraces were part of a large depositional system in which rivers in this region had high deposition and avulsion rates in an aggradational pattern (Hoorn et al., 2010; Pupim et al., 2019; Wilkinson et al., 2010). During this stage, flooded environments covered much larger areas, because they were not restricted to incised valleys as is currently the case. During the late Pleistocene, 50 most likely because of an adjustment of the river's long profile to a lowering of the base level or a change in discharge, the main rivers started to incise to reach their modern position. This phase of incision was associated with a widening of the valleys, which was accomplished through channel migration and lateral bank erosion (Merritts et al., 1994). This long-term period of fluvial erosion was accompanied, at a shorter time scale, by multiple cycles of 55 incision and aggradation, which resulted in the formation of both cut terraces and cut-and-fill sequences with various depositional ages and elevations (Pupim et al., 2019). Sea level changes have been proposed as the controlling mechanisms for the formation of these cycles (Irion & Kalliola, 2010), however, the depositional ages show that the cycles of aggradation and incision occurred at a different frequency or phase than the cycles of sea-level changes 60 during the Pleistocene (Pupim et al., 2019). In fact, there is ample evidence for the occurrence of Cretaceous bedrock exposed at the bottom of the Amazon River next to the confluence with the Negro River (Ianniruberto et al., 2018; Gualtieri et al., 2020), implying no net aggradation since the last glacial maximum. Therefore, the Holocene sea-level rise cannot be used to explain the occurrence of incised floodplains upstream of the confluence 65 with the Negro River. Climate driven discharge variations is another mechanism that has been used to explain the multiple phases of aggradation and erosion because such controls have the potential to impose changes on the course of the main and tributary rivers through changes in the sediment supply and the rivers' capacity to evacuate the supplied material (e.g., Tucker and Slingerland, 1999). There is indeed ample evidence that rainfall rates have


70 varied in central Amazonia during the last 200 kyrs (Cheng et al. 2013; Mertes and Dunne,
2007; Fritz et al., 2004). In addition, the pattern of terrace ages suggests that the related
cyclicality of sediment accumulation and erosion has been quite similar to that of rainfall
variations (Pupim et al., 2019). As yet, no quantitative model has been employed to test the
hypothesis of a climate driven control on the formation of the various terraces in central
75 Amazonia.

Here, we develop a landscape evolution model (LEM), which is tailored to
reproduce alluvial processes at a 10^4 yr temporal scale and for a spatial extent of several 10^6
 km^2 , which is comparable in size as the Sacek et al. (2014) model. This new model also
introduces a simplified approach to reproduce lateral erosion and the formation of cut
80 terraces. We start with a model that is calibrated with current field-based data from central
Amazonia including the relationships between water and sediment discharge, and the modern
topography along the Solimões River in an area between the confluence with the Iça and
Negro Rivers (Fig. 1). Using this calibrated model, we consider the landscape response to
different conditions of water discharge, sediment flux, and base level changes. We focus on
85 these parameters because of their high frequency variability, which is compatible with the
dynamics of the terrace formation in central Amazonia (Cheng et al. 2013; Mertes and
Dunne, 2007; Fritz et al., 2004 ; Pupim et al., 2019). We then compare the model results with
the modern topography. The ultimate scope is to quantitatively explore possible controls on
the formation of terrace levels in central Amazonia, which has occurred at a frequency of c.
90 10^4 years.

2 Local Setting

The study area (Fig. 1) is situated in central Amazonia to the west of the city of
Manaus, which is near the confluence between the Negro and the Solimões Rivers. The study
95 region, which is c. $300,000 \text{ km}^2$ large, additionally hosts 4 major tributary streams of the
Solimões River, which are from the West to the East: the Jutáí River, the Juruá River, the
Japurá River, and the Purus River. While the Solimões and the Japurá Rivers are sourced
from the Andean orogen, the Negro River is sourced from the Guiana shield, and the other
streams have their headwaters in the Fitzcarrald arch (Filizola and Guyot, 2009; Regard et al.,
100 2009). At Manaus downstream of the confluence with the Negro River, the mean water
discharge of the Amazon River is c. $130,000 \text{ m}^3/\text{s}$ and the mean annual flux of suspended
sediment amounts to c. $500 \times 10^6 \text{ t yr}^{-1}$ (Ianniruberto et al., 2018; Filizola and Guyot, 2011).
In the tributary rivers, mean water discharge values vary from c. $2000 \text{ m}^3/\text{s}$ for the Jutáí River

to approximately 30,000 m³/s for the Negro River. Except for the Jutaí and Japurá Rivers,
 105 gauging records of suspended sediment loads also exist for the other major streams in the
 study area. The values range from 25 x 10⁶ t yr⁻¹ for the Juruá River to c. 100 x 10⁶ t yr⁻¹ for
 the Purus River (Filizola and Guyot, 2009).



110 **Figure 1** Topography of central Amazonia with key geographical designations such as cities, rivers, and
 geological and geomorphic features including terrace levels (Lowlands), floodplains (Wetlands) and drainage
 network. The red square illustrates the location of the city of Manaus. Red dotted rectangle represents the region
 that is reproduced in the model and that represents the model grid. This illustration is based on a Digital
 Elevation Model from the Shuttle Radar Topography Mission.

115

The morphology of the study region is characterized by the occurrence of multiple
 terrace levels and floodplains. Conventionally, these are referred to as wetlands for the
 floodplains, and lowlands for the higher-elevated terraces (see Fig. 1 and Räsänen et al.,
 1990). Estimates of the elevation differences between the highest terrace level (e.g., green
 120 colored lowland surfaces in Fig. 1) and the floodplain (blue colored wetlands) range from c.
 10 to >30 m over a distance of hundreds of meters. This estimate is based on a SRTM digital
 elevation model, which, however, has not been corrected for the tree coverage. Note that this

could possibly add a bias in the modelling, which we address in the discussion section. In addition, several low-altitude terrace levels, referred to as mid-lower terraces, have been mapped in the study area inside the wetlands in Fig. 1 (Pupim et al., 2019). Estimates of the heights of the mid-lower terraces are associated with large uncertainties, but because they are at a lower elevation than the higher terraces their heights are less than 10 meters relative to the floodplain at flood stage. The floodplain widths vary between c. 15 km (Jutaí River) and 60 km (Solimões River). In addition, multiple local streams originate in the terrace levels and display a dendritic drainage pattern (Fig. 1).

Ages for some of these terrace levels have been established with optically luminescence dating techniques and measurements of ^{14}C concentrations in organic matter. As shown by Pupim et al. (2019), these analyses yielded ages between 50 and >200 kyrs for the highest terrace levels, and ages of <50 kyrs for the lower terrace levels adjacent to the floodplains. These results are summarized in Fig. 2 with OSL ages of the studied sedimentary profiles across fluvial terraces in central Amazonia and published ^{14}C and OSL ages from late Quaternary deposits from other Amazonian sites in Fig. 2 a. The ages are presented in groups (rows) related to river systems to highlight the regional variability of the sedimentary processes. Sample location, sedimentary and dating details, and references can be found in Pupim et al. (2019). The main domains are modern floodplains (green squares); mid-lower terraces (yellow squares); higher terraces (red squares). Subsurface sediments (gray squares) are remnants of older deposits at the base of a younger terrace. In Fig. 2 b there is speleothem ^{18}O records from Cueva del Diamante cave (purple line) indicating precipitation changes in western Amazonia (Cheng et al., 2013). The reported ages, the elevations and the spatial occurrences of the mid-lower terraces do not show a distinct pattern nor a clustering or a trend in any direction (Pupim et al., 2019). Although these ages reveal a large scatter and the sampling is still very sparse, we suggest that they record two distinct phases of landscape evolution. The first one is recorded by the highest terrace levels, which formed when the streams in the study area accumulated material over a larger area as revealed by the data of Pupim et al. (2019) in Fig. 2. This accumulation of fluvial deposits is superseded by a phase of erosional recycling. This second period started sometime during the late Pleistocene, and was associated with a shift towards a period with short-term (10^4 years) variations of climate (Cheng et al., 2013; Fritz et al., 2004; Mertes and Dunne, 2007). It has been proposed that this shift in paleoclimate eventually resulted in the formation of cut-and-fill terrace sequences and cut terraces in central Amazonia (see Fig. 4 in Pupim et al., 2019).

3 Methods

Following the scope of this paper, we developed a landscape evolution model which allows us to simulate fluvial processes of erosion, transport and deposition of sediments in alluvial landscapes (e.g., Tucker and Slingerland, 1999; Braun, 2006; Sacek, 2014). This model, which is referred to as SPASE (Sedimentary Processes and Alluvial Systems Evolution) is implemented in Python. The source code is openly accessible and can be downloaded from the GitHub repository (https://github.com/Ariel-H-Prado/SPASE_Model.git). In the context of the Late Pleistocene to Holocene terrace systems in central Amazonia, the search for the controls on the depositional and erosional cycles requires a conceptual model that considers short-term (10^4 years) variations of base level and climate changes (Cheng et al. 2013; Fritz et al., 2004; Mertes and Dunne, 2007), leading to the formation of cut-and-fill terrace sequences and cut terraces in central Amazonia. This model also needs to be able to simulate the erosion of the valley margins in response to the lateral migration of channels. In fact, such a process has been considered as essential to understand the formation and the widening of the in-valley floodplains in central Amazonia (Merritts et al., 1994).

3.1 Model setup and governing equations

To simulate fluvial erosion and mass transport, previously developed LEMs, such as CAESAR, SIBERIA and others (Beaumont et al., 1992; Braun, 2006; Hancock et al., 2010; Sacek, 2011; 2014), considered both fluvial processes, such as the long-distance advection of material by rivers and slope processes, such as local mass transport by hillslope diffusion for slopes flatter than a threshold, and advection for steep slopes (Tucker and Slingerland, 1999; Braun, 2006; Sacek, 2014). Since the Amazon alluvial system has very gentle slopes and because the area bordering the floodplains and terraces is occupied by a dense vegetation cover, we do not consider that hillslope processes largely contribute to the redistribution of sediment in the channel network inside the model grid. In fact, all material in the model domain is considered to result from either accumulation and/or erosion by fluvial and channelized processes only. However, the sediment supply from outside of the model grid could indeed be produced by hillslope processes in a steeper landscape such as e.g., the Andes, which is considered as a boundary condition (see section 3.3).




Figure 2 Chronology of sediment deposition representing the formation of terraces in Amazonian lowlands compared with Quaternary climate records from Pupim et al. (2019). (a) OSLand¹⁴C ages from late Quaternary deposits (b) Speleothem ¹⁸O records.

Upon modeling fluvial dynamics, we proceed following Beaumont et al. (1992) and employ a stream power law where, for a given channel, the carrying capacity of sediment, Q_{eqb} , which is equivalent to the maximum volume of sediment per unit time [$L^3 T^{-1}$] a stream can carry, is given by the equation:

$$Q_{eqb} = K_f Q_r^m S^n \quad (1)$$

Here, Q_r is the water discharge in the channel [$L^3 T^{-1}$], and S is the slope [$L L^{-1}$]. Note that in order to maintain the consistency of the equation, we set the dimension of the constant K_f to [$T L^{-3}$] ^{$m-1$} . This constant K_f is the dimensional transport coefficient that adjusts the carrying capacity (Q_{eqb}). The constants m and n determine whether the discharge and slope have a linear ($m = n = 1$) or a power ($m \neq 1$ and $n \neq 1$) relationship with Q_{eqb} . Here, we follow the experimental results and field observations of Whipple and Tucker (2002) and set

205 $m \geq 1$ and fixed $n = 1$. We then estimate the values for the parameters K_f and m (Table 1) through an ordinary least squares fitting on Eq. (1). This is accomplished using modern water and suspended sediment discharge data for multiple Amazonian rivers obtained from Filizola and Guyot (2009), and using slope values that we obtained from the SRTM digital elevation model. In this context, we infer that the sediment discharge in the current Amazonia rivers is
 210 at capacity. A table with the used modern values is presented in the Supplementary Material (Table S3).

Similar to Braun (2006), the entrainment, deposition and transport of sediment in a channel (in suspension or as bedload) is computed by Eq. (2), where the erosion/deposition rate is considered proportional to the difference between the carrying capacity (Q_{eqb}) and
 215 sediment discharge, Q_{sed} [$L^3 T^{-1}$]:

$$\frac{\delta h}{\delta t} = -\frac{1}{W L_f} (Q_{eqb} - Q_{sed}) \quad (2)$$

Here, W denotes the channel width [L] and L_f is the erosion/deposition length scale
 220 [L], which is a material property. From Eq. (2) it follows that if $Q_{sed} < Q_{eqb}$, sediment entrainment will occur. In this case, L_f takes the length-scale property of the river bed load, L_a . Alternatively, the model will simulate the occurrence of deposition if $Q_{sed} > Q_{eqb}$. In this case, L_f takes the length-scale property of the suspended load in the river, L_s , where $L_s < L_a$.

To simulate lateral erosion we infer that there is no beakrock confinement for the
 225 channels and that the eroded material is homogeneous in composition and strength. Therefore, the volume of sediment B [L^3] that a channel can laterally erode at a particular site is computed through:

$$\frac{\delta B}{\delta t} = A_{le} [(h_{terrace} - h_{channel}) + h_{depth}] \quad (3)$$

230

where $h_{terrace}$ is the height of the terrace adjacent to the channel [L], $h_{channel}$ is the height of the channel surface [L], h_{depth} is the depth of the channel [L]. A_{le} is the rate at which an area is laterally eroded per unit time [$L^2 T^{-1}$]. This rate is defined as:

$$235 \quad A_{le} = \alpha_1 Q_{eqb}^{\alpha_2} \quad (4)$$

Here, α_1 and α_2 are the lateral erosion coefficients, which depend on the erosional resistance of the lateral margins of the channels. These coefficients are defined through an iterative process where different values are assigned to the α_1 and α_2 parameters and where the impact on the model floodplains are then analyzed. We finally accepted those values for the α_1 and α_2 parameters where the simulated floodplain widths of the Solimões and Juruá Rivers correspond to the current situation and where the lateral migration of the model streams will not result in an obvious increase of the floodplains' widths. As another constraint, we inferred that this development towards the modern nearly stable condition had to occur within less than 20 kyrs, mainly because this corresponds to the timescales recorded by the terrace levels. An example of the model behavior with a different value for α_1 is shown in the Supplementary Material (Figure S9 and S10).

Because one model cell can potentially contain both a part of a channel belt-floodplain and a portion of a bordering terrace, it is necessary to record two elevation values upon modelling lateral erosion (Eq. (3) and (4)). One elevation variable, referred to as $h_{channel}$ in Eq. 3, represents the elevation of the active channel and the floodplain, and the other elevation parameter $h_{terrace}$ in Eq. (3) considers the elevation of a possibly older and higher terrace within the same cell. All maps in this paper illustrate the elevation of terraces ($h_{terrace}$) inside each cell. Additionally, the coefficient h_{depth} in Eq. (3) plays an important role upon modelling the lateral migration of the channels since it simulates the recycling of sediments that is associated with this process. In this context, accumulation of sediment (where $Q_{sed} > Q_{eqb}$) might raise the channel floor within a particular model cell. This can initiate a lateral shift of the active channel to a neighboring cell if its elevation is lower than the cell hosting the active channel. For a detailed description of the numerical solutions for the equations, their implementation in the model and the way of how the drainage network and the lakes are calculated the reader is referred to the Supplementary Material (S1).

3.2 Setup of the initial scenario

We use the modern topography as the initial condition. This surface was reconstructed over a regular mesh with 129 x 62 points, where each point corresponds to a center of a regular hexagonal cell. This input topography was obtained from the Shuttle Radar Topography Mission (SRTM) digital elevation model (DEM) for the area of interest (Fig. 3 a). The DEM was smoothed and adjusted for the grid of the model through the QGIS openly

accessible software. As mentioned before and addressed in the discussion, a limitation in the
 270 use of the SRTM is that the tree canopy heights are included in the elevation model.

Table 1 Fixed parameters in all simulated scenarios.

Parameter	Description	Value
k_f	Erosional coefficient due to fluvial processes	0.026
m	Stream power discharge exponent	1.425
n	Stream power slope exponent	1.0
L_a	Erosion length scale of the sediments	10.0 km
α_1	Lateral erosion coefficient 1	15.73
α_2	Lateral erosion coefficient 2	0.3
	Rainfall rate - Current	3000.0 mm
	Runoff coefficient	50 %
	Hexagonal cell side	4 km


The initial landscape displays six main rivers entering the grid. These are the
 275 Solimões, Japurá, Juruá, Jutaí, Purus and Negro Rivers (Fig. 1 and 3b). For these streams,
 water flux values were obtained by the HydroRivers model for the locations where they enter
 the study area (Lehner and Grill, 2013). Also for these locations, the sediment discharge of
 these streams was then estimated by applying Eq. (1), thereby using the calibrated parameters
 of K_f and m and the slopes S that we measured along the corresponding reach (Table 2).
 280 Modern average rainfall rate in the region is 3000 mm/year (Espinoza et al., 2009), and the
 runoff coefficient (Effective proportion of rainfall) is set to 50% (Guimberteau et al. 2013).
 For simplification, all erosional and transport processes occur over the same material, here
 considered as sand. The bedrock at Manaus is considered as a rigid material that cannot be
 eroded. Therefore the elevation at Manaus is defined as the minimum base level in the model,
 285 locked at 0 m in the model (Fig. 3).

In order to make the initial topography more stable upon modelling, the numerical
 model was run using the smoothed DEM as initial topography with the model parameters of
 Table 1 and 2 during 10 kyr inside the model. The output topography for this scenario (Fig.
 3b) was then used as initial topography for all the climatic and base-level tests in this work.

290

Table 2 Current discharge, slope and sediment flux of the rivers outside of the grid. The sediment flux was calculated using the Eq. (1) and the parameters on Table 1.

River	Discharge ($\text{m}^3 \text{s}^{-1}$)	Slope (m m^{-1})	Sediment flux ($\text{m}^3 \text{s}^{-1}$)
Solimões	55522.754	0.000057	8.624734014
Juruá	4970.593	0.00015	0.7285283026
Japurá	14555.073	0.0001	2.245332418
Jutaí	2301.853	0.00014	0.2270145616
Purus	6650.166	0.00012	0.8824582715
Negro	29187.611	0.000022	1.331445249



295

Figure 3 (a) Shuttle Radar Topography Mission (SRTM) digital elevation model of central Amazonia, with a resolution of 90 m at the Equator. (b) Initial topography and configuration of the numerical simulations. The blue lines represent the channels and the thickness of the line is proportional to the discharge of the corresponding stream.

300 3.3 Boundary conditions for the modelled scenarios

For the model scenarios where different climate conditions are taken into account, it is necessary to consider the variation of the rainfall rate outside of the model grid, which induces a change of the water discharge of the six main rivers entering the model grid and consequently impacts the sediment flux. We used the following relationship in order to estimate the sediment flux, Q_{sed} . For the rivers entering the grid, sediment discharge Q_{sed} for a given water discharge Q_r is computed through:

$$Q_{sed} = \alpha \cdot Q_r^\gamma \quad (5)$$

310 The value for γ is obtained through a regression of Eq. (5) using data from Filizola and Guyot (2009) about water discharge and sediment flux of the main rivers in the Amazonian lowlands as constraints. This regression yields a value of c. 0.5 for the exponent γ on Eq. (5) (see also Figure S7 on Supplementary Material). The parameter α varies for each river and is defined as:

$$315 \quad \alpha = Q_{sed_current} \cdot Q_{r_current}^{-\gamma} \quad (6)$$

in which $Q_{sed_current}$ and $Q_{r_current}$ are the current sediment and water flux of the rivers listed in Table 2 (Figure S8 on Supplementary Material shows this relation for the six main rivers). We note that this relationship between sediment and water flux, scaled through γ based on modern observations, is considered as a constant value during all model runs. Accordingly, we infer that sediment discharge in the major Amazonian rivers occurs at near capacity conditions, which we consider as a valid assumption.

325 Additionally, all cells on the right edge of the grid are considered as sink cells. They also control the elevation of the base-level of the grid. For the generation of the initial topography, the sink cells are kept at a fixed position. This corresponds to the elevation of the bedrock at Manaus, which we set to 0 m (see above).

4 Results

330 The controls of climatic variations on the formation of terraces are explored in the framework of two scenarios, the results of which are detailed in the following sections. In scenario one (Fig. 4), a dryer period is simulated by a 30% decrease of the current water input during the first 10 kyr, which is then followed, until 20 kyr, by an instantaneous increase in

the water supply up to the same level as the current water input. Scenario two is similar but considers first a 30% increase of the water input during the first 10 kyr (Fig. 5), thereby
335 simulating a wetter period, which is superseded by a 10 kyr-long model run during which the
water supply is instantaneously reduced to the current level. The 30% increase or decrease in
rainfall on central Amazonia during the Holocene is based on estimatives proposed by van
Breukelen et al. (2008) and Cheng et al. (2013). In both scenarios, after shifting to a drier or
wetter period, the model returns to the modern climate setting to evaluate what would be
340 preserved in the landscape from such climatic shifts. In these models, a change in
precipitation rates will affect the water input into the model, which is the sum of the water
flux where the stream enters the grid and the runoff that is generated through rain on the grid
itself. Both scenarios start with the same initial topography (Fig. 3b), and the grid base-level
is locked at a bedrock elevation of 0 m, thereby considering the occurrence of a bedrock in
345 the Amazon trunk stream at the downstream end of our model. The consequence of such a
fixed base level is that sediment does not get eroded or deposited after the confluence of the
Solimões and Negro Rivers although that the high water supply of the Negro River lowers the
sediment to water discharge ratio, which, in turn, would acutely promote a phase of erosion.
Following these model runs, we also explore a third scenario where we allow the base level
350 first to increase and then to decrease to the modern situation.

4.1 Scenario one: Shift to drier and then wetter conditions

The model results of scenario one, which commences with a dry period, show that
sediment starts to accumulate in the upstream part, and the wave of accumulation progresses
355 from upstream to downstream. As a consequence, the average slope of the Solimões River
and its main tributaries increases in pace with the accumulation of sediment. This change
towards steeper channel gradients occurs during the first 5 kyr, after which a nearly steady
state situation is established with constant topographic gradients. At this stage, a total of
approximately 13 meters of sediment has accumulated along the Solimões River upstream of
360 the Juruá River confluence (Fig. 4, at 2, 5 and 10 kyr). Also during the first 10 kyr with drier
conditions, the course of the Solimões trunk river is deflected to the North at the confluence
with the Purus River. The Purus River flows from the south and experiences, similar to the
other main tributaries in the model, a phase of sediment accumulation in the upstream part. In
the confluence area, this northward-directed shift of the Solimões River initiates a wave of
365 erosion along the northern banks of this stream (Fig. 4). The model simulates the occurrence
of lateral bank erosion also along those river segments where sediment accumulation occurs

(Fig. 4). This is mainly caused by the meandering of the channel's course to the lateral floodplain boundaries, which then initiates a phase of lateral erosion as long as higher terrace levels form cut banks. These results document that the SPASE model reproduces one of the first order features of a meandering stream such as cut banks despite the simple model architecture.

During the subsequent 10 kyr-long period of scenario one, the modelled increase in water input causes a wave of erosional recycling of the previously deposited sediments. An example can be seen in Fig. 4 where new terrace levels are formed e.g., around the Japura River (see location A in Figure). This phase of downcutting is associated with a phase of valley widening mainly along the main rivers, thus forming cut-terraces (Fig. 4, at 15 and 20 kyr). At the end of the model run, just a fraction of the previously deposited sedimentary material is preserved (red cells on Fig. 4 at 20 kyr). The model results predict that such terrace fragments mainly occur in the upstream part of the Solimões River and next to the Japurá, Juruá, Jutaí and Purus Rivers.

4.2 Scenario two: Shift to wetter and then drier conditions

In model run two, which corresponds to a scenario where water supply is increased during the first 10 kyr and then decreased to modern conditions, the landscape response is different. In particular, the increase in water runoff initiates a wave of erosion from upstream to downstream, which is also associated with a widening of the floodplain through lateral bank erosion. Erosion and valley lowering results in a decrease of the energy gradient, and in the formation of new floodplains at a lower elevation (Fig. 5, at 2, 5 and 10 kyr) of c. 10 meters. During the last 10 kyr when the model water input is the same as the modern one, the model rivers adapt an aggradational pattern, but the rivers' thalwegs do not reach the same initial elevations. At the end, no new terrace levels are formed (Fig. 5, at 15 and 20 kyr). Finally, during the evolution of this model scenario, the main rivers do not show any preferential direction of lateral bank erosion. Similar to the previous model run, the topography response to each climate change occurs mainly in the first 5 kyr, after which nearly steady state conditions are reached. Note that in comparison to scenario 1 (Fig. 4), the changes in the landscape are much less (Fig. 5).

Scenario 1: - 30 % Rainfall until 10 kyr





Figure 4 Results of the landscape evolution of scenario 1. The left figures show the difference of the model topography in relation to the initial topography, with colors saturated in red for values in elevation differences higher than 10 m, and in blue for values below - 10 m. The right column shows the model topography itself. The occasional blue stains on the map represent lakes (local minima). Location A indicates new terrace levels being formed around the Japura River.

Scenario 2: + 30 % Rainfall until 10 kyr



405

Figure 5 Results of the landscape evolution of scenario 2. The left figures show the difference of the model topography in relation to the initial topography, with colors saturated in red for values in elevation differences higher than 10 m, and in blue for values below - 10 m. The right column shows the topography itself. The occasional blue stains on the map represent lakes (local minima).

410 4.3 Scenario three: Topography response due to base-level variations

To simulate the consequences of a hypothetical base-level variation on the topography of Central Amazonia, we consider a scenario where the base level gradually increases until 35 m during the first 10 kyr, which is followed by a rapid base level drop to the elevation of the bedrock at 0 m (Fig. 6). The water input is equal to the current water input during the
415 entire scenario.


During the first 10 kyr, the main rivers turn to an aggradational pattern and thus adjust their long-stream profile to the increasing base level, even accumulating new sedimentary material above older terraces near the interfluvial area of the Solimões and Negro Rivers (Red cells on Fig. 6 at 2, 5 and 10 kyr). The smaller tributaries, with a lower sediment discharge,
420 are not able to accumulate sediment in response to the base-level rise at the same pace as the main rivers, thus forming shallow lakes within their own valleys (inundated cells on Fig. 6 at 2, 5 and 10 kyr). During the second period of the model run the main rivers initiate a phase of erosion, which occurs in response to the drop in the base level. This downwearing is associated with a period of valley widening. In addition, the model predicts that streams can
425 adapt to a new course. This is exemplified by a northward shift of the Solimões River that takes the course of the Negro River towards the end of the model grid (Fig. 6 at 15 and 20 kyr). Such shifts in the course of the major streams is due to the random position of the avulsive rivers during the aggradational stage and the preservation of their last configuration during the incision stage.

430

5 Discussion

The numerical model (SPASE) presented in this work simulates the evolution of alluvial landscapes. It is capable of reproducing the formation of floodplains and terraces at conditions similar to central Amazonia. The tested scenarios allow us to quantitatively
435 explore the responses of the central Amazonia topography to climatic and base-level changes.

Scenario 3: - Base level variation



440 **Figure 6** Model results showing the landscape evolution of scenario 3. The model water input is constant and equal to the current water input. The left figures show the difference of the model topography in relation to the initial topography, with colors saturated in red for values in elevation differences higher than 10 m, and in blue for values below - 10 m. The right column shows the topography itself. The occasional blue stains on the map represent lakes (local minima).

5.1 Model behavior

445 On the floodplains, the occurrence of erosion or sediment accumulation in the modelled scenarios are inherently related to the difference between the sediment transport capacity, which depends on water discharge and slope (Eq. (1) in the *Methods*), and the sediment discharge (Eq. (2) in the *Methods*). Accordingly, as we start our calculations with the conditions where sediment discharge in the model streams is at capacity, then any
450 changes in either water or sediment discharge will initiate a phase of channel steepening or flattening in order to re-establish the at-capacity conditions. Accordingly, by applying Eq. 5 we infer that the rivers in our model are at capacity upstream the model grid during the model runs. Since most of these rivers have their sediment sources either in the high Andes or in the elevated plateaus with a generally thick regolith cover (an exception is the Rio Negro), we
455 anticipate that sufficient sediment is available to be eroded by overland flow erosion so that sediment flux of the major streams is at capacity. Accordingly, if we maintain sediment discharge at a constant value and only change water supply, then an increase or a decrease in water discharge will respond in a flatter or steeper channel gradient, respectively, in order to reach a condition where sediment discharge equals the sediment transport capacity. In
460 addition, because data on modern sediment and water discharge in the Amazonian streams (Filizola and Guyot, 2009) suggests that sediment discharge relates to water discharge with an exponential factor of $\gamma \sim 0.5$ (Eq. (5) in the *Methods*, Figure S7 Supplementary Material), then changes in the sediment transport capacity are less than shifts in water discharge. Therefore, provided that sediment discharge in the current Amazonian rivers is at capacity, as
465 has been inferred upon setting up the initial conditions (Fig. 3), then a 30% drop in water supply will decrease the sediment transport capacity by c. 16%. In this case and according to Eq. (1) the channel and thus the entire floodplain will steepen, depending on the combined effect of the deposition rate of sediment (Eq. (2)) and the width of the floodplain. Sediment accumulation will then proceed from upstream to downstream because the modelled change
470 in the water budget is considered as downstream directed forcing. In the same sense, a 30% increase in water discharge will increase the sediment supply by the same amount, i.e. 16%. Because an increase in water runoff shifts the model streams to a condition where sediment flux is at under capacity, the model rivers will start to entrain sediment, thereby flattening the valley floor, as revealed by the model results (Fig. 5). Because in either scenario, erosion will
475 continue on the higher elevated terraces adjacent to the trunk streams, the sediment contribution in the tributary streams will first be high and then decrease with time. As a consequence, the elevation of the main valley floor will not be the same as the initial situation

after several model cycles (Fig. 5). The model response to changes in water discharge and precipitation thus ultimately reflects the effect of the sediment rating curve, which yields a
480 non-linear relationship ($\gamma = 0.5$) between water and sediment discharge.

The model shows that terrace levels form not only where the sediment transport capacity outpaces the sediment discharge but also where the meanders' cut banks approach the lateral margin of a floodplain. A terrace level might thus even form during a period when sediment accumulation occurs, particularly where the model streams start to erode the lateral
485 banks of a floodplain (Fig. 4). The combination of cut bank erosion and sediment accumulation and/or erosion on the floodplains result in a pattern where the terrace levels appear as randomly distributed within the Amazonian landscape, and at various elevations. As outlined in the following section, the model results thus offer an explanation for why published ages and elevation of terrace levels record a pattern which is rather difficult to
490 correlate with specific changes in external forcings.

We also conduct model runs where we change the initial precipitation conditions (see Fig. S11 to S19 in Supplementary File). The model predicts that the erosion of the elevated plateaus adjacent to the trunk rivers becomes larger as we increase the precipitation rates (1000 mm to 3000 mm as initial condition). However, the patterns at which the low-elevated
495 terraces form will not change. Therefore, a change of the initial precipitation rates will not alter the main conclusions.

5.2 Spatial distribution of mid-lower terrace levels

The model outputs predict what appears to be a random distribution of mid-lower
500 terrace levels, particularly if the dry-to-wet climate change scenario is considered. This pattern is caused by lateral shifts of the main rivers during both the aggradation and incision phases. These lateral shifts occur where the sediment flux exceeds the sediment transport capacity, which forces the streams to accumulate a fraction of the transported sediment. The models show that such conditions occur randomly along most of the main streams.
505 Exceptions are observed at the confluence of streams where the tributary stream is in an aggradational state. Such aggradation locally increases the elevation in the tributary floodplain relative to that of the trunk stream, with the consequence that it reduces the possibility and thus the probability of the trunk stream to laterally shift towards the tributary river. In this sense, the trunk river is pushed towards the opposite floodplain margin where
510 erosion can form a cut terrace.

The model results also show that climatic variations that occur at the scale of thousands of years in central Amazonia (Cheng et al., 2013; Fritz et al., 2004; Mertes and Dunne, 2007) can have a large influence on the cycles of sediment aggradation and incision. The main differences in the topographic responses between the two climatic conditions are
515 mainly observed within the floodplains, because the channels steepen during dryer periods in response to sediment accumulation, and they flatten during wetter periods as sediment is entrained in the floodplains. Both scenarios result in the formation of terrace levels at different locations and elevations. Whereas the drier-to-wetter climate change scenario results in the formation of new mid-lower terrace levels that have quite a large preservation potential
520 and that could eventually be mapped over a larger area, the wetter-to-drier climate change scenario returns model results where new terrace levels are poorly preserved. Following these model results, we thus propose that the mid-lower terrace systems in Central Amazonia mainly reflect the response to climate cycles from drier to wetter in comparison to modern conditions. These results could offer a solution of why published terrace ages neither show at
525 distinct clustering of ages nor a particular spatial pattern. We explain this by the short response times in combination with the high frequency of climate change and the rather stochastic pattern at which channels migrate laterally.

It is important to note that the initial topography that we used for our model was based on the SRTM digital elevation model (DEM) where the elevation of trees was maintained
530 upon calculating the DEM. This introduces a bias in the sense that the difference in elevations between the terraces and the floodplains are larger in the DEM than they actually should be. This implies that the spatial extents where sediment accumulations during the drier periods could be even larger than our model suggests. This implies that our modeling results can be considered as conservative scenarios of landscape change.

535

5.3 Response of low-gradient landscapes to climate changes and contrasts to mountainous areas

The model results show that climatic variations that occur at the scale of thousands of years in central Amazonia (Cheng et al., 2013; Fritz et al., 2004; Mertes and Dunne, 2007)
540 can have a large influence on the cycles of sediment aggradation and incision, and that the mid-lower terrace systems in Central Amazonia mainly reflect the response to climate cycles from drier to wetter in comparison to modern conditions. This is different to the cyclicity recorded by terrace systems e.g., in the Andean mountain belt valleys where terrace levels have been related to wetter-to-drier climate cycles, also relative to the modern climate (e.g.,

545 Steffen et al., 2009; Veit et al., 2016). The main difference between these environments is
that in mountain belts, hillslopes have been considered as an additional landscape component
where sediment can be stored during dry periods, whereas the material gets mobilized and
supplied to the drainage network during wet climates, resulting in a temporary accumulation
of material in the valley floor. During dry conditions, the hillslopes are considered as stable,
550 which allows the trunk streams to recycle the previously deposited sediment in the valley
floor, thereby forming a terrace level. Such a mechanism at work in mountainous areas has
also been reproduced with numerical models (e.g., Tucker and Slingerland, 1999; Norton et
al., 2016). This is, however, different to our case where the buffering effect of hillslopes is
missing, since the central Amazonian landscape is mostly flat and far from the Andean
555 mountain belt. Therefore, we mainly see the channel's responses to climate change, which
thus have an aggradation-incision cyclicity and which thus contrasts to that in mountain
valleys.

5.4 Controls of base level changes on the formation of the highest terrace levels

560 The scenario where we model the landscape response to an increase and then a
decrease in the local base-level (Fig. 6) provides an explanation why the topography of
central Amazonia has preserved terrace levels at a high elevation (Fig. 5 at 10 kyr). Such a
scenario has the potential to form wide floodplains at higher elevations. As the base-level
drops, the rivers start to recycle previously deposited material. In addition, the rivers can take
565 new courses. The model results thus predict that major changes in the channel network can
preferentially be initiated following a phase of major sediment accumulation, possibly
controlled by a higher base level at the downstream end of the model. Published ages of the
higher terraces suggest that these geomorphic features are also made up of multiple terrace
sequences with a stochastic pattern of ages and elevations, which is a feature that is similar to
570 the mid-lower terrace levels but at a higher elevation (Fig. 2). Accordingly, using the model
results of scenarios 1 and 2, we suggest that the formation of the higher terrace levels,
possibly conditioned by a higher local base level, was superimposed by high-frequency
climate changes. Accordingly, it is very likely that at such a stage of a higher base level,
climate cycles would also cause the formation of cut-and-fill terrace sequences with a similar
575 stochastic distribution of ages as those of the mid-lower terraces. This suggests that we can
expect a superposition of low-level, high-frequency terraces on terrace deposits occurring at
high levels, yielding a complex pattern in the landscape architecture.

6 Conclusions

580 According to the scenarios described herein, the current configuration of the terraces and floodplains in central Amazonia could be explained by aggradational and incisional processes controlled by high-frequency climatic variations and local base level changes, which however occurred at a lower frequency. In particular, the higher terraces were deposited in a condition of a higher base-level for the basins upstream of the confluence
585 between the Solimões and Negro Rivers. The subsequent decrease in the base level initiated a phase of gradual incision, thereby resulting in the current fluvial architecture. The high-frequency climate changes then yielded in the construction of mid-low terraces at various elevations, which however, are all situated at a lower elevation than the higher terrace levels. Our model shows that dry-to-wet shifts in climate, in relation to the modern situation, returns
590 a landscape architecture where mid-lower terrace levels are better preserved than wet-to-dry changes in climate, again if the current situation is considered as reference.

Our results also show that fast and widespread landscape changes possibly occurred in response to high-frequency climate changes in central Amazonia, at least since the Late Pleistocene, with great implications for the distribution and the connectivity of different
595 biotic environments in the region. Because of this short time scale of response to external perturbations, we suggest that the Central Amazonian streams in the study area, and also at a broader scale, possibly respond in a rapid and sensitive way to human perturbations. This could be, for instance, the ongoing deforestation with implications for sediment flux and hydrology, and particularly the planned construction of multiple hydropower dams
600 (Latrubesse et al., 2017; Best, 2018) with major implications on water discharge and local base levels as we have modeled in our contribution.

7 Code availability

The open-source landscape evolution model SPASE is freely available at
605 https://github.com/Ariel-H-Prado/SPASE_Model.git.

8 Author contribution

RPA designed the study, together with AHP. AHP developed the model with support by VS and RPA. CPG prepared the geological framework and synthesized data on water
610 discharge, channel widths and sediment flux. AHP analyzed the model results together with RPA, CPG, VS and FS. AHP wrote the manuscript with support from FS and RPA. All authors approved the current version.

9 Competing Interests

615 The author declares that there is no conflict of interest.

10 Acknowledgements

The authors were funded by the São Paulo Research Foundation (FAPESP) through research grants 2016/03091-5 and 2018/23899-2, and FAPESP student scholarships
620 2017/06874-3 and 2018/02197-0, as well as by the National Research Foundation of Korea (NRF) through research grant 2019R1A6A1A10073437. Part of the contribution of do Prado has been supported by various sources from the Institute of Geological Sciences of the University of Bern, and by the Innovative Training Network S2S project 860383. We also thank CNPq (#305218/2009-3 and #426654/2018-8), The Royal Society - Newton Advanced
625 Fellowship (NAF\R2\192188), and CAPES-PPGG (demanda social) for research support and student scholarships. SRTM data provided by <http://srtm.csi.cgiar.org>.

12 References

- Beaumont, C., Fullsack and P. Hamilton, J.: Erosional control of active compressional
630 orogens, in : Thrust Tectonics: London, edited by: McClay, K.R., Springer-Science+Business Media, B.V., 1–18., <https://doi:10.1007/978-94-011-3066-0>, 1992.
- Best, J.: Anthropogenic stresses on the world's big rivers, *Nature Geoscience* 12, 7–21, <https://doi.org/10.1038/s41561-018-0262-x>, 2019.
- Bicudo, T. C., Sacek, V., Almeida, R. P., Bates, J. M. and Ribas, C. C.: Andean Tectonics
635 and Mantle Dynamics as a Pervasive Influence on Amazonian Ecosystem, *Sci. Rep.* 9, 16879, <https://doi.org/10.1038/s41598-019-53465-y>, 2019.
- Braun, J.: Recent advances and current problems in modelling surface processes and their interaction with crustal deformation, *Geological Society of London Special Publications*, 253, 307–325, <https://doi:10.1144/GSL.SP.2006.253.01.16>, 2006.
- 640 Cheng, H., Sinha, A., Cruz, F.W., Wang, X.F., Edwards, R.L., d'Horta, F.M., Ribas, C.C., Vuille, M., Stott, L.D. and Auler, A.S.: Climate change patterns in Amazonia and biodiversity, *Nat. Commun.* 4, 1411, <https://doi.org/10.1038/ncomms2415>, 2013.

- Espinoza J.C., Ronchail J., Guyot J.L., Cochonneau G., Naziano F., Lavado W., de Oliveira E., Pombosa R. and Vauchel P.: Spatio-temporal rain-fall variability in the Amazon basin countries (Brazil, Peru, Bolivia, Colombia, and Ecuador), *Int. J. Climatol.* 29(11): 1574–1594, <https://doi.org/10.1002/joc.1791>, 2009.
- 645 Filizola, N. and Guyot, J. L.: Suspended sediment yields in the Amazon basin: an assessment using the Brazilian national data set, *Hydrol. Processes*, 23, 3207–3215, <https://doi.org/10.1002/hyp.7394>, 2009.
- 650 Filizola, N. and Guyot, J. L.: Fluxo de sedimentos em suspensão nos rios da Amazônia, *Rev. Bras. Geosci.*, 41 (4), 566–576, <https://doi.org/10.25249/0375-7536.2011414566576>, 2011.
- Fritz, S. C., Baker, P. A., Lowenstein, T. K., Seltzer, G. O., Rigsby, C. A., Dwyer, G., Tapia, P. M., Arnold, K. K., Ku, T. and Luo, S.: Hydrologic variation during the last 170,000 years in the Southern Hemisphere tropics of South America. *Quat. Res.* 61:95–104, <https://doi.org/10.1016/j.yqres.2003.08.007>, 2004.
- 655 Gualtieri, C., Martone, I., Filizola Junior, N.P. and Ianniruberto, M.: Bedform Morphology in the Area of the Confluence of the Negro and Solimões-Amazon Rivers, Brazil, *Water*, 12, 1630, <https://doi.org/10.3390/w12061630>, 2020.
- 660 Guimberteau, M., Ronchail, J., Espinoza, J.C., Lengaigne, M., Sultan, B., Polcher, J., Drapeau, G., Guyot, J.L., Ducharne, A. and Cialis, P.: Future changes in precipitation and impacts on extreme streamflow over Amazonian sub-basins, *Environmental Research Letters*, 8, 1-13, <https://doi:10.1088/1748-9326/8/1/014035>, 2013.
- Hancock, G. R., Lowry, J.B.C., Coulthard, T.J., Evans, K.G. and Moliere, D.R.: A catchment scale evaluation of the SIBERIA and CAESAR landscape evolution models, *Earth Surface Processes and Landforms*, 35(8), 863-875, <https://doi.org/10.1002/esp.1863>, 2010.
- 665 Hoorn, C. , Wesselingh, F. P., Ter Steege, H., Bermudez, M. A., Mora, A. , Sevink, J., Sanmartín, I., Sanchez-meseguerA., Anderson, C. L., Figueiredo, J. P., Jaramillo, C., Riff, D., Negri, F. R., Hooghiemstra, H., Lundberg, J., Stadler, T., Särkinen, T. and Antonelli, A.: Amazonia Through Time: Andean uplift, climate change, landscape evolution, and biodiversity, *Science*, 330, 927–931, <https://doi:10.1126/science.1194585>, 2010.

- Ianniruberto, M., Trevethan, M., Pinheiro, A., Andrade, J. F., Dantas, E., Filizola, N., Santos, A. and Gualtieri, C.: A field study of the confluence between Negro and Solimões Rivers. Part 2: bed morphology and stratigraphy, *Compt. Rendus Geosci.* 350 (1), 43–54, <https://doi.org/10.1016/j.crte.2017.10.005>, 2018.
- 675
- Irion, G., Kalliola, R.: Long-term landscape development processes in Amazonia, in: *Amazonia: Landscape and Species Evolution: A Look into the Past*, edited by: Hoorn, C., Wesselingh, F., Oxford, Wiley-Blackwell, 185–197, <https://doi.org/10.1002/9781444306408.ch22>, 2010.
- 680
- Latrubesse, E., Arima, E., Dunne, T., Park, E., Baker, V. R., d’Horta, F. M., Wight, C., Wittmann, F., Zuanon, J., Baker, P. A., Ribas, C. C., Norgaard, R. B., Filizola, N., Ansar, A., Flyvbjerg, B. and Stevaux J. C.: Damming the rivers of the Amazon basin. *Nature*, 546, 363–369, <https://doi.org/10.1038/nature22333>, 2017.
- 685
- Lehner, B. and Grill G.: Global river hydrography and network routing: baseline data and new approaches to study the world’s large river systems, *Hydrological Processes*, 27(15): 2171–2186. Data is available at www.hydrosheds.org, 2013.
- Merritts, D. J., Vincent, K.R. and Wohl, E. E.: Long river profiles, tectonism, and eustasy: a guide to interpreting fluvial terraces, *Journal of Geophysical Research*, 99, 14031–14050, <https://doi.org/10.1029/94JB00857>, 1994.
- 690
- Mertes, L.A.K. and Dunne, T.: Effects of tectonism, climatic change, and sea-level change on the form and behaviour of the modern Amazon River and its floodplain, in: *Large Rivers: Geomorphology and Management*, edited by: Gupta, A., Wiley, Chichester, 115–144, 2007.
- 695
- Norton, K. P., Schlunegger, F., and Litty, C.: On the potential for regolith control of fluvial terrace formation in semi-arid escarpments, *Earth Surf. Dynam.*, 4, 147–157, <https://doi.org/10.5194/esurf-4-147-2016>, 2016.
- Pupim, F. N., Sawakuchi, A. O., Almeida, R. P., Ribas, C. C., Kern, A. K., Hartmann, G. A., Chiessi, C. M., Tamura, L. N., Mineli, T. D., Savian, J. F., Grohmann, C. H., Bertassoli Jr., D. J., Stern, A. G., Cruz, F. W. and Cracraft, J.: Chronology of terra firme formation in Amazonian lowlands reveals a dynamic Quaternary landscape. *Quaternary*
- 700

- Science Reviews, 210, 154– 163, <https://doi.org/10.1016/j.quascirev.2019.03.008>, 2019.
- 705 Räsänen, M.E., Salo, J.S., Jungnert, H. and Pittman, L.R.: Evolution of the western Amazon lowland relief: impact of Andean foreland dynamics. *Terra Nova*, 2(4), pp.320-332, <https://doi.org/10.1111/j.1365-3121.1990.tb00084.x>, 1990.
- Regard, V., Lagnous, R., Espurt, N., Darrozez, J., Baby, P., Roddaz, M., Calderón, Y. and Hermoza, W.: Geomorphic evidence for recent uplift of the Fitzcarrald Arch (Peru): a response to the Nazca Ridge subduction, *Geomorphology*, 107, 107–117, <https://doi.org/10.1016/j.geomorph.2008.12.003>, 2009.
- 710
- Ribas, C. C., Aleixo, A., Nogueira, A. C. R., Miyaki, C. Y. and Cracraft, J.: A paleobiogeographic model for biotic diversification within Amazonia over the past three million years, *Proceedings of the Royal Society B, Biological Sciences*, 279, 681-689, <https://doi:10.1098/rspb.2011.1120>, 2012.
- 715
- Ribas, C. C. and Aleixo, A.: Diversity and evolution of Amazonian birds: implications for conservation and biogeography, *Anais da Academia Brasileira de Ciências*, 91, 3, 1-9, <https://doi:10.1590/0001-3765201920190218>, 2019.
- Rossetti, D. F., Toledo, P. M. and Góes, A. M.: New geological framework for Western Amazonia (Brazil) and implications for biogeography and evolution, *Quat. Res.* 63, 78–89, <https://doi:10.1016/j.yqres.2004.10.001>, 2005.
- 720
- Sacek, V.: Modelagem numérica conjunta de processos sedimentares e tectônicos em bacias sedimentares, Ph.D. thesis, São Paulo, Universidade de São Paulo, <https://doi:10.11606/T.14.2011.tde-21062013-151844>, 2011.
- 725
- Sacek, V.: Drainage reversal of the amazon river due to the coupling of surface and lithospheric processes, *Earth and Planetary Science Letters*, 401, Supplement C, 301–312, <https://doi:10.1016/j.epsl.2014.06.022>, 2014.
- Sioli, H.: The Amazon and its main afluent: hydrography, morphology of the river courses, and river types, in: *The Amazon*, edited by: Sioli, H., Springer, Dordrecht, 127-165, https://doi.org/10.1007/978-94-009-6542-3_5, 1984.
- 730

- Steffen, D., Schlunegger, F. and Preusser, F.: Drainage basin response to climate change in the Pisco valley, Peru, *Geology* 37, 491–494, <https://doi.org/10.1130/G25475A.1>, 2009.
- Tucker, G.E. and Slingerland, R.: Drainage basin responses to climate change, *Water Resources Research* 33, 2031–2047, <https://doi.org/10.1029/97WR00409>, 1997.
- 735 van Breukelen, M. R., Vonhof, H. B., Hellstrom, J. C., Wester, W. C. G. and Kroon, D.: Fossil dripwater in stalagmites reveals Holocene temperature and rainfall variation in Amazonia, *Earth and Planetary Science Letters*, 275, 54–60, <https://doi.org/10.1016/j.epsl.2008.07.060>, 2008.
- 740 Veit, H., May, J.-H., Madella, A., Delunel, R., Schlunegger, F., Szidat, S. and Capriles, J. M.: Palaeo-geoecological significance of Pleistocene trees in the Lluta Valley, Atacama Desert, *J. Quat. Sci.* 31 (3), 203–213, <https://doi.org/10.1002/jqs.2857>, 2016.
- Whipple, K. and Tucker, G.: Implications of sediment-flux-dependent river incision models for landscape evolution, *Journal of Geophysical Research*, 107, 1-20, <https://doi:10.1029/2000JB000044>, 2002.
- 745 Wilkinson, M. J., Marshall, L. G., Lundberg, J. G. and Kreslavsky, M. H.: Megafan environments in northern South America and their impact on Amazon Neogene aquatic ecosystems, in: *Amazonia, Landscape and Species Evolution: A Look into the Past*, edited by: Hoorn, C. and Wesselingh, F. P., Blackwell, 162–184, 2010.