

Spatio-temporal variability and controlling factors for postglacial denudation rates in the Dora Baltea catchment (western Italian Alps)

Elena Serra^{1,2}, Pierre G. Valla^{3,1,2}, Romain Delunel⁴, Natacha Gribenski^{1,2}, Marcus Christl⁵, Naki Akçar¹

¹Institute of Geological Sciences, University of Bern, Bern, 3012, Switzerland

5 ²Oeschger Centre for Climate Change Research, University of Bern, Bern, 3012, Switzerland.

³University Grenoble Alps, University Savoie Mont Blanc, CNRS, IRD, IFSTTAR, ISTERre, Grenoble, 38000, France

⁴Université Lumière Lyon 2, CNRS, UMR 5600 EVS, F-69635, France

⁵Laboratory of Ion Beam Physics, Swiss Federal Institute of Technology Zurich (ETHZ), Zurich, 8093, Switzerland

Correspondence to: Elena Serra (elena.serra@geo.unibe.ch)

10 **Abstract.** Disentangling the influence of **bedrock lithology** from the respective roles of climate, topography and tectonic forcing on catchment denudation is often challenging in mountainous landscapes due to the diversity of geomorphic processes in action and of spatial/temporal scales involved. The Dora Baltea catchment (western Italian Alps) ~~appears~~ the ideal setting for such investigation, since its large drainage system, extending from the Mont Blanc Massif to the Po Plain, cuts across different major litho-tectonic units of the western Alps, whereas this region has experienced relatively homogeneous climatic

15 conditions and glacial history throughout the Quaternary. We acquired new ¹⁰Be-derived catchment-wide denudation rates from 18 river-sand samples collected both along the main Dora Baltea river and at the outlet of its main tributaries. The inferred denudation rates vary between 0.2 and 0.9 mm/yr, consistent with previously-published values across the European Alps. Spatial variability in denudation rates was statistically compared with topographic, environmental and geologic metrics. ¹⁰Be-derived ~~denudation does~~ not correlate with modern precipitation and rock geodetic uplift. We find, rather, that catchment

20 topography, in turn conditioned by bedrock ~~structuration~~ and erodibility (litho-tectonic origin) and glacial overprint, ~~has the main influence on~~ ¹⁰Be-derived denudation patterns. We calculated the highest denudation rate for the Mont Blanc Massif, whose granitoid rocks and long-term tectonic uplift support high elevations, steep slopes and high relief and thus favour intense glacial/periglacial processes and recurring rock-fall events. Finally, our results, in agreement with modern sediment budgets, demonstrate that the high sediment input from the Mont Blanc catchment dominates the Dora Baltea sediment flux, explaining

25 the constant low ¹⁰Be concentrations measured along the Dora Baltea course even downstream the multiple junctions with tributary catchments.

1 Introduction

The use of *in-situ* ¹⁰Be concentrations measured in river sediments to quantify catchment-wide denudation rates over centennial to millennial time scale is now well-established (e.g. Brown et al., 1995; Granger et al., 1996; Bierman and Steig, 1996;

30 Granger and Schaller, 2014). ¹⁰Be concentrations are measured at the outlet of the studied basin and are inversely correlated ~~to~~ mean catchment denudation rate (von Blanckenburg, 2005). Widespread research has used this technique to estimate catchment denudation around the globe (see reviews in Portenga and Bierman, 2011; Willenbring et al., 2013; **Codilean et al.,**


35 2018) and specifically in mountain belts such as the European Alps (Delunel et al., 2020 and references therein), with the aim of illustrating the controlling mechanisms on recent (10²-10⁵ years) denudation dynamics and assessing the respective roles of climate and tectonics forcing.

In mountainous areas, the climatic imprint on the Earth's surface denudation has been recognized over both long and short timescales. Over the Late Cenozoic to Quaternary, temperature fluctuations, increased precipitation and glaciations have participated in a worldwide increase in erosion rates (e.g. Peizhen et al., 2001; Herman et al., 2013). In the European Alps, climate forcing has led to significant topographic modification through increased slope steepness and relief (e.g. Valla et al.,

40 2011), which in turn triggered a postglacial erosional response (Norton et al., 2010; Valla et al., 2010; Glotzbach et al., 2013;

Dixon et al., 2016). Over recent timescales (10^2 - 10^3 years), climate also exerts a control on denudation rates through precipitation and associated runoff (Moon et al., 2011; Bookhagen and Strecker, 2012) and by governing temperature/precipitation-dependent glacial and periglacial erosion processes (e.g. Delunel et al., 2010; Deline et al., 2014). Alternatively, other studies have shown a dominant litho-tectonic control on denudation rates (e.g. Cruz Nunes et al., 2015).
45 Rock-uplift and denudation rates are strongly coupled, with (1) erosional unloading driving uplift through isostatic rebound (Wittmann et al., 2007; Champagnac et al., 2009), and (2) tectonic rock uplift itself conditioning denudation rate by building new topographic gradient and increasing channel incision (e.g. Burbank et al., 1996; Montgomery and Brandon, 2002; Binnie et al., 2007; Godard et al., 2014). In transient landscapes such as recently deglaciated alpine settings, the topographic relief has not reached a steady state equilibrium between rock-uplift and denudation (e.g. Schlunegger and Hinderer, 2003; Delunel et al., 2020) and bedrock lithology may exert a significant control on millennial catchment denudation rates through its
50 structuration and erosional resistance (erodibility, Kühni and Pfiffner, 2001). More resistant lithologies have contrasting potential controls on denudation, (1) either decreasing denudation rates because of rock-mechanical strength (Scharf et al., 2013), (2) or promoting higher denudation rates by sustaining steep topography (Norton et al., 2011).

In the European Alps, the large-scale compilation of catchment-wide denudation rates by Delunel et al. (2020) highlighted (1)
55 the first-order correlation between denudation rate and mean catchment slope (derived from glacial imprint on Alpine topography), (2) the absence of relationship between modern climate and denudation patterns and (3) a significant correlation between rock uplift and denudation for >100 -km² catchments. This compilation also pointed at a rather weak control of bedrock lithology on denudation, with the lowest rates in the low-elevation foreland areas (with clastic sedimentary lithologies) and highest rates in the high-elevation crystalline parts (with gneissic, granitic or metamorphic lithologies) within the core of the
60 Alps. This trend, however, was not investigated further, since, at the scale of the European Alps, it appeared difficult to disentangle the relative influence of bedrock erodibility, topography and tectonic forcing on denudation rate as these are closely interrelated. Our study thereby aims to further explore the potential links and controls between climatically-driven topography, tectonic uplift and bedrock erodibility on the efficiency of denudation processes by investigating spatial variability of ¹⁰Be-derived denudation rates within the Dora Baltea (DB) catchment (western Italian Alps; Fig. 1). The DB catchment appears the
65 ideal setting for this investigation, since its large drainage system, extending from the Mont Blanc Massif (4808 m a.s.l.) to the Po Plain (around 200 m a.s.l.), cuts across the main litho-tectonic units of the western Alps (Fig. 2). Relatively similar climatic gradients and glacial history but variable bedrock lithology and geodetic uplift within the DB catchment and its tributaries allow us to assess how spatial variability in bedrock erodibility between litho-tectonic units may participate in controlling catchment topography and ¹⁰Be-derived denudation rates.



70

Figure 1: Study area with investigated Dora Baltea and main tributary river catchments (mosaic DEM from Regione Autonoma Valle d'Aosta, Regione Piemonte, swisstopo, and Institut Géographique National). Red and light blue circles indicate locations of river-sediment samples collected along the Dora Baltea river and at the outlet of the main river tributaries, respectively (for DB01 red-light blue circle as both along the Dora Baltea and considered as an individual tributary). Solid yellow lines delimit the catchments upstream of each sampling location (sample names indicated in white box). Present-day glaciers (GlaRiskAlp Project, <http://www.glariskalp.eu>), main topographic peaks and dams are indicated. Inset shows location of the DB catchment (red open box) within the European Alps.

75

2 Study area

The Dora Baltea (DB) catchment is a large drainage system of ~3900 km² located in the western Italian Alps (Fig. 1). Over a 170-km long distance, the DB river flows NW-SE from the Mont Blanc Massif to the Po Plain, and drains several tributaries connected to major >4000-m Alpine peaks (e.g. Mont Blanc, Monte Rosa, Matterhorn and Gran Paradiso; Fig. 1). Present-day mean annual temperatures range from -10°C in high-elevation zones to 15°C at valley bottoms (Regione Autonoma Valle d'Aosta, 2009). Precipitation varies between the semi-arid conditions prevailing at low elevations in the central part of the DB valley (mean annual precipitation of 400-500 mm/yr) and the wet conditions in the high-elevation internal valleys (Isotta et al., 2014). Higher mean annual precipitation values are observed in the Mont Blanc Massif (around 1800 mm/yr) compared to the north-western and southern sectors of the DB catchment (around 1400 mm/yr for Matterhorn and Monte Rosa area, and around 1150 mm/yr in the Grand Paradiso; Isotta et al., 2014). Present-day glaciers cover 3.6% of the total DB area, and are distributed within the upstream high-elevation parts of DB tributary catchments (Fig. 1), with terminus glacier elevations ranging from 2601 to 2800 m a.s.l. (data from 2005; Diolaiuti et al., 2012).


85

The geology of the DB catchment is complex, since its drainage network cuts across the main litho-tectonic units of the western Alps, recording the long-term collisional history between the European and Adriatic plates (e.g. Dal Piaz et al., 2008; Perello et al., 2008; Polino et al., 2008; Fig. 2). West of the Penninic Frontal Thrust, the European basement is exposed in the granitoid of the Mont Blanc External Massif and its Helvetic sedimentary cover. Bedrock units belonging to the thinned European crust (gneisses and schists of the Briançonnais basement and its terrigenous to carbonate metasedimentary cover, high-pressure gneisses of the Internal Massifs), the Tethyan oceanic crust (meta-ophiolite and calcschists of the Piedmont units) and the

90

95 Adriatic margin (Austroalpine gneisses and eclogitic micaschists) are exposed roughly from NW to SE across the axial belt, delimited by the Penninic Frontal Thrust to the NW and the Insubric Fault to the SE (Fig. 2). Long-term (10^6 - 10^7 years) exhumation rates from bedrock apatite fission-track data are higher in the western sector of the DB catchment (0.4-0.7 km/Myr for the External zones, west of Internal Houiller Fault; Fig. 2) than in the east (0.1-0.3 km/Myr for the Internal zones, i.e. between the Internal Houiller and the Insubric Faults, Malusà et al., 2005; Fig. 2). A similar pattern has been illustrated by short-term sediment budgets inferred from detrital apatite fission-track (Resentini and Malusà, 2012) and sediment gauge (Bartolini et al., 1996; Vezzoli, 2004; Bartolini and Fontanelli, 2009) data. Modern geodetic rock uplift also appears spatially variable within the DB catchment, with rates up to 1-1.6 mm/yr in the Monte Rosa, Mont Blanc and Ruitor areas, around 0.6-0.7 mm/yr in the axial belt and in the Gran Paradiso Massif, while decreasing to 0.2 mm/yr in the Po plain (Sternai et al., 2019). For the entire DB catchment, a ^{10}Be -derived denudation rate of 0.6 mm/yr was obtained by Wittmann et al. (2016, sample T12).

100 The DB catchment was repeatedly glaciated during the Quaternary, with major glaciers covering most of the catchment with the exception of the highest peaks ($\sim 3000 \text{ km}^2$, $>1000 \text{ m}$ thick; Serra et al., in press) and extending into the Po Plain during the Last Glacial Maximum (LGM, ca. 26-19 ka; Clark et al., 2009). The tributary glaciers already retreated in their upper valley catchments during the Lateglacial climatic oscillations (14-12 ka; Baroni et al., 2021; Serra et al., in press). As shown by the present-day topography (Fig. 1), postglacial fluvial dissection and hillslope processes have locally re-shaped the glacial landscape, with the development of V-shape valleys profiles and the deposition of large alluvial fans and infills along the main valleys.



115 **Figure 2.** Simplified litho-tectonic map of the study area with output catchment-wide denudation rates ($\pm 1\sigma$, mm/yr) reported at sampling locations (catchment boundaries in solid black lines). Major litho-tectonic domains and structural features (dashed lines) of the western European Alps are shown (modified map after Resentini and Malusà, 2012). Output catchment-wide denudation rates are corrected for topographic, LIA-glacier and snow shielding, and for quartz-content (see text and Table 1 for details).

3 Methods

3.1 ¹⁰Be-derived catchment-wide denudation rates

120 Eighteen river-sand samples were collected within the DB catchment, 5 along the main DB river and 13 at the outlet of the main tributaries (Fig. 1). One sample (DB06) was collected at the same location as sample T12 from Wittmann et al. (2012) to assess for the possible temporal variability of the in-situ ¹⁰Be signal exported by the DB river. Around 20-50 g of pure quartz were extracted from the 250-400 μm grainsize fraction, following sieving, magnetic separation and leaching in diluted HCl, H₃PO₄ and HF (detailed protocol reported in Akçar et al., 2017). The purified quartz was dissolved in concentrated HF after
125 addition of around 200 μg of ⁹Be carrier (Table S1), and Be extraction was performed through anion and cation exchange column chemistry (Akçar et al., 2017). Measurements of ¹⁰Be/⁹Be ratios were performed at ETH Zürich with the MILEA AMS system (Maxeiner et al., 2019), and normalized to the ETH in-house standards S2007N and S2010N (isotope ratios 28.1×10^{-12} and 3.3×10^{-12} , respectively; Christl et al., 2013). Calculated ¹⁰Be concentrations (Table 1) were corrected using a full process blank ¹⁰Be/⁹Be ratio of $2.96 \pm 0.32 \times 10^{-15}$.

130 Spatially-averaged ¹⁰Be production rates were derived from pixel-based calculations conducted with the Basinga ‘Production rates’ GIS tool (Charreau et al., 2019), using a 35-m resolution DEM from Regione Autonoma Valle d’Aosta and Regione Piemonte as input for catchment topography. For production rate calculations, we used the Lal/Stone time-dependent scaling model (Lal, 1991; Stone, 2000), which integrates corrections for atmospheric pressure and geomagnetic field fluctuations according to the ERA-40 reanalysis database (Uppala et al., 2005) and the Muscheler’s VDM database (Muscheler et al.,
135 2005), respectively. The total ¹⁰Be surface production rate at each DEM cell of the studied catchments was calculated based on a ¹⁰Be production rate at sea-level and high-latitude (SLHL) of 4.18 ± 0.26 at $\text{g}^{-1} \text{yr}^{-1}$ (Martin et al., 2017) considering relative contributions of 98.86, 0.87 and 0.27% by neutrons, slow muons, and fast muons, respectively (Charreau et al., 2019, after Braucher et al., 2011, and Martin et al., 2017). Attenuation length values of 160, 1500 and 4320 g cm^{-2} were used for neutrons, slow muons, and fast muons, respectively (Charreau et al., 2019, after Braucher et al., 2011), and a rock density of 2.7 g cm^{-3}
140 was assumed. Denudation-rate uncertainties (one-sigma external) were estimated only based on values and relative errors of ¹⁰Be concentrations and cosmogenic production rates from neutron and muons (Eq. 5 in Charreau et al., 2019). Integration times associated to the denudation rates (i.e. the time needed to erode the uppermost ~0.6 m of bedrock and ~1 m of soil; von Blanckenburg, 2005) were calculated as mean estimates (no uncertainty propagated).

Catchment-averaged production rates were corrected for (1) topographic shielding, (2) quartz-content, (3) Little Ice Age (LIA, 1250-1860 CE) glacier cover, and (4) snow shielding (Charreau et al., 2019). Catchment topographic shielding was computed
145 with the ‘toposhielding’ Topotoolbox function (Schwanghart and Scherler, 2014), following the method of Dunne et al. (1999) and Codilean (2006). We acknowledge the recent publication by DiBiase (2018) suggesting no need to correct for topographic shielding when calculating catchment-wide ¹⁰Be denudation rates. Our ¹⁰Be production rates were however corrected for topographic shielding to follow a conservative approach similar to the recent Alpine compilation study by Delunel et al. (2020).
150 As reported in Table 1, mean topographic shielding values obtained within the DB catchment are all very similar (~0.95), implying that neglecting the topographic-shielding correction would result in similar output rates generally within error estimates.

Based on the 1/100,000- and 1/250,000-scale digital geological maps from Regione Autonoma Valle d’Aosta and Regione Piemonte, respectively, we mapped and excluded from the ¹⁰Be production-rate calculation the catchment areas covered by
155 mafic and non-siliceous sedimentary (carbonate) bedrocks (Fig. S1), based on the assumption that they do not provide (or to a minor extent) quartz grains to the fluvial routing system. Crystalline bedrocks and Quaternary deposits (Fig. S1) were instead considered as quartz-bearing lithologies in our approach. In addition, we excluded areas with slope < 3°, assuming that they are likely not linked to the stream network or act as storage/transfer areas and therefore do not reflect catchment denudation (Fig. S1; Delunel et al., 2010). In order to estimate shielding correction due to glacier cover, ¹⁰Be production rates were set to

160 null for areas covered by LIA glaciers (GlaRiskAlp Project, <http://www.glariskalp.eu>; Fig. S1). This conservative approach
assumes sufficient ice thickness for complete cosmic-ray shielding (e.g. Delunel et al., 2010; Wittmann et al., 2007). Shielding
correction factors for snow cover were calculated as function of the average elevation for each individual catchment, by
applying an empirical model reported in Delunel et al. (2020) and predicting snow-shielding factors as a function of elevation
for the European Alps. The obtained average snow-shielding correction factors vary between 0.82 and 0.87 and were combined
165 to the topographic-shielding corrections as scaling factors for each sub-catchment. Catchment-wide denudation rates were then
obtained using the previously-calculated catchment-averaged ^{10}Be production rates and the measured ^{10}Be concentrations
(Table 1), with the Basinga ‘Denudation rates’ GIS tool (Charreau et al., 2019).

3.2 Topographic, environmental and geological metrics

We investigated potential drivers conditioning the observed spatial variability in DB catchment-wide denudation rates, and to
170 this aim we performed topographic analyses, and extracted environmental and geological variables of the quartz-bearing areas
(Fig. S1) for each investigated catchment through an ArcGIS-Matlab routine (Delunel et al., 2020).

Topographic analyses were conducted using a 35-m resolution DEM (Regione Autonoma Valle d'Aosta and Regione
Piemonte). We calculated drainage area, mean elevation, mean slope, percentage of slopes steeper than 40° , geophysical relief,
and hypsometric integral for the quartz-bearing areas of the individual catchments (Table 2). For slope analyses, the ‘gradient8’
175 Topotoolbox function was used (Schwanghart and Scherler, 2014), returning the steepest downward gradient of the 8-
connected neighbouring cells of the DEM. The percentage of catchment slope steeper than 40° was calculated as indicative of
the areal proportion of oversteepened threshold landscape (DiBiase et al., 2012). The geophysical relief (i.e. averaged elevation
differences between a surface connecting highest topographic points and the current topography; Small and Anderson, 1998)
was calculated in ArcGIS using a 5-km radius sampling window, and can be used as an indicator of past landscape change or
180 potential for locally increased erosion (see Champagnac et al., 2014 for discussion). The hypsometric integral was computed
based on Eq. 1 from Brocklehurst and Whipple (2004) and is inversely related to the stage of landscape evolution (i.e. more
evolved landscapes, whose high-elevation areas have been eroded, present low hypsometric integrals).

In addition, we extracted average values of the following environmental variables for the quartz-bearing areas of each
individual catchment. Average annual precipitation was obtained from the 5-km resolution grid of mean annual Alpine
185 precipitation from Isotta et al. (2014), in order to investigate the potential influence of modern precipitation/runoff on
denudation dynamics. Percentage of bare-rock area was estimated from the extent of class 30 (“bare bedrock”) of the 100-m
resolution CORINE Land Cover Inventory (2018), **to consider if catchment areas with null to low soil/vegetation cover are
more subjected to physical erosion and chemical weathering.** LIA-glacier areal cover was calculated based on the LIA-glacier
extent mapped within the GlaRiskAlp Project (<http://www.glariskalp.eu>), to assess the influence of modern to historical glacial
and periglacial processes on output ^{10}Be -derived denudation rates (Delunel et al., 2010). Mean LGM ice-thickness and areal
percentage of each catchment above the LGM Equilibrium Line Altitude (ELA) were estimated using the LGM paleo-glacier
reconstruction of the DB system (70-m resolution, LGM ELA at 2103 m a.s.l.; Serra et al., in press). Both metrics give
190 **indication on** the LGM glacial imprint on topography and subsequent potential for postglacial erosion response (Norton et al.,
2010; Salcher et al., 2014; Delunel et al., 2020). Catchment average temperature was not estimated since, at the relatively
constant latitude of the investigated catchments, temperature variability directly follows catchment hypsometric distribution
and thus relates to catchment elevation **which is already investigated** in the present study (topographic metric).

Lastly, we extracted geological variables for the studied catchments. Based on the simplified litho-tectonic map of the DB
catchment (Fig. 2), modified after Resentini and Malusà (2012), we estimated the relative proportion of the different litho-
tectonic units within the quartz-bearing areas of each catchment. Catchment-average geodetic uplift rates were ~~as well~~
200 considered using the 30-km resolution interpolation grid from Sternai et al. (2019), here downscaled to 600-m resolution grid
(Delunel et al., 2020).

4 Results

4.1 Spatial variability in catchment-wide denudation rates

Calculated catchment-wide denudation rates vary according to the applied production-rate correction factors (Table S2).
 205 Uncorrected denudation rates (i.e. including only mean catchment topographic shielding and excluding areas with slope $<3^\circ$)
 range between 0.27 ± 0.02 and 1.49 ± 0.13 mm/yr, while rates obtained by applying all corrections vary between 0.21 ± 0.02 and
 0.91 ± 0.08 mm/yr (Table 1 and Fig. 2). Significant production-rate corrections were obtained when taking into account snow
 shielding and LIA-glacier cover (up to 17 and 42% reduction compared to uncorrected ^{10}Be production/denudation rates,
 respectively), especially for catchments with high mean elevations and associated large LIA-glacier coverage (Table 2). Lower
 210 corrections were obtained when considering quartz-bearing areas (maximum 10% reduction in output ^{10}Be
 production/denudation for catchments DB08 and 11, where relatively abundant sedimentary and mafic bedrocks occur; Fig.
 S1). All corrections combined together lead to reduction in ^{10}Be production/denudation rates of 16-53% compared to the
 uncorrected estimates (Table S2). Hereafter, we consider ^{10}Be production/denudation rates obtained by applying all corrections
 (Table 1 and Fig. 2), in order to maintain a conservative approach as in the recent Alpine compilation study (Delunel et al.,
 215 2020).

Sample	Location WGS 84 (dd N/ dd E)	Elevation (m a.s.l.)	^{10}Be concentration ($\times 10^4$ at g^{-1})	Topographic shielding	Mean production rate (at $\text{g}^{-1} \text{yr}^{-1}$)		Denudation rate (mm/yr)		Integration time (yr)	
					Uncor.	Corr.	Uncor.	Corr.	Uncor.	Corr.
DB01	45.8040/ 6.9653	1230	1.29 ± 0.07	0.92	29.4	13.7	1.45 ± 0.11	0.68 ± 0.05	400	900
DB02	45.7167/ 7.1101	783	1.08 ± 0.07	0.94	25.1	15.3	1.49 ± 0.13	0.91 ± 0.08	400	700
DB03	45.6925/ 7.1935	699	2.35 ± 0.14	0.94	28.6	18.0	0.76 ± 0.06	0.48 ± 0.04	800	1300
DB04	45.7003/ 7.2019	664	2.20 ± 0.11	0.94	27.5	17.8	0.78 ± 0.06	0.51 ± 0.04	800	1200
DB05	45.7001/ 7.2337	638	2.05 ± 0.10	0.95	27.8	17.4	0.85 ± 0.06	0.53 ± 0.04	700	1100
DB06	45.5228/ 7.8375	251	1.54 ± 0.08	0.95	22.6	16.1	0.94 ± 0.07	0.68 ± 0.05	600	900
DB07	45.5962/ 7.7956	325	2.25 ± 0.26	0.95	22.2	15.3	0.61 ± 0.07	0.42 ± 0.05	1000	1400
DB08	45.6118/ 7.7310	373	4.85 ± 0.21	0.96	20.2	15.9	0.27 ± 0.02	0.21 ± 0.02	2300	2900
DB09	45.7352/ 7.6124	465	3.34 ± 0.18	0.96	24.4	17.5	0.46 ± 0.04	0.33 ± 0.03	1300	1800
DB10	45.7079/ 7.6713	375	1.35 ± 0.07	0.95	23.5	16.5	1.12 ± 0.09	0.79 ± 0.06	500	800
DB11	45.6830/ 7.7115	546	3.47 ± 0.20	0.96	24.5	16.0	0.44 ± 0.04	0.29 ± 0.02	1400	2100
DB12	45.7183/ 7.2651	594	1.26 ± 0.08	0.95	25.6	16.4	1.30 ± 0.11	0.84 ± 0.07	500	700
DB13	45.7482/ 7.3224	753	2.79 ± 0.12	0.95	24.4	17.2	0.56 ± 0.04	0.40 ± 0.03	1100	1500
DB14	45.7882/ 7.3061	600	2.83 ± 0.13	0.96	22.1	18.5	0.50 ± 0.04	0.42 ± 0.03	1200	1400
DB16	45.7386/ 7.4292	524	3.88 ± 0.35	0.94	23.6	18.7	0.38 ± 0.04	0.30 ± 0.03	1600	2000
DB17	45.7039/ 7.1622	689	2.71 ± 0.13	0.95	27.2	17.3	0.63 ± 0.05	0.40 ± 0.03	1000	1500
DB18	45.7039/ 7.1622	689	2.30 ± 0.12	0.94	27.0	17.2	0.75 ± 0.06	0.48 ± 0.04	800	1300

DB19	45.7619/ 6.9873	1005	4.19±0.15	0.96	25.8	16.0	0.39±0.03	0.25±0.02	1500	2400
------	--------------------	------	-----------	------	------	------	-----------	-----------	------	------

Table 1: River-sediment sample locations, measured ^{10}Be concentrations, calculated mean catchment ^{10}Be production rates, and output denudation rates and apparent ages. Sample coordinates are given in decimal degrees (dd), and sample names collected along the main DB river are written in italics. Production rate estimates (and derived denudation rates / apparent ages) are provided for (1) topographic shielding correction (column labeled “Uncor.”) and (2) including corrections for topographic shielding, snow and LIA-glacier shielding and for quartz-content (column labelled “Cor.”). Mean catchment ^{10}Be production rates (and derived catchment denudation rates) obtained by applying each individual correction are reported in Table S2.




Figure 3: Downstream evolution of river-sand ^{10}Be concentrations in the Dora Baltea (DB) catchment. Data are plotted versus distance from the main DB source (upper Val Veny, right tributary upstream DB01). In red are samples collected along the main DB river, in light blue are samples at the outlet of tributaries (Fig. 1 for locations). ^{10}Be concentration of sample T12 (red square) from Wittmann et al. (2016) is also shown for discussion. Samples DB03, 14 and 18 are omitted since they are in turn tributaries of catchments DB04 and DB13, respectively, and do not directly connect to the main DR river.

The highest denudation rates (0.7-0.9 mm/yr) were obtained from riverine samples collected along the main DB river (DB01, 02, 12, 10, 06), showing a slightly-decreasing trend in denudation rate with the river distance (apart from DB01; Fig. 2). ^{10}Be concentrations measured in these samples are the lowest and overall constant along the DB course (around 1.2×10^4 at/g; Fig. 3). Within tributary catchments, with the exception of sample DB01, ^{10}Be concentrations are higher (2.0 - 4.9×10^4 at/g; Fig. 3), and calculated denudation rates are lower, generally within 0.4-0.5 mm/yr and down to 0.2-0.3 mm/yr for some catchments (DB08, 09, 11, 16 and 19; Fig. 2).

In order to quantify at first-order the relative contribution of the Mont Blanc Massif (represented by the lowest ^{10}Be concentration of sample DB02; Fig. 3) to the ^{10}Be signal measured along the DB river, we followed the approach reported in Delunel et al. (2014). River-sediment ^{10}Be concentrations from tributaries and along the DB river have been first normalised to the SLHL ^{10}Be production rate (i.e. 4.18 ± 0.26 at $\text{g}^{-1} \text{yr}^{-1}$), implying that variations in normalised ^{10}Be concentrations represent the variability in denudation rates only. We then estimated the respective contributions of the Mont Blanc Massif and different tributaries through a mixing model (mass-balance model involving catchment ^{10}Be concentrations and contributing areas; Delunel et al., 2014) considering (A) the normalised ^{10}Be concentration for river materials exported from the Mont Blanc catchment, (B) the averaged normalised ^{10}Be concentration from the upstream tributaries contributing to each sampling points along the main DB river and (C) the normalised ^{10}Be concentration at the sampling points along the main DB river (DB12, 10, 06). Between our two most upstream DB river samples (DB01 and DB02), we based our model on DB02,

245 which provides a more conservative estimate of the contribution of the Mont Blanc Massif to the ^{10}Be signal measured along the DB river (i.e. the potential contributions of the tributaries are maximized). By applying this simple model, we find that the Mont Blanc Massif (upstream catchment DB02) contributes to 90, 87 and 77% of the river-sediment ^{10}Be signal measured respectively at locations DB12, 10 and 06, in line with overall constant ^{10}Be concentrations measured along the DB course (Fig. 3).

250 4.2 Catchment metrics and denudation rates


Results of catchment topographic analyses, along with estimates of environmental and geological metrics, are reported in Table 2. As for calculated catchment-wide denudation rates, DB01 (Mont Blanc Massif) also appears as an end-member with maximum values in most of the reported metrics (Table 2).

Catchment	Total drainage area (km ²)	Relative quartz-bearing area (%)	Mean elevation (m a.s.l.)	Mean slope (°)	Relative abundance of slopes > 40° (%)	Geophysical relief (5-km, m)	Hypsometric integral	Mean annual precipitation (mm)	Relative abundance of bare-rock (%)	LIA glacier cover (%)	Mean LGM ice-thickness (m)	Basin area above LGM ELA (%)	Mean geodetic rock uplift (mm yr ⁻¹)
<i>DB01</i>	189	88.7	2600	32.2	32.7	1632	0.39	1762	34.6	37.3	474	77	1.08
<i>DB02</i>	496	87.3	2288	28.5	28.2	1297	0.38	1475	24.6	19.8	577	62	1.18
DB03	147	80.3	2499	28.8	29.6	1246	0.54	1088	35.0	16.9	441	77	0.89
DB04	279	78.0	2387	28.5	30.4	1219	0.51	1071	32.5	13.0	476	71	0.97
DB05	257	71.9	2419	29.5	28.2	1191	0.52	1045	30.1	16.3	427	71	0.68
<i>DB06</i>	3321	73.6	2096	27.7	25.6	1214	0.40	1147	19.4	10.3	526	53	0.95
DB07	278	85.7	2030	30.2	26.9	1076	0.41	1352	11.1	8.7	304	43	0.67
DB08	108	56.4	1827	29.3	29.3	978	0.52	1164	4.1	0.3	282	36	0.48
DB09	207	62.0	2228	24.4	18.6	1232	0.45	1092	21.1	10.8	317	61	1.32
<i>DB10</i>	2464	77.9	2156	28.1	26.0	1234	0.40	1130	22.3	11.3	553	57	1.03
DB11	226	48.5	2060	25.5	18.2	1211	0.40	1051	13.3	7.8	377	47	0.96
<i>DB12</i>	1310	80.7	2284	28.2	26.9	1226	0.40	1228	25.2	15.3	534	64	1.03
DB13	450	90.6	2224	29.0	28.0	1148	0.46	1144	26.4	10.2	558	61	1.10
DB14	141	92.4	2090	26.8	20.0	1092	0.45	1269	12.4	0.8	568	51	0.93
DB16	54	53.4	2154	27.8	25.2	1146	0.62	949	18.7	2.1	369	58	0.68
DB17	158	79.4	2335	27.6	23.7	1107	0.55	1047	26.3	13.9	467	70	1.24
DB18	277	89.4	2406	30.9	34.0	1137	0.48	1123	36.9	16.4	493	72	1.22
DB19	148	80.3	2313	24.5	21.1	914	0.53	1251	23.8	18.0	506	69	1.39


255 **Table 2:** Topographic, environmental and geological metrics extracted for the studied catchments (upstream of sampling river locations for ^{10}Be analysis). Names of sample collected along the main DB river are given in italics. With the exception of the total drainage area, all the metrics were calculated using the quartz-bearing area of each catchment. See text for details.

We compared the ^{10}Be -derived catchment denudation rates against topographic, environmental and geological metrics and evaluated the statistical significance of linear correlations (p-value and r^2 ; Figs. 4 and 5, Table S3). Samples along the main DB river (downstream of DB01, i.e. DB02, 12, 10, 06; Fig. 2) were excluded from the investigated correlations since their apparent denudation rates are potentially affected by cumulative drainage and sediment mixing along the DB course. Correlations were calculated both including and excluding sample DB01. Cook's distance values were also calculated in order to assess whether DB01 strongly influences the derived correlations as a potential outlier. We selected a threshold value of 3 times for DB01 Cook's distance compared to the data mean Cook's distance. As a consequence, DB01 appears as an outlier in different investigated correlations between catchment denudation rates and topographic, environmental or geological metrics (Table S3). Nevertheless, significant linear correlations (i.e. p-value <0.05) both with and without DB01 were obtained between catchment denudation rates and mean elevation (Fig. 4A), 5-km geophysical relief (Fig. 4C), the relative abundance of bare bedrock (Fig. 5B), and the percentage of area above the LGM ELA (Fig. 5D). Significant linear correlations between

270 catchment denudation rates and mean slopes (Fig. 4B), proportions of oversteepened slopes (Table S3) and relative area covered by LIA glaciers (Fig. 5C) were instead only found when including DB01. Non-significant correlations (p -value ≥ 0.05) were observed between catchment denudation rates and drainage areas (Table S3), hypsometric integrals (Table S3), mean annual precipitation values (Fig. 5A), mean geodetic uplift rates (Fig. 4D) and mean LGM ice-thickness (Table S3).



275 **Figure 4:** Relationships between tributary-catchment denudation rates and mean catchment (A) elevation, (B) slope, (C) 5-km geophysical relief, and (D) geodetic uplift. Linear correlations have been calculated including or not sample DB01 (red and black lines, respectively; see main text for discussion). Correlation coefficients (p -value and r^2) are reported for each linear regression with statistical significance (p -value < 0.05). r^2 is not reported for non-significant correlations (p -value > 0.05). Asterisks indicate linear correlations for which DB01 has been considered as an outlier (based on Cook's distance, Table S3).



280

Figure 5: Relationships between tributary-catchment denudation rates and catchment (A) mean annual precipitation, (B) relative bare-bedrock area, (C) relative area covered by LIA glaciers, and (D) relative area above LGM ELA (2103 m a.s.l.). Correlations have been calculated including or not sample DB01 (red and black lines, respectively; see text for discussion). Correlation coefficients (p-value and r^2) are reported for each linear regression with statistical significance (p-value < 0.05). r^2 is not reported for non-significant correlations (p-value > 0.05). Asterisks indicate linear correlations for which DB01 has been considered as an outlier (based on Cook's distance, Table S3).

285


4.3 Litho-tectonic units and denudation rates

In addition to catchment metrics, we explored the potential influence of bedrock properties on the postglacial evolution of the DB catchment by analysing the correlation between tributary-catchment denudation rates and the spatial distribution of litho-tectonic units (Figs. 2 and 6; only quartz-bearing areas considered, Fig. S1). The highest denudation rates are observed for tributaries with widespread bedrock exposure of granites of the Mont Blanc External Massif and its Helvetic terrigenous to carbonate sedimentary cover (85%; DB01: 0.68 ± 0.05 mm/yr). Moderate denudation rates around 0.4-0.5 mm/yr are observed for catchments with abundant gneisses of the Gran Paradiso Internal Massif (41-73%; DB03, 04, 05: average denudation rate of 0.51 ± 0.02 mm/yr), with dominant Austroalpine gneisses and eclogitic micaschists (58-85%; DB07, 13, 18: average denudation rate of 0.43 ± 0.03 mm/yr, with the exception of DB08 at 0.21 ± 0.02 mm/yr) or with abundant gneisses and schists of the Briançonnais basement (63-89%; DB14, 17: average denudation rate of 0.41 ± 0.01 mm/yr). The lowest denudation rates were obtained for tributaries dominated by meta-ophiolites and calcschists of the Piedmont units (73-100%; DB09, 11, 16: average denudation rate of 0.31 ± 0.02 mm/yr), and by the terrigenous to carbonate Briançonnais metasedimentary cover (87%; DB19: 0.25 ± 0.02 mm/yr). It is however to be noted that for these tributaries (DB09, 11 and 16), the quartz-bearing areas

290

295

300 considered in our calculations and analysis mainly consist of Quaternary deposits rather than exposed bedrocks (Fig. 2 and S1).



305 **Figure 6:** Tributary-catchment denudation rates (A) and relative proportion of litho-tectonic units within the quartz-bearing areas of the individual catchments (B). Colour code in (A) refers to the most abundant litho-tectonic unit in each individual catchment (see Figure 2 for spatial distribution of the different litho-tectonic units).

310 Finally, we also assessed the potential influence of bedrock lithological properties on catchment morphometry, by evaluating the distribution of elevation, slope and 5-km geophysical relief for the quartz-bearing areas of each individual litho-tectonic unit (Fig. 7). Higher elevations are observed for the External and Internal Massifs (median of 2500-2700 m a.s.l.) compared to the other litho-tectonic units (median of 1900-2200 m a.s.l.; Fig. 7A). The slope distributions appear slightly higher for the External Massif, the Austroalpine units and the Briançonnais basement (median of 31-32°) than for the other units (median of 23-26°; Fig. 7B). The External Massif present the highest geophysical reliefs (median of 1700 m), while all the other litho-tectonic units have similar geophysical relief with medians varying between 1000 and 1300 m (Fig. 7C).

315




Figure 7: Box-and-whisker plots for the spatial distribution of elevation (A), slope (B), and 5-km geophysical relief (C) within the entire DB catchment, classified by individual litho-tectonic unit. Red horizontal line represents the median of each distribution, bottom and top of each box are the 25th and 75th percentiles. Whiskers extend up to 1.5 times the interquartile range, outliers (red crosses) are observations beyond the whiskers.

5. Discussion

5.1 Correction factors for catchment-wide ^{10}Be production and denudation rates

As reported in Table S2, the different correction factors for quartz-content, LIA-glacier cover and snow-shielding lead to 16-
325 53% decrease in catchment ^{10}Be production and inferred denudation rates compared to uncorrected estimates (i.e. including
only catchment-averaged topographic shielding and excluding areas with slope $<3^\circ$). Such correction factors build on several
assumptions and have different implications for our catchment-wide denudation rate results that are discussed hereafter.
However, we should also note that the investigated correlations between denudation rates and topographic, environmental and
geological metrics (Fig. 4-5) remain similar when using non-corrected denudation rates.

330 First, assuming Quaternary deposits as quartz-bearing lithologies is a first-order approximation, since deposits derived from
mafic and carbonate-sedimentary bedrocks would bring no or minor quartz to the sediment routing system. For some tributaries
dominated by these lithologies (DB09, 11 and 16, as reported in section 4.3), we nevertheless considered that Quaternary
deposits may bear quartz given that the upper part of their catchments drain crystalline bedrock (Fig. S1). However,
distinguishing deposit provenance/lithology in this Alpine environment, with complex glacial/periglacial systems, would
335 require detailed field investigation and mapping, which is beyond the scope of this work. Moreover, our calculations show that
correction for quartz-bearing area has only a minimal effect on catchment-averaged ^{10}Be production and denudation rates, with
only up to 10% difference between uncorrected and corrected results thus overlapping within uncertainties (Table S2).

Second, correction factors for LIA-glacier cover and snow shielding lead instead to significant decrease in catchment-averaged
 ^{10}Be production and thus denudation rates (up to 42 and 17%, respectively). Since sediments in sub-/proglacial environments
340 can derive from periglacial erosion from bedrock walls/peaks and/or re-mobilization of previously exposed material (with non-
zero ^{10}Be concentration, e.g. moraine deposits; Wittmann et al., 2007; Delunel et al., 2014; Guillon et al., 2015), assuming null
 ^{10}Be concentration input from areas covered by LIA glaciers might lead to overcorrections of our denudation rate estimates.
Uncertainties are related also to the snow-shielding correction approach. The snow-shielding vs. elevation model reported by
Delunel et al. (2020) has been calibrated on snow-water equivalent records of the Swiss and French Alps, which are wetter
345 regions compare to the DB catchment (Isotta et al., 2014). Therefore, LIA-glacier cover and snow-shielding corrections may
be overestimated for the DB catchments, especially for high-elevation tributaries. In particular, catchment DB01 shows the
maximum corrections for both LIA-glacier cover and snow shielding (42 and 17% respectively; Table S2) and consequently
relatively low output denudation rate compared to estimates obtained for catchments downstream along the main DB river
(DB02, 12, 10; Fig. 2), despite similar ^{10}Be concentrations (Fig. 3). We therefore acknowledge that our corrected catchment-
350 averaged ^{10}Be production and denudation rates (Table 1 and Fig. 2) should be considered as minimum estimates, given the
correction factors for LIA-glacier cover and snow shielding, in line with the recent compilation over the entire European Alps
(Delunel et al., 2020).

Finally, we need to assess the impact of LGM glacial erosion on our ^{10}Be -derived denudation rates (Glotzbach et al., 2014;
Dixon et al., 2016), since our study area has been largely glaciated during the LGM (Serra et al., in press). Deep glacial erosion
355 may have largely to completely zeroed ^{10}Be concentration on bedrock surfaces, with non steady-state ^{10}Be concentration depth
profiles during postglacial surface exposure leading to apparent overestimate in denudation rates from ^{10}Be concentrations in
river sands (Glotzbach et al., 2014). However, given the deglaciation history of the DB catchment (i.e. largely deglaciated by
14-12 ka; Baroni et al., 2021; Serra et al., in press) and the range of our ^{10}Be -derived denudation rates (0.2-0.9 mm/yr, Table
1 and Fig. 2), we can **estimate an overestimate** of our ^{10}Be -derived denudation rates by maximum 10-15%, similar to the
360 proposed estimate of Dixon et al. (2016) in the Eastern Alps, with a maximum 9% overestimate for slower ^{10}Be -derived
denudation rates (~ 0.2 mm/yr). We thus are confident in the validity of our ^{10}Be -derived denudation rates (Table 1), and can

exclude any potential strong bias influencing the spatial pattern (Fig. 2) and interpretation with regards to topographic, environmental and geological metrics (Figs. 4-5).

365 5.2 Propagation of ^{10}Be signal along the DB course

Our results highlight the strong ^{10}Be -dominance of the Mont Blanc Massif (represented by sample DB02, see section 4.1 for discussion) on downstream sediment samples collected along the DB course, below the tributary junctions (Fig. 3, Table 1). The relatively constant low ^{10}Be concentrations measured for samples DB01, 02, 12, 10, 06 (around 1.2×10^4 at/g, Fig. 3) compared to the tributaries ($2.0\text{--}4.9 \times 10^4$ at/g), and the outcomes of our mixing model indicate unequal sediment contribution
370 (non-balanced sediment budget; Savi et al., 2014) between the main DB stream and its tributaries. The Mont Blanc Massif appears to govern the sediment yield along the main DB river, contributing to $>77\%$ of the river-sediment ^{10}Be signal carried all along the DB river.

A key factor governing the mixing and flux balance of ^{10}Be concentrations between river streams is the quartz flux from each stream, which is in turn influenced by (1) catchment denudation rate, (2) drainage area, (3) catchment quartz content (Carretier et al., 2015), (4) sediment storage (e.g. dams, lakes, floodplains reducing mass flux but not changing the ^{10}Be concentration; Wittmann et al., 2016). Our results show significantly higher denudation rate for catchment DB01 compared to other DB
375 tributaries (Fig. 2, Table 1). While the Mont Blanc Massif (upstream DB02 catchment) represents only a minor fraction of the total DB catchment area ($\sim 18\%$), its quartz-bearing surface area appear 5-90% larger than for other tributaries (Table 2). Likewise, the sediment-provenance studies of Vezzoli et al. (2004) and the sediment-yield estimates of Vezzoli (2004)
380 highlighted that river sands from the Mont Blanc catchment (analogous catchment to DB02) have up to $\sim 20\%$ higher quartz content compared to some other DB tributaries (analogous to DB09, 11, 16; Table S4) and contribute to $\sim 62\%$ of the quartz flux of the entire DB catchment (analogous to DB06; Table S4). Since the occurrence of dams is limited to few catchments (Fig. 1), the high quartz flux and ^{10}Be -signal dominance of the Mont Blanc Massif along the DB course could derive from (1) its high denudation rate (Fig. 2 and Table 1), (2) its large quartz-bearing drainage area and (3) the high quartz content of the
385 Mont Blanc granitoid (Vezzoli, 2004). Between this three potential causes, we propose that the ^{10}Be -signal dominance of the Mont Blanc Massif along the DB course is mainly driven by its high denudation rather than quartz fertility or area coverage, as illustrated by the similar trend of modern denudation rates derived from sediment gauging (Hinderer et al., 2013; see also discussion in section 5.4). The high rock-slope instability and glaciogenic sediment production in the Mont Blanc Massif supply abundant low ^{10}Be concentration quartz to the river system, being therefore efficient in diluting the ^{10}Be concentration
390 in the downstream course of the DB river. Controlling factors explaining the high denudation rate of the Mont Blanc Massif are further discussed below (section 5.3).

For the entire DB catchment, we can note that the ^{10}Be concentration is $\sim 30\%$ higher for sample T12 ($1.99 \pm 0.14 \times 10^4$ at/g; Wittmann et al., 2016) compared to DB06 ($1.52 \pm 0.08 \times 10^4$ at/g; Fig. 3), both collected at the same location (DB catchment outlet, Fig. 1) but at different time periods. The observed difference is probably related to a stochastic change in sediment
395 sources (Lupker et al., 2012), with potentially a temporary dominant sediment input from a DB tributary catchment with higher ^{10}Be concentration (e.g. DB07, close location and similar ^{10}Be concentration as T12; Fig. 3) than from the Mont Blanc catchment. By comparing our results to the Po catchment (Wittmann et al., 2016), which drains several main river systems from the south-western Alps in addition to the DB river basin, it emerges that the low ^{10}Be concentration signal deriving from the Mont Blanc Massif, remaining overall constant along the DB course, increases significantly soon after the DB flows into
400 the Po river. The high ^{10}Be concentrations measured by Wittmann et al. (2016) in Po river-sediment samples, immediately downstream the DB confluence (samples P1 and P3: around 3.6×10^4 at/g), show that the Po river is dominated in its initial lowland flow by the high ^{10}Be concentration inputs from other south-western Alpine catchments (Wittmann et al., 2016).

5.3 Controlling factors and processes on ¹⁰Be-derived catchment denudation rates

405 Our ¹⁰Be-derived denudation rates, varying between 0.2 and 0.9 mm/yr, fit within the values obtained over the European Alps, where 95% of the considered catchments yield denudation rate values <1.2 mm/yr and rates for the Western European Alps range between 0.1 and 1.2 mm/yr (Delunel et al., 2020). Correlations with topographic, environmental and geologic metrics allowed us to identify and discuss potential controlling mechanisms for denudation-rate variability within the DB catchment, that we present here in comparison with studies conducted in other Alpine sectors.

410 While precipitation and rock uplift have been recognized as main drivers for Alpine denudation rates, especially for the Central Alps (e.g. Chittenden et al., 2014; Wittmann et al., 2007, respectively), their respective influence on denudation-rate variability within the DB catchment is not significant (Figs. 5A and 4D). Interestingly, it can be observed that catchment geodetic rock uplift is overall higher (20-80%) than ¹⁰Be denudation in all the investigated tributary catchments, suggesting a net surface uplift of the DB area for recent timescales, in line with other observations across the European Alps (Norton et al., 2011; Delunel et al., 2020).

Catchment topography appears instead to have a major role in controlling the observed spatial variability in DB denudation rates. First, denudation rates are positively correlated with catchment-averaged elevation (Fig. 4A). Elevation influences denudation rates through periglacial (i.e. frost-cracking; Delunel et al., 2010) and glacial erosive processes, both increasing with altitude due to their temperature dependency, as well as by modifying soil and vegetation cover, with bare-rock exposure being positively correlated with denudation rates (Fig. 5B). Second, correlations with slope and geophysical relief need to be considered (Figs. 4B and C). It has been previously proposed that topographic slope and geophysical relief are positively correlated to catchment denudation until a threshold slope angle of 25-30° (Montgomery and Brandon, 2002; Champagnac et al., 2014; Delunel et al., 2020). Below this threshold, denudation was shown to respond to a slope-dependent equilibrium between regolith cover production through weathering and its downslope diffusion. In oversteepened catchments, denudation rates are instead controlled by mass wasting processes (i.e. rockfalls, debris flows, landslides) which stochastically influence river-sediment ¹⁰Be concentrations. While the potential effect of slope alone is here challenging to evaluate as all the tributaries exhibit similar averaged slope values, between ~25-30° (with the exception of DB01 with average slope of ~32°, Fig. 4B), denudation rate exhibits a clear correlation with geophysical relief (Fig. 4C), which is function of both slope and elevation difference (Small and Anderson, 1998; Champagnac et al., 2014). We suggest that slope differences between the investigated catchments, while not significant, are nevertheless still close to threshold values (Fig. 4B; Delunel et al., 2020), which, when combined with elevation differences between catchments, would explain the significant relationship observed between geophysical relief and denudation rates (Fig. 4C).

Our results also show a correlation between catchment denudation rates and bedrock litho-tectonic classification (Fig. 6), which has been proposed to drive denudation through rock mechanical strength (erodibility; Kühni and Pfiffner, 2001). Similar to what has been suggested based on DB modern sediment provenance (Vezzoli et al., 2004), we observe a general trend with the highest denudation rates in catchments dominated by apparent “low erodibility” bedrocks (granite and gneiss), and the lowest rates in catchments with apparent “high erodibility” bedrocks (sedimentary and terrigenous rocks; erodibility classes according to Kühni and Pfiffner, 2001). This trend has already been observed locally in the Eastern and Southern Alps (Norton et al., 2011) as well as at the scale of the entire European Alps (Delunel et al., 2020). Such observations were interpreted to be related to the influence of bedrock resistance on catchments morphometry (in turn connected to denudation dynamics), with the most resistant lithologies located at highest elevations and sustaining the steepest slopes/highest reliefs (Kühni and Pfiffner, 2001; Stutenbecker et al., 2016). Our results are in line with this interpretation, with the “low-erodibility” granite of the Mont Blanc External Massif supporting the highest elevation and reliefs and slightly steeper slopes (Fig. 7), where efficient geomorphic processes promote the highest catchment denudation rate (Fig. 6). On the other hand, the “high erodibility” rocks of the Briançonnais cover and of the Piedmont units present low elevation, relief and slope values, and are associated with low denudation rates. High elevation sustained by gneisses and granite of the Internal Massifs (2700 m a.s.l.; Fig. 7A) and slightly

steeper slopes supported by gneisses and micaschists of the Austroalpine units and of the Briançonnais basement (30-31°; Fig. 7B) would also drive the moderate denudation rates observed in these three litho-tectonic domains (Fig. 6). Moreover, the different long-term tectonic histories between the litho-tectonic domains could also explain some of the observed variability in catchment denudation between areas west and east of the Penninic Frontal Thrust (Fig. 2). Bedrock tectonic fracturing (Molnar et al. 2007) may influence subsequent erodibility and denudation, facilitated by the exhumation of more fractured bedrock units such as the crystalline units of the Mont Blanc External Massif and its Helvetic sedimentary cover (no deep Eocene subduction during Alpine orogeny), compared to deeply-exhumed rocks of the Internal Massifs and Piedmont units (Schmid et al., 2004). Additionally, higher late-Miocene uplift rates in the Mont Blanc Massif compared to the rest of the DB catchment (Malusà et al., 2005) could have sustained high-elevations in the Mont Blanc Massif, which in turn would also promote efficient geomorphic processes and high denudation rates.

Lastly, we consider the potential connection between landscape glacial imprint and catchment denudation rates. Statistically significant correlation between catchment denudation rates and catchment area proportion above LGM ELA (Fig. 5D) suggests an impact of LGM/large Quaternary glaciations on ¹⁰Be-derived denudation rates. Most of the catchments have >50% of their area above the LGM-ELA, indicating that large glaciers persisted during the LGM (and potentially older Quaternary glacial stages), with a significant impact on catchment topography characterized by steep slopes and high reliefs (Fig. 4A-C, Pedersen and Egholm, 2013). We hence tentatively interpret the significant correlation between denudation rates and high elevation / pronounced geophysical relief (Figs. 4B and C) to be indicative of a long-term glacial topographic control on the postglacial erosional response, as suggested by previous studies (Norton et al., 2010a; Glotzbach et al., 2013; Dixon et al., 2016). The glacial pre-conditioning of the topography has been also enhanced during postglacial times with coupled fluvial incision and hillslope processes increasing Alpine valley slopes and reliefs locally (Korup and Schlunegger, 2007; Valla et al., 2010; van den Berg et al., 2012). Over a shorter term, the positive correlation between catchment denudation rates and LIA glacial cover (only when including DB01, Fig. 5C) suggests also an important role of Holocene to modern glacial processes in influencing catchment denudation, by contributing to high-sediment delivery (Stutenbecker et al., 2018).

By considering the above-mentioned controlling mechanisms for catchment denudation, we propose the following interpretation for the high denudation rate obtained for catchment DB01 compared to other DB tributaries (Figs. 2 and 4-5). Catchment DB01 has maximum values for most of the investigated metrics (Figs. 4 and 5, Table 2). Its location in the high-elevation core of the Alps (Mont Blanc Massif, long- and short-term high uplift rate) was the site of intense Quaternary glaciations (large catchment area above the LGM ELA), which deeply modified the landscape as illustrated by the high geophysical relief of this catchment. Thanks to the highly-resistant granitoid lithology, steep slopes and high reliefs deriving from glacial erosion could be maintained, in turn promoting high millennial to present-day denudation rates in this catchment. Finally, the supply of sediments by retreating glaciers and active periglacial processes, and the contribution of frequent rockfall events triggered by abundant precipitations (Fig. 5A) and present-day permafrost degradation (Ravanel et al., 2010; Akçar et al., 2012; Deline et al., 2015) participate to the significant sediment yield in the DB01 catchment. Catchment DB01 thus supplies material with highly depleted ¹⁰Be concentrations to the river system, which is in turn capable to significantly dilute the ¹⁰Be signal along the DB course (Fig. 3).

5.4 Long- and short-term DB denudation rates

Our ¹⁰Be-derived denudation rates show a general trend of higher millennial denudation rates in the Mont Blanc Massif compared to other DB tributaries (Fig. 2), which is overall in agreement, albeit with different absolute values, with denudation rate estimates on different timescales.

Long-term (10⁶-10⁷ yr) exhumation rates estimated from bedrock apatite fission-track data (Malusà et al., 2005) show higher values (0.4-0.7 km/Myr) in the western sector of the DB catchment (west of Internal Houiller Fault, Fig. 2) than in the east (around 0.2 km/Myr, between the Internal Houiller and the Insubric Faults, Fig. 2). Likewise, results from detrital apatite

fission-track (Resentini and Malusà, 2012) indicate that short-term (10^2 - 10^5 years) denudation rates are higher (around 0.5 mm/yr) in the Mont Blanc External Massif and its sedimentary cover (west of the Penninic Frontal Thrust) than in the axial belt, east of the Penninic Frontal Thrust (around 0.1 mm/yr; Fig. 2). Similarly to what has been shown by Glotzbach et al. (2013), the external Alps catchments (west of the Penninic Frontal Thrust; Fig. 2) appear to have equivalent long-term (apatite fission-track derived) and short-term (^{10}Be -derived) denudation rates, while internal Alps catchments (east of the Penninic Frontal Thrust; Fig. 2) show higher short-term than long-term denudation rates. This has been tentatively explained by potential differences in driver mechanisms of denudation before and during the Quaternary (Glotzbach et al., 2013). Tectonic forcing dominated Neogene denudation rates, with fast exhuming External Massifs having steeper rivers and higher reliefs and therefore eroding faster than the slowly-exhuming Internal Alpine Massifs. During the Quaternary, instead, climate fluctuations and associated glaciations modified both the Internal and External Alps morphology and topographic reliefs, also resulting in increased denudation rates for the Internal Alps. Our ^{10}Be -derived catchment denudation rates for the DB catchment are therefore not totally reflecting long-term exhumation rates over Myr timescales but most probably highlight Quaternary erosion dynamics.




Figure 8: Comparison of ^{10}Be - and modern river yield-derived catchment-wide denudation rates. For all tributary catchments and DB locations, modern denudation rates are derived from river-bedload empirical estimates (Vezzoli et al., 2004; red and light blue dots for DB and tributary catchments, respectively). For catchment DB06 and DB011, also modern denudation rates derived respectively from sediment gauging and sediment trapping are plotted with green diamonds (Hinderer et al., 2013, after Bartolini et al., 1996 and Bartolini and Fontanelli, 2009). Errors are represented only for ^{10}Be -derived denudation rates, since they are not reported for the modern rates.

Modern denudation rates, obtained from sediment-yield estimates (all DB catchments; Vezzoli et al., 2004; based on Gavrilovic empirical formula, Gavrilovic, 1988) and measurements (sediment gauging for DB06 and sediment trapping for DB11; Hinderer et al., 2003, after Bartolini et al., 1996 and Bartolini and Fontanelli, 2009), display higher values for samples along the DB (0.07-0.73 mm/yr) compared to DB tributaries (0.01-0.08 mm/yr). While such a pattern is consistent with our ^{10}Be -derived records, millennial denudation rates are 2 to 50 times greater than modern denudation estimates, with the exception of sample DB01 for which modern and ^{10}Be -derived denudation rates are roughly similar (Fig. 8). Equivalent order of discrepancy between modern sediment-yield and ^{10}Be -derived denudation rates has been observed by several studies (e.g.

515 Kirchner et al., 2001; Schaller et al., 2001; Wittmann et al., 2007, 2016; Stutenbecker et al., 2018; Pitlick et al., 2021). Among
other factors (including the exclusion of non quartz-bearing areas for ^{10}Be denudation rates, see Table 2 and discussion in
section 5.1), this discrepancy was interpreted to point to the separate or combined effects of (1) incorporation of high-
magnitude low-frequency erosion events in the ^{10}Be -derived but not in the sediment-yield denudation rates, (2) contribution
520 of bedload and chemical weathering to ^{10}Be -derived but not to sediment-yield denudation rates, (3) preferential postglacial
erosion of material with low ^{10}Be concentration, increasing ^{10}Be -derived denudation rates, through fluvial linear dissection of
the landscape and subglacial sediment export, (4) sediment traps (e.g. lakes, dams), changing the flux measured by sediment
gauging but less probably the ^{10}Be concentrations which are averaged over longer timescales. The first and third hypotheses
could be the most plausible for our results. Modern denudation rates are potentially not capturing the occurrence of large
sporadic erosional events (Kirchner et al., 2001; Schaller et al., 2001), with the exception of catchment DB01 (and therefore
525 DB02, 06, 10, 12 along the main DB course), where major erosional events have been occurring during the Holocene towards
present-day (i.e. rockfall events; Deline et al., 2012, 2015) and therefore potentially included in the 10^1 - 10^2 yr integration time
of the modern denudation rates. Alternatively, low ^{10}Be -concentration sediment input in the river system, coming from linear
fluvial incision and subglacial sediment export, could explain the mismatch between modern and millennial denudation rates,
with ^{10}Be -derived denudation rates being potentially overestimated (Stutenbecker et al., 2018).

530 **Conclusions**

Our ^{10}Be -derived catchment-wide denudation rates obtained in the Dora Baltea (DB) catchment (western Italian Alps) vary
between 0.2 and 0.9 mm/yr and fit within literature values across the European Alps (Delunel et al., 2020). Correlation of
output denudation rates with topographic, environmental and geologic metrics excludes any significant control of precipitation
and rock uplift on the observed variability in denudation rates within the DB catchment. Our results instead highlight the main
535 influence of catchment bedrock structuration and erodibility (litho-tectonic origin) and resulting topographic metrics on
denudation rate variability among the 13 main tributaries. As previously supposed for some other parts of the Alps, our study
shows that the most resistant lithologies (granite and gneiss) support high-elevation and high-relief catchments where glacial
and slope processes are more intense and denudation rates are higher than in low-elevation/relief catchments, dominated by
“high erodibility” bedrocks (carbonate and terrigenous rocks).

540 This litho-tectonic control on catchment denudation is exemplified by the tributary catchment draining the Mont Blanc Massif,
which has the highest ^{10}Be -derived denudation rate from our dataset and appears as an end-member for most of the investigated
metrics. Located in the long-term actively-uplifting core of the European Alps, the Mont Blanc Massif also experienced intense
Quaternary glaciations which deeply modified the landscape. Steep slopes and high reliefs could be supported by the highly-
resistant granitoid lithology, which in turn have been influencing the millennial to present-day high denudation of the
545 catchment, governed by intense glacial/periglacial processes and recurring rockfall events. In addition, our results also show
that the high sediment input from the Mont Blanc catchment dominates the DB sediment flux, contributing to >77% of the
 ^{10}Be signal carried by river sediments along the DB main river, even downstream of multiple tributary junctions. This suggests
unequal sediment contribution between tributary fluxes along the DB catchment.

Finally, our ^{10}Be -derived denudation rates allow for comparison with long-term (10^6 - 10^7 yr, from thermochronology) and
550 modern (10^1 - 10^2 yr, from sediment budget) denudation rates, showing that, albeit different absolute values, the spatial trend in
catchment denudation is overall in agreement over different timescales, with higher millennial denudation rates in the Mont
Blanc Massif compared to the rest of the DB catchment.

Data availability

The data used in this study is available upon request, supplementary information are available online at [XX](#).

555 **Author contributions**

ES, PGV and RD designed the study. ES, PGV and NG performed field investigations and sample collection. ES performed ^{10}Be cosmogenic sample preparation under supervision of NA, ^{10}Be production/denudation rate calculation and analyses. MC performed ^{10}Be measurements. RD performed ^{10}Be -derived sediment budget calculations. ES wrote the manuscript with input from all co-authors.

560 **Acknowledgements**

The authors warmly thank F. Magrani for the help during fieldwork. J. Krbanjevic is thanked for support for ^{10}Be sample preparation. This study was supported by the Swiss National Science Foundation SNSF (Grant PP00P2_170559) and the French ANR-PIA programme (ANR-18- MPGA-0006). We thank three anonymous reviewers and the Associate Editor S. Mudd for insightful comments that helped to improve the manuscript.

565 **Competing interests**

The authors declare no competing interests.

References

- Akçar, N., Deline, P., Ivy-Ochs, S., Alfimov, V., Hajdas, I., Kubik, P. W., Christl, M. and Schlüchter, C.: The AD 1717 rock avalanche deposits in the upper Ferret Valley (Italy): A dating approach with cosmogenic ^{10}Be , *J. Quat. Sci.*, 27(4), 383–392, doi:10.1002/jqs.1558, 2012.
- 570 Akçar N., Ivy-Ochs S., Alfimov V., Schlunegger F., Claude A., Reber R., Christl M., Vockenhuber C., Dehnert A., Meinert R., and Schlüchter C.: Isochron-burial dating of glaciofluvial deposits: primary results from the Alps. *Earth Surface Processes and Landforms*, 42, 2414–2425, <https://doi.org/10.1002/esp.4201>, 2017.
- Baroni, C., Gennaro, S., Cristina, M., Ivy-ochs, S., Christl, M., Cerrato, R. and Orombelli, G.: Last Lateglacial glacier advance in the Gran Paradiso Group reveals relatively drier climatic conditions established in the Western Alps since at least the Younger Dryas, *Quat. Sci. Rev.*, 255, 106815, doi:10.1016/j.quascirev.2021.106815, 2021.
- 575 Bartolini, C. and Fontanelli, K.: Present versus long term sediment yield to the Adriatic Sea and the reliability of gauging stations data, *Bollettino della Società Geologica Italiana* 128(3), 655–667, doi:10.3301/IJG.2009.128.3.655, 2009.
- Bartolini, C., Caputo, R. and Pieri, M.: Pliocene-Quaternary sedimentation in the Northern Apennine foredeep and related denudation, *Geol. Mag.*, 133(3), 255–273, doi:10.1017/s001675680009006, 1996.
- 580 Bierman, P. and Steig, E.J.: Estimating rates of denudation using cosmogenic isotope abundances in sediment. *Earth Surf. Process. Landforms*, 21(2), 125-139, [https://doi.org/10.1002/\(SICI\)1096-9837\(199602\)21:2<125::AID-ESP511>3.0.CO;2-8](https://doi.org/10.1002/(SICI)1096-9837(199602)21:2<125::AID-ESP511>3.0.CO;2-8), 1996.
- Binnie, S.A., Phillips, W.M., Summerfield, M.A. and Fifield, L.K.: Tectonic uplift, threshold hillslopes, and denudation rates in a developing mountain range, *Geology*, 35(8), 743-746, <https://doi.org/10.1130/G23641A.1>, 2007.
- 585 von Blanckenburg, F.: The control mechanisms of erosion and weathering at basin scale from cosmogenic nuclides in river sediment, *Earth Planet. Sci. Lett.*, 237(3–4), 462–479, doi:10.1016/j.epsl.2005.06.030, 2005.
- Bookhagen, B. and Strecker, M. R.: Spatiotemporal trends in erosion rates across a pronounced rainfall gradient: Examples from the southern Central Andes, *Earth Planet. Sci. Lett.*, 327–328, 97–110, doi:10.1016/j.epsl.2012.02.005, 2012.
- 590 Brocklehurst, S. H. and Whipple, K. X.: Hypsometry of glaciated landscapes, *Earth Surf. Process. Landforms*, 29(7), 907–926, doi:10.1002/esp.1083, 2004.

- Brown, E.T., Stallard, R.F., Larsen, M.C., Raisbeck, G.M. and Yiou, F.: Denudation rates determined from the accumulation of in situ-produced ^{10}Be in the Luquillo Experimental Forest, Puerto Rico. *Earth Planet. Sci. Lett.*, 129(1-4), 193-202, [https://doi.org/10.1016/0012-821X\(94\)00249-X](https://doi.org/10.1016/0012-821X(94)00249-X), 1995.
- 595 Burbank, D.W., Leland, J., Fielding, E., Anderson, R.S., Brozovic, N., Reid, M.R., Duncan, C.: Bedrock incision, rock uplift and threshold hillslopes in the northwestern Himalayas. *Nature* 379, 505-510, doi:10.1038/379505a0, 1996.
- Carretier, S., Regard, V., Vassallo, R., Martinod, J., Christophoul, F., Gayer, E., Audin, L. and Lagane, C.: A note on ^{10}Be -derived mean erosion rates in catchments with heterogeneous lithology: Examples from the western Central Andes, *Earth Surf. Process. Landforms*, 40(13), 1719–1729, doi:10.1002/esp.3748, 2015.
- 600 Champagnac, J. D., Schlunegger, F., Norton, K., von Blanckenburg, F., Abbühl, L. M. and Schwab, M.: Erosion-driven uplift of the modern Central Alps, *Tectonophysics*, 474(1–2), 236–249, doi:10.1016/j.tecto.2009.02.024, 2009.
- Champagnac, J. D., Valla, P. G. and Herman, F.: Late-Cenozoic relief evolution under evolving climate: A review, *Tectonophysics*, 614, 44–65, doi:10.1016/j.tecto.2013.11.037, 2014.
- Charreau, J., Blard, P. H., Zumaque, J., Martin, L. C. P., Delobel, T. and Szafran, L.: Basinga: A cell-by-cell GIS toolbox for computing basin average scaling factors, cosmogenic production rates and denudation rates, *Earth Surf. Process. Landforms*, 44(12), 2349–2365, doi:10.1002/esp.4649, 2019.
- 605 Chittenden, H., Delunel, R., Schlunegger, F., Akçar, N. and Kubik, P.: The influence of bedrock orientation on the landscape evolution, surface morphology and denudation (^{10}Be) at the Niesen, Switzerland, *Earth Surf. Process. Landforms*, 39(9), 1153–1166, doi:10.1002/esp.3511, 2014.
- 610 Christl, M., Vockenhuber, C., Kubik, P. W., Wacker, L., Lachner, J., Alfimov, V. and Synal, H. A.: The ETH Zurich AMS facilities: Performance parameters and reference materials, *Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 294, 29–38, doi:10.1016/j.nimb.2012.03.004, 2013.
- Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica, J.X., Hostetler, S.W., and McCabe, A.M.: The Last Glacial Maximum. *Science* 325, 710–4. <https://doi.org/10.1126/science.1172873>, 2009
- 615 Codilean, A.T., 2006. Calculation of the cosmogenic nuclide production topographic shielding scaling factor for large areas using DEMs. *Earth Surf. Process. Landf.* 31(6), 785-794. <https://doi.org/10.1002/esp.1336>, 2006
- Dal Piaz, G. V., Gianotti, F., Monopoli, B., Pennacchioni, G., Tartarotti, P. and Schiavo, A.: Carta Geologica d'Italia - Foglio 091, Chatillon, , 1–152, 2008.
- Deline P., Akçar N., Ivy-Ochs S., and Kubik P.W.: Repeated Holocene rock avalanches onto the Brenva Glacier, Mont Blanc massif, Italy: A chronology. *Quaternary Science Reviews*, 126, 186-200, <https://doi.org/10.1016/j.quascirev.2015.09.004>, 2015.
- 620 Deline, P., Kirkbride, M. P., Ravello, L. and Ravello, M.: The Tré-la-Tête rockfall onto the Lex Blanche Glacier, Mont Blanc Massif, Italy, in September 2008, *Geogr. Fis. e Din. Quat.*, 31(2), 251–254, 2008.
- Deline, P., Gruber, S., Delaloye, R., Fischer, L., Geertsema, M., Giardino, M., Hasler, A., Kirkbride, M., Krautblatter, M., Magnin, F., McColl, S., Ravello, L. and Schoeneich, P.: Ice Loss and Slope Stability in High-Mountain Regions, Elsevier Inc., 2014.
- 625 Delunel, R., Van der Beek, P.A., Bourlès, D.L., Carcaillet, J. and Schlunegger, F. Transient sediment supply in a high-altitude Alpine environment evidenced through a ^{10}Be budget of the Etages catchment (French Western Alps). *Earth Surface Processes and Landforms*, 39(7), 890-899, <https://doi.org/10.1002/esp.3494>, 2014.
- 630 Delunel, R., van der Beek, P. A., Carcaillet, J., Bourlès, D. L. and Valla, P. G.: Frost-cracking control on catchment denudation rates: Insights from in situ produced ^{10}Be concentrations in stream sediments (Ecrins-Pelvoux massif, French Western Alps), *Earth Planet. Sci. Lett.*, 293(1–2), 72–83, doi:10.1016/j.epsl.2010.02.020, 2010.

- Delunel, R., Schlunegger, F., Valla, P. G., Dixon, J., Glotzbach, C., Hippe, K., Kober, F., Molliex, S., Norton, K. P., Salcher, B., Wittmann, H., Akçar, N. and Christl, M.: Late-Pleistocene catchment-wide denudation patterns across the European Alps, *Earth-Science Rev.*, 211, doi:10.1016/j.earscirev.2020.103407, 2020.
- DiBiase, R. A., Heimsath, A. M. and Whipple, K. X.: Hillslope response to tectonic forcing in threshold landscapes, *Earth Surf. Process. Landforms*, 37(8), 855–865, doi:10.1002/esp.3205, 2012.
- Diolaiuti, G. A., Bocchiola, D., Vagliasindi, M., D’Agata, C. and Smiraglia, C.: The 1975-2005 glacier changes in Aosta Valley (Italy) and the relations with climate evolution, *Prog. Phys. Geogr.*, 36(6), 764–785, doi:10.1177/0309133312456413, 2012.
- Dixon, J. L., Von Blanckenburg, F., Stüwe, K. and Christl, M.: Glaciation’s topographic control on Holocene erosion at the eastern edge of the Alps, *Earth Surf. Dyn.*, 4(4), 895–909, doi:10.5194/esurf-4-895-2016, 2016.
- Dunne, J., Elmore, D. and Muzikar, P.: Scaling factors for the rates of production of cosmogenic nuclides for geometric shielding and attenuation at depth on sloped surfaces, *Geomorphology*, 27(1–2), 3–11, doi:10.1016/S0169-555X(98)00086-5, 1999.
- Glotzbach, C., Van Der Beek, P., Carcaillet, J. and Delunel, R.: Deciphering the driving forces of erosion rates on millennial to million-year timescales in glacially impacted landscapes: An example from the Western Alps, *J. Geophys. Res. Earth Surf.*, 118(3), 1491–1515, doi:10.1002/jgrf.20107, 2013.
- Granger, D.E., Kirchner, J.W. and Finkel, R.: Spatially averaged long-term erosion rates measured from in situ-produced cosmogenic nuclides in alluvial sediment. *The Journal of Geology*, 104(3), 249-257, <https://doi.org/10.1086/629823>, 1996.
- Hinderer, M., Kastowski, M., Kamelger, A., Bartolini, C. and Schlunegger, F.: River loads and modern denudation of the Alps - A review, *Earth-Science Rev.*, 118, 11–44, doi:10.1016/j.earscirev.2013.01.001, 2013.
- Isotta, F. A., Frei, C., Weilguni, V., Perčec Tadić, M., Lassègues, P., Rudolf, B., Pavan, V., Cacciamani, C., Antolini, G., Ratto, S. M., Munari, M., Micheletti, S., Bonati, V., Lussana, C., Ronchi, C., Panettieri, E., Marigo, G. and Vertačnik, G.: The climate of daily precipitation in the Alps: Development and analysis of a high-resolution grid dataset from pan-Alpine rain-gauge data, *Int. J. Climatol.*, 34(5), 1657–1675, doi:10.1002/joc.3794, 2014.
- Kirchner, J. W., Finkel, R. C., Riebe, C. S., Granger, D. E., Clayton, J. L., King, J. G. and Megahan, W. F.: Mountain erosion over 10 yr, 10 k.y., and 10 m.y. time scales, *Geology*, 29(7), 591–594, doi:10.1130/0091-7613(2001)029<0591:MEOYKY>2.0.CO;2, 2001.
- Korup, O. and Schlunegger, F.: Bedrock landsliding, river incision, and transience of geomorphic hillslope-channel coupling: Evidence from inner gorges in the Swiss Alps. *Journal of Geophysical Research: Earth Surface*, 112(F3), <https://doi.org/10.1029/2006JF000710>, 2007.
- Kober, F., Hippe, K., Salcher, B., Ivy-Ochs, S., Kubik, P. W., Wacker, L. and Hählen, N.: Debris-flow-dependent variation of cosmogenically derived catchment-wide denudation rates, *Geology*, 40(10), 935–938, doi:10.1130/G33406.1, 2012.
- Kühni, A. and Pfiffner, O. A.: The relief of the Swiss Alps and adjacent areas and its relation to lithology and structure: Topographic analysis from a 250-m DEM, *Geomorphology*, 41(4), 285–307, doi:10.1016/S0169-555X(01)00060-5, 2001.
- Lupker, M., Blard, P. H., Lavé, J., France-Lanord, C., Leanni, L., Puchol, N., Charreau, J. and Bourlès, D.: ¹⁰Be-derived Himalayan denudation rates and sediment budgets in the Ganga basin, *Earth Planet. Sci. Lett.*, 333–334, 146–156, doi:10.1016/j.epsl.2012.04.020, 2012.
- Malusà, M. G., Polino, R., Zattin, M., Bigazzi, G., Martin, S. and Piana, F.: Miocene to Present differential exhumation in the Western Alps: Insights from fission track thermochronology, *Tectonics*, 24(3), 1–23, doi:10.1029/2004TC001782, 2005.
- Martin, L. C. P., Blard, P.-H., Balco, G., Lavé, J., Delunel, R., Lifton, N. and Laurent, V.: The CREp program and the ICE-D production rate calibration database: A fully parameterizable and updated online tool to compute cosmic-ray exposure ages, *Quat. Geochronol.*, 38, 25–49, doi:10.1016/J.QUAGEO.2016.11.006, 2017.

- 675 Maxeiner, S., Synal, H. A., Christl, M., Suter, M., Müller, A. and Vockenhuber, C.: Proof-of-principle of a compact 300 kV multi-isotope AMS facility, *Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 439(November 2018), 84–89, doi:10.1016/j.nimb.2018.11.028, 2019.
- Montgomery, D.R., and Brandon, M.T.: Topographic controls on erosion rates in tectonically active mountain ranges, *Earth Planet. Sci. Lett.*, 201 (3–4), 481–489, doi:10.1016/S0012-821X(02)00725-2, 2002.
- 680 Moon, S., Page Chamberlain, C., Blisniuk, K., Levine, N., Rood, D. H. and Hilley, G. E.: Climatic control of denudation in the deglaciated landscape of the Washington Cascades, *Nat. Geosci.*, 4(7), 469–473, doi:10.1038/ngeo1159, 2011.
- Muscheler, R., Beer, J., Kubik, P. W. and Synal, H. A.: Geomagnetic field intensity during the last 60,000 years based on ^{10}Be and ^{36}Cl from the Summit ice cores and ^{14}C , *Quat. Sci. Rev.*, 24(16–17), 1849–1860, doi:10.1016/j.quascirev.2005.01.012, 2005.
- 685 Norton, K. P., Abbühl, L. M. and Schlunegger, F.: Glacial conditioning as an erosional driving force in the Central Alps, *Geology*, 38(7), 655–658, doi:10.1130/G31102.1, 2010.
- Norton, K. P., von Blanckenburg, F., DiBiase, R., Schlunegger, F. and Kubik, P. W.: Cosmogenic ^{10}Be -derived denudation rates of the Eastern and Southern European Alps, *Int. J. Earth Sci.*, 100(5), 1163–1179, doi:10.1007/s00531-010-0626-y, 2011.
- Perello, P., Gianotti, F. and Monopoli, B.: *Carta Geologica d'Italia - Foglio 089*, Courmayeur, 2008.
- 690 Pedersen, V.K. and Egholm, D.L.: Glaciations in response to climate variations preconditioned by evolving topography. *Nature*, 493(7431), 206–210, <https://doi.org/10.1038/nature11786>, 2013.
- Polino, R., Malusà, M. G., Martin, S., Carraro, F., Gianotti, F. and Bonetto, F.: *Carta Geologica d'Italia - Foglio 090*, Aosta, (3), 2008.
- Pitlick, J., Recking, A., Liebault, F., Misset, C., Piton, G., and Vazquez-Tarrio, D.: Sediment production in French Alpine rivers, *Water Resources Research*, 57, e2021WR030470. <https://doi.org/10.1029/2021WR030470>, 2021.
- 695 Portenga, E. W. and Bierman, P. R.: Understanding earth's eroding surface with ^{10}Be , *GSA Today*, 21(8), 4–10, doi:10.1130/G111A.1, 2011.
- Ravanel, L., Allignol, F., Deline, P., Gruber, S. and Ravello, M.: Rock falls in the Mont Blanc Massif in 2007 and 2008, *Landslides*, 7(4), 493–501, doi:10.1007/s10346-010-0206-z, 2010.
- 700 Resentini, A. and Malusà, M. G.: Sediment budgets by detrital apatite fission-Track dating (Rivers Dora Baltea and Arc, Western Alps), *Spec. Pap. Geol. Soc. Am.*, 487(08), 125–140, doi:10.1130/2012.2487(08), 2012.
- Sarmiento, J. L., Gruber, N., Brzezinski, M. A. and Dunne, J. P.: High-latitude controls of thermocline nutrients and low latitude biological productivity, *Nature*, 427(6969), 56–60, doi:10.1038/nature02127, 2004.
- Savi, S., Norton, K., Picotti, V., Brardinoni, F., Akçar, N., Kubik, P. W., Delunel, R. and Schlunegger, F.: Effects of sediment mixing on ^{10}Be concentrations in the Zielbach catchment, central-eastern Italian Alps, *Quat. Geochronol.*, 19, 148–162, doi:10.1016/j.quageo.2013.01.006, 2014.
- 705 Schaller, M., Von Blanckenburg, F., Hovius, N. and Kubik, P. W.: Large-scale erosion rates from in situ-produced cosmogenic nuclides in European river sediments, *Earth Planet. Sci. Lett.*, 188(3–4), 441–458, doi:10.1016/S0012-821X(01)00320-X, 2001.
- 710 Schlunegger, F., and Hinderer, M.: Pleistocene/Holocene climate change, reestablishment of fluvial drainage network and increase in relief in the Swiss Alps, *Terra Nova*, 15, 88–95, doi: 10.1046/j.1365-3121.2003.00469.x, 2003.
- Schmid, S.M., Fügenschuh, B., Kissling, E., Schuster, R.: Tectonic map and overall architecture of the Alpine orogen. *Eclogae Geologicae Helvetiae*, 97(1), 93–117, doi: 10.1007/s00015-004-1113-x, 2004.
- Schwanghart, W. and Scherler, D.: Short Communication: TopoToolbox 2 - MATLAB-based software for topographic analysis and modeling in Earth surface sciences, *Earth Surf. Dyn.*, 2(1), 1–7, doi:10.5194/esurf-2-1-2014, 2014.
- 715 Serra, E., Valla, P.G., Gribenski, N., Carcaillet, Deline, P., in press. Post-LGM glacial and geomorphic evolution of the Dora Baltea valley (western Italian Alps). *Quat. Sci. Rev.* <https://doi.org/10.1016/j.quascirev.2022.107446>

- Small, E. E. and Anderson, R. S.: Pleistocene relief production in Laramide mountain ranges, western United States., *Geology*, 26(2), 123–126, doi:10.1130/0091-7613(1998)026<1150:PRPILM>2.3.CO;2, 1998.
- 720 Sternai, P., Sue, C., Husson, L., Serpelloni, E., Becker, T. W., Willett, S. D., Faccenna, C., Di Giulio, A., Spada, G., Jolivet, L., Valla, P., Petit, C., Nocquet, J. M., Walpersdorf, A. and Castellort, S.: Present-day uplift of the European Alps: Evaluating mechanisms and models of their relative contributions, *Earth-Science Rev.*, 190(July 2018), 589–604, doi:10.1016/j.earscirev.2019.01.005, 2019.
- Stutenbecker, L., Costa, A. and Schlunegger, F.: Lithological control on the landscape form of the upper Rhône Basin, Central Swiss Alps, *Earth Surf. Dyn.*, 4(1), 253–272, doi:10.5194/esurf-4-253-2016, 2016.
- 725 Stutenbecker, L., Delunel, R., Schlunegger, F., Silva, T. A., Šegvić, B., Girardclos, S., Bakker, M., Costa, A., Lane, S. N., Loizeau, J. L., Molnar, P., Akçar, N. and Christl, M.: Reduced sediment supply in a fast eroding landscape? A multi-proxy sediment budget of the upper Rhône basin, Central Alps, *Sediment. Geol.*, 375, 105–119, doi:10.1016/j.sedgeo.2017.12.013, 2018.
- 730 Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., da Costa Bechtold, V., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, I., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J. F., Morcrette, J. J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo,
- 735 P. and Woollen, J.: The ERA-40 re-analysis, *Q. J. R. Meteorol. Soc.*, 131(612), 2961–3012, doi:10.1256/qj.04.176, 2005.
- Valla, P. G., van der Beek, P. A. and Carcaillet, J.: Dating bedrock gorge incision in the French Western Alps (Ecrins-Pelvoux massif) using cosmogenic ¹⁰Be, *Terra Nov.*, 22(1), 18–25, doi:10.1111/j.1365-3121.2009.00911.x, 2010.
- Valla, P. G., Shuster, D. L. and Van Der Beek, P. A.: Significant increase in relief of the European Alps during mid-Pleistocene glaciations, *Nat. Geosci.*, 4(10), 688–692, doi:10.1038/ngeo1242, 2011.
- 740 Van den Berg, F., Schlunegger, F., Akçar, N. and Kubik, P.: ¹⁰Be-derived assessment of accelerated erosion in a glacially conditioned inner gorge, Entlebuch, Central Alps of Switzerland. *Earth Surface Processes and Landforms*, 37(11), 1176–1188, <https://doi.org/10.1002/esp.3237>, 2012.
- Vezzoli, G.: Erosion in the Western Alps (Dora Baltea Basin): 2. Quantifying sediment yield, *Sediment. Geol.*, 171(1–4), 247–259, doi:10.1016/j.sedgeo.2004.05.018, 2004.
- 745 Vezzoli, G., Garzanti, E. and Monguzzi, S. Erosion in the western Alps (Dora Baltea basin): 1. Quantifying sediment provenance. *Sedimentary Geology*, 171(1-4), 227-246, <https://doi.org/10.1016/j.sedgeo.2004.05.017>, 2004.
- Wittmann, H., von Blanckenburg, F., Kruesmann, T., Norton, K. P. and Kubik, P. W.: Relation between rock uplift and denudation from cosmogenic nuclides in river sediment in the Central Alps of Switzerland, *J. Geophys. Res. Earth Surf.*, 112(4), 1–20, doi:10.1029/2006JF000729, 2007.
- 750 Wittmann, H., Malusà, M. G., Resentini, A., Garzanti, E. and Niedermann, S.: The cosmogenic record of mountain erosion transmitted across a foreland basin: Source-to-sink analysis of in situ ¹⁰Be, ²⁶Al and ²¹Ne in sediment of the Po river catchment, *Earth Planet. Sci. Lett.*, 452, 258–271, doi:10.1016/j.epsl.2016.07.017, 2016.
- Miller, B. B. and Carter, C.: The test article, *J. Sci. Res.*, 12, 135–147, doi:10.1234/56789, 2015.
- Smith, A. A., Carter, C., and Miller, B. B.: More test articles, *J. Adv. Res.*, 35, 13–28, doi:10.2345/67890, 2014.