

1 **Supplementary material**

2 **Table S1.** Additional analytical details on ^{10}Be measurements of river-sediment samples collected
3 in the present study. Sample locations are reported in Table 1 in the main text.

Sample	Quartz dissolved (g)	Be carrier (mg)	$^{10}\text{Be}/^9\text{Be}$ not blank corrected (10^{-14})	$^{10}\text{Be}/^9\text{Be}$ blank corrected (10^{-14}) ^a	^{10}Be concentration (10^4 at g^{-1})
DB01	50.2785	0.1997	5.16±0.26	4.87±0.25	1.29±0.07
DB02	49.8504	0.1967	4.39±0.25	4.10±0.24	1.08±0.07
DB03	49.7478	0.1977	9.15±0.53	8.85±0.51	2.35±0.14
DB04	49.9469	0.1968	8.64±0.40	8.34±0.38	2.20±0.11
DB05	49.9931	0.1872	8.49±0.40	8.20±0.39	2.05±0.10
DB06	50.1976	0.1939	6.25±0.29	5.96±0.28	1.54±0.08
DB07	20.0443	0.1914	3.82±0.40	3.53±0.37	2.25±0.26
DB08	49.6077	0.1962	18.66±0.80	18.37±0.79	4.85±0.21
DB09	37.7201	0.1975	9.86±0.52	9.56±0.51	3.34±0.18
DB10	50.0416	0.1868	5.69±0.28	5.39±0.27	1.35±0.07
DB11	32.9841	0.1969	9.00±0.50	8.70±0.49	3.47±0.20
DB12	50.1821	0.1903	5.26±0.32	4.97±0.30	1.26±0.08
DB13	49.7747	0.1930	11.07±0.46	10.77±0.45	2.79±0.12
DB14	51.4029	0.1881	11.85±0.53	11.55±0.52	2.83±0.13
DB16	17.8550	0.2000	5.47±0.46	5.18±0.44	3.88±0.35
DB17	49.9150	0.1904	10.94±0.51	10.64±0.50	2.71±0.13
DB18	49.8754	0.1788	9.89±0.49	9.59±0.48	2.30±0.12
DB19	46.7700	0.1975	15.12±0.51	14.83±0.50	4.19±0.15

4

5 ^a Calculated ^{10}Be concentrations were corrected for full process blank $^{10}\text{Be}/^9\text{Be}$ ratio of 2.96

6 $\pm 0.32 \times 10^{-15}$.

7 **Table S2.** Mean catchment ^{10}Be production rates and denudation rates calculated by applying different correction factors (see main text
8 for each correction description). Uncorrected results refer to values obtained by including only mean catchment topographic shielding.
9 The last two columns report the results obtained by applying all the corrections and used in the main text.

Catchment	<i>Uncorrected</i>		<i>Snow correction</i>		<i>Lithology correction</i>		<i>LIA-glacier correction</i>		<i>All corrections</i>	
	Mean production rate (at $\text{g}^{-1} \text{yr}^{-1}$) ^a	Denudation rate (mm yr^{-1}) ^b	Mean production rate (at $\text{g}^{-1} \text{yr}^{-1}$)	Denudation rate (mm yr^{-1})	Mean production rate (at $\text{g}^{-1} \text{yr}^{-1}$)	Denudation rate (mm yr^{-1})	Mean production rate (at $\text{g}^{-1} \text{yr}^{-1}$)	Denudation rate (mm yr^{-1})	Mean production rate (at $\text{g}^{-1} \text{yr}^{-1}$)	Denudation rate (mm yr^{-1})
DB01	29.4	1.45±0.11	24.3	1.19±0.09	30.0	1.47±0.12	17.1	0.85±0.07	13.7	0.68±0.05
DB02	25.1	1.49±0.13	21.2	1.26±0.11	25.1	1.49±0.13	18.3	1.10±0.09	15.3	0.91±0.08
DB03	28.6	0.76±0.06	23.7	0.63±0.05	28.5	0.76±0.06	22.7	0.60±0.05	18.0	0.48±0.04
DB04	27.5	0.78±0.06	22.9	0.65±0.05	26.4	0.75±0.06	21.4	0.61±0.05	17.8	0.51±0.04
DB05	27.8	0.85±0.06	23.1	0.70±0.05	27.2	0.83±0.06	21.9	0.67±0.05	17.4	0.53±0.04
DB06	22.6	0.94±0.07	19.4	0.81±0.06	22.6	0.95±0.07	18.8	0.79±0.06	16.1	0.68±0.05
DB07	22.2	0.61±0.07	19.2	0.53±0.06	22.0	0.61±0.07	18.5	0.51±0.06	15.3	0.42±0.05
DB08	20.2	0.27±0.02	17.5	0.23±0.02	18.4	0.24±0.02	19.9	0.26±0.02	15.9	0.21±0.02
DB09	24.4	0.46±0.04	20.8	0.39±0.03	24.5	0.46±0.04	20.6	0.39±0.03	17.5	0.33±0.03
DB10	23.5	1.12±0.09	20.1	0.96±0.08	23.4	1.12±0.09	19.3	0.93±0.07	16.5	0.79±0.06
DB11	24.5	0.44±0.04	20.8	0.38±0.03	22.1	0.40±0.03	20.0	0.36±0.03	16.0	0.29±0.02
DB12	25.6	1.30±0.11	21.6	1.10±0.10	25.1	1.28±0.11	19.8	1.01±0.09	16.4	0.84±0.07
DB13	24.4	0.56±0.04	20.7	0.47±0.03	24.2	0.55±0.04	20.5	0.47±0.03	17.2	0.40±0.03
DB14	22.1	0.50±0.04	19.0	0.43±0.03	21.9	0.50±0.04	21.7	0.49±0.04	18.5	0.42±0.03
DB16	23.6	0.38±0.04	20.0	0.32±0.03	22.8	0.37±0.04	22.8	0.37±0.04	18.7	0.30±0.03
DB17	27.2	0.63±0.05	22.6	0.52±0.04	25.6	0.59±0.05	19.9	0.46±0.04	17.3	0.40±0.03
DB18	27.0	0.75±0.06	22.5	0.62±0.05	26.9	0.74±0.06	20.9	0.58±0.05	17.2	0.48±0.04
DB19	25.8	0.39±0.03	21.6	0.33±0.02	25.3	0.39±0.03	19.2	0.29±0.02	16.0	0.25±0.02


10 ^a Catchment-averaged ¹⁰Be production rates were calculated with Basinga (Charreau et al., 2019), based on SLHL total ¹⁰Be production rate of
11 4.18±0.26 at g⁻¹ yr⁻¹ (Martin et al., 2017) and the Lal/Stone time-dependent scaling model (Lal, 1991; Stone, 2000).

12 ^b ¹⁰Be-derived catchment denudation rates were calculated with Basinga (Charreau et al., 2019), using default attenuation length values of 160, 4320,
13 and 1500 g cm⁻², for neutrons, fast muons, and slow muons, respectively (Charreau et al., 2019, after Braucher et al., 2011), and assuming a rock
14 density of 2.7 g cm⁻³. Denudation-rate uncertainties are estimated only based on values and relative errors of ¹⁰Be concentrations and cosmogenic
15 production rates from neutron and muons (Eq. 5 in Charreau et al., 2019).

16 **Table S3.** Statistical significance of investigated linear correlations between tributary-catchment
 17 denudation rates and metrics. Correlations were calculated both including (in red) and excluding
 18 (in black) sample DB01 (see text for details). Linear trends were considered significant (bold) if
 19 $p\text{-value} < 0.05$. Some representative correlations are represented in Figures 3 and 4.

Catchment metric	<i>p-value (with DB01)</i>	<i>R² (with DB01)</i>	<i>p-value (no DB01)</i>	<i>R² (no DB01)</i>
Drainage area (km ²)	0.28	0.10	0.13	0.20
Mean elevation (m)	0.01	0.49	0.03	0.37
Mean slope (°)	0.02	0.38	0.10	0.23
Relative abundance of slopes > 40° (%)	0.05	0.28	0.20	0.13
Geophysical relief (5-km, m)	0.00	0.62	0.02	0.39
Hypsometric integral	0.12	0.19	0.61	0.02
Mean annual Precipitation (mm)	0.08	0.23	0.74	0.01
Relative abundance of bare-rock (%)	0.02	0.40	0.02	0.38
Basin area covered by LIA glaciers (%)	0.01	0.43	0.18	0.16
Mean LGM ice- thickness (m)	0.08	0.24	0.11	0.22
Basin area above LGM ELA (2103 m a.s.l.) (%)	0.06	0.27	0.10	0.23
Mean geodetic rock uplift (mm yr ⁻¹)	0.89	0.00	0.92	0.00

20



21
 22 **Figure S1.** Spatial map of lithological/geomorphic correction factors employed for ^{10}Be
 23 production/denudation rate calculation, in addition to topographic and snow shielding. Crystalline
 24 rocks and Quaternary deposits (1/100,000- and 1/250,000-scale digital geological maps from
 25 Regione Autonoma Valle d'Aosta and Regione Piemonte) were considered as quartz-bearing
 26 lithologies and included in catchment production rate calculation. Mafic and sedimentary rocks
 27 were instead excluded based on assumption of no to minor quartz content. Low-lying areas (slope
 28 $< 3^\circ$) were excluded from the calculations (no geomorphic link with stream network; Delunel et
 29 al., 2010). For LIA-glacier correction, ^{10}Be production rates were assumed to be null for areas
 30 covered by LIA glaciers (GlaRiskAlp Project, <http://www.glariskalp.eu>).