



# Multisensor monitoring and data integration reveal cyclical destabilization of Äußeres Hochebenkar Rock Glacier

Lea Hartl<sup>1, 6</sup>, Thomas Zieher<sup>1</sup>, Magnus Bremer<sup>1, 2</sup>, Martin Stocker-Waldhuber<sup>1</sup>, Vivien Zahs<sup>3</sup>, Bernhard Höfle<sup>3, 4</sup>, Christoph Klug<sup>2</sup>, and Alessandro Cicoira<sup>5</sup>

<sup>1</sup> Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences, Innrain 25, 3. OG 6020 Innsbruck, Austria

<sup>2</sup> Department of Geography, University of Innsbruck, Innrain 52f, 6020 Innsbruck, Austria

<sup>3</sup> Institute of Geography, Heidelberg University, Im Neuenheimer Feld 368, 69120 Heidelberg, Germany

<sup>4</sup> Interdisciplinary Center for Scientific Computing (IWR), Heidelberg University, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany

<sup>5</sup> Department of Geography, University of Zurich, Winterthurerstr. 190, 8057 Zurich, Switzerland

<sup>6</sup> Alaska Climate Research Center, Geophysical Institute, University of Alaska Fairbanks, 2156 Koyukuk Dr, Fairbanks, AK 99775, United States

**Correspondence:** Lea Hartl (Lea.Hartl@oeaw.ac.at)

**Abstract.** This study investigates rock glacier destabilization based on the results of a unique in situ and remote sensing-based monitoring network focused on the kinematics of the rock glacier in Äußeres Hochebenkar (Austrian Alps). We consolidate, homogenize, and extend existing time series to generate a comprehensive dataset consisting of 14 digital surface models covering a 68 year time period, as well as in situ measurements of block displacement since the early 1950s. The digital surface models are derived from historical aerial imagery and, more recently, airborne and uncrewed aerial vehicle-based laser scanning (ALS, ULS). Since 2017, high-resolution 3D ALS and ULS point clouds are available at annual temporal resolution. Additional terrestrial laser scanning data collected in bi-weekly intervals during the summer of 2019 is available from the rock glacier front. Using image correlation techniques, we derive velocity vectors from the digital surface models, thereby adding rock glacier-wide spatial context to the point scale block displacement measurements. Based on velocities, surface elevation change, analysis of morphological features, and computations of the bulk creep factor and strain rates, we assess the combined datasets in terms of rock glacier destabilization. To additionally investigate potential rotational components of the movement of the destabilized section of the rock glacier, we integrate in situ data of block displacement with ULS point clouds and compute changes in the rotation angles of single blocks during recent years. The time series shows two cycles of destabilization in the lower section of the rock glacier. The first lasted from the early 1950s until the mid 1970s. The second began around 2017 after approximately two decades of more gradual acceleration and is currently ongoing. Both destabilization periods are characterized by high velocities and the development of morphological destabilization features on the rock glacier surface. Acceleration in the most recent years has been very pronounced, with velocities reaching 20-30m/a in 2020/21. These values are unprecedented in the time series and suggest highly destabilized conditions in the lower section of the rock glacier, which shows signs of translational as well as rotational, landslide-like movement. Due to the length and granularity of the time series, the cyclic destabilization process at Äußeres Hochebenkar rock glacier is well resolved in the dataset. Our study highlights the



importance of interdisciplinary, long-term and continuous, high-resolution 3D monitoring to improve process understanding and model development related to rock glacier rheology and destabilization.

## 1 Introduction

Rock glaciers are lobate or tongue shaped landforms supersaturated by ice and generated by the former or current creep of frozen ground and widespread features of the mountain cryosphere (Whalley, 1974; Barsch, 1992; Haeberli et al., 2006; RGIK, 2022). As such, they serve as proxies for former and current permafrost occurrence, play an important role in the hydrological system, and form part of the sediment cascade (Wahrhaftig and Cox, 1959; Barsch, 1996; Haeberli et al., 2006; RGIK, 2022). Rock glacier kinematics have increasingly become of interest during approximately the last two decades, as accelerating trends became apparent at many rock glaciers in the Alps (Roer, 2005; Delaloye et al., 2008, 2010; Kellerer-Pirklbauer et al., 2018; Fleischer et al., 2021). These general trends are broadly attributed to rising air and ground temperatures, although interannual variations of rock glacier velocities often cannot be explained with variations in mean annual or summer air temperatures alone, indicating a multifaceted and scale-dependent relationship between climate forcing and rock glacier response (Delaloye et al., 2008, 2010; Sorg et al., 2015; Hartl et al., 2016b; Kellerer-Pirklbauer et al., 2017; Müller et al., 2016; Cicoira et al., 2019a; Fleischer et al., 2021). Sub-seasonal monitoring of rock glacier movement has shown correlations between short-term velocity fluctuations and water input from snow melt or precipitation events, indicating that the presence of liquid water within the rock glacier strongly affects the material properties relevant to its rheology and, hence, kinematics (Krainer and He, 2006; Wirz et al., 2016; Kenner et al., 2017; Buchli et al., 2018; Cicoira et al., 2019b, 2020; Fleischer et al., 2021).

In recent years, the term *destabilization* has gained traction in the context of rock glacier acceleration. While there is no universal, exact definition of rock glacier destabilization to date, it is generally understood to mean anomalous “landslide-like behaviour” (Marcer et al., 2021) characterized by a sudden, strong, often localized increase in velocity and the concurrent appearance of morphological destabilization signs like cracks, crevasses, and scarps (Roer, 2005; Roer et al., 2008; Delaloye et al., 2013). Destabilization appears to require favourable terrain, i.e., a relatively steep slope angle, and onset often occurs at convexities or terrain steps (Marcer et al., 2019, 2021). Destabilization signs and velocity discontinuities develop in these areas and the rock glacier is essentially split into a destabilized (usually lower) section and a comparatively unaffected (usually upper) section. Features like crevasses and deep cracks allow rain or melt water to enter the rock glacier, further contributing to acceleration and changes in rheology and kinematics (Roer, 2005; Delaloye et al., 2013; Buchli et al., 2018; Eriksen et al., 2018; Vivero and Lambiel, 2019). In terms of rheology, it is assumed that sliding on shear horizons, which may be basal or located within the structure of the rock glacier, dominates the destabilized movement, while “normal” rock glacier movement is driven by viscous creep of the ice-rich permafrost core (Arenson et al., 2002; Roer et al., 2008; Krainer et al., 2015; Schoeneich et al., 2015; Marcer et al., 2019; Cicoira et al., 2019b, 2020). The underlying causes of destabilization may be climatic (rheological changes due to increased temperature and/or liquid water input, e.g., Delaloye et al. (2013)), mechanical (local overloading due to rock fall events and subsequent propagation of compressive processes, e.g., Scotti et al. (2017)), or a combination thereof.



We follow the terminology of Marcer et al. (2021) regarding rock glacier destabilization and refer to their publication for a more comprehensive overview of the conceptual framework around different phases of destabilization.

55 Relatively little is known about regional distribution of destabilized rock glaciers. Marcer et al. (2019) find evidence of destabilization at 10% of active rock glaciers in France based on topographic data from 2000 to 2013. Interestingly, they note that destabilizing rock glaciers tend to be pebbly as opposed to bouldery (Ikeda and Matsuoka, 2006) and none of the destabilizing rock glaciers identified in their study are in crystalline lithology. Expanding their analysis further into the past, Marcer et al. (2021) identify a small number of rock glaciers in France that experienced a form of destabilization around the middle of the 20th century, then returned to normal behaviour, and began destabilizing once again in the last two to three decades. Case studies have detailed the destabilization of rock glaciers throughout the Alps in Switzerland, France, Italy, and Austria (Avian et al., 2005; Delaloye et al., 2013; Scotti et al., 2017; Vivero and Lambiel, 2019; Kaufmann et al., 2021; Bearzot et al., 2022). While not all accelerating rock glaciers are necessarily destabilized, pronounced or unusual acceleration patterns may hint at destabilization processes at the respective sites. This applies to the Alps as well as other mountain regions, even though related studies do not always explicitly use the term destabilization to describe the observed changes (Delaloye et al., 60 2010; Hartl et al., 2016b; Eriksen et al., 2018; Käab et al., 2021).

In most cases, destabilization is followed by degradation (Cicoira et al., 2020), i.e., deceleration and eventual inactivity of the previously destabilized section of the rock glacier. In rare cases, the complete collapse of the destabilized section followed by a rapid mass movement of substantial parts of the affected rock glacier has been observed. Such events tend to be preceded 70 by wet conditions (significant precipitation events and/or longer periods with anomalous amounts of precipitation, water input through snow melt) and high temperatures and morphological destabilization signs prior to the collapse are reported in cases for which detailed analyses of collapse events exist (Krysiiecki et al., 2008; Bodin et al., 2012, 2017; Marcer et al., 2020; Kofler et al., 2021). In some cases, e.g., Grabengufer rock glacier in Switzerland, the combination of very high velocities, obvious signs of destabilization, and downstream topography favourable for mass movements led to concern but did not produce debris 75 flows (Delaloye et al., 2013). This suggests that the underlying processes are complex and depend on a variety of internal (composition of the rock glacier, internal hydrological processes) and external (meteorological and climatological forcing, surrounding terrain, ...) factors (Marcer et al., 2019, 2021).

### 1.1 Äußeres Hochebenkar rock glacier

In the following, we present data from an interdisciplinary monitoring network at the rock glacier in Äußeres Hochebenkar 80 (HEK) in the Austrian Alps (Fig. 1), and discuss the behaviour of the lower section of the rock glacier through the lens of destabilization. HEK is a fast moving, tongue-shaped talus rock glacier with a long history of scientific study (Table 1). It is one of 556 intact rock glaciers in the Ötztal mountain range (Wagner et al., 2020) and covers an area of about 0.4 km<sup>2</sup>. The root zone reaches a maximum elevation of about 2870 m and the terminus currently extends down to about 2370 m. The rock glacier flows over a terrain step at around 2570 m, below which the terrain steepens. The terminus funnels into a drainage gully, 85 which is located directly above an access road leading to mountain huts further up-valley. The road has recently been affected by rock fall from the terminus and is threatened by any further destabilization.



Using refraction seismics, Haeberli and Patzelt (1982) found the mean thickness of the rock glacier to be about 40 m and estimated an ice content of 50%. More recent ground penetrating radar (GPR) surveys found comparable values for the mean depth of the bedrock, with considerable variation in depth throughout the rock glacier area, and a several metres (up to >10 m) thick surface layer of ice-free, coarse debris (Nickus et al., 2015; Hartl et al., 2016a). The surface debris has an average grain size of 35 cm to about 60 cm, with some blocks reaching diameters of up to a few metres (Nickus et al., 2015). A layer of finer material below the coarse surface debris is exposed at the steep frontal slope of the rock glacier. The surrounding lithology is part of the crystalline Ötztal-Stubai complex and the bedrock is composed mainly of paragneiss and mica shists (Nickus et al., 2015).

Surveys of the temperature at the bottom of the winter snow cover (BTS) carried out in 1976 (Haeberli and Patzelt, 1982) and 2010 indicate a reduction in the extent of the discontinuous permafrost surrounding the rock glacier during this period, particularly at the margins of the rock glacier and directly below the terminus (Nickus et al., 2015). The rock glacier terminus was at the lower end of the discontinuous permafrost margin even in the 1970s and has advanced downwards since, hence moving further below the permafrost margin. Zahs et al. (2019) found no permafrost in an electrical resistivity tomography (ERT) profile beside the rock glacier at an elevation of about 2470 m and reported isolated ice-lenses in a profile at similar elevation on the rock glacier terminus.

Regular measurements of block displacements on the surface of HEK rock glacier began in the early 1950s and continue to this day. Since the 2000s, numerous remote sensing-based studies have used a variety of methods and analysis techniques to quantify surface elevation change and horizontal displacement at HEK, providing spatial context to the in situ monitoring of block movement. See Table 1 for an overview of the respective literature. Summarising broadly, previous studies show that vertical surface elevation changes are typically small or slightly negative in the upper part of the rock glacier, while the lower section is extending downwards and thinning.

Morphological observations in early studies and the long time series of block velocities indicate processes of destabilization at HEK in the 1950s and '60s: During surveys in the 1970s, Haeberli and Patzelt (1982) observed large crevasses in the lowest section of the rock glacier tongue. They noted one crevasse crossing the entire width of the rock glacier at around 2540 m and speculated that the lowest section of the terminus below this crevasse had previously undergone a process of separation or decoupling from the upper part. Schneider and Schneider (2001) interpreted the kinematics of the lower section of the rock glacier in a similar manner, agreeing with the idea that the terminus area had separated from the main body of the rock glacier, which they describe as being in a normal state (“*Normalzustand*”) and healthy (“*gesund*”), whereas the behaviour of the presumably separated part is considered extraordinary (“*außergewöhnlich*”). They speculated that the process of separation of the entire section below the terrain step was completed by the early 1970s, and that it might repeat in the future given continued advancement of the rock glacier over the terrain step. The study further suggests that the lowest section of the terminus became inactive after this destabilization phase due to the loss of ice-rich permafrost in the terminus area. Schneider and Schneider (2001) attributed the anomalous behaviour in the lower section to the steep slope angle and basal gliding processes as opposed to the permafrost creep governing the upper section, describing the same general destabilization processes as more recent work (Roer et al., 2008; Schoeneich et al., 2015; Marcer et al., 2019; Cicoira et al., 2020).



After this early period of high velocities and destabilization, the rock glacier returned to its “normal” state until the onset of a second phase of acceleration in the mid-1990s. The lower section below the terrain step and the crevasse observed by Haerberli and Patzelt (1982) also accelerated again with a peak in velocities in 2004 and a stronger, currently ongoing acceleration after  
125 2010 (Hartl et al., 2016b).

## 1.2 Objectives

Our overarching goal is to contribute to the developing scientific discussion around rock glacier destabilization by making available and, for the first time, consolidating the in situ and remote sensing data basis from our study site. We provide a comprehensive overview of two separate destabilization periods at the same rock glacier by combining the most recent multi-  
130 sensor and multi-method monitoring results with data from previous studies and long-term observations (Table 1). Specifically, we

- Homogenize and update a time series of 14 digital surface models (DSMs) derived from aerial imagery (Klug, 2011), and LiDAR point clouds, also referred to as laser scanning (time series: 1953-2021).
- Present the most recent data from the long-term in situ time series of DGNS-based block displacement, updating the  
135 dataset presented in Hartl et al. (2016b) (time series: 1952-2021).
- Compute a time series of the bulk creep factor (BCF) as a metric of destabilization (Cicoira et al., 2020) to aid interpretation of velocity change
- Extract additional information about the rock glacier’s recent surface change from very high-resolution (cm point spacing) 3D point clouds. We derive (a) 3D change information between two epochs based on corresponding planar boulder  
140 faces at the rock glacier front (Zahs et al., 2022a, b) and (b) a time series of rotating single blocks.

Following an overview of the different datasets and methods, the results are structured chronologically, moving from the first period of destabilization to the subsequent “stable” period and the start of the ongoing renewed acceleration and destabilization. For each period, we describe kinematic and morphological changes based on the in situ and remote sensing datasets. For the most recent years, we present additional results based on the fusion of high-resolution laser scanning data and block  
145 displacement, as well as data on sub-seasonal 3D topographic change at the rock glacier front for the summer of 2019. In the discussion, we then consider uncertainties and limitations related to the different datasets and methods, and offer an interpretation of the observed changes in the general context of rock glacier destabilization, and specifically in terms of transitions between normal behaviour and different phases destabilization.

Table 1: Overview of previous studies at HEK rock glacier, grouped by thematic focus and listed chronologically per thematic group.

### Interdisciplinary overview publications

---



|  |  |
|--|--|
| Permafrost mapping based on BTS survey, descriptions of morphological features on the rock glacier surface, refraction seismics.   | Haeberli and Patzelt (1982)  |
| Geological setting, BTS data, presentation of first GPR survey, stream flow data and analysis of conductivity and chemical composition of rock glacier runoff.   | Nickus et al. (2015)   |
| <b>Displacement of block profiles</b>  |  |
| Details on establishment of block profiles and first 2 decades of measurements.  | Vietoris (1958, 1972); Pillewizer (1957)   |
| Digitisation of early data, homogenisation of time series, systematic assessment of mean profile velocities and single blocks, analysis of surface elevation change at the block locations.  | Schneider (1999a, b)   |
| Overview of block movement since beginning of measurements and consolidation of results of Schneider (1999a, b); considerations on role of climate parameters as drivers of block movement.  | Schneider and Schneider (2001)   |
| Update to time series of block profiles including profile 0 (established in 1997). Statistical analysis of correlation between cumulative anomalies of movement and annual and seasonal means of climate parameters.                                   | Hartl et al. (2016b)   |
| <b>Remote sensing</b>  |  |
| Terrestrial photogrammetric surveys of the rock glacier tongue (Advance of 1.1m between 1953-55 at the main, orographic left lobe); No movement detected at the smaller, orographic right lobe for 1956-66 (unpublished data cited in Vietoris (1972)) | Vietoris (1972)  |
| Terrestrial photogrammetric surveys of the lower section of the rock glacier in 1986, 1999, 2003, 2008, and detailed analysis thereof. Computation of 3D flow vectors. Overview and analysis of historic cartographic and photogrammetric data.        | Kaufmann and Ladstaedter (2002); Kaufmann and Ladstaedter (2002); Ladstaedter and Kaufmann (2005); Kaufmann (2012) |
| Analysis of orthophotos to generate digital surface models (DSM) and horizontal flow vectors (orthophotos available for: 1953, 1969, 1971, 1977, 1990, 1997)   | Klug (2011); Klug et al. (2012)  |
| Analysis of airborne laser scanning (ALS) data to generate DEM and flow vectors (ALS data for 2006, 2009, 2010, 2011)  | Bollmann et al. (2012); Klug et al. (2012)   |



|   |  |
|---|--|
| Multi-source (ALS, terrestrial laser scanning (TLS), UAV-borne laser scanning (ULS), and UAV-borne photogrammetry-based dense image matching (DIM)) 3D topographic change analysis for different time spans in the period 2006-2021, including bi-weekly monitoring in summer 2019. | Zahs et al. (2019); Ulrich et al. (2021); Zahs et al. (2022b, a) |
| <b>Geophysics</b>   |  |
| Refraction seismics: Surveys carried out in 1975 and 1977. Mean ice content of about 50% at surveyed profiles.  | Haeberli and Patzelt (1982)                                      |
| Ground penetrating radar: Surveys carried out in 2000, 2008, 2013 with two different radar systems at different locations on the rock glacier. Mean depth of bedrock between 34 and 45m depending on survey year.   | Nickus et al. (2015); Hartl et al. (2016a)                       |
| Electrical resistivity tomography: Survey carried out in 2016 at two profile lines next to and on the margin of the rock glacier tongue. No indications of permafrost beside the rock glacier, isolated ice lenses identified in the profile on the tongue.                         | Zahs et al. (2019)   |

## 2 Data and methods

### 150 2.1 Long-term geodetic surface displacement monitoring

Since 1954, surface displacement of HEK rock glacier has been measured geodetically along three cross profiles (Vietoris, 1958, 1972). See Fig.1 for an overview of the profile locations. Measurements were initially carried out annually but the time series has substantial gaps from the 1960s until 1997, when a fourth cross profile and a longitudinal profile were installed in the lowest section and the monitoring efforts were revitalized. Schneider (1999a, b); Schneider and Schneider (2001), and  
 155 Niederwald (2009) give details on the homogenization of the time series and early measurement techniques. Since 2008, displacement measurements have been carried out with a Topcon HiperPro real-time kinematic differential global navigation satellite system (DGNSS), replacing a theodolite and tachymeter previously used for the same purpose. Hartl et al. (2016b) describe the current measurement system and give an overview of the kinematics of HEK rock glacier up to 2015. Historically, reporting focused on mean displacement values for the cross profiles. The mean value was computed as the arithmetic mean  
 160 of all blocks in the profile and displacement refers to the absolute, three-dimensional distance moved. Given issues related to averaging and changing numbers of blocks in the profiles, Hartl et al. (2016b) estimate an uncertainty of between  $\pm 0.2$  m/a and  $\pm 0.5$  m/a for the mean profile velocities. We adhere to the same method of using 3D trajectories of single blocks as part of a profile when referring to profile means in order to be consistent with the long-term time series. Measurements are carried out annually in summer, usually in late August or early September. Annual mean values are derived from the absolute displacement  
 165 and the number of days between measurement campaigns. The tachymeter and DGNSS displacement measurements for single blocks are considered accurate to  $\pm 3$  cm vertically and horizontally (Niederwald, 2009; Nickus et al., 2015). We refer to the



existing literature for further details on the time series of block displacement. (Tab. 1; in particular Schneider and Schneider (2001); Niederwald (2009); Hartl et al. (2016b)).

170 The dynamics of destabilized rock glaciers are characterized by velocity discontinuities between the faster, destabilized (usu-  
ally lower) section and the slower, non-destabilized (usually upper) section. As this discontinuity becomes more pronounced  
under ongoing destabilization, the surface strain between the two sections also increases (Marcer et al., 2021). Hence, changes  
in surface strain rates can serve as indicators of destabilization onset in specific sections of a rock glacier. To quantify these  
changes for recent years at HEK, we use the positions of individual blocks in P1 and P2 to compute surface strain rates across  
the terrain step and the velocity discontinuity located between P1 and P2, i.e., between the currently destabilized lower section  
175 and the non-destabilized upper section. The surface strain rate between a pair of blocks (b1, b2) in P1 and P2 is given by the  
difference in velocity between b1 and b2 divided by the distance between b1 and b2, following Marcer et al. (2021).

As part of the HEK monitoring network, an automatic weather station (AWS) was installed at 2565 m.a.s.l. in 2010 di-  
rectly beside the rock glacier (Stocker-Waldhuber et al., 2013). (Note: Unless otherwise specified, all elevation values in this  
manuscript refer to orthometric heights; EVRF2000 Austria height, EPSG:9274). Long-term meteorological data is available  
180 from a semi-automatic weather station in Obergurgl. This station is located about 4 km down-valley from the rock glacier at  
1941 m.a.s.l. and data is available from the data portal of the Austrian Central Institution for Meteorology and Geodynamics  
(ZAMG), which operates the station.


## 2.2 Remote-sensing-based area-wide monitoring of topographic change

### 2.2.1 Long-term change monitoring

185 Photogrammetrically reconstructed surface topography based on historical aerial imagery (Klug, 2011) is available for multiple  
timesteps from 1953 until 1997. In more recent years, various airborne (ALS), terrestrial (TLS) and UAV-based laser scanning  
(ULS) surveys were carried out at HEK. Table 2 gives an overview of all topographic data used for this study, which includes  
14 DSMs covering a time period of 68 years, as well as orthophotos from UAV surveys and TLS-based point clouds of the  
front and lower terminus area.

190 The photogrammetrically reconstructed topography and the 3D point clouds derived from ALS and ULS were co-registered  
within bedrock outcrops assumed to be stable throughout the established time series. For this purpose, the iterative closest  
point algorithm (ICP, Besl and McKay (1992)) was used, with the 2017 ALS data (Table 2) serving as reference. All datasets  
are projected according to the Austria GK West definition (EPSG 31254, with a vertical datum of ...). Digital surface models  
(DSMs) with a spatial resolution of 1m were computed for each epoch of the multi-temporal point clouds, aggregating the aver-  
195 age elevation of all points per raster cell. For deriving area-wide displacement vectors based on an image correlation technique  
(IMCORR, Scambos et al. (1992)), shaded reliefs were computed from the DSMs with the ambient occlusion method (Tarini  
et al., 2006), preventing cast shadows which may hinder pattern matching. The employed IMCORR algorithm, implemented  
in SAGA GIS (Conrad et al., 2015), included the DSMs for adding the vertical displacement component, resulting in 2.5D  
vectors.





**Figure 1.** Location of the study area in the Austrian Alps (top right) and detail of the geographical context of the Hochebenkar Rock Glacier in the Ötztal Alps (top left). 3D visualization of the study site (data from SRTM and © GoogleEarth, bottom left) and slope angle above hillshade generated from the 2017 DSM (bottom right). Black dots mark the positions of the blocks in the 4 cross profiles and the longitudinal profile for 2015–2021. Also shown are the central flowline and the lines between the fixed reference points that define the positions of the cross profiles.

200 The image correlation was conducted at points in a regular grid-like pattern with a distance of 5 m. From the resulting displacements, mean annual velocities (m/a) between the individual epochs were calculated by dividing the 2.5D lengths by the respective time period. The vectors were then scaled according to the mean velocity and filtered semi-automatically, only considering downslope movement. In the vicinity of the monitored blocks the mean velocities derived from the image correlation analysis were spatially aggregated to allow comparing the results with those of the DGNS monitoring. To assess  
205 the uncertainty of the derived velocities, the individual directional components of the pseudo-vectors were analysed within stable bedrock outcrops near the rock glacier, following a similar approach as, e.g., Fleischer et al. (2021). Arbitrary directions of the vectors within the stable area reflect sufficient quality of the data and its registration, while a non-random distribution indicates higher errors. Additionally, differential digital surface models (DDSMs) were computed by subtracting the DSMs of two consecutive epochs (Williams, 2012) to show patterns of elevation gain and loss. DDSM uncertainty was assessed  
210 by computing the 2.5% and 97.5% quantile within stable bedrock outcrops, providing an estimate of the inherent noise and




constituting the detection threshold for obtaining significant surface changes beyond measurement noise (Table 2). The analyses of the multi-temporal DSMs and the displacement vectors were conducted with the R statistical programming language (R Core Team, 2021).

### 2.2.2 Recent high-resolution monitoring of 3D topographic change

215 In 2018, 2019, 2020 and 2021, ULS campaigns were conducted with a Riegl RiCopter UAV, carrying a VUX-1LR kinematic laser scanner, run with a 336° field of view and combined with an Applanix AP20 inertial measurement unit (IMU) (Bremer et al., 2019). Additionally, a Nadir-looking Sony Alpha 6000 RGB camera was mounted for additional orthophoto creation. In 2018, only the lower terminus area was captured. The following flight campaigns were extended to the middle part of the rock glacier. All acquisitions followed the same flight plan: Aside from small connection lines, the rock glacier was captured  
220 by a set of parallel flight lines with a horizontal spacing of 100 m, oriented perpendicular to the flow direction of the rock glacier. The average flying height above ground level (AGL) was between 70 m and 120 m. The flight speed was 8 m/s, the pulse repetition rate (PRR) was 820 kHz and the angular resolution was 0.0476°. Following standard procedures such as flight trajectory post-processing, point extraction, geo-referencing and strip-adjustment, the resulting point clouds were co-registered to the ALS 2017 dataset by using the ICP algorithm to minimize distances between the point clouds in stable areas of bedrock  
225 outcrops. Additionally, 0.1 m and 1 m resolution DSMs were created. For both resolutions, image correlation (IMCORR) and DDSMs were analysed.

To quantify sub-seasonal changes at the rock glacier front and lower terminus, a bi-weekly (June 24 to August 30) time series of six terrestrial laser scanning (TLS) point clouds was captured in 2019. This temporally highly resolved dataset complements an annual TLS time series starting in 2015 (Ulrich et al., 2021; Zahs et al., 2022a). Bi-weekly change in 2019 was computed  
230 using the correspondence-driven plane-based M3C2 (CD-PB M3C2) algorithm (Zahs et al., 2022b). The CD-PB M3C2 is tailored to quantifying small-magnitude (<0.1 m) 3D topographic change with low uncertainties and, thus, allows confident detection of change in natural landscapes even with complex surface dynamics, i.e. locally planar but overall rough morphology, and changes of low magnitudes. The algorithm quantifies 3D change between two epochs  $t_1$  and  $t_2$  in terms of the distance between corresponding planar areas (plane pairs). Change is thereby computed along the normal vector of the plane in  $t_1$ . When  
235 applied to rock glaciers, the method can make use of the surface of a rock glacier being most planar at the scale of faces of individual boulders and of these boulders moving with the general creep of the rock glacier rather than independently (Ulrich et al., 2021). Moreover, boulders can be re-identified in successive epochs due to their moderate magnitudes of movement at the monitored interval (Ulrich et al., 2021). We used these corresponding planar boulder faces for change analysis of the rock glacier front and lower terminus between the epochs considering plane-based change in flow direction of the rock glacier,  
240 in vertical direction, and in horizontal direction. The CD-PB M3C2 additionally estimates the uncertainty associated with quantified change and therefore allows confident analysis of change by separating significant change (magnitude of change > uncertainty) from non-significant or no change (magnitude of change  $\leq$  uncertainty).



**Figure 2.** Principle of block shape matching in the ULS point clouds (grey dots) between consecutive time steps (2018-2021). The iterative closest point algorithm (ICP) was used to match the shape of a block in one epoch onto its shape in another epoch. A rotation in the opposite direction of the translation can be recognized for the block shapes. The figure shows examples of downslope translation and upslope rotation.

### 2.2.3 Data fusion approach to generate time series of block rotation

The movement of destabilized rock glaciers is described as landslide-like and may have translational as well as rotational components (Buchli et al., 2018; Marcer et al., 2021). While translational movement is relatively well documented in kinematic  
245 rock glacier monitoring, few data on rotational movement are available. To assess potential rotational movement in the recently destabilized section of HEK, we analyzed the rotational movement of individual blocks in the profile lines. In 2021, the DGNSS-measurements of block profiles and the ULS campaign were conducted only three days apart. This temporal proximity of the measurements made it possible to identify individual blocks from the profiles in 3 cm resolution ortho-images generated  
250 from the 2021 UAV data. Unique block identifiers were manually assigned to the distinct block shapes in the respective ULS point cloud. For each of these selected point ID groups, the following analyses were carried out:

1. We applied an IMCORR image correlation at the centre position of a given block in the 2021 data using the 0.1 m shaded reliefs of 2021 and 2020, in order to reconstruct the position of the block in the previous epoch (2020). Starting from the resulting 2020 position, the same procedure was repeated to find the 2019 and 2018 positions. For each interval, this led  
255 to three shift parameters describing the estimated block movement between two epochs.
2. Considering this translation for the initialization of a 4-by-4 transformation matrix, we used the ICP algorithm on a block-by-block basis in order to optimize the alignment of the block shapes between two consecutive epochs. This was done by a 6-parameter transformation optimizing translational and rotational components. The resulting 4-by-4 transformation matrix describes the full-3D transformation of a block (assumed to be a rigid body).
- 260 3. Finally, the derived matrix allowed decomposing both the translational and rotational components of the transformation. For better interpretation, the rotational components were derived relative to the translation: rotation in the same or opposite direction of translation and rotation around the translation vector (Fig. 2).



### 2.3 Bulk creep factor

To interpret the observed rock glacier velocities from a dynamic perspective, we computed the Bulk Creep Factor (BCF) as described by Cicoira et al. (2020). This dimensionless parameter provides a quantitative basis for the analysis of the rheological properties of the rock glacier material, disentangling the geometrical component from the velocity signal. The calculation of the BCF is based on a modified version of Glen’s flow law (Glen, 1955) adapted for rock glaciers (Arenson and Springman, 2005; Arenson et al., 2002; Cicoira et al., 2020), in which the strain rates are a function of the rock glacier’s slope angle, thickness, and mechanical properties. In general terms, high BCF values (roughly above 10) indicate destabilized rock glaciers, with anomalous short-term deformation processes in the shear horizon dominating over long-term secondary creep in the permafrost material. However, the BCF includes both components (ice-rich core and shear horizon) and remains a proxy value for the heterogeneous properties of rock glacier material, which should be discussed in detail case by case.

We computed the BCF for the four cross profiles shown in Fig. 1 using the time series of the respective mean profile velocities as obtained through the block displacement measurements (Section 2.1). The slope angle at each profile is given by the mean angle along the line between the fixed points that define the respective profile. The slope angle was extracted from the DSM time series data (Section 2.2.1) resampled to a 10 m x10 m grid in order to avoid variability at the scale of surface features (such as furrows and ridges or single blocks), which are not representative of the rock glacier geometry with regards to dynamics. The slope angle between DSM epochs was linearly interpolated for years in which DGNSS block measurements are available but DSMs are not. Rock glacier thickness is given by a map of the rock glacier’s bedrock extrapolated from GPR data and presented in Hartl et al. (2016a). For parameter calibration, we used the same values suggested in Cicoira et al. (2020), which were calibrated for a dataset of rock glaciers mostly in the French Alps. This approach allows us to directly compare our data with their results.

Table 2: Metadata for the presented topographic data.

| Long-term DSM time series |             |                  |   |   |           |
|---------------------------|-------------|------------------|---|---|-----------|
| Acquisition date          | Data source | Spatial coverage | DDSM uncertainty (values refer to the DDSM pair of the given and the previous year, in m/a) | Median velocities in stable areas in x, y, z direction (values refer to the DSM pair of the given and the previous year in m/a) | Reference |
|                           |             |                  |   |   |           |



|            |   |                        |               |                      |  |
|------------|---|------------------------|---------------|----------------------|--|
| 1953-08-31 | Aerial photographs (analogue aerial stereoscopic pairs, Federal Office of Metrology and Surveying-BEV)                            | Entire glacier<br>rock |               |                      | Klug (2011)  |
| 1971-08-18 | Aerial photographs (analogue aerial stereoscopic pairs, BEV)  | Entire glacier<br>rock | -0.07 / +0.08 | -0.01 / 0.04 / -0.03 | Klug (2011)  |
| 1977-09-07 | Aerial photographs (analogue aerial stereoscopic pairs, BEV)  | Entire glacier<br>rock | -0.21 / +0.17 | -0.03 / -0.16 / 0.03 | Klug (2011)  |
| 1990-10-10 | Aerial photographs (analogue aerial stereoscopic pairs, BEV)  | Entire glacier<br>rock | -0.11 / +0.12 | 0.01 / 0.03 / -0.01  | Klug (2011)  |
| 1997-09-11 | Aerial photographs (analogue aerial stereoscopic pairs, BEV)  | Entire glacier<br>rock | -0.18 / +0.21 | -0.01 / 0.05 / -0.01 | Klug (2011)  |
| 2006-08-23 | ALS flight campaign, government of Tyrol  | Entire glacier<br>rock | -0.19 / +0.13 | 0.04 / -0.01 / 0.02  | Land Tirol<br>Abteilung<br>Geoin-<br>formation<br>(2011, 2019) |
| 2009-09-09 | ALS flight campaigns carried out within the ACRP (Austrian Climate Research Programme) project C4AUSTRIA (project no:384 A963633) | Entire glacier<br>rock | -0.13 / +0.18 | -0.01 / 0.00 / 0.00  | Bollmann et al. (2012); Klug et al. (2012)                     |
| 2010-10-09 | ALS flight campaigns carried out within the project MUSICALS of the Centre of climate change adaption strategies(alpS)            | Entire glacier<br>rock | -0.33 / 0.31  | 0.01 / -0.09 / 0.02  | Roncat et al. (2013a, b)                                       |



|            |   |                     |               |                       |   |
|------------|---|---------------------|---------------|-----------------------|---|
| 2011-10-03 | ALS flight campaigns carried out within the ACRP (Austrian Climate Research Programme) project C4AUSTRIA (project no:384 A963633) | Entire rock glacier | -0.33 / +0.35 | -0.01 / 0.06 / -0.01  | Bollmann et al. (2012, 2015); Zahs et al. (2019)          |
| 2017-09-15 | ALS flight campaign, government of Tyrol  | Entire rock glacier | -0.06 / +0.06 | 0.01 / 0.00 / 0.00    | Land Tirol Abteilung Geoinformation (2011); Rieger (2019) |
| 2018-07-30 | ULS   | Terminus            | -0.16 / +0.28 | 0.07 / 0.15 / -0.02   | This study  |
| 2019-08-30 | ULS   | Lower section       | -0.21 / +0.11 | 0.00 / -0.04 / -0.01  | This study  |
| 2020-09-18 | ULS   | Lower section       | -0.09 / +0.08 | 0.00 / -0.02 / -0.01  | This study  |
| 2021-08-13 | ULS   | Lower section       | -0.07 / +0.16 | -0.04 / -0.01 / -0.01 | This study  |

**Other data**

|                                  |                  |               |   |   |                        |
|----------------------------------|------------------|---------------|---|---|------------------------|
| 2019, biweekly during the summer | TLS              | Terminus      | Alignment error between point clouds: 0.011 m - 0.013 m (*) | / | Zahs et al. (2022b, a) |
| 2020-09-18                       | UAV ortho photos | Lower section | /   | / | This study             |
| 2021-08-13                       | UAV ortho photos | Lower section | /   | / | This study             |

\*Alignment error is assessed by calculating the standard deviation of M3C2 distances on stable bedrock outcrops distributed around the rock glacier (Fey and Wichmann, 2017; Zahs et al., 2022a)



### 3 Results

#### 285 3.1 Historical timeline of terminus destabilization

The first result we focus on is an update to the time series of surface displacement at HEK and the homogenization and extension of the DSM time series for the site, which now covers a time period of 68 years. In the following, we present analyses of the multi-temporal DSMs alongside the block displacement measurements. The early data shows a destabilization phase of the rock glacier from onset to deceleration, with a peak in the 1960s (Section 3.1.1). This phase is followed by a period of relative stability, which includes the onset of renewed acceleration in the mid-1990s (Section 3.1.2). Detailed data on the block profiles for the early years of the time series have been presented in previous studies (Schneider, 1999a; Schneider and Schneider, 2001; Hartl et al., 2016b). We include the mean profile velocities here again in brief to contextualise the in situ data with the multi-decadal DSM time series and morphological observations based on the DSMs. The high-frequency, high-resolution 3D data collected in the past 5 years are presented separately in Section 3.2.

##### 295 3.1.1 First period of destabilization: 1953 to 1977

From 1956/57 onwards, the rock glacier entered a period of acceleration. Considering the block profiles (Fig. 3), P1 and P2 arguably showed irregular behaviour in the few available years prior to that, but this signal is less clear and cannot be contextualised historically. Velocity vectors derived from the 1953 and 1971 DSMs reach values of 1-2 m/a in roughly the lower half of the rock glacier, up to the area between P2 and P3 (visible in the velocity vector maps in Fig. 4). It should be noted that this DSM pair does not resolve the terminus well and likely does not capture the full range of velocities in the lowest part of the rock glacier. In 1971-77, velocity vectors of more than 5 m/a were recorded at the terminus. In the upper part of the rock glacier, the measurements show a decrease in velocity compared to the 1953-71 period. The 1971-77 DSM pair shows that the fastest moving part of the rock glacier was below P1 during this time (Fig. 4).

Comparing the velocities obtained from the DSMs with the mean block profile velocities, the velocity of profile P1 seems low in comparison with the maximum values of the velocity vectors because the block profiles did not capture changes in the lowest section of the terminus. The highest mean profile velocities during this first period of acceleration were recorded in 1961/62 at P1 and 2. Single blocks reached a maximum velocity of 6.6 m/a at P1 and 2.2 m/a at P2 in this measurement year (Schneider, 1999a). The next measurement was carried out in 1970. Between these two measurements, mean profile velocities decreased from 3.9 m/a to just under 1.8 m/a for P1, and from 1.5 m/a to 1 m/a at P2 (Schneider, 1999a). P3 experienced a slight increase in velocity at the same time as the lower two profiles, as well as a slight decrease after 1969/70, but changes were minor and, in contrast to P1 and P2, do not clearly stand out from the later years of the time series. Similarly, both P1 and P2 show large BCF values during the 1960s and early 1970s (12.8 for P1), but BCF at P3 was only marginally higher during the period of accelerated movement than during the subsequent, more stable period (Fig. 3).

From a geomorphological perspective, the first signs of destabilization are already visible in the earliest data. In the 1953 DSM, an isolated scarp can be seen at around 2580 m.a.s.l. (close to zone "A" in Fig. 5 and Fig. 6). In the 1971 DSM, the scarp is a few tens of metres further downhill and has notably increased its size in the centre of the rock glacier body. Below this




area, several other destabilization signs are visible from 2580 to 2480 m.a.s.l. The most prominent scarp (Fig. 5, "B") appears at about 2520 m.a.s.l., with a width of more than 150 m across and almost 300 m following the crown. The highest elevation difference is observable on the orographic left side, with more than 30 vertical metres between the crown and the top of the destabilized body. Nearby, several other cracks and scarps are clearly visible. In the following years, the destabilization signs rapidly change in size and geometry and move downslope. By 1977, the scarp at "A" is almost stable, while the lower area below "B" continues its evolution, especially towards the rock glacier front, where a series of profound scarps develop in a short time span, concurrent with a notable advance of the landform (Fig. 6, Fig. 5, Fig. S1). At the front, the oversteepened slope shows the last destabilization signs of the first phase with the onset of the lowermost scarp in 1977. In total, the front advanced roughly  $115 \text{ m} \pm 10 \text{ m}$  horizontally and 50 m vertically between 1953 and 1977. The 1953-71 and 1971-77 DSM pairs clearly show a pattern of elevation loss below area "B" and concurrent elevation gain at the lowest end of the advancing terminus (Fig. 7).

### 3.1.2 Intermediate period of relative stability: 1977 to 2017

From the mid-1970s until the later half of the 1990s, the displacement rates at the surface of the rock glacier stagnated in a narrow range with low variability. The mean block profiles and the velocity vectors derived from the 1977-90 DSM pair show similar values (between 0.3 m/a and 0.7 m/a for the mean block profile velocities and between 0.2 m/a and 1.7 m/a for the 1977-90 DSM pair at the profile locations, Fig. 3). There are larger discrepancies between the mean profile velocities and the DSM derived velocities at the profile locations in the first years of the time series due to lower performance of the DSM matching algorithm and the low density of the velocity vectors in the area around the profiles (Fig. 4). During the following periods (1990-onwards), the results from the DSM matching improve in the area of the profiles (Fig. 4) and so does the correspondence with mean block profile velocities (Fig. 3). Starting in the late 1990s, the velocities at all four profiles show a clear increasing trend, accompanied by large inter-annual variations. In the 1997-2006 DSM pair, the area of elevated velocity values extends approximately from the terrain step at 2570 m.a.s.l. to P3, with maximum values of up to about 3.5 m/a recorded near P1 on the orographic right side of the rock glacier. Velocities at the terminus did not noticeably rise compared to the 1990-97 period. The 2006-09 DSM pair shows a very similar velocity pattern (Fig. 4). In the following year (2009-10), velocities increase in the area slightly below and above the terrain step up to close to P3. This trend continues in the 2010-11 DSM pair, with elevated velocities of over 1 m/a recorded above P3 on both lobes of the rock glacier. For all the profiles, two velocity peaks were reached in 2004 and in 2015-16, with 3.3 m/a and 6.4 m/a respectively at P1. A velocity minimum was reached in 2007 at P0 and 2008 at P1, P2, and P3 (0.3 m/a at P0, 1.7 m/a at P1).

The BCF remained relatively constant at P1, P2, and P3 from the mid-1970s until the mid-1990s, at what may be considered a baseline value for the respective profiles in stable conditions (Fig. 3). This stable BCF value is about 2 at P2 at about 1 at P1 and P3 based on the mean profile velocities. P0 was not established yet during this time, but computing the BCF from the DSM-derived velocities for the area of P0 indicates a higher stable BCF of 4-5 in this section of the rock glacier. The 2004 velocity peak and subsequent decrease in velocity translates to a similar pattern of BCF, which is least pronounced at P3.






**Figure 3.** Time series of: a) mean profile velocities (m/a) at the 4 block profiles (Uncertainty estimate for the block profiles:  $\pm 0.2 - \pm 0.5$  m/a depending on the year, see Hartl et al. (2016b)). b) Mean velocities at the locations of the stone profiles derived from image correlation of DSM pairs, vertical dashed lines indicate years for which DSMs are available. See Fig. S2, S3, S4 (supplement) for corresponding boxplots of velocity uncertainties. c) Bulk creep factor (BCF) computed for the mean block profile velocities (solid lines) and the DSM velocities (dotted lines).


350 Interestingly, BCF at P2 is very close to BCF at P1 during the 2004 peak, although velocities are substantially higher at P1. During the 2015-16 peak, BCF at P1 rose to values comparable to the first period of destabilization in the 1960s.

The morphological signs of destabilization remained mostly unchanged until the 2011 DSM, as visible in Fig. 5 and in the movie in the supplement. However, some surface evolution is visible, especially in the central part of the destabilized area, where the terrain is steep and relatively large surface velocities were recorded even in this period. Between 1977 and 1990,  
355 the upper scarp shifted only minimally further downhill. The large, secondary scarp in area "B" and the smaller scarps lower down on the terminus became less prominent over time, reducing their length and height over the years (see Fig. 6 and Fig. 8). When analyzing the elevation difference between the top of the displaced material and the low point of the secondary




**Figure 4.** Velocity vectors (m/a) for the time series of DSM pairs, plotted over hillshades of the later DSM of each pair. Reference lines defining block profiles P0-P3 added for orientation. (Coordinate grid: EPSG: 31254)

scarp (Fig. 8), it is evident that the smoothing process develops slowly up to the 1997 DSM. The scarp height decreases by about 3 m in almost 40 years (Fig 8, b). Between 2011 and 2017, scarp height decreases by about another 2 m, as velocities  
 360 increase in the area above the scarp, and the scarp moves downhill shifting over 10 m in terms of horizontal distance (Fig 8, b). Similarly, the majority of the other destabilization signs on the terminus gradually smoothen (Fig. 6). The advance of the terminus in this period was small compared to the previous, more unstable period. The 1977-90 and 1990-97 DSM pairs show moderate elevation gain at the very end of the terminus and elevation loss directly above this area (Fig. 7). Later, in 1997, 2006, 2009, 2010, and 2011, an area of positive elevation change can be seen between about the elevation of P1 and just above the  
 365 terrain step. The upper part of the rock glacier mostly shows slightly negative elevation change, with the magnitude of negative



**Figure 5.** Hillshades of DSMs from 1953 to 2021. A and B mark upper and lower sections of morphological destabilization signs as referred to in the text. An animated version of the time series of hillshades is available in the video supplement. (Coordinate grid: EPSG: 31254)

change increasing in the later DSM pairs. The signal of elevation change in the terminus area is characterized by small scale



**Figure 6.** Surface elevation along the flow line as extracted from the DSMs. Estimated bedrock profile (dashed line) from Hartl et al. 2016b. Annotations show positions of the cross profiles. A and B mark upper and lower zones of morphological destabilization signs, as in Fig. 5.


variations around individual morphological destabilization features shifting downhill (Fig. 7). The front of the smaller, less active orographic right lobe shows positive elevation change due to a gradual advance.

### 3.2 Recent destabilization observed from high-resolution monitoring

370 In this section we take a closer look at the data since 2017, for which a much higher temporal and spatial resolution is available, extending the previous analysis and adding an additional focus on the movement and rotation of single blocks from 3D point clouds, and sub-seasonal changes at the rock glacier front. High-resolution 3D point clouds were derived from state-of-the-art close-range sensing techniques (ALS, ULS, TLS). These datasets are characterized by a spacing of the individual point measurements in the order of centimeters and, hence allow a detailed study of the kinematics and geomorphological evolution  
375 of the rock glacier. Since 2017, surface change at the rock glacier is characterized by unusually large displacement rates and by the onset and development of a second destabilization phase in the lower part.

#### 3.2.1 Kinematics


In measurement year 2018/19, velocities at the block profiles started increasing again. P1 was consistently the fastest profile throughout the time series and remained so with a mean profile velocity of 12.6 m/a in 2020/21. As of 2020/21, P0 was second  
380 fastest with 7.4 m/a. The high velocity at P0 is particularly noteworthy compared to previous years: Since its establishment in 1997, P0 was the slowest of the 4 profiles - often by a substantial margin - suggesting advanced degradation of the rock glacier in the terminus area. However, in the 2019/20 and 2020/21 measurement years, P0 was faster than P2 and P3, deviating strongly from the pattern of the previous 2 decades (Fig. 3). BCF values reflect the shift in behaviour at P0 and jump to over



**Figure 7.** Vertical surface elevation change (m/a) beyond the individual uncertainty (Table 2) for the time series of DSM pairs, plotted over hillshades of the latter DSM of each pair. High change values in the steep rock face above the root zone of the rock glacier are artefacts from the photogrammetric processing due to shadows and/or snow cover in the images and are included for the sake of transparency. Reference lines defining block profiles P0-P3 added for orientation. (Coordinate grid: EPSG: 31254)

20 in the 2020/21. P1 shows a similarly sharp increase in BCF. At P2 and P3, BCF is also elevated in recent years but remains  
 385 in the same range as during the 2015-16 peak. At P1 and P0, the 2020/21 profile mean velocities represent 567% and 582% of  
 the time series mean, respectively. At P2 and P3 the anomaly is pronounced but less extreme (224% and 336%, respectively)

Considering the movement of single blocks in the profiles (Fig. 9), it is apparent that the large increase in mean profile  
 velocities at P0 and P1 is driven by blocks in the central and orographic right section of the profiles. In 2020/21, the maximum




**Figure 8.** a) Surface elevation along the flow line as extracted from the DSMs, zoomed in on the scarp in zone "B", and approximate locations of the bottom of the scarp and the high point of the displaced material. The scarp is not yet present in 1953. b) Height and length of the scarp in different years (height is the vertical distance between the respective grey and red dots in a); length is the horizontal distance).

block velocities of P0 and P1, respectively, were 13.6 m/a and 20.6 m/a. Blocks on the orographic left and at the margins of  
390 the profiles show a far more gradual acceleration without the extreme jump in velocities in the most recent two years.

The highest annual 2.5D velocity detected in the DSM time series at a derived IMCORR-vector is 32.1 m/a in 2020/21 just  
below the terrain step (Fig. 4). This could vary slightly if the correlation analysis was performed at different nodes, so some  
caution is required when interpreting vector magnitudes. Considering only the 99.9 quantile of velocity vectors derived from  
the 2017/18-2020/21 DSM pairs, the highest value is 25.9 m/a in the same area of the rock glacier. The velocity vectors in the  
395 DSM pairs of 2011, 2017, 2018, 2019, 2020, and 2021 show a similar pattern of velocity distribution as the previous years,  
but with significantly and progressively higher velocities each year in the area at and below the terrain step. This is a strong  
indication that this section is central to the renewed destabilization process.

### 3.2.2 Destabilization signs

In the 2017 DSM, a large new crevasse is visible shortly upstream of the terrain step where the first destabilization phase started  
400 (zone "A"). A large section of the slope below this area was displaced by about 10 vertical metres between 2011 and 2017  
(Fig. 5, Fig. 6, Section 3.1.2). Between 2017 and 2021, the crevasse developed into a scarp concomitant with the appearance of  
new destabilization signs in its vicinity (cracks and crevasses). Some of the cracks closer to the terrain step, initially confined  
towards the left and right margins of the rock glacier, get connected by a long crevasse in 2019. This feature rapidly develops  
towards a scarp in the following years. By 2021, another large and very convex scarp appears just a few metres downslope. The  
405 appearance of these destabilization signs coincides with the large velocities illustrated in the previous section. While the scarps  
in zone "A" widened and deepened, the large scarp in zone "B" (Fig. 8) that developed during the first destabilization cycle



**Figure 9.** Block velocities for the four cross profiles (P0-P3) and for the longitudinal profile (P long.) in m/a for 2015/16 - 2020/21. The x-axis shows the distance from the orographic left reference point for P0, 1, 2, and 3, respectively. For P long., the x-axis shows the elevation of the blocks in each year. In P0, the third block from the right could not be located after 2018, presumably because it rotated so that the marking is obscured. In P1, a block was similarly lost in 2021. For P long., only blocks that were found every year since 2016 are shown.


continued decreasing in size and moved downslope, as described in the previous section. Recent TLS data from the lowest section of the terminus also show continuous smoothing of individual morphological features in recent years (Fig. S8).

The distribution of elevation change (Fig. 7) shows a similar pattern. Starting in 2011, pronounced surface elevation gain in the lower part and elevation loss in the upper part are visible. The signal of positive elevation change rapidly (within 4 years) propagates towards the front (Fig. 6, Fig. 7).

Zone “A” lies between profiles 1 and 2 and the surface strain rates between the two profiles also reflect the rapid changes in this part of the rock glacier in recent years. Strain rates were in a relatively low range of up to about  $0.005 \text{ a}^{-1}$  until 2011 and then showed a strong increase for blocks in the central and orographic right part of the rock glacier. Until 2019, strain rates in this section ranged from roughly  $0.01$  to  $0.02 \text{ a}^{-1}$ . Then, a large jump occurred and strain rates reached values around  $0.03 \text{ a}^{-1}$  in 2020 and between  $0.04$  and  $0.05 \text{ a}^{-1}$  in 2021. Strain rates remained low on the left margin of the rock glacier throughout and even decreased there slightly since about 2017 (Fig. 10).

### 3.2.3 Rotation of blocks

Individual blocks in profiles 0, 1, 2 and in the longitudinal profile were identified in high-resolution UAV orthophotos and tracked in the associated ULS point clouds for 2018-2021, yielding rotation angles of the blocks along the axis of the flow




**Figure 10.** Strain rates for the blocks in P1 and P2 (excluding the furthest left and furthest right blocks in P2) for the measurement years 2009/10 to 2020/21. Panel a) shows the location of the blocks in each year (red dots) over the 2021 hillshade. Panel b) shows the strain rate per block pair, counting the blocks in the profile lines from west to east.

direction. Rotation angles are generally low (roughly  $\pm 1^\circ$ ) at P2 and higher at P1, P0, and the longitudinal profile (Fig. 11). The blocks on the orographic left side of P1 show little rotational movement or tend to tilt forwards, while the blocks further to the right in the faster moving sector tend to tilt backwards in 2018/19 and 2019/20. In 2020/21, the rotational movement at P1 appears reduced overall. Blocks in P0 and in the lower part of the longitudinal profile where it intersects P0 tend to tilt backwards in all three years, with highest rotation angles ( $>5^\circ$ ) in the central part of the terminus near a morphological destabilization feature (Fig. 11). From 2018/19-2020/21, patterns of positive and negative surface elevation change around this feature became more pronounced. The strongly positive rotation angles are predominantly found in an area above the feature with positive surface elevation change.

### 3.2.4 Sub-seasonal displacement (2019)

430 For the bi-weekly TLS time series of summer 2019 we quantified 3D topographic change between two epochs by calculating CD-PB M3C2 distances between corresponding planar boulder faces (plane pairs). We considered CD-PB M3C2-based surface change in flow direction, vertical direction and horizontal direction. We quantified significant change in flow direction for 58,074 to 62,138 plane pairs, in vertical direction for 38,779 to 40,543 plane pairs, and in horizontal direction for 27,227 to 30,078 plane pairs. In this study, the uncertainty associated with quantified change ranges between 0.014 m and 0.015 m. This  
435 allows significant surface change to be quantified in all three directions in 73.95% to 75.28% of the total area for which change





**Figure 11.** Pitch angle of each DGNSS block identified in the ULS data for the years 2018/19, 2019/20, and 2020/21, plotted over the elevation change layer of the same time period and a hillshade of the later DSM. Black lines show the displacement vectors for the cross profiles during the 3 time periods.


quantification is applied (i.e. where corresponding boulder faces are identified). In Figures 12, S6, and S7 (supplementary material) only significant surface change is shown.

Our analysis reveals short-term variations in the magnitude of the movement along the flow direction at the rock glacier front from 0.12 m in 13 days to 0.21 m in 15 days (Fig. 12). These values correspond to average movement of 0.009 m/day in early July and August and 0.014 m/day during the second half of July and August, respectively. Standard deviation of the magnitudes is relatively constant at 0.07-0.08 m for each time step (Figures S6, S7). The magnitude of surface elevation change per time period also varies, with the largest values recorded in the first measurement period between the last week of June and the first week of July. Vertical change between plane pairs at the rock glacier front is predominantly negative (Fig. S6).

## 4 Discussion

### 4.1 Kinematic data and change detection: Uncertainties and challenges

We follow the method of previous studies for the calculation of mean profile velocities from the locations of the marked blocks, i.e. the profile mean is the mean of the available blocks per profile (Schneider and Schneider, 2001; Hartl et al., 2016b). This ensures the consistency of the time series, but we acknowledge that it is arguably a simplistic approach, as it does not account in detail for missing data from blocks that can no longer be found, the periodic repainting and repositioning of single blocks or profile lines, missing years, or the slight year-to-year differences in measurement dates. The magnitudes of the velocity vectors derived from the DSM pairs and the general distribution pattern of increase and decrease in velocities agree well with previous



**Figure 12.** Magnitudes of surface change in flow direction derived from the correspondence-driven plane-based M3C2 for 2-week timespans. A hillshade derived from airborne laser scanning data is used as baselayer in all subfigures. As the CD-PB M3C2 algorithm favors confident detection of small-magnitude changes over full point-wise change quantification, change information is more dense in at the steep rock glacier front where high point density, spatial coverage and spatial overlap between point clouds of two epochs could be achieved with TLS-based data acquisition.

studies, with minor discrepancies due to different node placements and methodological adjustments (Klug, 2011; Klug et al., 2012).

455 Based on the analysis of resulting vectors within stable areas around the rock glacier, the uncertainties inherent to the change detection between DSM pairs is in a similar range as was found by comparable studies at other rock glacier sites (e.g., Bodin et al. (2018); Fleischer et al. (2021); Kummert et al. (2021)). However, we individually consider the directions of the resulting pseudo-vectors within the stable areas to additionally analyze directional biases of the included data. This provides further information about data and registration quality (see Table 2 and Fig. S2, S3, and S4 in the supplement). In east-west



direction, the median velocity within the stable areas shows only minor deviations (e.g. +0.07 m/a in 2017-2018). In north-south  
460 direction however, distinct deviations emerge for the periods 1971-1977 (-0.16 m/a), 2009-2010 (-0.09 m/a) and 2017-2018  
(+0.15 m/a). This may result from the fact that within the area of interest mainly east- and west-facing slopes are covered by  
the data with sufficient quality, allowing for a more robust registration in the east-west direction. The deviations of median  
velocity within stable areas in the vertical direction are generally negligible, with the highest deviation in the period 1954-  
1971 (-0.03 m/a). In terms of data acquisition and processing techniques, the DSM pairs photogrammetrically derived from  
465 scanned analogue historical aerial imagery show the highest uncertainty. However, considering the longer periods between the  
acquisition campaigns (e.g. 1953-1971, 1977-1990) the respective velocity uncertainty reduces distinctly. The photogrammetric  
DSM pairs covering shorter periods (1971-1977, 1990-1997) show an accordingly higher velocity uncertainty. The DSM  
pairs marking the transitions between acquisition methods (1997-2006: photogrammetry-ALS; 2017-2018: ALS-ULS) are also  
associated with higher velocity uncertainty. This may be the result of the different quality in representing the terrain, which  
470 is typically lower using photogrammetric techniques based on historical aerial imagery. Furthermore, the spatial resolution  
increases drastically from photogrammetry to ALS and ULS. The higher spatial resolution of the ALS and ULS data includes  
more topographic details which will affect the aggregation of DSMs with 1m cell size, chosen as a compromise between the  
historical and recent data. Furthermore, in the case of the DSM pair 2017-2018, the reduced spatial coverage of the 2018  
ULS dataset limits the selection of stable areas for the co-registration with the 2017 ALS dataset. This particularly affects the  
475 uncertainty in both east-west and north-south directions. All in all, we consider the resulting velocity uncertainty adequate for  
the used data acquisition/processing techniques and they are well below the derived velocities.

The TLS dataset of the rock glacier front allows the quantification of short-term (bi-weekly) variations of 3D surface change.  
As magnitudes of rock glacier surface change tend to be small (<0.1 m) at such monitoring intervals, sophisticated methods for  
3D change detection capable of quantifying 3D surface change with low uncertainties are required. We were able to confidently  
480 quantify 3D surface change in different directions (flow direction, vertical direction, horizontal direction) and to reveal related  
sub-seasonal variations for a large number of corresponding planar boulder faces in two epochs. The sub-seasonal dataset  
presented in this study mainly covers the rock glacier front, thus limiting the comparability of this data with the long-term  
DSM time series. Future high-frequency monitoring set-ups might integrate point clouds derived from TLS as well as UAV-  
borne 3D sensing techniques (UAV-borne laser scanning, UAV-borne photogrammetry) as the latter offers increased coverage  
485 and a more uniform point distribution compared to ground-based sensing techniques (Zahs et al., 2022a). This will allow  
detailed study of sub-seasonal variations of 3D surface change for larger parts of the rock glacier.

Combining the in-situ DGNS data of the block profiles with the remote sensing data leverages the advantages of both  
methods and partially compensates for their disadvantages: On the one hand, the analysis of the DSM pairs during the first  
acceleration period shows high velocities in the terminus area that were not recorded in the time series of the blocks, since  
490 there was no block profile at the terminus during this time. The velocity patterns in the DSM analysis provide further spatial  
context to the point measurements of the blocks, yielding a very high-resolution overview of the spatial progression of the  
destabilization process. On the other hand, the higher temporal resolution of the block time series resolves shorter-term vari-  
ability - notably the 2004 and 2015-16 velocity peaks - which is not apparent in the DSM data. The fusion of DGNS data



and high-resolution 3D point clouds further allows the extraction of rotational information for individual blocks, aiding the  
495 interpretation of the rotational movement of the destabilized rock glacier terminus (see following section) and highlighting  
the potential of multi-method monitoring. In the future, such analysis might be extended by integrating change information  
obtained between corresponding planar boulder faces with the CD-PB M3C2 algorithm.

## 4.2 Interpretation of results in a destabilization context

In the following we discuss the long time series of surface displacement at HEK from a geomorphological perspective. The  
500 temporal resolution achieved with the 14 DSMs, complemented by surface velocities from block profiles from 42 measurement  
years allows for a comprehensive assessment of the two phases of destabilization (first DSM from 1953; annual DSMs since  
2017; first block data from P1 in 1952 and from P2 and 3 in 1955; annual data from 4 profiles since about 1997).

In literature, the evolution of a rock glacier with regard to the destabilization process is dynamically and geomorphologi-  
cally divided into four phases: normal activity (phase 1), destabilization onset (phase 2), destabilization peak (phase 3), and  
505 deceleration/inactivation (phase 4, Cicoira et al. (2020); Marcer et al. (2021)). We follow this approach and divide the two  
destabilization cycles observed in our dataset on the basis of a combination of displacement and surface strain rates, geomor-  
phological signs, and the values of the BCF for different areas of the rock glacier. The earliest data in our time series coincide  
with the onset of the first destabilization cycle, with clear geomorphological destabilization signs rapidly developing. Despite  
the relatively low temporal resolution of the first part of our dataset, the time series represents an unprecedented result for this  
510 period with three DSMs and 14 years of block velocity measurements documenting the first destabilization cycle between the  
early 1950s and late 1970s. As a result, we are able to distinctly map and follow the geomorphic signs as they evolve downslope  
and are able to resolve the acceleration and destabilization phase in the kinematic signal.

The two distinct cycles of destabilization of HEK are in general accordance with other publications that have analysed  
landform evolution over multi-decadal time series (e.g., Kääb et al. (2021); Marcer et al. (2021)). From these studies, the  
515 first cycle took place around the middle of the last century and the second cycle is reported approximately after the 1990s,  
with locally variable timing of the destabilization onset. For HEK, the onset of a second cycle of destabilization can first be  
observed in the 2017 data, after a longer but more gradual acceleration phase. The first destabilization cycle at the HEK started  
in 1953 and lasted until the late 1970s, affecting mostly the area of the rock glacier below the terrain step and leaving relatively  
undisturbed the terrain above. This observation is in line with the previous assessments of Haeberli and Patzelt (1982) and  
520 Schneider and Schneider (2001), who described the same general process and timeline in terms of a separation of the tongue  
below the terrain step from the upper, “healthy” part of the rock glacier. The surface elevation change along the flow line  
between 1953 and 1971 and 1977 as extracted from the respective DSMs (Fig. 7) also agrees well with the data on terminus  
advance and elevation change presented and discussed in Schneider and Schneider (2001).

During the first few decades of our time series, the integration of data from DSMs and block measurements shows destabi-  
525 lization phases 2 (onset) to 3 (peak) and 4 (deceleration, return to normal conditions). Phase 3 is characterized by high velocities  
and values of the BCF close to 40 in the lowest section of the rock glacier. During the subsequent period of relative stability  
and lower velocity values, the BCF values at P1 and P3 remain close to 1, which can be considered these profiles’ “stable



BCF" value corresponding to normal conditions rather than the extraordinary (Schneider and Schneider, 2001) conditions of destabilization. The renewed onset of acceleration in the 1990s did not coincide with the appearance of new morphological destabilization signs, nor with a clear and drastic dynamic decoupling between the two upper and lower sections of the rock glacier, as indicated by the low values of the BCF in both regions. After a long period of relatively continuous acceleration, clear signs of destabilization onset (phase 2) appeared in 2017, marking the beginning of the second destabilization cycle in terms of morphological destabilization signs. This was followed by a drastic increase in the velocities and in the BCF at the lower profiles in the most recent years of the time series. At P0 in particular, the dramatic rise in velocity and BCF at P0 since 2018 suggests a fundamental shift in underlying processes. The lowest section of the terminus appeared strongly degraded and showed comparatively little activity even during the 2004 velocity peak. The recent increase in velocities at P0 is likely due to the accelerating and increasingly destabilized upper section of the terminus pushing this lower section in the flow direction. In general, the velocities of the destabilized section seem entirely decoupled from the velocity patterns in the upper part and on the margins of the rock glacier and are significantly higher than at other rock glaciers in the region (e.g., Fleischer et al. (2021)). The increasing surface strain rates across the terrain step (between blocks in P1 and P2) and in particular the very high values in the most recent years are yet another indicator of highly destabilized conditions in the lower sector of the rock glacier compared to the relatively stable upper part. The magnitudes of the strain rates are in close agreement with findings by Marcer et al. (2021), who calculated surface strain rates for destabilized and non-destabilized rock glaciers and found that strain rates are about an order of magnitude larger in the destabilized cases. The large jumps in strain rates at HEK in 2020 and 2021 across zone "A" are likely caused by the formation and evolution of the newly developed scarps in this area.

Cicoira et al. (2020) and Bearzot et al. (2022) show that destabilization is characterized by discontinuities in BCF throughout the rock glacier area and argue that such discontinuities in BCF and surface velocity are stronger indicators of destabilization than the magnitude of either parameter. HEK exhibits a pronounced pattern break just below the terrain step, where the destabilization - or "separation" as per Schneider and Schneider (2001) and Haeberli and Patzelt (1982) - of a part of the terminus is occurring (Fig. 6, Fig S5). In contrast, the upper section is much more homogenous. Compared to Plator rock glacier (Italy) (Bearzot et al., 2022), HEK shows fewer distinct zones of similar BCF values but rather one main discontinuity.

Comparing absolute values of BCF at HEK to the larger dataset of Cicoira et al. (2020), we note that during highly destabilized conditions (phase 3), BCF at HEK as well as the corresponding velocities are higher than at any of the 340 rock glaciers assessed in their study. This could be explained by the difference between a landform-wide approach and the detailed investigations of single areas or even individual boulders (see also Bearzot et al. (2022)), and, in a more general sense, ties into the complexities of the dependence of spatial variability on scale. Under stable conditions, P1 and P3 have a BCF of about 1. In Cicoira et al. (2020), the rheological parameters were calibrated so that the peak of the distribution under normal conditions would match the value of 1. In this sense, HEK and its bulk creep behaviour are in line with most other rock glaciers analysed previously.

By extracting topographic information from the multi-temporal DSM dataset, we can extend the analysis presented in previous studies (Cicoira et al., 2020; Bearzot et al., 2022) and adjust the computation of the BCF for changes in slope angle and thickness over time. However, we do not account for changes in internal properties, i.e., friction angle, cohesion, shear resis-



tance, etc., which can vary considerably over space and time and strongly depend on temperature (Cuffey and Paterson, 2010; Moore, 2014; Cicoira et al., 2019a; Millstein et al., 2022). The lack of detailed subsurface information limits the possibilities  
565 for modelling the movement of the rock glacier. We use the bedrock estimate by Hartl et al. (2016a) to calculate the BCF, which implies that we assume a unique shear horizon at the depth of the bedrock for the computation. In terms of rheology, this is a substantial assumption that likely deviates from reality in some if not all parts of the rock glacier. In general, using bulk parameters to describe rock glacier rheology - as in the computation of the BCF - is a strong simplification of complex processes and by definition does not resolve local variations in rock glacier composition, or external factors, which have to be  
570 taken into account by means of ad-hoc parameterisations. However, as shown, it is possible to use such bulk parameters for the definition of dynamic phases and to highlight different dynamic behaviours, which can then be investigated in more detail, especially when a multi-temporal, good quality data basis is available. Although the number of studies incorporating the relatively new concept of BCF into their analysis is limited, the correspondence between the HEK results from a geomorphological perspective and the more dynamic, BCF focussed approach are increasing our confidence in the validity of this method.

575 The geomorphological destabilization signs in combination with the analysis of elevation differences and pitch angles indicate a landslide-like behaviour with roto-translational kinematics. Decreasing surface elevation change at rock glaciers can typically be expected in cases of permafrost thawing and extensional flow. In contrast, positive elevation change is most often linked to the advance of a front, the re-stabilization of a scarp, or to compressive flow. The large elevation changes ( $> 2$  m/a) observed during the destabilization phases at HEK appear to be related to the development of shearing surfaces and the conse-  
580 quent rotational and/or translational movement of the unstable permafrost masses. The change rates around the morphological destabilization features are orders of magnitude larger than in the area not affected by the destabilization process, and surface elevation change during highly destabilized conditions is generally higher than during the intermediate phase with normal creep behaviour. While change rates in the stable areas of HEK are in the range reported for similar landforms by other studies, the values in the destabilized section are considerably higher (e.g. Käab et al. (2003); Cusicanqui et al. (2021); Fleischer et al.  
585 (2021) and Wee and Delaloye (2022)).

A detailed visual inspection of the DSMs and the hillshade maps reveals that the destabilization process develops from higher to lower elevations, indicating an enlarging landslide pattern. For both destabilization phases, the first signs to appear are cracks at the sides of what will develop into a larger scarp. Other (subsequent) cracks appear further downslope as the ice-debris mixture loses cohesion, develops new sliding surfaces, and accelerates downslope. Springman et al. (2013) report  
590 a similar pattern as we observe in the HEK boulder rotations for Grabengufer rock glacier in Switzerland, where they found backwards tilting at the terminus and interpret this as evidence of “rotational failures at the foot of the rock glacier combined with slumping”. In one of the most comprehensive in situ investigations of rock glacier destabilization to date, Buchli et al. (2018) showed the existence of multiple shear horizons within the body of the Furggwanghorn rock glacier (Switzerland) and that rotational as well as translational movement can be associated with individual shear zones. We consider our results to be  
595 additional supportive evidence of such processes, although further geomorphological analysis and detailed numerical modelling experiments are clearly needed for a more detailed assessment. Generally speaking, the evidence of rotational movement at destabilized rock glaciers suggests that the underlying processes may resemble slow moving landslides after rupture (past the



tertiary creep phase) more closely than rock glacier creep. We hope the presented dataset will contribute to advancing our process understanding in this direction.

### 600 4.3 Meteorological and climatological setting

Considering long-term changes, the last approximately three decades of profile velocities at HEK show two distinct peaks and subsequent slowdowns in 2004 and 2015-16, respectively. Nickus et al. (2015) suggest that the 2004 peak and deceleration at HEK may have been related to the very warm summer of 2003. Summer temperature anomalies in 2003 stand out as an exceptionally warm outlier in the past two decades at the automatic weather station in Obergurgl (Fig. S10). 2015 was less  
605 extreme in terms of temperature anomalies but marks the start of a series of anomalously warm years that extends unbroken until 2021. The summers of 2018 and 2019 were the third and second warmest summers since 2003, respectively. 2015-2016, the years of the velocity plateau, saw relatively dry summers. In contrast, 2017 was unusually wet. It is likely that high temperatures and liquid water input through precipitation both contributed to the renewed acceleration from 2017/18 onwards, but rock glacier change monitoring at high temporal resolution in combination with modelling efforts is needed to gain a clearer  
610 picture of these connections. A number of rock glaciers in the Swiss, French, and Italian Alps also show a velocity peak in 2015 followed by 2-3 slower years and renewed acceleration from 2018 onwards, which strongly suggests a common climate forcing at the regional scale (Beutel et al., 2019; Bearzot et al., 2022; Thibert and Bodin, 2022; Wee and Delaloye, 2022).

The sub-seasonal TLS time series for the summer of 2019 primarily shows that movement rates at the front of the rock glacier fluctuate considerably on relatively short time scales, which is in keeping with other studies (e.g., Wirz et al. (2016);  
615 Buchli et al. (2018)). The magnitude of change per day (dividing total change by the length of the two week time span) is comparable to values reported for the front of Furggwanghorn rock glacier by Buchli et al. (2018). Short-term fluctuations of change at the rock glacier front are likely related to meteorological forcing but the approximately two week resolution of the TLS data does not resolve potential correlations with individual precipitation events and correlations are not linear. A speculative interpretation of the TLS data suggests that cumulative effects play a role for both temperature and precipitation on  
620 the time scales of the 2019 dataset. The first two week observation period (June 24 - July 6) was the warmest of the summer, but rock glacier movement was moderate. The third observation period, in which maximum velocities were reached, was not as warm as the first, but saw a lot of precipitation and a significant temperature increase from close to freezing to about 15°C (Fig. S9). A similar jump in temperatures between periods 4 and 5 also coincides with an increase in velocity. Buchli et al. (2018) observed localized, high daily change rates in sections of the front that experience water outflows and link the occurrence of  
625 such outflows to hydrological processes, e.g. water flow due to snow melt. At HEK, large changes in temperature combined with liquid water in the rock glacier system seem plausible as general drivers of the short-term changes measured during the 2019 summer season, but continuous monitoring of rock glacier movement is clearly needed to improve understanding of rock glacier response to short-term meteorological input.

High-resolution monitoring is also needed to better assess the hazard potential of the destabilized section of the rock glacier.  
630 localized rock fall from the steep rock glacier front has already led to temporary closures of the access road and is very likely to continue given the terrain and the movement of the front. The possibility of a complete collapse of the destabilized section



of the rock glacier and a subsequent rapid mass movement cannot be ruled out given what we know about the internal and external contributing factors. However, large uncertainties remain about the composition of the rock glacier both in the lower, destabilized section as well as in the upper part and it is unclear how much material is affected by the destabilization, nor is it currently possible to predict the likelihood of a collapse.

## 5 Conclusions

We present an updated and extended dataset of in situ and remote sensing-based change monitoring from Äußeres Hochebenkar rock glacier consisting of 14 DSMs covering a time span of 68 years (photogrammetry using historical aerial imagery for 1953, 1971, 1977, 1990, and 1997, airborne laser scanning for 2006, 2009, 2010, 2011, and 2017, and UAV-borne laser scanning for 2018, 2019, 2020, and 2021), as well as 42 individual measurement years of block displacement on the rock glacier surface (starting in the early 1950s, annual since about 1997), and a short-term time series of high-resolution 3D topographic change at the rock glacier front throughout the summer of 2019. Integrating the time series block displacement, DSM derived velocities, and geomorphological analysis allows for a detailed assessment of two cycles of destabilization at HEK and shows that while recent velocities measured in the lower sector of the rock glacier are unprecedented in the long time series, the destabilization process driving these kinematic changes is not. The highest velocities were recorded in the most recent measurement year (2020/21) and reach values between 20 m/a (DGNSS, single blocks) and about 30 m/a (image correlation from ULS data) in the highly destabilized lower section of the rock glacier.

Combining in situ data of block displacement and high-resolution 3D point clouds allows for method development based on data fusion approaches, expanding the change detection capacity of the monitoring network to new parameters, such as block rotation. Interdisciplinary monitoring and the coordinated integration of data and methods have the potential to shed light on the processes controlling rock glacier dynamics through geomorphological analysis and numerical modelling based on the resulting data. Such advances are only practicable when the data basis shows high granularity in terms of temporal and spatial resolution as well as accuracy.

We wish to highlight the importance of maintaining the monitoring network at Äußeres Hochebenkar rock glacier and, as a future goal, expanding it to subsurface monitoring. Well-studied sites like HEK are essential for the development and testing of new methods, as well as for advances in fundamental process understanding: The extensive historical data basis extending nearly 70 years into the past and the cyclic destabilization behaviour of the rock glacier provide a promising starting point for the development, calibration, and validation of numerical models pertaining to rock glacier movement in general and the destabilization process in particular. We are confident the dataset presented in this study (freely available through data repositories, see section Data Availability) will contribute to such efforts.

*Code and data availability.* – The DGNSS time series of block velocities is available on the pangaea data repository and updated yearly (Stocker-Waldhuber et al., 2021).





- The multi-temporal digital surface models, shaded reliefs and differential digital surface models as well as the derived velocity vectors are available on the Zenodo data repository (Zieher et al., 2022).
- 665 – The TLS data for summer 2019 and the source code for the correspondence-driven plane-based M3C2 method are available at: Zahs et al. (2021).
- The code to derive the velocity vectors and for the visualisations in this paper can be found at [https://github.com/thomaszieher/HRG\\_reanalysis](https://github.com/thomaszieher/HRG_reanalysis)
- The code to make the figures in this manuscript can be found at: [https://github.com/LeaHartl/Hochebenkar\\_figures](https://github.com/LeaHartl/Hochebenkar_figures)

*Video supplement.* An animated version of the time series of DSM hillshades is currently available as a .gif file in the supplementary material.  
670 *Please note: We are in the process of uploading this to a citable repository and will adjust this section once we have a doi.*

*Author contributions.* TZ, MB, and CK generated the DSM time series from source data. TZ homogenized the DSM dataset and derived the velocity time series. MB carried out the analysis related to the rotational movement of blocks. MS provided data and analysis related to the DGNSS block displacement and coordinates the in situ monitoring program. VZ and BH provided TLS data of the rock glacier front and carried out detailed 3D change analysis based on this data. LH and AC conceptualized the paper and carried out the kinematic and  
675 geomorphological analyses of the time series. LH wrote the paper with contributions from all co-authors.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* The contributions of Heral and Britta Schneider to the historical time series of block displacement are invaluable. In recent times, the maintenance of the block displacement time series and related monitoring efforts are supported by the “Verein Gletscher und Klima” and carried out by a team consisting mostly of volunteers. Without their dedication, continued long-term monitoring would not  
680 be possible.

Open source software was essential to the data processing, analyses, and visualizations in this study, in particular: QGIS, SAGA, and numerous python packages (QGIS.org, 2010; Oliphant, 2006; Hunter, 2007; GeoPandas, 2013-2019; Gillies et al., 2013; Conrad et al., 2015; Harris et al., 2020).



## References

- 685 Arenson, L., Hoelzle, M., and Springman, S.: Borehole deformation measurements and internal structure of some rock glaciers in Switzerland, *Permafrost and Periglacial Processes*, 13, 117–135, 2002.
- Arenson, L. U. and Springman, S. M.: Triaxial constant stress and constant strain rate tests on ice-rich permafrost samples, *Canadian Geotechnical Journal*, 42, 412–430, 2005.
- Avian, M., Kaufmann, V., and Lieb, G. K.: Recent and Holocene dynamics of a rock glacier system: The example of Langtalkar (Central Alps, Austria), *Norsk Geografisk Tidsskrift-Norwegian Journal of Geography*, 59, 149–156, 2005.
- 690 Barsch, D.: Permafrost creep and rockglaciers, *Permafrost and Periglacial Processes*, 3, 175–188, 1992.
- Barsch, D.: Indicators for the present and former geoecology in high mountain environments, Springer Berlin, Heidelberg, 1996.
- Bearzot, F., Garzonio, R., Colombo, R., Crosta, G. B., Di Mauro, B., Fioletti, M., Morra Di Cella, U., and Rossini, M.: Flow Velocity Variations and Surface Change of the Destabilised Plator Rock Glacier (Central Italian Alps) from Aerial Surveys, *Remote Sensing*, 14, 635, 2022.
- 695 Besl, P. J. and McKay, N. D.: Method for registration of 3-D shapes, in: *Sensor fusion IV: control paradigms and data structures*, vol. 1611, pp. 586–606, Spie, 1992.
- Beutel, J., Delaloye, R., Hilbich, C., Mollaret, C., Nötzli, J., Pellet, C., and Phillips, M.: PERMOS 2019. Permafrost in Switzerland 2014/2015 to 2017/2018, *Glaciological Report (Permafrost)*, 16, 2019.
- 700 Bodin, X., Krysiecki, J.-M., and Anocona, P. I.: Recent collapse of rock glaciers: two study cases in the Alps and in the Andes, in: *12th Congress, Protection of living spaces from natural hazards, Interpraevent 2012*, vol. 1, pp. p–409, 2012.
- Bodin, X., Krysiecki, J.-M., Schoeneich, P., Le Roux, O., Lorier, L., Echelard, T., Peyron, M., and Walpersdorf, A.: The 2006 collapse of the Bérard rock glacier (Southern French Alps), *Permafrost and Periglacial Processes*, 28, 209–223, 2017.
- Bodin, X., Thibert, E., Sanchez, O., Rabatel, A., and Jaillet, S.: Multi-annual kinematics of an active rock glacier quantified from very high-resolution DEMs: An application-case in the French Alps, *Remote Sensing*, 10, 547, 2018.
- 705 Bollmann, E., Klug, C., Sailer, R., Stötter, J., and Abermann, J.: Quantifying Rock glacier Creep using Airborne Laserscanning. A case study from two Rock glaciers in the Austrian Alps, in: *10th International Conference on Permafrost. The Northern Publisher, Salekhard, Russia*, pp. 49–54, 2012.
- Bollmann, E., Girmair, A., Mitterer, S., Krainer, K., Sailer, R., and Stötter, J.: A rock glacier activity index based on rock glacier thickness changes and displacement rates derived from airborne laser scanning, *Permafrost and Periglacial Processes*, 26, 347–359, 2015.
- 710 Bremer, M., Wichmann, V., Rutzinger, M., Zieher, T., and Pfeiffer, J.: SIMULATING UNMANNED-AERIAL-VEHICLE BASED LASER SCANNING DATA FOR EFFICIENT MISSION PLANNING IN COMPLEX TERRAIN., *International Archives of the Photogrammetry, Remote Sensing & Spatial Information Sciences*, 2019.
- Buchli, T., Kos, A., Limpach, P., Merz, K., Zhou, X., and Springman, S. M.: Kinematic investigations on the Furggwanhorn rock glacier, Switzerland, *Permafrost and Periglacial Processes*, 29, 3–20, 2018.
- 715 Cicoira, A., Beutel, J., Faillettaz, J., Gärtner-Roer, I., and Vieli, A.: Resolving the influence of temperature forcing through heat conduction on rock glacier dynamics: a numerical modelling approach, *The Cryosphere*, 13, 927–942, 2019a.
- Cicoira, A., Beutel, J., Faillettaz, J., and Vieli, A.: Water controls the seasonal rhythm of rock glacier flow, *Earth and Planetary Science Letters*, 528, 115 844, 2019b.



- 720 Cicoira, A., Marcer, M., Gärtner-Roer, I., Bodin, X., Arenson, L. U., and Vieli, A.: A general theory of rock glacier creep based on in-situ and remote sensing observations, *Permafrost and Periglacial Processes*, 32, 139–153, 2020.
- Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., Wehberg, J., Wichmann, V., and Böhner, J.: System for automated geoscientific analyses (SAGA) v. 2.1. 4, *Geoscientific Model Development*, 8, 1991–2007, 2015.
- Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, Academic Press, 2010.
- 725 Cusicanqui, D., Rabatel, A., Vincent, C., Bodin, X., Thibert, E., and Francou, B.: Interpretation of volume and flux changes of the Laurichard rock glacier between 1952 and 2019, French Alps, *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006 161, 2021.
- Delaloye, R., Perruchoud, E., Avian, M., Kaufmann, V., Bodin, X., Hausmann, H., Ikeda, A., Käab, A., Kellerer-Pirklbauer, A., Krainer, K., et al.: Recent interannual variations of rock glacier creep in the European Alps, 2008.
- Delaloye, R., Lambiel, C., and Gärtner-Roer, I.: Overview of rock glacier kinematics research in the Swiss Alps, *Geographica Helvetica*, 65,  
730 135–145, 2010.
- Delaloye, R., Morard, S., Barboux, C., Abbet, D., Gruber, V., Riedo, M., and Gachet, S.: Rapidly moving rock glaciers in Mattertal, Mattertal–ein Tal in Bewegung, edited by: Graf, C., *Publikation zur Jahrestagung der Schweizerischen Geomorphologischen Gesellschaft*, 29, 21–31, 2013.
- Eriksen, H. Ø., Rouyet, L., Lauknes, T. R., Berthling, I., Isaksen, K., Hindberg, H., Larsen, Y., and Corner, G. D.: Recent acceleration  
735 of a rock glacier complex, Adjet, Norway, documented by 62 years of remote sensing observations, *Geophysical Research Letters*, 45, 8314–8323, 2018.
- Fey, C. and Wichmann, V.: Long-range terrestrial laser scanning for geomorphological change detection in alpine terrain—handling uncertainties, *Earth Surface Processes and Landforms*, 42, 789–802, 2017.
- Fleischer, F., Haas, F., Piermattei, L., Pfeiffer, M., Heckmann, T., Altmann, M., Rom, J., Stark, M., Wimmer, M. H., Pfeifer, N., et al.:  
740 Multi-decadal (1953–2017) rock glacier kinematics analysed by high-resolution topographic data in the upper Kaunertal, Austria, *The Cryosphere*, 15, 5345–5369, 2021.
- GeoPandas: GeoPandas 0.8.0, <https://geopandas.org/>, last access: 2022, 2013–2019.
- Gillies, S. et al.: Rasterio: Geospatial raster i/o for Python programmers, Mapbox, <https://github.com/mapbox/rasterio>, last access: 2022, 2013.
- 745 Glen, J. W.: The creep of polycrystalline ice, *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 228, 519–538, 1955.
- Haeberli, W. and Patzelt, G.: Permafrostkartierung im Gebiet der Hochebenkar-Blockgletscher, Obergurgl, Ötztaler Alpen, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 18, 127–150, 1982.
- Haeberli, W., Hallet, B., Arenson, L., Elconin, R., Humlum, O., Käab, A., Kaufmann, V., Ladanyi, B., Matsuoka, N., Springman, S., et al.:  
750 Permafrost creep and rock glacier dynamics, *Permafrost and periglacial processes*, 17, 189–214, 2006.
- Harris, C. R., Millman, K. J., Van Der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., et al.: Array programming with NumPy, *Nature*, 585, 357–362, 2020.
- Hartl, L., Fischer, A., Klug, C., and Nicholson, L.: Can a simple numerical model help to fine-tune the analysis of ground-penetrating radar data? Hochebenkar rock glacier as a case study, *Arctic, Antarctic, and Alpine Research*, 48, 377–393, 2016a.
- 755 Hartl, L., Fischer, A., Stocker-waldhuber, M., and Abermann, J.: Recent speed-up of an alpine rock glacier: an updated chronology of the kinematics of outer hochebenkar rock glacier based on geodetic measurements, *Geografiska Annaler: Series A, Physical Geography*, 98, 129–141, 2016b.



- Hunter, J. D.: Matplotlib: A 2D Graphics Environment, *Computing in Science & Engineering*, 9, 90–95, <https://doi.org/10.1109/mcse.2007.55>, 2007.
- 760 Ikeda, A. and Matsuoka, N.: Pebbly versus bouldery rock glaciers: Morphology, structure and processes, *Geomorphology*, 73, 279–296, 2006.
- Kääb, A., Kaufmann, V., Ladstädter, R., and Eiken, T.: Rock glacier dynamics: implications from high-resolution measurements of surface velocity fields, in: *Eighth International Conference on Permafrost*, vol. 1, pp. 501–506, Permafrost, 2003.
- Kääb, A., Strozzi, T., Bolch, T., Caduff, R., Trefall, H., Stoffel, M., and Kokarev, A.: Inventory and changes of rock glacier creep speeds in  
765 Ile Alatau and Kungöy Ala-Too, northern Tien Shan, since the 1950s, *The Cryosphere*, 15, 927–949, 2021.
- Kaufmann, V.: The evolution of rock glacier monitoring using terrestrial photogrammetry: the example of Äusseres Hochebenkar rock glacier (Austria), *Austrian Journal of Earth Sciences*, 105, 2012.
- Kaufmann, V. and Ladstaedter, R.: Spatio-temporal analysis of the dynamic behaviour of the Hochebenkar rock glaciers (Oetztal Alps, Austria) by means of digital photogrammetric methods, *Proceedings of the 6th International Symposium on High Mountain Remote  
770 Sensing Cartography. Grazer Schriften der Geographie und Raumforschung*, 37, Institute of Geography and Regional Science, University of Graz, 119–139., 2002.
- Kaufmann, V. and Ladstaedter, R.: Monitoring of active rock glaciers by means of digital photogrammetry, *International Archives of Photogrammetry Remote Sensing and Spatial Information Sciences*, 34, 108–111, 2002.
- Kaufmann, V., Kellerer-Pirklbauer, A., and Seier, G.: Conventional and UAV-Based Aerial Surveys for Long-Term Monitoring (1954–2020)  
775 of a Highly Active Rock Glacier in Austria, *Frontiers in Remote Sensing*, 2, 732744, 2021.
- Kellerer-Pirklbauer, A., Lieb, G. K., and Kaufmann, V.: The Dösen Rock Glacier in Central Austria: A key site for multidisciplinary long-term rock glacier monitoring in the Eastern Alps, *Austrian Journal of Earth Sciences*, 110, 16, 2017.
- Kellerer-Pirklbauer, A., Delaloye, R., Lambiel, C., Gärtner-Roer, I., Kaufmann, V., Scapozza, C., Krainer, K., Staub, B., Thibert, E., Bodin, X., et al.: Interannual variability of rock glacier flow velocities in the European Alps, in: *Proceedings of the EUCOP5 5th European  
780 Conference on Permafrost, Chamonix-Mont Blanc, France*, vol. 23, pp. 396–397, 2018.
- Kenner, R., Phillips, M., Beutel, J., Hiller, M., Limpach, P., Pointner, E., and Volken, M.: Factors controlling velocity variations at short-term, seasonal and multiyear time scales, Ritigraben rock glacier, Western Swiss Alps, *Permafrost and Periglacial Processes*, 28, 675–684, 2017.
- Klug, C.: Blockgletscherbewegungen in den Stubai (Reichenkar) und Ötztal (Äußeren Hochebenkar) Alpen : Methodenkombination aus digitaler Photogrammetrie und Airborne Laserscanning. Diploma thesis., 2011.
- 785 Klug, C., Bollmann, E., Kääb, A., Krainer, K., Sailer, R., and Stötter, J.: Monitoring of permafrost creep on two rock glaciers in the Austrian eastern Alps: combination of aerophotogrammetry and airborne laser scanning, in: *Proceedings of the 10th international conference on permafrost*, vol. 1, pp. 215–220, 2012.
- Kofler, C., Mair, V., Gruber, S., Todisco, M. C., Nettleton, I., Steger, S., Zebisch, M., Schneiderbauer, S., and Comiti, F.: When do rock glacier fronts fail? Insights from two case studies in South Tyrol (Italian Alps), *Earth Surface Processes and Landforms*, 46, 1311–1327,  
790 2021.
- Krainer, K. and He, X.: Flow velocities of active rock glaciers in the austrian alps, *Geografiska Annaler: Series A, Physical Geography*, 88, 267–280, <https://doi.org/10.1111/j.0435-3676.2006.00300.x>, 2006.
- Krainer, K., Bressan, D., Dietre, B., Haas, J. N., Hajdas, I., Lang, K., Mair, V., Nickus, U., Reidl, D., Thies, H., et al.: A 10,300-year-old permafrost core from the active rock glacier Lazaun, southern Ötztal Alps (South Tyrol, northern Italy), *Quaternary Research*, 83,  
795 324–335, 2015.



- Krysiecki, J.-M., Bodin, X., Schoeneich, P., et al.: Collapse of the Bérard rock glacier (southern French Alps), in: Proc. 9th Int. Conf. Permafrost, Ext. Abstr, pp. 153–154, 2008.
- Kummert, M., Bodin, X., Braillard, L., and Delaloye, R.: Pluri-decadal evolution of rock glaciers surface velocity and its impact on sediment export rates towards high alpine torrents, *Earth Surface Processes and Landforms*, 46, 3213–3227, 2021.
- 800 Ladstädter, R. and Kaufmann, V.: Terrestisch-photogrammetrische Dokumentation des Blockgletschers im Äußeren Hochebenkar, Chesi, G., and Weinold, T. (eds.), *Internationale Geodätische Woche Obergurgl*, Herbert Wichmann Verlag, Heidelberg, 2005.
- Land Tirol Abteilung Geoinformation: Landesweite Laserscanbefliegung Tirol 2006-2010, Available at [https://www.tirol.gv.at/fileadmin/themen/sicherheit/geoinformation/Laserscandaten/ALS\\_Tirol\\_Onlinebericht.pdf](https://www.tirol.gv.at/fileadmin/themen/sicherheit/geoinformation/Laserscandaten/ALS_Tirol_Onlinebericht.pdf) (2022/08/19), 2011.
- Land Tirol Abteilung Geoinformation: Laserscandaten, Available at <https://www.tirol.gv.at/sicherheit/geoinformation/geodaten/laserscandaten/> (2022/08/19), 2019.
- 805 Marcer, M., Serrano, C., Brenning, A., Bodin, X., Goetz, J., and Schoeneich, P.: Evaluating the destabilization susceptibility of active rock glaciers in the French Alps, *The Cryosphere*, 13, 141–155, 2019.
- Marcer, M., Ringsø Nielsen, S., Ribeyre, C., Kummert, M., Duvillard, P.-A., Schoeneich, P., Bodin, X., and Genuite, K.: Investigating the slope failures at the Lou rock glacier front, French Alps, *Permafrost and Periglacial Processes*, 31, 15–30, 2020.
- 810 Marcer, M., Cicoira, A., Cusicanqui, D., Bodin, X., Echelard, T., Obregon, R., and Schoeneich, P.: Rock glaciers throughout the French Alps accelerated and destabilised since 1990 as air temperatures increased, *Communications Earth & Environment*, 2, 1–11, 2021.
- Millstein, J. D., Minchew, B. M., and Pegler, S. S.: Ice viscosity is more sensitive to stress than commonly assumed, *Communications Earth & Environment*, 3, 1–7, 2022.
- Moore, P. L.: Deformation of debris-ice mixtures, *Reviews of Geophysics*, 52, 435–467, 2014.
- 815 Müller, J., Vieli, A., and Gärtner-Roer, I.: Rock glaciers on the run—understanding rock glacier landform evolution and recent changes from numerical flow modeling, *The Cryosphere*, 10, 2865–2886, 2016.
- Nickus, U., Abermann, J., Fischer, A., Krainer, K., Schneider, H., Span, N., and Thies, H.: Rock Glacier Äußeres Hochebenkar (Austria)—recent results of a monitoring network, *Zeitschrift für gletscherkunde und glazialgeologie*, 47, 43–62, 2015.
- Niederwald, T.: Festpunktbestimmung mit GPS für Gletscher-Monitoring-Projekte in den Ötztaler Alpen (Tirol), 2009.
- 820 Oliphant, T. E.: *A guide to NumPy*, vol. 1, Trelgol Publishing USA, 2006.
- Pillewizer, W.: Untersuchungen an Blockströmen der Otztaler Alpen, *Geomorphologische Abhandlungen des Geographischen Institutes der Freien Universität Berlin (Otto-Maull-Festschrift)*, 5, 37–50, 1957.
- QGIS.org: QGIS Geographic Information System. Open Source Geospatial Foundation Project, <http://qgis.org>, 2010.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, (last accessed: 2022/08/22), 2021.
- 825 RGIK: Towards standard guidelines for inventorying rock glaciers: baseline concepts (version 4.2.2)., IPA Action Group Rock glacier inventories and kinematics, 13 pp., 2022.
- Rieger, W.: ALS Tirol 2017-18 Abschlussbericht, Tech. rep., 2019.
- Roer, I.: Rockglacier kinematics in a high mountain geosystem, Ph.D. thesis, Geographisches Institut, Universität Bonn, 2005.
- 830 Roer, I., Haerberli, W., Avian, M., Kaufmann, V., Delaloye, R., Lambiel, C., and Kääh, A.: Observations and considerations on destabilizing active rock glaciers in the European Alps, 2008.



- Roncat, A., Wieser, M., Briese, C., Bollmann, E., Sailer, R., Klug, C., and Pfeifer, N.: Analysing the suitability of radiometrically calibrated full-waveform lidar data for delineating Alpine rock glaciers, *ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, II-5/W2, 247–252, 2013a.
- 835 Roncat, A., Wieser, M., Briese, C., Bollmann, E., Sailer, R., and Pfeifer, N.: Digital surface model, hillshade and Lambertian reflectance model of the rock glaciers Oelgrube and Aeusseres Hochebenkar (Oetztal Alps, Tyrol, Austria), <https://doi.org/10.1594/PANGAEA.816225>, supplement to: Roncat, A et al. (2013): Analysing the suitability of radiometrically calibrated full-waveform lidar data for delineating Alpine rock glaciers. *ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, II-5/W2, 247–252, <https://doi.org/10.5194/isprsannals-II-5-W2-247-2013>, 2013b.
- 840 Scambos, T. A., Dutkiewicz, M. J., Wilson, J. C., and Bindschadler, R. A.: Application of image cross-correlation to the measurement of glacier velocity using satellite image data, *Remote sensing of environment*, 42, 177–186, 1992.
- Schneider, B.: Die Bewegungsmessungen am Blockgletscher im Äußeren Hochebenkar (Öztaler Alpen, Tirol) seit 1938, 1999a.
- Schneider, B.: Climate data and velocity of the Äußeres Hochebenkar (Öztal, Tyrolian Alps, Austria), <https://doi.org/10.1594/PANGAEA.836621>, 1999b.
- 845 Schneider, B. and Schneider, H.: Zur 60 jährigen Messreihe der kurzfristigen Geschwindigkeitsmessungen am Blockgletscher im Äußeren Hochebenkar, Öztaler Alpen, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 37, 1–33, 2001.
- Schoeneich, P., Bodin, X., Echelard, T., Kaufmann, V., Kellerer-Pirklbauer, A., Krysiecki, J.-M., and Lieb, G.: Velocity changes of rock glaciers and induced hazards, in: *Engineering Geology for Society and Territory-Volume 1*, pp. 223–227, Springer, 2015.
- Scotti, R., Crosta, G. B., and Villa, A.: Destabilisation of creeping permafrost: The Plator rock glacier case study (Central Italian Alps),
- 850 *Permafrost and Periglacial Processes*, 28, 224–236, 2017.
- Sorg, A., Kääh, A., Roesch, A., Bigler, C., and Stoffel, M.: Contrasting responses of Central Asian rock glaciers to global warming, *Scientific reports*, 5, 1–6, 2015.
- Springman, S. M., Yamamoto, Y., Buchli, T., Hertrich, M., Maurer, H., Merz, K., Gärtner-Roer, I., and Seward, L.: Rock glacier degradation and instabilities in the European Alps: a characterisation and monitoring experiment in the Turtmantal, CH, in: *Landslide science and practice*, pp. 5–13, Springer, 2013.
- 855 Stocker-Waldhuber, M., Emprechtinger, M., Hartl, L., and Fischer, A.: Continuous meteorological observations at weather station HEK (Hochebenkar), Öztal Alps, Austria, in 2012, <https://doi.org/10.1594/PANGAEA.808806>, 2013.
- Stocker-Waldhuber, M., Fischer, A., Hartl, L., Abermann, J., and Schneider, H.: Flow velocity records at Rock Glacier Outer Hochebenkar (Äußeres Hochebenkar), Öztal, Tyrolian Alps, Austria, 1997 et seq, <https://doi.org/10.1594/PANGAEA.928244>, 2021.
- 860 Tarini, M., Cignoni, P., and Montani, C.: Ambient occlusion and edge cueing for enhancing real time molecular visualization, *IEEE transactions on visualization and computer graphics*, 12, 1237–1244, 2006.
- Thibert, E. and Bodin, X.: Changes in surface velocities over four decades on the Laurichard rock glacier (French Alps), *Permafrost and Periglacial Processes*, 33, 323–335, 2022.
- Ulrich, V., Williams, J. G., Zahs, V., Anders, K., Hecht, S., and Höfle, B.: Measurement of rock glacier surface change over different
- 865 timescales using terrestrial laser scanning point clouds, *Earth Surface Dynamics*, 9, 19–28, 2021.
- Vietoris, L.: Der Blockgletscher des äußeren Hochebenkars, *Gurgler Berichte*, 1, 41–45, 1958.
- Vietoris, V.: Über den Blockgletscher des Äußeren Hochebenkars, *Z. Gletscherk. Glazialgeol.*, 8, 169–188, 1972.
- Vivero, S. and Lambiel, C.: Monitoring the crisis of a rock glacier with repeated UAV surveys, *Geographica Helvetica*, 74, 59–69, 2019.



- Wagner, T., Pleschberger, R., Kainz, S., Ribis, M., Kellerer-Pirklbauer, A., Krainer, K., Philippitsch, R., and Winkler, G.: The first consistent  
870 inventory of rock glaciers and their hydrological catchments of the Austrian Alps., *Austrian Journal of Earth Sciences*, 113, 2020.
- Wahrhaftig, C. and Cox, A.: Rock glaciers in the Alaska Range, *Geological Society of America Bulletin*, 70, 383–436, 1959.
- Wee, J. and Delaloye, R.: Post-glacial dynamics of an alpine Little Ice Age glacitectonized frozen landform (Aget, western Swiss Alps),  
Permafrost and Periglacial Processes, 2022.
- Whalley, W. B.: Rock glaciers and their formation as part of a glacier debris-transport system, University of Reading, Department of Geog-  
875 raphy, 1974.
- Williams, R.: DEMs of difference, *Geomorphological Techniques*, 2, 2012.
- Wirz, V., Gruber, S., Purves, R. S., Beutel, J., Gärtner-Roer, I., Gubler, S., and Vieli, A.: Short-term velocity variations at three rock glaciers  
and their relationship with meteorological conditions, *Earth Surface Dynamics*, 4, 103–123, 2016.
- Zahs, V., Hämmerle, M., Anders, K., Hecht, S., Sailer, R., Rutzinger, M., Williams, J. G., and Höfle, B.: Multi-temporal 3D point cloud-based  
880 quantification and analysis of geomorphological activity at an alpine rock glacier using airborne and terrestrial LiDAR, *Permafrost and  
Periglacial Processes*, 30, 222–238, 2019.
- Zahs, V., Winiwarter, L., Anders, K., Williams, J. G., Rutzinger, M., Bremer, M., and Höfle, B.: Correspondence-driven plane-based M3C2  
for quantification of 3D topographic change with lower uncertainty [Data and Source Code], <https://doi.org/10.11588/data/TGSVUI>, 2021.
- Zahs, V., Winiwarter, L., Anders, K., Bremer, M., Rutzinger, M., Potčková, M., and Höfle, B.: EVALUATION OF UAV-BORNE PHO-  
885 TOGRAMMETRY AND LASER SCANNING FOR 3D TOPOGRAPHIC CHANGE ANALYSIS AT AN ACTIVE ROCK GLACIER,  
*The International Archives of Photogrammetry, Remote Sensing and Spatial Information Sciences*, 43, 1109–1116, 2022a.
- Zahs, V., Winiwarter, L., Anders, K., Williams, J. G., Rutzinger, M., and Höfle, B.: Correspondence-driven plane-based M3C2 for lower  
uncertainty in 3D topographic change quantification, *ISPRS Journal of Photogrammetry and Remote Sensing*, 183, 541–559, 2022b.
- Zieher, T., Bremer, M., Klug, C., and Hartl, L.: Multisensor monitoring data of Hochebenkar Rock Glacier,  
890 <https://doi.org/10.5281/zenodo.7010292>, 2022.