

1 **Feedbacks between the formation of secondary minerals and the** 2 **infiltration of fluids into the regolith of granitic rocks in different** 3 **climatic zones (Chilean Coastal Cordillera)**

4 Ferdinand J. Hampl¹, Ferry Schipperski¹, Christopher Schwerdhelm², Nicole Stroncik³, Casey Bryce⁴,
5 Friedhelm von Blanckenburg^{3,5}, Thomas Neumann¹

6
7 ¹Department of Applied Geochemistry, Technische Universität Berlin, Ernst-Reuter-Platz 1, 10587 Berlin, Germany

8 ²Geomicrobiology Group, Eberhard-Karls-University Tübingen, Schnarrenbergstrasse 94-96, 72076 Tübingen, Germany

9 ³Earth Surface Geochemistry, GFZ German Research Centre for Geosciences, Telegrafenberg, 14473 Potsdam, Germany

10 ⁴School of Earth Sciences, University of Bristol, Wills Memorial Building, Queens Road, Bristol BS8 1RJ, United Kingdom

11 ⁵Institute of Geological Sciences, Freie Universität Berlin, Malteserstrasse 74-100, 12249 Berlin, Germany

12

13 *Correspondence to:* Ferdinand J. Hampl (ferdinand.j.hampl@tu-berlin.de)

14

15 **Abstract.** Subsurface fluid pathways and the climate-dependent infiltration of fluids into the subsurface jointly control the
16 intensity and depth of mineral weathering reactions. The products of these weathering reactions (secondary minerals), such as
17 Fe(III) oxyhydroxides and clay minerals, in turn exert a control on the subsurface fluid flow and hence on the development of
18 weathering profiles.

19 We explored the dependence of mineral transformations on climate during the weathering of granitic rocks in two 6 m deep
20 weathering profiles in Mediterranean and humid climate zones along the Chilean Coastal Cordillera. We used geochemical
21 and mineralogical methods such as (micro-) X-ray fluorescence, oxalate/dithionite extractions, X-ray diffraction and electron
22 microprobe mapping to elucidate the transformations involved during weathering. In the profile of the Mediterranean climate
23 zone, we found a low weathering intensity affecting the profile down to 6 m depth. In the profile of the humid climate zone,
24 we found a high weathering intensity. Based on our results, we propose mechanisms that can intensify the progression of
25 weathering to depth. The most important is weathering-induced fracturing (WIF) by Fe(II) oxidation in biotite and precipitation
26 of Fe(III) oxyhydroxides, and by swelling of interstratified smectitic clay minerals that promotes the formation of fluid
27 pathways. We also propose mechanisms that mitigate the development of a deep weathering zone, like the precipitation of
28 secondary minerals (e.g., clay minerals) and amorphous phases that can impede the subsurface fluid flow. We conclude that
29 the depth and intensity of primary mineral weathering in the profile of the Mediterranean climate zone is significantly
30 controlled by WIF. It generates a surface-subsurface connectivity that allows fluid infiltration to great depth and hence
31 promotes a deep weathering zone. Moreover, the water supply to the subsurface is limited in the Mediterranean climate and
32 thus most of the weathering profile is generally characterized by a low weathering intensity. The depth and intensity of
33 weathering processes in the profile of the humid climate zone, on the other hand, are controlled by an intense formation of

34 secondary minerals in the upper section of the weathering profile. This intense formation arises from pronounced dissolution
35 of primary minerals due to the high water infiltration (high precipitation rate) into the subsurface. The secondary minerals, in
36 turn, impede the infiltration of fluids to great depth and thus mitigate the intensity of primary mineral weathering at depth.
37 These two settings illustrate that the depth and intensity of primary mineral weathering in the upper regolith are controlled by
38 positive and negative feedbacks between the formation of secondary minerals and the infiltration of fluids.

39

40 **Keywords:** Coastal Cordillera, feedback, weathering-induced fracturing, secondary minerals, Critical Zone, fluid flow

41 **1 Introduction**

42 The formation of weathered material (regolith) from unweathered rock (bedrock) is a key process for shaping Earth's surface.
43 It is of major importance for making mineral-bound nutrients accessible to the biosphere of the Critical Zone (e.g., Dawson et
44 al., 2020) and to supply rocks and minerals to the sediment cycle. In this process the in-situ disaggregation and chemical
45 depletion of weathered rock (saprock) to saprolite plays an essential role. This transformation is a result of fracturing and
46 mineral dissolution (e.g., Navarre-Sitchler et al., 2015). Both are associated with chemical, physical (e.g., Goodfellow et al.,
47 2016), and biological weathering processes (e.g., Drever, 1994; Lawrence et al., 2014; Napieralski et al., 2019). These
48 processes are linked to climate-related parameters such as precipitation rate, fluid flow (water and gases), and biological
49 activity. Apart from that, the weathering processes and hence the saprolite formation also depend on primary fractures (e.g.,
50 Molnar et al., 2007; Hynek et al., 2017; Kim et al., 2017; Holbrook et al., 2019; Hayes et al., 2020; Krone et al., 2021; Hampl
51 et al. 2022a), discontinuity density and tortuosity (Israeli et al., 2021), thermoelastic relaxation (e.g., Nadan and Engelder,
52 2009) as well as the topographic surface profile (e.g., Rempe and Dietrich, 2014; St. Clair et al., 2015). However, one of the
53 most fundamental parameters for the regolith formation is the mineral content of the bedrock. The weathering of some of these
54 primary minerals and the consequent formation of secondary minerals can lead to an amplification of the depth and intensity
55 (i.e., the parameter describing the elemental loss and relative amount of secondary minerals) of primary mineral weathering
56 (e.g., Fletcher et al., 2006; Lebedeva et al., 2007; Buss et al., 2008; Behrens et al., 2015; Hampl et al. 2022a). Such mechanisms
57 comprise (1) a forcing process like the formation of secondary minerals that is triggering (2) a responsive process such as more
58 intense infiltration of fluids to depth. The latter process reinforces the initial forcing process of secondary mineral formation.
59 Such a mechanism is therefore called positive feedback between (1) and (2). The formation of secondary minerals can also
60 have a weathering-impeding effect (e.g., Lohse and Dietrich, 2005; Navarre-Sitchler et al., 2015; Kim et al., 2017; Gerrits et
61 al., 2021) causing a mitigation of the weathering depth and -intensity. Such mechanisms comprise (1) a forcing process like
62 the formation of secondary minerals and (2) a responsive process such as reduced infiltration of fluids to depth. The latter
63 process damps the initial forcing process of secondary mineral formation, and the mechanism is therefore called negative
64 feedback between (1) and (2).

65 Deciphering the relationship between the formation of secondary minerals and the climatic conditions they were formed under
66 is a prerequisite for understanding the weathering system. It allows to determine whether feedbacks between the formation of
67 secondary minerals and the infiltration of fluids affect the intensity and depth of primary mineral weathering. We hypothesize
68 that a positive feedback loop results in a deep weathering depth, as secondary minerals form fluid pathways by fracturing due
69 to volume increase. On the other hand, we think that a negative feedback loop leads to a shallow weathering depth, as the
70 precipitation of secondary minerals seals fluid pathways.

71 To explore such connections and to elucidate the impact of secondary minerals on the development of weathering systems in
72 different climatic zones, we investigated two 6 m deep weathering profiles in the Chilean Coastal Cordillera. One profile is
73 located in a Mediterranean and another in a humid climate zone, and both developed from weathering of granitic rock. Both
74 sites are eroding and the surfaces in the locations are thus constantly turned over (see compilation of rates and environmental
75 parameters in Oeser and von Blanckenburg (2020) and references therein). The profiles were sampled in soil pits and
76 complemented with rock samples obtained by deep wireline rotary drilling close to the soil pits. Samples were investigated by
77 a combination of analytical techniques such as X-ray fluorescence and micro-X-ray fluorescence as well as oxalate-/dithionite
78 extraction to characterize the geochemical composition, and X-ray diffraction, magnetic susceptibility measurements, electron
79 microprobe as well as light microscopy to identify the mineral assemblages. The combined results of these techniques are used
80 to derive weathering-intensifying and -mitigating processes during subsurface weathering and to elucidate how these processes
81 influence the depth and intensity of weathering in the different climate zones.

82 **2 Study sites**

83 **2.1 La Campana (LC)**

84 The soil pit (-33.02833° N, -71.04370° E, 894 m) and the drilling site some 15 m next to it (-33.02833° N, -71.04354° E,
85 898 m) are located south of the La Campana National Park approximately 60 km NW of Santiago de Chile (Fig. 1a). They are
86 situated on a ridge with steep slope dip angles of 20–30°.

87 The vegetation can be characterized as Mediterranean sclerophyllous forest with *Cryptocarya alba* and *Lithraea caustica* as
88 dominant plants (Luebert and Plissock, 2006; Oeser et al., 2018; Fig. 1b,c). The annual precipitation rate (measured from April
89 2016 to April 2020) is 346 mm yr⁻¹ (Übernicker et al., 2020) and the Holocene net primary production is 280 ± 50 g C m⁻² yr⁻¹
90 (Werner et al., 2018; Oeser and von Blanckenburg, 2020).

91 The regolith profile developed on top of Upper Cretaceous intrusions of mainly granodiorites and tonalites with subordinate
92 quartz monzodiorites (Gana et al., 1996). The depths of the soil horizons are A: 0–30 cm, B: 30–83 cm and C (saprolite): >83
93 cm (Fig. 1d). Uplift rates for the north of Santiago de Chile vary between 0.01 and 0.23 mm yr⁻¹ with a general mean value of
94 0.13 ± 0.04 (Melnik, 2016). The soil denudation rate in the nearby La Campana National Park is 53.7 ± 3.4 (S-facing slope)
95 to 69.2 ± 4.6 t km⁻² yr⁻¹ (N-facing slope; Oeser et al., 2018).


96 2.2 Nahuelbuta (NA)

97 The investigated soil pit (-37.79371° N, -72.95065° E, 1113 m) and the drilling site next to it (-37.79381° N, -72.95043° E,
98 1114 m) are located approximately 20 km west of Angol (Region IX (Araucanía), Province Malleco) in southern Chile (Fig.
99 1a). The borehole was located on a plateau-like ridge with gently dipping slopes (ca. 10°).

100 The pre-land-use vegetation in the study area resembled the recent vegetation found in the Nahuelbuta National Park which
101 can be characterized as temperate forest with *Araucaria araucana* as the dominant tree (Luebert and Plissock, 2006; Fig. 1e).
102 However, extensive modern pastoral farming (cow grazing) and fires have converted the ecosystem in the study area to a
103 sparse forest of deciduous trees such as *Nothofagus obliqua* (see Oeser et al., 2018; Fig. 1f). Numerous signs of burning can
104 be observed in the field and charcoal is an integral component of the soil down to 25 cm (A horizon). The precipitation rate
105 (measured from end of March 2016 to April 2020) is 1927 mm yr⁻¹ (Übernicker et al., 2020) and the Holocene net primary
106 production is 520 ± 130 g C m⁻² yr⁻¹ (Werner et al., 2018; Oeser and von Blanckenburg, 2020).

107 The regolith profile developed on top of granitoid rocks of the Nahuelbuta central pluton which contains heterogenous
108 lithological portions (Hervé, 1977; Ferraris, 1979). It is part of the Nahuelbuta Batholith which in turn belongs to the late
109 Carboniferous Chilean Coastal Batholith (Steenken et al., 2016; Deckart et al., 2013). The depths of the soil horizons are A:
110 0–25 cm, B: 25–90 cm, and C (saprolite): >90 cm (Fig. 1g). Today's exhumation rates in NA are high (>0.2 mm yr⁻¹; Glodny
111 et al., 2008b), whereas the catchment-wide denudation rate is small (27.4 ± 2.4 mm kyr⁻¹; van Dongen et al., 2019) compared
112 to LC. The soil denudation rate in the nearby Nahuelbuta National Park ranges between 17.7 ± 1.1 (N-facing slope) to 47.5 ±
113 3.0 t km⁻² yr⁻¹ (S-facing slope; Oeser et al., 2018). Tectonic fractures in NA can be related to the Lanalhue Fault Zone (see
114 Glodny et al., 2008a).

115



116 **Figure 1: Overview of the study sites and soil profiles.** (a) Position of La Campanana (LC) and Nahuelbuta (NA) in Chile. Modified
 117 map data from [OpenStreetMap](https://www.openstreetmap.org/) (© [OpenStreetMap](https://www.openstreetmap.org/) contributors). (b) Original vegetation in LC (i.e., before human intervention;
 118 La Campanana National Park). (c) Vicinity of the soil pit and drilling site in LC and (d) the first 2 m of the soil profile in LC with
 119 inscribed soil horizons (A-C). A prominent discontinuity (dashed line) can be found in the depth interval 120–140 cm. (e) The original
 120 vegetation in NA (i.e., before human intervention; Nahuelbuta National Park) in comparison to (f) the recent vegetation in the
 121 vicinity of the soil pit and drilling site. (g) The first 2 m of the soil profile in NA with inscribed soil horizons (A-C).

122 3 Materials and methods

123 3.1 Soil pit sampling, drilling, and sample preparation

124 The sampled 6 m deep soil profiles were located close to the main boreholes at the respective sites. Bulk samples were collected
 125 in 20 intervals in each soil pit and weighed around 3 kg. Corestones were not encountered in the soil pit profiles of LC and
 126 NA. By using a rotary splitter (type PT, Retsch) the bulk samples were separated into aliquots (see Hampl et al., 2022b).
 127 During the drilling campaigns, up to 1.5 m long core runs were recovered by wireline diamond drilling (~80 mm core diameter)
 128 using potable water as drilling fluid (see Krone et al. (2021) for a detailed description of the drilling technique). Rock samples
 129 were separated from the core by mechanical methods (angle grinder, hammer and chisel), cut (diamond saw), impregnated
 130 with blue artificial resin filling the porosity, and subsequently thin-sectioned. Representative bedrock samples were separated
 131 from the core (diamond saw) and crushed (jaw crusher).

132 3.2 Analytical methods and calculations

133 A detailed description of the analytical methods can be found in the accompanying data publication of this study (Hampl et
134 al., 2022b).

135 3.2.1 X-ray fluorescence (XRF) and micro-X-ray fluorescence (μ -XRF)

136 Soil pit samples were ground with an agate disc mill and annealed (950°C for 1h) before adding a lithium borate flux to produce
137 glass beads in platinum crucibles. The element composition of the glass beads was analysed with a Thermo Scientific ARL
138 PERFORM[®]X X-ray fluorescence sequential spectrometer (WD-XRF; Thermo Fisher Scientific Inc., U.S.A.). Additional
139 powder pellets were produced by mixing the ground air-dried samples with wax. The mixtures were pressed and analysed with
140 a SPECTRO XEPOS energy dispersive X-ray fluorescence spectrometer (ED-XRF, SPECTRO Analytical Instruments GmbH,
141 Germany). Polished sample slabs of bedrock (Fig. 2) were mapped for the spatial distribution of elements with a μ -XRF
142 spectrometer M4 Tornado (Bruker, Germany).

143

144 Geochemical calculations

145 Zr contents obtained from the XRF element analyses on powder pellets were used as an immobile element for the calculation
146 of the chemical depletion fraction (CDF; Riebe et al., 2003; Eq. 1), and the mass transfer coefficient (τ ; Anderson et al., 2002;
147 Eq. 2).

148

$$149 \text{ CDF} = 1 - \frac{\text{Zr}_N^b \cdot (\text{sum}^w - \text{LOI}^w)}{\text{Zr}_N^w \cdot (\text{sum}^b - \text{LOI}^b)}, \quad (1)$$

150

$$151 \tau = \frac{X^w \cdot \text{Zr}^b}{X^b \cdot \text{Zr}^w} - 1, \quad (2)$$

152

153 LOI^b = loss on ignition of the bedrock, LOI^w = loss on ignition of the weathered sample, sum^b = measured total sum (wt.%)
154 of the bedrock, sum^w = measured total sum (wt.%) of the weathered sample, X^b = concentration of element X in the bedrock,
155 X^w = concentration of element X in the weathered sample, Zr^b = concentration of Zr in the bedrock, Zr_N^b = zirconium content
156 of the bedrock normalized to a LOI-free sum of 100 %, Zr^w = concentration of Zr in the weathered sample, Zr_N^w = zirconium
157 content of the weathered sample normalized to a LOI-free sum of 100 %.

158

159 The chemical index of alteration (CIA; Nesbitt and Young, 1982) was modified to Δ CIA (Eq. 3).

160

$$161 \Delta \text{CIA} = \left[\left(\frac{\text{Al}_2\text{O}_3^w}{\text{Al}_2\text{O}_3^w + \text{CaO}^w + \text{Na}_2\text{O}^w + \text{K}_2\text{O}^w} \right) - \left(\frac{\text{Al}_2\text{O}_3^b}{\text{Al}_2\text{O}_3^b + \text{CaO}^b + \text{Na}_2\text{O}^b + \text{K}_2\text{O}^b} \right) \right] \cdot 100, \quad (3)$$

162

163 $w =$ in the weathered sample, $b =$ in the bedrock.

164

165 **3.2.2 Oxalate- and dithionite extraction**

166 Air-dried bulk samples of <2 mm (dry-sieved) were used for oxalate- and dithionite extractions. The solutions thus obtained
167 were measured with an ICP-OES iCAP 6300 DUO (Thermo Fisher Scientific, USA) to determine the extractable Fe, Al, and
168 Si contents. The oxalate extraction employed targets the easily extractable mainly X-ray amorphous Fe(III) oxyhydroxides and
169 (poorly) crystalline Al-containing minerals (see review by Rennert (2019) and references therein). The dithionite extraction
170 dissolves crystalline and amorphous iron oxides (McKeague and Day, 1966). In doing so it can (partly) attack Al-bearing
171 (mineral) phases (see review by Rennert (2019) and references therein).

172 The oxalate extractions were performed after Schwertmann (1964) with an oxalic acid-/oxalate-extraction solution (0.2 M, pH
173 3.0). After the addition of the solution to the sample and shaking for 2 h in the dark (over-head shaker), the solution was filtered
174 in a darkened room and immediately measured. The cold dithionite extractions were performed based on Holmgren (1967)
175 with an extraction solution (mixture of 0.2 M NaHCO₃ and 0.24 M trisodium citrate) and sodium dithionite under oxic
176 conditions. The resulting mixture of chemicals and sample was shaken for 16 h and centrifuged before the supernatant was
177 filtered and immediately measured. Additional reference samples, blanks and calibration solutions were also prepared and
178 measured like the soil pit samples. The results of the samples presented here are the mean of duplicate measurements performed
179 on two individually extracted sample aliquots.

180 **3.2.3 Grain size determination**

181 Sample aliquots were suspended in de-ionized water (<10 µS m⁻¹) and dispersed in a rotating overhead shaker (approx. 15 h)
182 and a subsequent ultrasonic bath before vibrational wet sieving. The >63 µm sieving fractions were dried (50°C, approx. 24
183 h) and their weight percentages were measured. The clay and silt contents were determined using the <63 µm suspension and
184 a pipette method. Organic-rich samples were treated with H₂O₂ to decompose organic matter and sodium pyrophosphate was
185 used as a dispersion agent to prevent coagulation. Clay (<2 µm) was separated from the <63 µm fraction slurry via
186 centrifugation.

187 **3.2.4 X-ray diffraction (XRD)**

188 Untreated air-dried aliquots of bulk samples were crushed in a porcelain mortar and afterwards processed with a micronisation
189 XRD-mill McCrone (Retsch, Germany) to obtain a final powder of <10 µm. These powders were mounted to XRD sample
190 holders by back-loading and X-ray diffraction measurements were performed with a Rigaku SmartLab equipped with a 9 kW
191 rotating Cu-anode and a HyPix-3000 detector in Bragg-Brentano geometry (3–80° 2θ, scan step: 0.01°, scan speed: 1° min⁻¹,
192 and 60 rpm sample rotation). For the identification and semi-quantitative analyses, the software SmartLab Studio II and the
193 mineral database PDF-4 Minerals 2021 including reference intensity ratio (RIR) factors were used. Image processing (imageJ;

194 version 1.53a; Schneider et al., 2012) performed on the μ -XRF element distribution maps in Fig. 2 was used to get a rough
195 compositional information of the mineral content in the sampled bedrock. These analyses were used as a supporting basis for
196 the semi-quantitative XRD analyses with RIR factors.

197 Clay mineral contents in the samples were quantitatively estimated by combining the results of the grain size determination
198 with the semi-quantitative results of the XRD analyses. The clay-size fraction ($<2\ \mu\text{m}$) of which the mass was determined by
199 sieving/pipetting, was assumed to represent the entire clay mineral content of the sample, while the other size fractions were
200 considered to be free of clay minerals. This assumed clay mineral content (in wt.%) was combined with the XRD-semi-
201 quantitative weight percentages of the primary minerals in the same sample to approximate the mineral composition of the
202 whole soil pit sample (summarized to 100 wt.%). Despite the assumption that only the $<2\ \mu\text{m}$ grain size fraction contains clay
203 minerals, this estimate appears to be the most accurate because there are no matching files in the mineral database used here
204 that would accurately semi-quantify the identified interstratified clay minerals.

205 The separated clay-size fractions were measured as oriented clay films (texture preparation). A D2 Phaser XRD device
206 (Bruker) equipped with a Cu-anode was utilized for the measurements. The diffractograms were recorded in Bragg-Brentano
207 geometry in the range of $3\text{--}35^\circ\ 2\theta$ (step width: $0.01^\circ\ 2\theta$, 0.5 seconds per step). The samples were measured after air-drying,
208 during ethylene glycol saturation and after a thermal treatment at 550°C for 1 h. Selected samples were also treated with
209 glycerol and KCl (1 M) to characterize the clay minerals in more detail. The identification was supported by a clay mineral
210 identification chart (Starkey et al., 1984).

211 **3.2.5 Magnetic susceptibility measurements**

212 The magnetic susceptibility was measured on all twenty-one McCrone-milled bulk samples of the LC profile with a KLY-3
213 Kappabridge (AGICO, Czechia). Measurements were performed in triplicates at room temperature, a frequency of 875 Hz and
214 a peak magnetic field of $300\ \text{A m}^{-1}$.

215 To obtain the magnetite content of the bedrock, a representative $60\times 60\ \text{mm}$ sample slab (Fig. 2a) was mapped with the μ -XRF
216 spectrometer M4 Tornado. The μ -XRF map that depicts only the maximum Fe content was used as an approximation of the
217 magnetite content since magnetite is the mineral with the highest Fe concentration in the rock. Finally, the map was analysed
218 with the image processing program imageJ (version 1.53a; Schneider et al., 2012) to quantify the magnetite content. The
219 obtained value was equalled to the measured magnetic susceptibility of the same sample and used to convert the magnetic
220 susceptibility results of the LC soil pit samples into approximated magnetite contents by the rule of three. The investigated
221 bedrock of NA contains no magnetite.

222 **3.2.6 Light microscopy and electron microprobe analysis (EMPA)**

223 Thin sections were investigated with the light microscope DM750P (Leica, Wetzlar, Germany) equipped with a microscope
224 camera (Euromex, The Netherlands). Electron microprobe element distribution maps of selected areas were obtained for Al,

225 Ca, Fe, K, and Mg by using standard wavelength dispersive techniques on a JEOL Superprobe JXA-8230 fitted with a W-
226 emitter electron gun (accelerating voltage: 15 kV, beam current: 20 nA, beam diameter and step width: 1 µm).

227 **4 Results**


228 The data tables (cited as Table S1–S5) are included in the accompanying data publication (Hampl et al., 2022b).

229 **4.1 Bedrock**

230 According to the Streckeisen nomenclature the bedrock of LC can be described as granodiorite and the investigated bedrock
231 of NA can be described as granite. However, the drill core revealed that the bedrock of NA occasionally contains more mafic
232 sections. The most abundant minerals in the fine-grained bedrock of LC are plagioclase, quartz, microcline, hornblende, biotite,
233 and chlorite (Fig. 2a,b). The latter occurs solely and abundantly along with (former) biotite crystals as their hydrothermal
234 transformation products (i.e., chloritization; e.g., Kogure and Banfield, 2000). Magnetite is a ubiquitous accessory mineral
235 (Fig. 2c; <1 vol.%) in LC and shows no signs of alteration to hematite (martitisation). Pyrite and chalcopyrite are also observed
236 in much smaller abundance than magnetite. Mafic xenoliths can frequently be found in the granodiorite of LC.

237 In the coarse-grained Nahuelbuta granite, quartz, plagioclase, microcline, biotite, and chlorite are the main constituents (Fig.
238 2d,e). In contrast to LC, amphiboles can only be found as an accessory mineral (<1 vol.%) in the investigated bedrock of NA.
239 Like in LC, biotite is often chloritized. Magnetite and sulfides could not be identified in the investigated rock samples of NA.
240 Variations in the biotite content, the occurrence of amphibole crystals, differences in fabric (microcline of a few centimetres),
241 the alternation with mafic portions and the presence of pegmatites in the core make the overall lithology of NA far more
242 heterogenous compared to LC.

243



244 **Figure 2: Bedrock of the investigated profiles. (a) Bedrock from La Campana (IGSN: GFFJH0095) with (b) a corresponding μ -XRF**
 245 **map reflecting the spatial mineral distribution. (c) μ -XRF map of the maximum Fe content (black dots) representing the magnetite**
 246 **crystals in the bedrock sample slab of La Campana. (d) Typical unweathered granite from Nahuelbuta (IGSN: GFFJH00H0) and**
 247 **(e) a μ -XRF map reflecting the mineral content of the same.**


248

249 4.2 Regolith

250 4.2.1 Incipient weathering in rock

251 Weathered rock from the borehole of LC shows abundant indications of weathering-induced fracturing (WIF) due to Fe(II)
 252 oxidation in biotite, like fanned-out edges or opening due to dilatation (Fig. 3a,b). Secondary minerals like Fe(III)
 253 oxyhydroxides are subordinate and are mostly associated with biotite. They are detectable as Fe-enrichments at the edge of
 254 biotite crystals and within the cracks encompassing biotite (Fig. 3c,d). To a minor degree, Fe(III) oxyhydroxides are also
 255 associated with hornblende. Nevertheless, most micro-fractures in feldspar and quartz of the investigated thin sections are
 256 solely filled with blue resin and are bare of any secondary minerals.

257




258 **Figure 3: Rock weathering in La Campana (LC; porosity is represented by blue-dyed resin).** (a) Thin section image (transmitted
 259 **light)** of a weathered rock sample obtained from approx. 27 m depth in the LC drill core (IGSN: GFFJH00HY). (b) A detail image
 260 **of biotite showing signs of dilatation** (dashed lines indicate cleavage planes). (c) Secondary minerals in cracks around biotite. (d) The
 261 **electron microprobe map of the contact zone between biotite and quartz/feldspar displays Fe-enrichments at the interface.** Bt =
 262 **biotite.**

263

264 Indications of WIF around biotite are also present in weathered rock of NA (Fig. 4a,b). However, the cracks are often filled
 265 and covered with Fe(III) oxyhydroxides and clay minerals as observed with light microscopy (Fig. 4c) and electron microprobe
 266 investigations. Unlike LC, weathered rock in NA is characterized by distinct Ca-depletion and Al-enrichment in plagioclase
 267 which indicates partial dissolution (Fig. 4d–f). These alteration sites host secondary minerals covering the newly formed
 268 surfaces which were formed by the dissolution of the plagioclase.

269



270 **Figure 4: Rock weathering in Nahuelbuta (NA).** (a) Thin section image (transmitted light) of weathered rock obtained from approx.
 271 6 m depth in the NA drill core (note that the porosity (blue) is largely associated with weathered plagioclase; IGSN: GFFJH00HX).
 272 (b) Indications of WIF in quartz (backscattered electron image, EMP). (c) Thin section image (transmitted light) of a crack covered
 273 with brown Fe(III) oxyhydroxides from approx. 12 m depth (IGSN: GFFJH00J2). (d) Backscattered electron image (EMP) of partly
 274 dissolved plagioclase and (e) the respective Ca and (f) Al map of the section (IGSN: GFFJH00HX). Qz = quartz, Bt = biotite, Pl =
 275 plagioclase, Kfs = potassium feldspar.

276

277 **4.2.2 Saprolite and soil**


278 **Chemical alteration**

279

280 The mass transfer coefficient τ indicates moderate depletion below 80 cm (not smaller than -0.2) in the LC soil pit profile, but
281 clear depletion in the uppermost few decimetres where Na, K, Mg, Ca, Si and P can reach up to $\tau = -0.5$ and -0.6 (Fig. 5; Table
282 S1). A pronounced P depletion can be detected down to 1.4 m depth in LC. The chemical depletion fraction (CDF) of LC and
283 the bedrock-normalized chemical index of alteration (Δ CIA) indicate a weak chemical weathering degree below ca. 0.5–1 m,
284 but minor chemical depletion was analysed down to the bottom of the 6 m deep profile of LC (see Δ CIA; Fig. 5).

285 In contrast, Nahuelbuta is characterized by distinct chemical depletion of Ca and Na (up to $\tau = -0.9$; Fig. 5). K is depleted to a
286 depth of approximately 5 m, Si to a depth of ~6 m and Mg shows moderate depletion ($\tau \geq -0.3$) throughout the profile. P is
287 strongly depleted between ca. 2–6 m ($\tau \sim -0.6$) but the P content gradually increases from approx. 3 m depth towards the
288 surface and is enriched in the uppermost ~20 cm of the soil (A horizon; Fig. 5). The CDF values of NA indicate depletion
289 down to the bottom of the profile at 6 m depth. The Δ CIA of the profile underpins strong chemical alteration compared to the
290 bedrock (Fig. 5). However, overall chemical depletion decreases towards the bottom of the soil profile and according to the τ -
291 values in 550–600 cm only Na, Ca and P seem to be significantly depleted at >6 m depth.

292




293 **Figure 5:** τ -values of Na, K, Mg, Ca, Si, and P as well as the CDF (all based on Zr) and Δ CIA values of the soil pit profiles in La
 294 **Campana (LC) and Nahuelbuta (NA; Table S1).** Note that the scales are equal for the individual indices of LC and NA. dep. =
 295 **depletion, enr. = enrichment.**

296

297 Since many secondary minerals are formed via a metastable or amorphous precursor (e.g., Steefel and van Cappellen, 1990;
 298 Hellmann et al., 2012; Behrens et al., 2021), we assume that the extractable Fe, Si, and Al contents are indicative for recent
 299 weathering of primary minerals (Fig. 6; see chapter 3.2.2 for an assignment of the extractable elements to the minerals they
 300 likely originate from).

301



302 **Figure 6: Oxalate- and dithionite-extractable Fe, Si and Al contents divided by the respective total element contents of the bulk soil**
 303 **pit samples of La Campana (LC) and Nahuelbuta (NA; Table S2). The elevated ratio at 120–140 cm in (f) (arrow) coincides with the**
 304 **position of a discontinuity in the profile (Fig. 1d). Note that the scales for LC and NA are equal in (a)–(d). The scale in (e) and (f) is**
 305 **one order of magnitude larger for NA compared to LC.**

306

307 Extractable contents of Fe in LC are moderately elevated in the uppermost meter of the profile (up to $Fe_{dit}/Fe_{tot} \sim 14\%$)
 308 compared to the other depth intervals which show low contents ($Fe_{dit}/Fe_{tot} < 1\text{ m}$: $\sim 4\text{--}5\%$; Table S2; Fig. 6a,b). The extractable
 309 Si-contents show no clear pattern (Fig. 6c,d) whereas oxalate-/dithionite-extractable Al-contents are variable in the profile of
 310 LC (Fig. 6e,f). The elevated Al_{dit}/Al_{tot} value in the depth interval 120–140 cm in LC ($\sim 0.5\%$; Fig. 6f) coincides with a
 311 discontinuity in the saprolite (Fig. 1d) and may indicate more secondary crystalline and amorphous Al-bearing phases in this
 312 section. The profile in NA is characterized by high amounts of extractable Fe, Si and Al contents which are especially elevated
 313 in the uppermost meter of the profile (Fe_{dit}/Fe_{tot} up to $\sim 40\%$, Si_{dit}/Si_{tot} up to $\sim 0.14\%$, Al_{dit}/Al_{tot} up to $\sim 12\%$). The extractable
 314 contents rapidly decrease from the surface towards the bottom of the NA profile and starting at approx. 2 m they are similar
 315 down to 6 m (Fig. 6).

316 The Fe₂O₃ content in the investigated bedrock of LC is more than twice as high as that of the NA bedrock, but the oxalate- and
317 dithionite-extractable Fe contents (and hence the amount of the respective secondary minerals) are far higher in NA (Fig. 6a,b).
318 The difference between LC and NA is even more pronounced for the extractable Al contents as values in NA can be 10 times
319 higher than in LC (Fig. 6e,f). The extractable contents in the profiles of both study sites are generally within the range of
320 previous investigations on soil samples from the La Campana and Nahuelbuta National Parks, but the Fe_{dit}/Fe_{tot} contents in the
321 upper profile section of NA in this study are much higher (up to 40 %) than those measured in the Nahuelbuta National Park
322 (<25 %; Oeser et al., 2018).

323

324 **Mineral content and grain sizes**


325

326 The sieving results of LC show a gradual decrease in particle size from the bottom of the profile towards the surface and a
327 relatively constant sand-size content ranging from 65–80 wt.% with similar portions of the individual sand-size fractions (Fig.
328 7a). The small geochemical depletion below the uppermost ~2 m of the LC profile (Fig. 5) is also reflected in the little changing
329 mineral composition of the profile (Fig. 7b). Only the plagioclase (Ca-albite) content slightly decreases from approx. 1 m depth
330 towards the surface. A small decrease of biotite in the depth interval 120–140 cm coincides with the mentioned discontinuity
331 of this profile section (Fig. 1d). The abundant chlorite of the investigated bedrock in LC (~5 wt.%) is completely weathered
332 and absent from the soil pit samples (Fig. 7b).

333 Significant alteration of magnetite (e.g., martitisation) could not be observed in ore microscopic investigations of the magnetic
334 particles in soil pit samples of LC. Thus, the magnetic susceptibility directly reflects the magnetite content of the samples (e.g.,
335 Ferré et al., 2012). A relative magnetite enrichment was detected in the uppermost 40 cm of the LC profile (1–1.6 %) whereas
336 the rest of the profile shows approximately constant magnetite contents (mean ~0.9 %) close to the value of the investigated
337 bedrock (0.94 %; Fig. 7c). This almost consistent magnetite content underlines the homogeneity of the bedrock that was
338 weathered in the 6 m deep soil pit (i.e., no mafic dykes, pegmatites, or major xenoliths).

339 The soil pit profile of NA is characterized by a much higher pebble- and silt/clay-size content compared to LC (Fig. 7d). This
340 reflects the more heterogeneous grain size distribution of the investigated bedrock in NA compared to the bedrock of LC (see
341 Fig. 2). High clay contents can be detected in the uppermost meter of the NA profile (partly >20 wt.%) and the identified
342 mineral content of the soil pit samples differs significantly from the mineral content of the investigated bedrock (Fig. 7e). The
343 plagioclase (Ca-albite) content distinctly decreases from the bottom of the profile towards the surface and the bedrock content
344 of ~28 wt.% partly decreases down to 1 wt.% in the soil pit. The microcline content on the other hand is relatively uniform.
345 Just as in LC, the chlorite of the bedrock analysed here (~1 wt.%) is completely weathered in the NA soil pit profile and is
346 absent from the samples.

347




348 **Figure 7: Sieving/pipetting results, semi-quantitative XRD results and approximated magnetite contents of the investigated soil pit**
 349 **samples in La Campana (LC) and Nahuelbuta (NA). (a) Grain size distribution based on wet-sieving and pipetting, (b) semi-**
 350 **quantitative XRD and (c) magnetic susceptibility results converted to approximate magnetite contents of the LC profile. (d) Wet-**
 351 **sieving combined with pipetting results and (e) semi-quantitative XRD results of the NA samples. Semi-quantitative XRD results of**
 352 **the investigated bedrock samples (see Fig. 2) are given below the results of the soil pit samples. In (a) and (d): 1: ≤ 2000 to >1000 μm ,**
 353 **2: ≤ 1000 to >500 μm , 3: ≤ 500 to >250 μm , 4: ≤ 250 to >125 μm , 5: ≤ 125 to >63 μm .**

354

355 The mineral content of the clay-size fraction in LC differs significantly from that in NA (Fig. 8). La Campana is characterized
 356 by abundant expandable clay minerals (interstratified chlorite-smectite and interstratified mica-smectite) which can largely be
 357 traced back to the weathering of chlorite and biotite (Fig. 8a). Kaolinite can be found throughout the LC profile whereas
 358 interstratified mica-vermiculite only occurs in the depth interval of the discontinuity (120–140 cm; see Fig. 1d). The
 359 expandable portion of the interstratified minerals gradually decreases from the profile bottom towards the surface and cannot
 360 be detected in the uppermost centimetres of the LC profile. Only mica and kaolinite constitute the clay-size fraction of the
 361 uppermost part of the profile in LC. The mineral content in the clay-size fraction of NA is characterized by small amounts of
 362 interstratified mica-vermiculite below 1 m depth and ubiquitous kaolinite which shows small expandable portions below 2 m
 363 depth. Hydroxy-interlayered vermiculite (HIV) and gibbsite can first be detected in 400–450 cm depth and the content
 364 increases towards the surface. The main minerals of the clay-size fraction in the uppermost part of the profile are HIV, kaolinite,
 365 and gibbsite (Fig. 8b).

366



367 **Figure 8: Minerals in the clay-size fraction of the soil pit profiles in La Campana (LC) and Nahuelbuta (NA).** (a) The profile in LC
 368 features abundant expandable clay minerals. (b) NA is characterized by the presence of gibbsite and vermiculite but very minor
 369 amounts of expandable clay minerals. C-S = interstratified chlorite-smectite, G = gibbsite, HIV = hydroxy-interlayered vermiculite,
 370 K = kaolinite, K+ = kaolinite with expandable portions, M-S = interstratified mica-smectite, M-V = interstratified mica-vermiculite.

371 5 Discussion

372 5.1 Climate-dependent mineral transformations

373 Chemical depletion and mineral transformations are far more pronounced in the profile of NA compared to the profile of LC
 374 even though the bedrock of LC contains more minerals with higher solubility compared to NA (more plagioclase, biotite,
 375 chlorite or hornblende in LC than in NA where quartz and potassium feldspar dominate; see e.g., Wilson, 2004; Bandstra et
 376 al., 2008). The high chemical depletion ($\tau[\text{Na}, \text{Ca}]$ up to -0.9 and ΔCIA up to 22; Fig. 5) and the occurrence of gibbsite in NA
 377 are indicative of distinct dissolution of primary minerals (esp. plagioclase; Fig. 7e) and solute removal of alkali and alkaline
 378 earth metals while immobile Al remains as hydroxide ($\text{Al}(\text{OH})_3$ = gibbsite). This depletion is assumed to be the result of more
 379 water infiltration into the subsurface of NA (more precipitation due to humid climate) compared to LC (less precipitation due
 380 to Mediterranean climate). The measured $\tau[\text{P}]$ distribution in NA is a clear indication for biologically controlled nutrient uplift


381 and recycling within the topsoil (Jobbágy and Jackson, 2004). We assume that the high precipitation rate in NA leads to more
382 biomass production by plants, which in turn implies more litter production and a stimulation of biogenic decay that supplies
383 plants with nutrients. Thus, we concur with the hypothesis that the ecosystem in NA is thriving on nutrient recycling rather
384 than on an uptake of nutrients that were released by biogenic weathering at depth (Oeser and von Blanckenburg, 2020). Apart
385 from Ca, Na, and P (τ -values in Fig. 5), the chemical depletion successively decreases from the surface towards the bottom
386 part of the investigated profile in NA. To account for this shallow chemical depletion, we propose that a secondary-mineral-
387 controlled impeding of the fluid infiltration to depth is playing an important role for the depth of mineral transformations in
388 NA.

389 Chemical depletion can be detected throughout the investigated profile in LC, but the chemical weathering degree is very low
390 between 2–6 m depth (Fig. 5) and the mineral transformations in this section of the profile are only minor (Fig. 7b). On the
391 other hand, distinct mineral dissolution and removal of solutes is testified by the higher magnetic susceptibility values in the
392 uppermost decimetres of the LC profile. This can be related to a residual accumulation of weathering-resistant magnetite while
393 other minerals like plagioclase dissolve. The strong chemical depletion in this part of the profile is also reflected by the low τ -
394 and elevated CDF-/ Δ CIA-values. To account for the detected weak but deep chemical weathering in LC, we propose that a
395 secondary-mineral-controlled formation of fluid pathways is facilitating the fluid infiltration to depth and is thus an important
396 control on the chemical weathering reactions in the subsurface.

397 The difference between the profiles is also displayed by the oxalate- and dithionite-extractable Fe, Si, and Al contents. While
398 high extractable contents especially within the uppermost 2 m of the NA profile are interpreted to indicate considerable
399 ongoing (recent) transformations of primary to secondary minerals, LC shows comparatively little indications in this regard.
400 This difference underlines the higher degree of mineral transformations in NA compared to LC which is also reflected in the
401 mineral content of the clay-size fraction (see Fig. 9). That oxalate- frequently exceeds dithionite-extractable Al contents is
402 indicative for amorphous phases since oxalate is more effective at extracting amorphous forms of Al (McKeague and Day,
403 1966). Moreover, the highest contents of the clay-size fraction in the profiles are in good correlation with the elevated
404 extractable Fe, Si and Al contents and highlight the pronounced mineral transformation in the uppermost part of the profiles.
405 This size fraction hosts most of the products of primary silicate weathering. Clay-size minerals of NA mainly correspond to
406 distinct weathering of plagioclase and biotite, whereas in LC they can mainly be associated with chlorite and biotite weathering
407 (Fig. 9). Feldspar weathers to kaolinite and gibbsite in NA and biotite weathers to hydroxy-interlayered vermiculite (HIV).
408 Chlorite completely dissolved in the NA profile, whereas both chlorite and biotite in LC weather via interstratified clay
409 minerals to smectite. Finally, smectite and feldspar likely weather to kaolinite in LC (Fig. 9). The mineral composition of the
410 clay-size fraction (Fig. 8) is dependent on the bedrock composition (e.g., more chlorite in LC) and the climate-dependent
411 mineral dissolution (see Fig. 9) in the study sites. However, we argue that the amount of secondary minerals is largely a
412 function of the climatic conditions that control the weathering intensity via water availability in the study sites.

413 As this study does not consider the entire weathering profile in LC and NA, the interplay between erosion rate and weathering
414 advance rate (Lebedeva and Brantley, 2020) is not addressed here. However, the different denudation rates in the study areas

415 (mean soil denudation rate in LC: ~ 61, in NA: ~ 33 t km⁻² yr⁻¹; Oeser et al., 2018) likely affect the weathering intensity. Due
 416 to the higher denudation rates in LC compared to NA (Oeser et al., 2018; van Dongen et al., 2019), we hypothesize that the
 417 residence time of weathered material in the regolith of LC is shorter than in NA. Thus, there is less time for chemical weathering
 418 in LC. In combination with the lower water availability in LC, this factor might contribute to the lower weathering intensity
 419 in the regolith of LC compared to NA. NA, on the other hand, is characterized by a longer residence time of weathered material.
 420 Together with the higher water availability, this factor might contribute to the high weathering intensity in the upper regolith
 421 of NA. The situation in LC may be comparable to an incompletely developed profile, and the situation in NA to a completely
 422 developed profile (Reis and Brantley, 2019).
 423



424 Figure 9: Schematic diagram showing the transformation of primary minerals to secondary minerals (clay minerals and aluminium
 425 hydroxide) depending on the activities of H⁺, Si, Al, K, Na, Mg and Ca. The depletion of the alkali and alkaline earth metals, and
 426 the increase of the Al activity are coupled to an increase of the mineral dissolution and the removal of solutes by a higher subsurface

427 **fluid flow. Elevated a_{H^+} -values (i.e., lower pH) increase the mineral solubility. Modified from Chesworth et al. (2008). HIV = hydroxy-**
428 **interlayered vermiculite, a = thermodynamic activity**

429

430 **5.2 Weathering-intensifying processes**

431 **5.2.1 Porosity increase by weathering-induced fracturing and its impact on the weathering depth**

432 Ferrous primary minerals of the LC granodiorite can frequently be identified as initiating locations of micro-cracks. This
433 observation can be related to weathering-induced fracturing (WIF) due to the increase in volume caused by the oxidation of
434 Fe(II) in Fe(II)-bearing silicates (e.g., Buss et al., 2008; Behrens et al., 2015; Kim et al., 2017) and the formation of secondary
435 Fe(III) oxyhydroxides (Fletcher et al., 2006; Lebedeva et al., 2007; Anovitz et al., 2021; Fig. 3d). This process generates and
436 increases surface areas of primary minerals and in turn accelerates weathering reactions (positive feedback between the
437 formation of secondary minerals and the infiltration of fluids (esp. O₂ and water) to depth; e.g., Røyne et al., 2008). These
438 weathering-induced fractures consequently facilitate the presence of surface-derived O₂ in the deep subsurface (Kim et al.,
439 2017) and the corresponding transport through the saprolite/soil is dominated by advection (Lebedeva et al., 2007). The
440 bedrock of LC is richer in Fe-bearing minerals than the investigated granite of NA (ca. 25 wt.% in LC and ca. 10 wt.% in NA)
441 and hosts biotite, hornblende, chlorite, and magnetite as Fe(II) sources. A considerable amount of the total Fe content is bound
442 in magnetite (roughly 0.7 wt.% of the total Fe₂O₃ content if the magnetite content of the bulk sample is 1 wt.%). However, we
443 found no microscopic evidence (no oxidation) nor indications in the magnetic susceptibility results that the Fe(II) in magnetite
444 is available for weathering reactions. Thus, we conclude that magnetite is stable under the environmental conditions of LC. Of
445 the three remaining Fe(II)-bearing minerals, biotite was found to be the most important one for the generation of WIF in LC
446 (see also Buss et al., 2008; Bazilevskaya et al., 2013, 2015) due to its volumetric expansion during weathering (e.g.,
447 Goodfellow et al., 2016). Although WIF also occurs in NA (Fig. 4b) it does not seem to significantly increase the permeability
448 of the rock which can be related to the low Fe(II) content of the dominant bedrock (Fe₂O₃ (total Fe): <3 wt.%; Table S1; see
449 Kim et al., 2017).

450 Other than that, chlorite is suggested to be an important mineral in the development of the investigated weathering profile in
451 LC. The original chlorite content of the bedrock in LC has been completely transformed into interstratified chlorite-smectite
452 in the soil pit profile. We suggest that this transformation plays a significant role for the development of the LC profile since
453 expandable clay minerals are known to disaggregate rock by swelling (e.g., Dunn and Hudec, 1966; Jiménez-González et al.,
454 2008). The ensuing fracturing also forms new fluid pathways and new access to reactive surfaces of primary minerals which
455 in turn fosters weathering reactions (positive feedback mechanism; see e.g., Røyne et al., 2008). Even though expandable clay
456 minerals can also cause sealing of the subsurface (Kim et al., 2017) we do not regard this effect as significant for LC since
457 clay contents are very low. However, a minor retardation of the fluid flow from surface to depth due to the expansion of the
458 interstratified clay minerals in LC (Fig. 8a) cannot be excluded (see Kim et al., 2017). In conclusion, we propose that small

459 amounts of expandable clay minerals like in LC can generate porosity whereas high amounts of expandable clay minerals can
460 reduce porosity.
461 The feedback mechanism of weathering-induced fracturing is presented here for granodiorite. However, the significance of
462 this mechanism is not restricted to plutonic rocks. Weathering-induced fracturing requires Fe(II)-bearing minerals such as
463 biotite and/or potentially the presence of expandable clay minerals that cause the formation of cracks by volume increase
464 during weathering. This feedback concept is thus transferable to all igneous, metamorphic and sedimentary rocks that contain
465 these minerals.

466 **5.2.2 Increase of weathering intensity by biogenic activity**

467 The formation of secondary minerals such as clay minerals and aluminium hydroxide is among other factors controlled by
468 biogenic activity since organic acids and an acidity-increase by elevated organic-derived CO₂ contents accelerate dissolution
469 rates of primary minerals (see e.g., Lucas, 2001; Lawrence et al., 2014). This effect needs to be considered for the organic-rich
470 and acidic subsurface of NA (see Bernhard et al., 2018). The acidity likely contributes to the high degree of mineral dissolution
471 in NA (see a_{H^+} in Fig. 9), which consequently leads to an increased formation of secondary minerals.

472 The depth interval 120–140 cm in LC is characterized by lower amounts of biotite and a different clay mineral composition
473 compared to the surrounding depth intervals (Fig. 7b; Fig. 8a). This depth interval coincides with a discontinuity crossing the
474 entire profile width (Fig. 1d). We interpret this plant-root-containing discontinuity in the saprolite as a fracture remnant since
475 there are no indications of a lithological heterogeneity in this zone (e.g., a significant change in the magnetic susceptibility or
476 of the primary mineral content; Fig. 7b,c). To explain the lower biotite content and the different clay composition in this part
477 of the profile, we propose an intensification of weathering reactions in the vicinity of the fracture fostered by the observed
478 plant roots (e.g., Fimmen et al., 2008; Pawlik et al., 2016; Nascimento et al., 2021). This weathering-promoting mechanism
479 might account for the increase in interstratified chlorite-smectite and the appearance of interstratified mica-vermiculite (Fig.
480 8a), while the amount of biotite decreases due to its transformation to secondary minerals (Fig. 7b).

481 **5.3 Weathering-mitigating processes**

482 **5.3.1 O₂ consumption by Fe-bearing silicates and its impact on the weathering depth and intensity**

483 The granodiorite of LC hosts an abundance of Fe(II)-bearing minerals (Fig. 7b). The Fe₂O₃ content of the LC bedrock after
484 subtraction of the inert magnetite-bound Fe₂O₃ fraction (since 100 % pure magnetite contains 69 % Fe₂O₃, 0.94 % magnetite
485 as analysed in the LC bedrock equals to 0.65 % magnetite-bound Fe₂O₃ which needs to be subtracted) is 5.34 wt.% (for
486 comparison: 2.33 wt.% Fe₂O₃ in NA). Since O₂ is reduced by the oxidation of mineral-bound Fe(II) (e.g., White and Yee,
487 1985; Perez et al., 2005) and the consequent formation of secondary minerals, the O₂ content and hence oxidative weathering
488 reactions are expected to decrease from surface to depth. A rapid decrease of the O₂ concentration to depth is characteristic for
489 weathering systems in which O₂ transport is dominated by diffusion (Behrens et al., 2015). Given the observed deep fracturing

490 due to Fe(II) oxidation (i.e., WIF) in LC and the consequent deep connectivity between the surface and the subsurface (Kim
491 et al., 2017), the O₂ transport in LC is most likely dominated by advection. As a consequence, diffusive O₂ transport is
492 insignificant in the upper regolith of LC and the O₂ consumption by Fe(II) oxidation is not limiting the regolith depth in LC
493 (compare Bazilevskaya et al., 2013). It has been argued that WIF and thus a thicker regolith is more likely when the ratio
494 pO₂/pCO₂ in soil water is greater than the ratio of the capacity for O₂ consumption to the capacity for CO₂ consumption in
495 bedrock (Stinchcomb et al., 2018). In the study sites, decomposition of organic matter is restricted to the topsoil, likely because
496 organic matter at depth becomes stabilized against microbial decomposition (Scheibe et al., 2023). Thus, we suggest that the
497 pCO₂ of water in the deeper profile part of LC is low (i.e., pO₂/pCO₂ is high), and O₂ is not being consumed by organic matter
498 decomposition but is available for Fe(II) oxidation and hence WIF. The WIF-controlled connectivity between the surface and
499 the subsurface results in an O₂ availability for oxidative weathering processes at great depth. On the other hand, the weak
500 chemical weathering in LC is in good agreement with the low precipitation rate (~350 mm yr⁻¹; Übernickel et al., 2020). The
501 low precipitation rate entails a small infiltration of water to depth and hence minor primary mineral dissolution and thus
502 chemical weathering at depth.

503 The cracks around weathered biotite in the investigated samples of LC are (mainly) filled with Fe(III) oxyhydroxides as
504 revealed by the high Fe-enrichment detected in electron microprobe maps (Fig. 3d). Newly formed weathering-induced
505 fractures make the biotite more accessible to surface inputs like water and O₂ which promotes the dissolution of biotite. The
506 solutes formed as a result migrate along the weathering-induced cracks and precipitate in the vicinity of the biotite crystal as
507 secondary phases (Fig. 3c). Thus, we propose that the reactive surface of biotite is partly shielded from weathering reactants
508 (water, O₂) due to the precipitation of secondary minerals (see e.g., Navarre-Sitchler et al., 2015; Vázquez et al., 2016; Gerrits
509 et al., 2020; 2021). Combined with the low subsurface water availability in LC causing a low mineral dissolution degree, this
510 shielding might contribute to the relatively stable biotite content throughout the LC profile (Table S4).

511 **5.3.2 Reduction of weathering intensity and -depth by damping of fluid flow**

512 The formation of secondary minerals such as clay minerals (via amorphous and poorly crystalline precursors; see Fig. 6) can
513 decrease the porosity (e.g., Bazilevskaya et al., 2015; Navarre-Sitchler et al., 2015) formed by WIF and dissolution. Al-rich
514 phases were found as precipitates in partly dissolved plagioclase of NA (Fig. 4d–f) and within cracks which can often be
515 identified as weathering-induced. We suggest that the abundant presence of clay minerals and gibbsite in NA restricts the fluid
516 flow through such fractures and pores. The clay-rich zone in the uppermost metre of the NA soil pit profile (around 50 cm
517 depth; Fig. 7d) likely acts as a (partially) shielding horizon (impeding vertical flow of surface inputs to the deep subsurface;
518 see e.g., Lohse and Dietrich, 2005). Clay-rich horizons can therefore influence the dynamic of the subsurface fluid flow and
519 thus mitigate mineral transformations and chemical weathering at depth. At the same time, these conditions foster a long fluid
520 residence time in the upper regolith and thus promote the precipitation of secondary minerals such as clay minerals that may
521 impel the weathering of primary minerals in the upper part of the weathering profile (see Maher, 2010). However, the seasonal


522 sealing of fractures and pore spaces due to an increase of soil moisture and an ensuing clay expansion (Kim et al., 2017) is not
523 assumed for NA as expandable secondary minerals barely occur in the clay-size fraction of NA (Fig. 8b).
524 The negative feedback mechanism presented here is demonstrated for granite. However, the concept is essentially based on
525 newly formed minerals such as clay minerals that inhibit the subsurface fluid flow by blocking pathways. This feedback
526 mechanism can thus be significant for weathering systems developing from all igneous, metamorphic and sedimentary rocks
527 where secondary minerals can block the permeable porosity formed by WIF and primary mineral dissolution.

528 **6 Conclusions**

529 In two 6 m deep weathering profiles formed on granitic rock in two climatic zones (Mediterranean and humid climate), we
530 found different degrees of elemental loss by chemical weathering, and different secondary minerals. Under Mediterranean
531 climate conditions (La Campana), Fe(II)-oxidation, precipitation of Fe(III) oxyhydroxide and clay swelling lead to fracturing
532 and the formation of fluid pathways. This weathering-induced fracturing (WIF) is likely one of the dominant controls on the
533 development of the upper regolith as it leads to a deep infiltration of surface inputs (esp. water and O₂) which in turn causes
534 further WIF. While the intensity of chemical weathering at the Mediterranean site is low, it was detected throughout the entire
535 6 m deep profile. This suggests that the weathering front is located at much greater depth in La Campana. The overall low
536 abundance of secondary minerals can be explained by the low climate-related subsurface water availability in La Campana.
537 The lack of large quantities of secondary minerals ensures that fractures and porosity generated by WIF remain accessible to
538 water and gases. Thus, we conclude that the development of the deep but weak chemical weathering in the upper regolith of
539 La Campana is significantly controlled by two mechanisms: (1) A positive feedback loop between the formation of secondary
540 minerals and the infiltration of fluids to depth induced by (mainly) biotite weathering (WIF) which leads to a deep surface-
541 subsurface connectivity for weathering reactants (in particular O₂). (2) Low subsurface water availability resulting in a low
542 amount of secondary minerals which would otherwise seal this connectivity.

543 Under humid climate conditions (Nahuelbuta), clay minerals, gibbsite as well as amorphous and poorly crystalline secondary
544 minerals largely formed due to intense plagioclase dissolution. We link this intense dissolution to the high climate-related
545 subsurface water availability in Nahuelbuta. The secondary minerals thus formed are suggested to impede the flow of surface
546 inputs to depth. Moreover, the generally lower amount of Fe(II)-bearing silicates in Nahuelbuta compared to La Campana
547 results in less WIF and thus fewer fluid pathways. Therefore, we conclude that the development of the weathering profile in
548 Nahuelbuta is predominantly governed by two mechanisms: (1) Considerable climate-related subsurface water availability and
549 high biogenic activity which lead to intense weathering of primary minerals in the upper part of the regolith. (2) A negative
550 feedback loop between the formation of secondary minerals and the infiltration of fluids to depth induced by (mainly)
551 plagioclase weathering and the ensuing formation of secondary minerals which leads to a poor surface-subsurface connectivity
552 for weathering reactants. The main findings and factors that are most relevant to the development of the different weathering
553 systems are summarized in Figure 10.

554 The relationship between precipitation and the degree of chemical weathering along the climate gradient of the Chilean Coastal
 555 Cordillera was found to be non-linear and non-systematic (Oeser and von Blanckenburg, 2020; Schaller and Ehlers, 2022).
 556 We argue that a systematic relationship is likely concealed by variations in the mineral content of the bedrocks and the
 557 associated feedback mechanisms. However, the investigated feedbacks provide a causal explanation for the depth of chemical
 558 weathering. This study illustrates how the formation of secondary minerals and the infiltration of surface-derived fluids to
 559 depth are interlinked by positive and negative feedback loops. We demonstrated that these feedback loops and the climatic
 560 conditions they occur in are important controls on the development of the upper regolith.
 561



562
 563 **Figure 10: Schematic summary of the two weathering systems. According to our findings, the regolith of La Campana (LC) is**
 564 **dominated by a positive feedback loop between weathering-induced fracturing (WIF) and the infiltration of fluids to depth. WIF**
 565 **creates deep-reaching pathways for fluids (water, O₂) and hence a good connectivity between the surface and the subsurface.**
 566 **Moreover, the low water availability in the Mediterranean climate inhibits the formation of large amounts of secondary minerals**
 567 **(i.e., low weathering intensity) that could seal these pathways. The high denudation rate in LC results in a short residence time of**
 568 **weathered material in the profile and could therefore contribute to the detected lower weathering intensity (i.e., less chemical**
 569 **weathering). The regolith of Nahuelbuta (NA), on the other hand, was found to be dominated by a negative feedback loop between**
 570 **the formation of secondary minerals and amorphous phases, and the infiltration of fluids to depth. These secondary solids are**
 571 **consequences of the high water availability in NA that results in intense chemical weathering (i.e., high weathering intensity). The**
 572 **high weathering intensity entails the formation of abundant secondary minerals and amorphous phases that reduce the connectivity**
 573 **between the surface and the subsurface. The lower denudation rate and thus longer residence time of weathered material in NA**
 574 **likely contributes to the more intense chemical weathering. dep. = depletion.**
 575

576 **Data availability**

577 Datasets related to this article can be found in the data publication Hampl et al. (2022b). The data publication is hosted at the
578 GFZ Data Services and can be downloaded by clicking on “Download data and description” in the field “Files”.
579

580 **Sample availability**

581 The IGSN-registered samples used in this article are deposited at the Department of Applied Geochemistry (Technische
582 Universität Berlin) and are listed in the data publication of this paper (Hampl et al., 2022b).
583

584 **Author contribution**

585 Ferdinand J. Hampl: conceptualization, methodology, investigation, writing – original draft preparation
586 Ferry Schiperski: methodology, supervision, writing – review & editing
587 Christopher Schwerdhelm: investigation, writing – review & editing
588 Nicole Stroncik: investigation, writing – review & editing
589 Casey Bryce: funding acquisition, writing – review & editing
590 Friedhelm von Blanckenburg: supervision, writing – review & editing
591 Thomas Neumann: funding acquisition, supervision, writing – review & editing
592

593 **Competing interests**

594 The authors declare that they have no conflict of interest.
595

596 **Acknowledgements**

597 This work was supported by the German Research Foundation (DFG) priority research program SPP-1803 “EarthShape: Earth
598 Surface Shaping by Biota” (grant number NE 687/9-1) and the EarthShape Coordination (EH 329/17-2, BL562/20-1). We are
599 grateful to Dr. Kirstin Übernickel for the management of the drilling campaigns and to Prof. Andreas Kappler for his support.
600 We would also like to thank Michael Facklam for his help in determining the clay content and Dr. Katja Emmerich for her
601 valuable hints on the clay mineralogy. The authors would also like to thank Prof. Dr. Peter Finke and Prof. Dr. Susan L.

602 Brantley for their valuable comments and suggestions that greatly improved the manuscript. Finally, we are grateful to Antonia
603 Roesrath for her help in registering the samples.
604

605 **References**

606 Anderson, S., Dietrich, W., and Brimhall, G.: Weathering profiles, mass-balance analysis, and rates of solute loss: Linkages
607 between weathering and erosion in a small, steep catchment, *Geol. Soc. Am. Bull.*, 114 (9), 1143–1158,
608 [https://doi.org/10.1130/0016-7606\(2002\)114<1143:WPMBAA>2.0.CO;2](https://doi.org/10.1130/0016-7606(2002)114<1143:WPMBAA>2.0.CO;2), 2002.

609 Anovitz, L.M., Cheshire, M.C., Hermann, R.P., Gu, X., Sheets, J.M., Brantley, S.L., Cole, D.R., Ilton, E.S., Mildner, D.F.R.,
610 Gagnon, C., Allard, L.F., and Littrell, K.C.: Oxidation and associated pore structure modification during experimental
611 alteration of granite, *Geochim. Cosmochim. Ac.*, 292, 532–556, <https://doi.org/10.1016/j.gca.2020.08.016>, 2021.

612 Bandstra, J. Z., Buss, H. L., Campen, R. K., Liermann, L. J., Moore, J., Hausrath, E. M., Navarre-Sitchler, A. K., Jang, J.-H.,
613 and Brantley, S. L.: Compilation of mineral dissolution rates, in: *Kinetics of Water-Rock Interaction*, edited by: Brantley, S.
614 L., Kubicki, J. D., and White, A. F., Springer New York, 737–823, <https://doi.org/10.1007/978-0-387-73563-4>, 2008.

615 Bazilevskaya, E., Lebedeva, M., Pavich, M., Rother, G., Parkinson, D. Y., Cole, D., and Brantley, S. L.: Where fast weathering
616 creates thin regolith and slow weathering creates thick regolith, *Earth Surf. Proc. Land.*, 38, 847–858,
617 <https://doi.org/10.1002/esp.3369>, 2013.

618 Bazilevskaya, E., Rother, G., Mildner, D. F., Pavich, M., Cole, D., Bhatt, M. P., Jin, L., Steefel, C. I., and Brantley, S. L.: How
619 Oxidation and Dissolution in Diabase and Granite Control Porosity during Weathering, *Soil Sci. Soc. Am. J.*, 79, 55–73,
620 <https://doi.org/10.2136/sssaj2014.04.0135>, 2015.

621 Behrens, R., Bouchez, J., Schuessler, J. A., Dultz, S., Hewawasam, T., and von Blanckenburg, F.: Mineralogical
622 transformations set slow weathering rates in low-porosity metamorphic bedrock on mountain slopes in a tropical climate,
623 *Chem. Geol.*, 411, 283–298, <https://doi.org/10.1016/j.chemgeo.2015.07.008>, 2015.

624 Behrens, R., Wirth, R., and von Blanckenburg, F.: Rate limitations of nano-scale weathering front advance in the slow-eroding
625 Sri Lankan Highlands, *Geochim. Cosmochim. Ac.*, 311, 174–197, <https://doi.org/10.1016/j.gca.2021.06.003>, 2021.

626 Bernhard, N., Moskwa, L.-M., Schmidt, K., Oeser, R. A., Aburto, F., Bader, M. Y., Baumann, K., von Blanckenburg, F., Boy,
627 J., van den Brink, L., Brucker, E., Büdel, B., Canessa, R., Dippold, M. A., Ehlers, T. A., Fuentes, J. P., Godoy, R., Jung, P.,
628 Karsten, U., Köster, M., Kuzyakov, Y., Leinweber, P., Neidhardt, H., Matus, F., Mueller, C. W., Oelmann, Y., Oses, R., Oses,

629 P., Paulino, L., Samolov, E., Schaller, M., Schmid, M., Spielvogel, S., Spohn, M., Stock, S., Stroncik, N., Tielbörger, K.,
630 Übernicker, K., Scholten, T., Seguel, O., Wagner, D., and Kühn, P.: Pedogenic and microbial interrelations to regional climate
631 and local topography: New insights from a climate gradient (arid to humid) along the Coastal Cordillera of Chile, CATENA,
632 170, 335–355, <https://doi.org/10.1016/j.catena.2018.06.018>, 2018.

633 Buss, H. L., Sak, P. B., Webb, S. M., and Brantley, S. L.: Weathering of the Rio Blanco quartz diorite, Luquillo mountains,
634 Puerto Rico: coupling oxidation, dissolution, and fracturing, *Geochim. Cosmochim. Ac.*, 72, 4488–4507,
635 <https://doi.org/10.1016/j.gca.2008.06.020>, 2008.

636 Chesworth W. et al.: Clay Mineral Formation, in: *Encyclopedia of Soil Science, Encyclopedia of Earth Sciences Series*, edited
637 by: Chesworth, W., Springer, Dordrecht, Figure C51, https://doi.org/10.1007/978-1-4020-3995-9_108, 2008.

638 Dawson, T. E., Hahm, W. J., and Crutchfield-Peters, K.: Digging deeper: what the critical zone perspective adds to the study
639 of plant ecophysiology, *New Phytol.*, 226, 666–671, <https://doi.org/10.1111/nph.16410>, 2020.

640 Deckart, K., Hervé, F., Fanning, C., Ramírez, V., Calderón, M., and Godoy, E.: U-Pb Geochronology and Hf-O Isotopes of
641 zircons from the Pennsylvanian Coastal Batholith, South-Central Chile, *Andean Geol.*, 41 (1), 49–82,
642 <https://doi.org/10.5027/andgeoV41n1-a03>, 2013.

643 Drever, J. I.: The effect of land plants on weathering rates of silicate minerals, *Geochim. Cosmochim. Ac.*, 58, 2325–2332,
644 [https://doi.org/10.1016/0016-7037\(94\)90013-2](https://doi.org/10.1016/0016-7037(94)90013-2), 1994.

645 Dunn, J. R., and Hudec, P. P.: Water, Clay and Rock Soundness, *Ohio J. Sci.*, 66 (2), 153–168, 1966.

646 Ferraris, F.: Hoja Los Angeles – Angol, Escala: 1:250.000, Carta Geológica de Chile N°5 (Preliminar), Instituto de
647 Investigaciones Geológicas, OCLC number: 1024799672, 1979.

648 Ferré, E., Michelsen, K., Ernst, W., Boyd, J., and Cañón-Tapia, E.: Vertical zonation of the Barcroft granodiorite, White
649 Mountains, California: Implications for magmatic processes, *Am. Mineral.*, 97, 1049–1059,
650 <http://dx.doi.org/10.2138/am.2012.4013>, 2012.

651 Fimmen, R., Richter, D., Vasudevan, D., Williams, M., and West, L.: Rhizogenic Fe-C redox cycling: A hypothetical
652 biogeochemical mechanism that drives crustal weathering in upland soils, *Biogeochemistry*, 87, 127–141,
653 <https://doi.org/10.1007/s10533-007-9172-5>, 2008.

654 Fletcher, R. C., Buss, H. L., and Brantley, S. L.: A spheroidal weathering model coupling porewater chemistry to soil
655 thicknesses during steady-state denudation, *Earth Planet. Sc. Lett.*, 244, 444–457, <https://doi.org/10.1016/j.epsl.2006.01.055>,
656 2006.

657 Gana, P., Wall, R., and Gutiérrez, A.: Mapa geológico del área de Valparaiso-Curacavi, Región de Valparaiso y Región
658 Metropolitana, Mapas Geológicos, N° 1, Escala 1:100 000, Servicio Nacional de Geología y Minería (Chile), OCLC number:
659 43901854, 1996.

660 Gerrits, R., Pokharel, R., Breitenbach, R., Radnik, J., Feldmann, I., Schuessler, J. A., von Blanckenburg, F., Gorbushina, A.
661 A., and Schott, J.: How the rock-inhabiting fungus *K. petricola* A95 enhances olivine dissolution through attachment, *Geochim.*
662 *Cosmochim. Ac.*, 282, 76–97, <https://doi.org/10.1016/j.gca.2020.05.010>, 2020.

663 Gerrits, R., Wirth, R., Schreiber, A., Feldmann, I., Knabe, N., Schott, J., Benning, L. G., and Gorbushina, A. A.: High-
664 resolution imaging of fungal biofilm-induced olivine weathering, *Chem. Geol.*, 559, 119902,
665 <https://doi.org/10.1016/j.chemgeo.2020.119902>, 2021.

666 Glodny, J., Echtler, H., Collao, S., Ardiles, M., Buron, P., and Figueroa, O.: Differential Late Paleozoic active margin evolution
667 in South-Central Chile (37° S–40° S) – the Lanalhue Fault Zone. *J. S. Am. Earth Sci.*, 26, 397–411,
668 <https://doi.org/10.1016/j.jsames.2008.06.001>, 2008a.

669 Glodny, J., Gräfe, K., Echtler, H., and Rosenau, M.: Mesozoic to Quaternary continental margin dynamics in south-central
670 Chile (36–42°S): The apatite and zircon fission track perspective, *Int. J. Earth Sci.*, 97, 1271–1291,
671 <https://doi.org/10.1007/s00531-007-0203-1>, 2008b.

672 Goodfellow, B. W., Hilley, G. E., Webb, S. M., Sklar, L. S., Moon, S., and Olson, C. A.: The chemical, mechanical, and
673 hydrological evolution of weathering granitoid, *J. Geophys. Res.-Earth*, 121, 1410–1435,
674 <https://doi.org/10.1002/2016JF003822>, 2016.

675 Hampl, F. J., Schiperski, F., Byrne, J. M., Schwerdhelm, C., Kappler, A., Bryce, C., von Blanckenburg, F., and Neumann, T.:
676 The role of iron-bearing minerals for the deep weathering of a hydrothermally altered plutonic rock in semi-arid climate
677 (Chilean Coastal Cordillera), *Chem. Geol.*, 604, 120922, <https://doi.org/10.1016/j.chemgeo.2022.120922>, 2022a.

678 Hampl, F. J., Schiperski, F., Schwerdhelm, C., Stroncik, N., Bryce, C., von Blanckenburg, F., and Neumann, T.: Mineralogical
679 and geochemical data of two weathering profiles in a Mediterranean and a humid climate region of the Chilean Coastal
680 Cordillera, GFZ Data Services [data set], <https://doi.org/10.5880/fidgeo.2022.035>, 2022b.

681 Hayes, N. R., Buss, H. L., Moore, O. W., Krám, P., and Pancost, R. D.: Controls on granitic weathering fronts in contrasting
682 climates, *Chem. Geol.*, 535, 119450, <https://doi.org/10.1016/j.chemgeo.2019.119450>, 2020.

683 Hellmann, R., Wirth, R., Daval, D., Barnes, J.-P., Penisson, J.-M., Tisserand, D., Epicier, T., Florin, B., and Hervig, R. L.:
684 Unifying natural and laboratory chemical weathering with interfacial dissolution–reprecipitation: A study based on the
685 nanometer-scale chemistry of fluid–silicate interfaces, *Chem. Geol.*, 294–295, 203–216,
686 <https://doi.org/10.1016/j.chemgeo.2011.12.002>, 2012.

687 Hervé, F.: Petrology of the crystalline basement of the Nahuelbuta Mountains, southcentral Chile, in: *Comparative Studies on*
688 *the Geology of the Circum-Pacific Orogenic Belt in Japan and Chile*, edited by: Ishikawa, T., and Aguirre, I., Japan Society
689 for the Promotion of Science, Tokyo, 1-51, 1977.

690 Holbrook, S., Marcon, V., Bacon, A., Brantley, S., Carr, B., Flinchum, B., Richter, D., and Riebe, C.: Links between physical
691 and chemical weathering inferred from a 65-m-deep borehole through Earth’s critical zone, *Sci. Rep.-UK*, 9, 4495,
692 <https://doi.org/10.1038/s41598-019-40819-9>, 2019.

693 Holmgren, G. G. S.: A Rapid Citrate-Dithionite Extractable Iron Procedure, *Soil Sci. Soc. Am. J.*, 31, 210–211,
694 <https://doi.org/10.2136/sssaj1967.03615995003100020020x>, 1967.

695 Hynek, S., Comas, X., and Brantley, S. L.: The effect of fractures on weathering of igneous and volcanoclastic sedimentary
696 rocks in the Puerto Rican tropical rain forest, *Proced. Earth Plan. Sc.*, 17, 972–975,
697 <https://doi.org/10.1016/j.proeps.2017.01.001>, 2017.

698 Israeli, Y., Salhov, E., and Emmanuel, S.: Impact of textural patterns on modeled rock weathering rates and size distribution
699 of weathered grains, *Earth Surf. Proc. Land.*, 46, 1177–1187, <https://doi.org/10.1002/esp.5093>, 2021.

700 Jiménez-González, I., Rodríguez-Navarro, C., and Scherer, G. W.: Role of clay minerals in the physicomaterial
701 deterioration of sandstone, *J. Geophys. Res.*, 113, F02021, <https://doi.org/10.1029/2007JF000845>, 2008.

702 Jobbágy, E. G., and Jackson, R. B.: The uplift of soil nutrients by plants: biogeochemical consequences across scales, *Ecology*,
703 85, 2380–2389, <https://doi.org/10.1890/03-0245>, 2004.

704 Kim, H., Stinchcomb, G., and Brantley, S.: Feedbacks among O₂ and CO₂ in deep soil gas, oxidation of ferrous minerals, and
705 fractures: A hypothesis for steady-state regolith thickness, *Earth Planet. Sc. Lett.*, 460, 29–40,
706 <https://doi.org/10.1016/j.epsl.2016.12.003>, 2017.

- 707 Kogure, T., and Banfield, J. F.: New insights into the mechanism for chloritization of biotite using polytype analysis, *Am.*
708 *Mineral.*, 85 (8–9), 1202–1208, <https://doi.org/10.1515/am-2000-8-913>, 2000.
- 709 Krone, L. V., Hampl, F. J., Schwerdhelm, C., Bryce, C., Ganzert, L., Kitte, A., Übernickel, K., Dielforder, A., Aldaz, S., Oses-
710 Pedraza, R., Perez, J. P. H., Sanchez-Alfaro, P., Wagner, D., Weckmann, U., and von Blanckenburg, F.: Deep weathering in
711 the semi-arid Coastal Cordillera, Chile, *Sci. Rep.-UK*, 11, 13057, <https://doi.org/10.1038/s41598-021-90267-7>, 2021.
- 712 Lawrence, C., Harden, J., and Maher, K.: Modeling the influence of organic acids on soil weathering, *Geochim. Cosmochim.*
713 *Ac.*, 139, 487–507, <https://doi.org/10.1016/j.gca.2014.05.003>, 2014.
- 714 Lebedeva, M. I. and Brantley, S. L.: Exploring an 'ideal hill': how lithology and transport mechanisms affect the possibility of
715 a steady state during weathering and erosion. *ESPL*, 45(3), 652-665, <https://doi.org/10.1002/esp.4762>, 2020.
- 716 Lebedeva, M. I., Fletcher, R. C., Balashov, V. N., and Brantley, S. L.: A reactive diffusion model describing transformation
717 of bedrock to saprolite, *Chem. Geol.*, 244, 624–645, <https://doi.org/10.1016/j.chemgeo.2007.07.008>, 2007.
- 718 Lohse, K. A., and Dietrich, W. E.: Contrasting effects of soil development on hydrological properties and flow paths, *Water*
719 *Resour. Res.*, 41, W12419, <https://doi.org/10.1029/2004WR003403>, 2005.
- 720 Lucas, Y.: The Role of Plants in Controlling Rates and Products of Weathering: Importance of Biological Pumping, *Annu.*
721 *Rev. Earth Pl. Sc.*, 29, 135–163, <https://doi.org/10.1146/annurev.earth.29.1.135>, 2001.
- 722 Luebert, F., and Plissock, P.: Sinópsis bioclimática y vegetal de Chile, Editorial Universitaria, Santiago de Chile, ISBN
723 956-11-1832-7, 2006.
- 724 Maher, K.: The dependence of chemical weathering rates on fluid residence time, *EPSL*, 294, 1–2, 101-110,
725 <https://doi.org/10.1016/j.epsl.2010.03.010>, 2010.
- 726 McKeague, J. A., and Day, J. H.: Dithionite- and oxalate-extractable Fe and Al as aids in differentiating various classes of
727 soils, *Can. J. Soil Sci.*, 46 (1), 13–22, <https://doi.org/10.4141/cjss66-003>, 1966.
- 728 Melnick, D.: Rise of the central Andean coast by earthquakes straddling the Moho, *Nat. Geosci.*, 9, 401–407,
729 <https://doi.org/10.1038/ngeo2683>, 2016.
- 730 Molnar, P., Anderson, R. S., and Anderson, S. P.: Tectonics, fracturing of rock, and erosion, *J. Geophys. Res.*, 112, F03014,
731 <https://doi.org/10.1029/2005JF000433>, 2007.

- 732 Nadan, B. J., and Engelder, T.: Microcracks in New England granitoids: A record of thermoelastic relaxation during
733 exhumation of intracontinental crust, *Geol. Soc. Am. Bull.*, 121, 80–99, <https://doi.org/10.1130/b26202.1>, 2009.
- 734 Napieralski, S. A., Buss, H. L., Brantley, S. L., Lee, S., Xu, H., and Roden, E. E.: Microbial chemolithotrophy mediates
735 oxidative weathering of granitic bedrock, *P. Natl. Acad. Sci. USA*, 116, 26394–26401,
736 <https://doi.org/10.1073/pnas.1909970117>, 2019.
- 737 Nascimento, D. L., Abrahão, A., Lambers, H., Teodoro, G. S., Ladeira, F. S. B., de Britto Costa, P., Oliveira, R. S., and de
738 Farias, C. H. B.: Biogeomorphological evolution of rocky hillslopes driven by roots in campos rupestres, Brazil,
739 *Geomorphology*, 395, 107985, <https://doi.org/10.1016/j.geomorph.2021.107985>, 2021.
- 740 Navarre-Sitchler, A., Brantley, S. L., and Rother, G.: How Porosity Increases During Incipient Weathering of Crystalline
741 Silicate Rocks, *Rev. Mineral. Geochem.*, 80 (1), 331–354, <https://doi.org/10.2138/rmg.2015.80.10>, 2015.
- 742 Nesbitt, H., and Young, G.: Early Proterozoic climates and plate motions inferred from major element chemistry of lutites,
743 *Nature*, 299, 715–717, <https://doi.org/10.1038/299715a0>, 1982.
- 744 Oeser, R. A., Stroncik, N., Moskwa, L.-M., Bernhard, N., Schaller, M., Canessa, R., van den Brink, L., Köster, M., Brucker,
745 E., Stock, S., Fuentes, J. P., Godoy, R., Matus, F. J., Osés Pedraza, R., Osses McIntyre, P., Paulino, L., Seguel, O., Bader, M.
746 Y., Boy, J., Dippold, M. A., Ehlers, T. A., Kühn, P., Kuzyakov, Y., Leinweber, P., Scholten, T., Spielvogel, S., Spohn, M.,
747 Übernichel, K., Tielbörger, K., Wagner, D., and von Blanckenburg, F.: Chemistry and microbiology of the Critical Zone along
748 a steep climate and vegetation gradient in the Chilean Coastal Cordillera, *CATENA*, 170, 183–203,
749 <https://doi.org/10.1016/j.catena.2018.06.002>, 2018.
- 750 Oeser, R. A., and von Blanckenburg, F.: Do degree and rate of silicate weathering depend on plant productivity?
751 *Biogeosciences*, 17, 4883–4917, <https://doi.org/10.5194/bg-17-4883-2020>, 2020.
- 752 Pawlik, Ł., Phillips, J. D., and Šamonil, P.: Roots, rock, and regolith: Biomechanical and biochemical weathering by trees and
753 its impact on hillslopes – A critical literature review, *Earth-Sci. Rev.*, 159, 142–159,
754 <https://doi.org/10.1016/j.earscirev.2016.06.002>, 2016.
- 755 Perez, J. R., Banwart, St. A., and Puigdomenech, I.: The kinetics of $O_{2(aq)}$ reduction by structural ferrous iron in naturally
756 occurring ferrous silicate minerals, *Appl. Geochem.*, 20, 2003–2016, <https://doi.org/10.1016/j.apgeochem.2005.07.001>, 2005.
- 757 Reis, F. D. A. A. and Brantley, S. L.: The impact of depth-dependent water content on steady state weathering and eroding
758 systems. *Geochim. Cosmochim. Acta*, 244, 40–55, <https://doi.org/10.1016/j.gca.2018.09.028>, 2019.

- 759 Rempe, D. M., and Dietrich, W. E.: A bottom-up control on fresh-bedrock topography under landscapes, *P. Natl. Acad. Sci.*
760 *USA*, 111 (18), 6576–6581, <https://doi.org/10.1073/pnas.1404763111>, 2014.
- 761 Rennert, T.: Wet-chemical extractions to characterise pedogenic Al and Fe species – a critical review, *Soil Res.*, 57, 1–16,
762 <https://doi.org/10.1071/SR18299>, 2019.
- 763 Riebe, C. S., Kirchner, J. W., and Finkel, R. C.: Long-term rates of chemical weathering and physical erosion from cosmogenic
764 nuclides and geochemical mass balance, *Geochim. Cosmochim. Ac.*, 67, 4411–4427, [https://doi.org/10.1016/S0016-](https://doi.org/10.1016/S0016-7037(03)00382-X)
765 [7037\(03\)00382-X](https://doi.org/10.1016/S0016-7037(03)00382-X), 2003.
- 766 Røyne, A., Jamtveit, B., Mathiesen, J., and Malthe-Sørenssen, A.: Controls on rock weathering rates by reaction-induced
767 hierarchical fracturing, *Earth Planet. Sc. Lett.*, 275, 364–369, <https://doi.org/10.1016/j.epsl.2008.08.035>, 2008.
- 768 Schaller, M., and Ehlers, T. A.: Comparison of soil production, chemical weathering, and physical erosion rates along a climate
769 and ecological gradient (Chile) to global observations, *Earth Surf. Dynam.*, 10, 131–150, [https://doi.org/10.5194/esurf-10-131-](https://doi.org/10.5194/esurf-10-131-2022)
770 [2022](https://doi.org/10.5194/esurf-10-131-2022), 2022.
- 771 Scheibe, A., Sierra, C. A., Spohn, M.: Recently fixed carbon fuels microbial activity several meters below the soil surface.
772 *Biogeosciences*, 20, 827-838, <https://doi.org/10.5194/bg-20-827-2023>, 2023.
- 773 Schneider, C. A., Rasband, W. S., and Eliceiri, K. W.: NIH Image to ImageJ: 25 years of image analysis, *Nat. Methods*, 9 (7),
774 671–675, <https://doi.org/10.1038/nmeth.2089>, 2012.
- 775 Schwertmann, U.: Differenzierung der Eisenoxide des Bodens durch Extraktion mit Ammoniumoxalat-Lösung, *Zeitschrift für*
776 *Pflanzenernährung, Düngung, Bodenkunde*, 105, 194–202, <https://doi.org/10.1002/jpln.3591050303>, 1964.
- 777 Starkey, H. C., Blackmon P. D., and Hauff P. L. The routine mineralogical analysis of clay-bearing samples, *U.S. Geological*
778 *Survey Bulletin*, 1563, <https://doi.org/10.3133/b1563>, 1984.
- 779 St. Clair, J., Moon, S., Holbrook, W. S., Perron, J. T., Riebe, C. S., Martel, S. J., Carr, B., Harman, C., Singha, K., and Richter,
780 D.: Geophysical imaging reveals topographic stress control of bedrock weathering, *Science*, 350 (6260), 534–538,
781 <https://doi.org/10.1126/science.aab2210>, 2015.
- 782 Steefel, C. I., and van Cappellen, P.: A new kinetic approach to modeling water-rock interaction: The role of nucleation,
783 precursors, and Ostwald ripening, *Geochim. Cosmochim. Ac.*, 54 (10), 2657–2677, [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-7037(90)90003-4)
784 [7037\(90\)90003-4](https://doi.org/10.1016/0016-7037(90)90003-4), 1990.

785 Steenken, A., Rabbia, O., and Hernández, L.: The Emplacement of the Nahuelbuta Batholith in an Active Continental Margin
786 (Central Chile), XVIII Congreso Peruano de Geología, Lima, Peru, 2016.

787 Stinchcomb, G. E., Kim, H., Hasenmueller, E. A., Sullivan, P. L., Sak, P. B., and Brantley, S. L.: Relating soil gas to weathering
788 using rock and regolith geochemistry. *Am. J. Sci.*, 318, 727-763, <https://doi.org/10.2475/07.2018.01>, 2018.

789 Übernickel, K., Ehlers, T. A., Ershadi, M. R., Paulino, L., Fuentes Espoz, J.-P., Maldonado, A., Oses-Pedraza, R., and von
790 Blanckenburg, F.: Time series of meteorological station data in the EarthShape study areas in the Coastal Cordillera, Chile,
791 GFZ Data Services [data set], <https://doi.org/10.5880/fidgeo.2020.043>, 2020.

792 van Dongen, R., Scherler, D., Wittmann, H., and von Blanckenburg, F.: Cosmogenic ^{10}Be in river sediment: where grain size
793 matters and why, *Earth Surf. Dynam.*, 7, 393–410, <https://doi.org/10.5194/esurf-7-393-2019>, 2019.

794 Vázquez, M., Ramírez, S., Morata, D., Reich, M., Braun, J.-J., and Carretier, S.: Regolith production and chemical weathering
795 of granitic rocks in central Chile, *Chem. Geol.*, 446, 87–98, <https://doi.org/10.1016/j.chemgeo.2016.09.023>, 2016.

796 Werner, C., Schmid, M., Ehlers, T. A., Fuentes-Espoz, J. P., Steinkamp, J., Forrest, M., Liakka, J., Maldonado, A., and Hickler,
797 T.: Effect of changing vegetation and precipitation on denudation – Part 1: Predicted vegetation composition and cover over
798 the last 21 thousand years along the Coastal Cordillera of Chile, *Earth Surf. Dynam.*, 6, 829–858, [https://doi.org/10.5194/esurf-](https://doi.org/10.5194/esurf-6-829-2018)
799 [6-829-2018](https://doi.org/10.5194/esurf-6-829-2018), 2018.

800 White, A. F., and Yee, A.: Aqueous oxidation-reduction kinetics associated with coupled electron transfer from iron-containing
801 silicates at 25°C, *Geochim. Cosmochim. Ac.*, 49, 1263–1275, [https://doi.org/10.1016/0016-7037\(85\)90015-8](https://doi.org/10.1016/0016-7037(85)90015-8), 1985.

802 Wilson, M. J.: Weathering of the primary rock-forming minerals: processes, products and rates, *Clay Miner.*, 39, 233–266,
803 <https://doi.org/10.1180/0009855043930133>, 2004.